less marked and the glucose uptake and lactic acid accumulation increased.

4. In unstimulated slices respiration, glucose consumption and lactic acid formation increased with rise of temperature (30.0–41.5°). In electrically stimulated preparations the functions mentioned above increased from 30° to 37.5° but decreased at 41.5°. Ethanol had no effect at 30°.

5. The relationship of these results to those obtained in earlier investigations with ethanol and other depressants is discussed. It is concluded that the action of ethanol is non-competitive and directed towards some process linked with the physiological functioning of nerve tissue.

The authors are indebted to Professor H. McIlwain, Dr Heikki Suomalainen, Dr K. O. Donner and Dr Olof Forsander for helpful criticism of the manuscript, to Mrs Kaija Salmela and Mr Ralph Lindbohm for technical assistance in the experimental work, and to Mr Peter Neuenschwander and Mr Keijo Bovellan for constructing special equipment.

REFERENCES

- Abood, L. G., Gerard, R. W. & Ochs, S. (1952). *Amer. J. Physiol.* **171**, 134.
- Ashford, C. A. & Dixon, K. C. (1935). *Biochem. J.* **29**, 157.
- Barker, S. B. & Summerson, W. H. (1941). *J. biol. Chem.* **138**, 535.
- Bartlett, G. R. & Barnet, H. N. (1949). *Quart. J. Stud. Alc.* **10**, 381.
- Batley, L. L., Heyman, A. & Patterson, J. L. (1953). *J. Amer. med. Ass.* **152**, 6.
- Beer, C. T. & Quastel, J. H. (1958a). *Canad. J. Biochem. Physiol.* **36**, 531.
- Beer, C. T. & Quastel, J. H. (1958b). *Canad. J. Biochem. Physiol.* **36**, 543.
- Bronk, D. W. & Brink, F. (1951). *Fed. Proc.* **10**, 19.
- Brooks, C. McC., Koizumi, K. & Malcolm, J. L. (1955). *J. Neurophysiol.* **18**, 205.
- Buchel, L. (1953). *Anésth. Analg.* **10**, 1.
- Burbridge, T. N., Sutherland, V. C., Hine, C. H. & Simon, A. (1959). *J. Pharmacol.* **126**, 70.
- Burns, B. D. (1958). *The Mammalian Cerebral Cortex*. London: Edward Arnold Ltd.
- Dickens, F. & Greville, G. B. (1935). *Biochem. J.* **29**, 1468.
- Field, J., Fuhrman, F. A. & Martin, A. W. (1944). *J. Neurophysiol.* **7**, 117.
- Fischer, E. (1957). *A.A.A.S. Publ.; Alcoholism*, no. 47 p. 19.
- Forda, O. & McIlwain, H. (1953). *Brit. J. Pharmacol.* **8**, 225.
- Fuhrman, F. A. & Field, J. (1948). *Proc. exp. Biol., N.Y.*, **69**, 331.
- Gallego, A. (1948). *J. cell. comp. Physiol.* **31**, 97.
- Ghosh, J. J. & Quastel, J. H. (1954). *Nature, Lond.*, **174**, 28.
- Green, D. E. (1956–57). *Harvey Lect.* **52**, 177.
- Himwich, H. E. (1956). In *Alcoholism*, p. 291. Ed. by Thompson, G. E. Springfield, Ill.: Charles C. Thomas.
- Hine, C. H., Shick, A. F., Margolis, L., Burbridge, T. N. & Simon, A. (1952). *J. Pharmacol.* **106**, 253.
- Kimura, Y. (1937). *Sci. Pap. Inst. phys. chem. Res., Tokyo*, **33**, 231.
- Kratzing, C. C. (1953). *Biochem. J.* **54**, 312.
- Larrabee, M. G. & Posternak, J. M. (1952). *J. Neurophysiol.* **15**, 91.
- Masoro, E. J., Abramovitch, H. & Birchard, J. R. (1953). *Amer. J. Physiol.* **173**, 37.
- McIlwain, H. (1951a). *Biochem. J.* **49**, 382.
- McIlwain, H. (1951b). *Biochem. J.* **50**, 132.
- McIlwain, H. (1953a). *Biochem. J.* **53**, 403.
- McIlwain, H. (1953b). *Biochem. J.* **55**, 618.
- McIlwain, H. (1953c). *J. Neurol. Psychiat.* **16**, 257.
- McIlwain, H. (1954). *Electroenceph. clin. Neurophysiol.* **6**, 93.
- McIlwain, H., Anguiano, G. & Cheshire, J. D. (1951). *Biochem. J.* **50**, 12.
- McIlwain, H. & Gore, M. B. R. (1953). *Biochem. J.* **54**, 305.
- McIlwain, H. & Ochs, S. (1952). *Amer. J. Physiol.* **171**, 128.
- Narayanaswami, A. & McIlwain, H. (1954). *Biochem. J.* **57**, 663.
- Quastel, J. H. (1952). *Curr. Res. Anesth.* **31**, 151.
- Somogyi, M. & Nelson, N. (1945). *J. biol. Chem.* **160**, 69.
- Sutherland, V. C., Hine, C. H. & Burbridge, T. N. (1956). *J. Pharmacol.* **116**, 469.
- Truitt, E. B., Bell, F. K. & Krantz, J. K. (1956). *Quart. J. Stud. Alc.* **17**, 594.
- Wallgren, H. (1960). *Acta physiol. scand.* (in the Press).
- Wright, E. B. (1947). *Amer. J. Physiol.* **148**, 174.

Biochem. J. (1960) **75**, 158

Purification of the Two Components of Leucocidin from *Staphylococcus aureus*

By A. M. WOODIN*

Sir William Dunn School of Pathology, University of Oxford

(Received 17 September 1959)

Staphylococcal leucocidin inhibits the respiration (Neisser & Wechsberg, 1901) and produces characteristic morphological changes (Gladstone &

* Member of the external staff, Medical Research Council.

van Heyningen, 1957) in the white blood cells of humans and rabbits. It has been shown that this activity results from the synergistic action of at least two substances, since the activity is lost when leucocidin preparations are separated on carboxy-

methylcellulose into two fractions but is regained when the fractions are mixed (Woodin, 1959). According to their rate of migration on carboxymethylcellulose these fractions were called the fast and the slow fraction of the leucocidin preparation. Chromatographically the fast fraction behaved as a single substance but immunologically several constituents could be distinguished. The electrophoretic behaviour of the fast fraction suggested the formation of complexes, and this may be responsible for the heterogeneous system behaving as a single substance on columns of cation-exchange resins. The slow fraction of leucocidin behaved essentially as a single substance contaminated with some of the constituents of the fast fraction.

The present paper describes the further purification of the fast and slow fractions of the leucocidin preparation. It is now clear that only one of the constituents of each is responsible for the activity and these are referred to below as the fast (F) and slow (S) components of leucocidin. Both components have been crystallized.

MATERIALS AND METHODS

Preparation of culture filtrates. Large-scale production of culture filtrates from the V8 strain of *Staphylococcus aureus* grown in the 'CCY' medium containing a diffusate from Oxoid Yeast extract (Woodin, 1959) was done by Mr R. Elsworth and Mr A. May at the Microbiological Research Establishment, Porton. Ammonium sulphate (750 g.) was stirred into each litre of culture filtrate and the solution left overnight. The bulk of the precipitate rose to the surface and, after the supernatant solution had been siphoned off, was concentrated by centrifuging. The precipitate is called the culture-filtrate slurry.

Fractionation of the culture-filtrate slurry. The 'displacement-eluate 1' and 'displacement-eluate 2' were derived by the method summarized in Fig. 6 of Woodin (1959) with the following modifications.

(a) The $(\text{NH}_4)_2\text{SO}_4$ concentration in the culture-filtrate slurry was reduced to M by adding 0.1 M-phosphate buffer, pH 6.7, at room temperature and the insoluble fraction rejected. With one exception subsequent operations were carried out at 4–7°.

(b) The extract in M- $(\text{NH}_4)_2\text{SO}_4$ was centrifuged at 25 000 rev./min. for 1 hr. and the sediment rejected. The supernatant was diluted to have a conductivity equal to that of 0.1 M-phosphate buffer, pH 6.7, and applied to a column of hydroxylapatite. This was eluted in turn with 0.1 M-phosphate buffer, pH 6.7, 0.14 M-phosphate buffer, pH 6.7, and then with 0.5 M-phosphate buffer, pH 6.7, at room temperature. The last effluent contained the bulk of the two components of leucocidin and was processed on carboxymethylcellulose to give the displacement-eluates 1 and 2 as already described. They were stored in saturated $(\text{NH}_4)_2\text{SO}_4$. The F and S fractions of the leucocidin preparation were prepared from displacement-eluate 2 as described by Woodin (1959) and stored in saturated $(\text{NH}_4)_2\text{SO}_4$.

Assay of leucocidin. To enable the measurement of the concentrations of the F and S components of leucocidin the

procedure has been modified. For each sample to be assayed dilutions differing by 10% are prepared. An excess (12.5 $\mu\text{g.}$) of the purified F component (see below) is added to a 1 ml. portion of each dilution and purified S component (2.5 $\mu\text{g.}$) is added to another 1 ml. portion of each dilution. Antitoxin (0.5 unit) is then added to each tube and the end points are determined by the inhibition of the respiration of macrophages as described previously (Woodin, 1959). The number of test doses (L +) per ml. of the original sample is equal to the dilution at the end point multiplied by the number of units of antitoxin added (i.e. 0.5). The result obtained in the presence of added F component is referred to below as the 'S titre' and that obtained in the presence of added S component as the 'F titre'.

When end points have also been determined by the observation of the morphological changes induced in polymorphonuclear leucocytes (Gladstone & van Heyningen, 1957) the results have not differed.

Staphylococcal antitoxin. This was a pepsin-refined antiserum CPP 76/63 kindly provided by Miss Mollie Barr of the Wellcome Research Laboratories. It was arbitrarily assigned the values 240 units of anti-S component and 240 units of anti-F component.

Assay of deoxyribonuclease. This was done viscometrically by a method similar to that of Laskowski & Seidel (1945). Deoxyribonucleic acid was prepared from calf thymus by the method of Kay, Simmons & Dounce (1952) and a 0.05% solution was prepared in 0.05 M-sodium borate buffer, pH 8.6, containing 0.01 M- CaCl_2 . This solution had a relative viscosity of 5 at 37° in a viscometer with a flow time of 18 sec. for 1 ml. of water. Preparations with deoxyribonuclease activity were diluted in 0.01 M- CaCl_2 -0.05 M-sodium borate buffer, pH 8.6, containing 0.2% of gelatin. A portion (0.25 ml.) of the deoxyribonuclease solutions was mixed with 1 ml. of the deoxyribonucleic acid solution at 37° and 1 ml. of the mixture was transferred to the viscometer. Deoxyribonuclease units were calculated by the method of Laskowski & Seidel (1945). For reproducible results it was necessary to dilute the deoxyribonuclease solutions so that they contained less than 6 units/ml.

Chromatographic methods. Dowex-2 ($\times 8$) was regenerated and set up in columns by the procedure described by Boman & Westlund (1956), except that after use the top 1 cm. of resin in the column was rejected. The other adsorbents were used in the way described by Woodin (1959).

Protein concentrations. In the partially purified fractions protein concentrations were assessed from the ultraviolet absorption at 280 m μ , assuming $E_{1\%}^{1\text{cm.}}$ 10. The data given in this paper relating to the properties of the F component and of the S component refer to the concentration calculated from the absorption at 280 m μ and the values $E_{1\%}^{1\text{cm.}}$ 16.2 and 15.8 for the F and S components respectively.

Nitrogen contents. These were determined by the micro-Kjeldahl method (Markham, 1942; Ma & Zuazaga, 1942). With the purified S component a solution was dialysed against distilled water for several days and dried at 110° in a weighed ampoule; the N content was then determined. With the F component the crystalline material was washed with water till the supernatant gave no precipitate with AgNO_3 , and dried at 110°. No corrections were made for residual ash content.

Ultracentrifuging. This was done in a Spinco model E analytical ultracentrifuge equipped with the RTIC unit and a phase plate in the schlieren-optical system. Sedimentation coefficients (S) are given in Svedberg units and were calculated from the plot of $\log x$ against t , where x is the distance from the centre of rotation to the peak of the gradient curve and t is the time of sedimentation. Apparent diffusion coefficients (D^*) of material sedimenting in the ultracentrifuge were calculated from the equation

$$D = \frac{1}{4\pi t} \left(\frac{A}{H} \right)^2 (1 - W^2 St),$$

where A is the area of the schlieren diagram and H its height, S the sedimentation coefficient in cm./sec./dyne/g., t the time during which diffusion has proceeded and W is the angular velocity. The values of D^* are in cm.²/sec. The concentration of material sedimenting in the ultracentrifuge was calculated from the areas of the gradient curves (corrected for the dilution effect due to the shape of the cell), the phase-plate angle and a calibration constant obtained by forming a boundary between solvent and solution in a synthetic boundary cell at low speed. The partial specific volume ($\bar{V}$) of the F component was calculated from density measurements on a 0.73% solution in 0.2 M-acetate buffer, pH 5.0. The values of $\bar{V}$ are in ml./g.

Boundary electrophoresis. This was done in an Antweiler microelectrophoresis apparatus.

Conductivity measurements. A Mullard conductivity bridge was used for these.

Immunological analysis. This was done by diffusion in agar, by the method of Ouchterlony (1949).

Buffers. Buffers used in the experiments described in this paper had the composition given by Woodin (1959); also 0.01 M-2-amino-2-hydroxymethylpropane-1.3-diol-HCl (tris-HCl) buffer, pH 7.5, prepared by adding HCl to 10 m-moles of tris to give pH 7.5, and diluting to 1 l.

RESULTS

Production of the two components of leucocidin by Staphylococcus aureus

Both the F and the S titres reached maximal values after 8–10 hr., when the culture contained 5–6 g. of organisms/l. The final titres in different batches of culture filtrate varied from 10 L+ of the F component/ml. and 50 L+ of the S component/ml. to 21 L+ of the F component/ml. and 150 L+ of the S component/ml. The latter values correspond

to 78 µg. of F component/ml. and 83 µg. of S component/ml.

Recovery of the two components in the preliminary purification

Table 1 lists the yields of the F and S components in the preliminary purification. At all stages the recovery of the S component is less than that of the F component. Little activity could be recovered in the fractions rejected and it is likely that the losses are due to irreversible precipitation.

The extent of separation of the F and S fractions of the leucocidin preparation in the gradient elution from carboxymethylcellulose depends upon the efficiency of the packing of the column, and where significant overlap occurred the overlapping material was processed again.

F component of leucocidin

Purification. The displacement-eluate 1 and the F fraction of the leucocidin preparation were combined. Both were found to contain deoxyribonuclease, some of the properties of which have been described by Privat de Garilhe, Fassina, Pochon & Pillet (1958). They showed that deoxyribonuclease can be adsorbed onto Amberlite CG-50 from a solution in 0.2 M-sodium phosphate buffer, pH 7.0. The F component of leucocidin is not adsorbed under these conditions. However, to achieve separation it is necessary to dialyse the displacement-eluate 1 and the F fraction of the leucocidin preparation against 0.2 M-sodium phosphate buffer, pH 7.0, to remove the (NH₄)₂SO₄, and this led to the precipitation of about 30% of the F component in a partly crystalline form. Crystallization at this stage is disadvantageous as the crystalline product contains many impurities.

The method used to purify the F component of leucocidin is shown in Fig. 1 and utilizes the finding that leucocidin can be separated from deoxyribonuclease at a lower pH than that used by Privat de Garilhe *et al.* (1958). Dowex-2 (× 8) is used to remove pigmented material. After passage through Dowex-2 (× 8) the effluent is adjusted to pH 5.0

Table 1. *Recovery of the two components of leucocidin in the preliminary purification*

The culture filtrate processed contained 1.3×10^6 L+ of F component and 8.7×10^6 L+ of S component.

Stage of purification	F component (% of starting material recovered)	S component (% of starting material recovered)
Fraction soluble in $\text{M}-(\text{NH}_4)_2\text{SO}_4$	54	35
Hydroxylapatite eluate in 0.5 M-phosphate buffer, pH 6.7	31	14
Displacement-eluate 1	19	0.2
Displacement-eluate 2	6	10
F fraction of the leucocidin preparation	4	0.3
S fraction of the leucocidin preparation	2	9

Fig. 1. Preparation of the crystalline F component of leucocidin.

Table 2. Recovery of the F component of leucocidin in the fractionation illustrated in Fig. 1

Stage in purification	Protein content (mg.)	F component content (L +)	Deoxyribonuclease content (Deoxyribonuclease units)
F fraction of the leucocidin preparation and displacement-eluate 1	120	8700	2 × 10 ⁷
Effluent from Dowex-2 (× 8)	84	8500	1.3 × 10 ⁷
Effluent from Amberlite CG-50 in 0.2 M-citrate buffer, pH 6.0	51	7500	(Not measured)
(NH ₄) ₂ SO ₄ -concentrate of the 0.2 M-citrate buffer effluent from Amberlite CG-50	42	5700	3200
Crystalline F component	20	3200	Less than 200

Fig. 2. The elution of the partially purified F component of leucocidin from Amberlite CG-50 by 0.2 M-citrate buffer, pH 6.0. ○, Total protein; ×, pH; ●, conductivity (arbitrary units); ▲, F component of leucocidin; ●, deoxyribonuclease.

and applied to a column of Amberlite CG-50. The F component is adsorbed and impurities are removed by elution with 0.1 M-citrate buffer, pH 5.0. When the ultraviolet absorption of the effluent has fallen to a constant value and the pH and conductivity of the effluent are those of the 0.1 M-citrate buffer, pH 5.0, the eluent is changed to 0.2 M-citrate buffer, pH 6.0. There is an increase in the conductivity of the effluent to a value approaching that of the inflowing buffer and this is followed by a gradual increase in the pH of the effluent. A single protein peak is eluted above pH 5.4. This stage in the purification is illustrated in Fig. 2. Table 2 gives the recovery of the F component at various stages.

Crystallization. The first 5% and the last 5% of the protein peak eluted from Amberlite CG-50 were rejected and the remainder was concentrated and precipitated by dialysis against saturated $(\text{NH}_4)_2\text{SO}_4$. The precipitate was dissolved in the smallest amount of 0.1 M-acetate buffer, pH 5.0,

and the solution was dialysed against 0.2 M-phosphate buffer, pH 6.7. In the course of 2–3 days needle-like crystals or, occasionally, hexagonal-plate-like crystals were deposited (Fig. 3). Re-crystallization could be effected by repeating this procedure after solution of the crystals in 0.1 M-acetate buffer, pH 5.0, which requires several hours. About 80% of the material crystallized in the course of a few days. The crystalline material was stored in 0.2 M-phosphate buffer, pH 6.7, at 4°.

Properties. The N content was 15.5%; $E_{1\text{cm}}^{1\%}$ at 280 m μ was 16.2; the antibody-combining power was 270 L+/mg. Immunological analysis by diffusion in agar showed a single precipitin band (Fig. 4). The supernatant from the first crystallization contained other antigens, indicating the value of the crystallization at this stage in removing trace contaminants.

Ultracentrifuging. All measurements were made at 22°. The twice-crystallized F component, suspended in phosphate buffer, was dissolved in 0.2 M-acetate buffer, pH 5.0, by dialysing a suspension of the crystals against this solvent. A 0.58% solution was centrifuged at 7000 rev./min. in the synthetic boundary cell and the constant relating the area of the gradient curve to the concentration was calculated. This, together with the optical constants and enlargement factors, gives 1.80×10^{-3} for the specific refractive increment. The apparent diffusion coefficient ($D_{20,w}^*$) was 8.5×10^{-7} .

A 1.32% (w/v) solution in 0.2 M-acetate buffer, pH 5.0, was centrifuged at 59780 rev./min. and had

Fig. 3. The two crystalline forms of the F component of leucocidin. (a) Hexagonal plates, dark-ground illumination; $\times 800$. (b) Needle-like form, light-ground illumination; $\times 480$.

Fig. 4. Immunological analysis of the F and S components of leucocidin. The centre well contained staphylococcal antitoxin CPP 76/63. Recorded after 3 days' diffusion at 37°. F₁, 300 μ g. of F component; F₂, 100 μ g. of F component; F₃, 33 μ g. of F component. S₁, 360 μ g. of S component; S₂, 120 μ g. of S component; S₃, 40 μ g. of S component.

$S_{20,w}$ 3.05. Fig. 5 shows the gradient curve. The concentration of the sedimenting material calculated from the areas of the gradient curves at various times was 95–100 % of the starting material. $D_{20,w}^*$ values were 7.14×10^{-7} – 7.42×10^{-7} , and showed no dependence upon the time of sedimentation. The values S 3.05, D 8.5×10^{-7} , $\bar{V}$ 0.721 give a molecular weight of 32 000.

Electrophoresis of the F component. A 1.5 % (w/v) solution of the crystalline material in 0.1M-acetate buffer, pH 5.0, showed a single component on boundary electrophoresis. A solution prepared from twice-crystallized material which had been washed with water and freeze-dried retained a slight opalescence after centrifuging at 7000 rev./min. for 20 min. and showed two components on boundary electrophoresis in 0.1M-acetate buffer,

pH 5.0 (Fig. 6). From the appearance of the descending boundary it was clear that the minor, faster component was responsible for the opalescence of the solution. It was removed by centrifuging at 60 000 rev./min. for 30 min. and the supernatant of this centrifuged solution showed a single component on boundary electrophoresis (Fig. 6). A solution prepared from crystalline material which had been stored in the frozen state also showed two components on boundary electrophoresis in 0.1M-acetate buffer, pH 5.0. It is probable that the minor component is produced from the crystallized F component by denaturation, and freezing or freeze-drying the crystalline material should be avoided.

S component of leucocidin

Purification. It was shown that the S fraction of the leucocidin preparation had a single peak on boundary electrophoresis at pH 5.0 and at pH 7.0 (Woodin, 1959). Immunological analysis by diffusion in agar showed a main precipitin band persisting to low antigen concentrations and other precipitin bands which were diluted out at much higher antigen concentrations and appeared to be identical with some of the constituents of the F fraction of the leucocidin preparation. To remove these the S fraction of the leucocidin preparation

Fig. 5. Schlieren diagrams of the two components of leucocidin sedimenting in the ultracentrifuge. (a) F component, 1.32 %, 64 min. after reaching full speed. (b) S component, 1.30 %, 85 min. after reaching full speed. In both cases the solvent was 0.2M-acetate buffer, pH 5.0.

Fig. 6. Boundary electrophoresis of the F component of leucocidin in 0.1M-acetate buffer, pH 5.0. (a) Freeze-dried material before high-speed centrifuging; (b) the same material after high-speed centrifuging.

Table 3. Recovery of the S component of leucocidin on fractionation by the scheme illustrated in Fig. 1

The analysis of the Amberlite CG-50 effluent is for material remaining after rejection of the first 10% and final 5% of the protein peak.

Stage in the purification	Protein content (mg.)	S component content (L +)	Deoxyribonuclease content (Deoxyribonuclease units)
S fraction of the leucocidin preparation	1000	9×10^5	5.4×10^7
Effluent from Dowex-2 ($\times 8$)	820	9.5×10^5	4.5×10^7
Effluent from Amberlite CG-50 in 0.2M-citrate buffer, pH 6.0	620	7.5×10^5	8.5×10^3

was put through the fractionation scheme illustrated in Fig. 1, as far as the elution from Amberlite CG-50 with 0.2M-citrate buffer, pH 6.0, and with the following modification. As the S component was not precipitated by dialysis at pH 7.0, before application of the material to the columns the salt concentrations were adjusted by dialysis and any insoluble material was centrifuged off. Air was excluded from the dialysis sac and the contents were agitated as little as possible, as agitation caused irreversible precipitation of some of the S component in a fibrous form. Fig. 3 of Woodin (1959) indicates a relative displacement of the F and S components on elution from Amberlite CG-50 with a pH gradient. To reduce the contamination of the S component by the F component the first 10% of the protein peak eluted by 0.2M-citrate buffer, pH 6.0, was rejected, and to decrease the deoxyribonuclease content the last 5% of the protein peak was also rejected. The remaining material was dialysed against distilled water and freeze-dried. Table 3 gives the recovery of the S component at various stages. Subsequent to the development of this fractionation scheme the conditions for the crystallization of the S component were discovered and are described below. The properties of the S component have been determined on the freeze-dried material before crystallization.

Properties. The N content was 15.2%, $E_{1\text{cm}}^{1\%}$ at 280 m μ was 15.8; the antibody-combining power was 1780 L+/mg. Immunological analysis by diffusion in agar showed a single precipitin band (Fig. 4). The S component has been stored in the freeze-dried form over P₂O₅. The freeze-dried solid is not completely soluble in water, 0.2M-acetate buffer, pH 5.0, or 0.1M-phosphate buffer, pH 6.7, and the amount remaining insoluble appears to increase with storage.

Ultracentrifuging. All measurements were made at 23°. A 0.47% solution of the S component, in 0.2M-acetate buffer, pH 5.0, was centrifuged at 7000 rev./min. and the constant relating the concentration to the area of the gradient curve was determined. This, together with the optical constants and the enlargement factors, gave the value

Fig. 7. Crystallized S component of leucocidin. Dark-ground illumination; $\times 600$.

1.87×10^{-3} for the specific refractive increment. $D_{20,w}^*$ under these conditions was 7.5×10^{-7} . A 1.30% solution of the S component in 0.2M-acetate buffer, pH 5.0, was centrifuged at 59 780 rev./min. and had $S_{20,w}$ 3.3. Fig. 5 shows the gradient curve, which is slightly asymmetric. The concentration of the sedimenting material calculated from the areas of the gradient curves at various times of sedimentation was 95–101% of that of the starting material. The apparent diffusion coefficients obtained from the gradient curves at different times were 9.2×10^{-7} – 9.8×10^{-7} and showed no dependence upon the time of sedimentation. The values S 3.3, D 7.5×10^{-7} , $\bar{V}$ 0.721 give a molecular weight of 38 000.

Electrophoresis. A 2.8% solution of the S component in 0.2M-acetate buffer, pH 5.0, showed a single boundary on electrophoresis.

Crystallization. To solutions of the S component, 2% (w/v) in 0.05M-phosphate buffer, pH 6.7, solid (NH₄)₂SO₄ was added to give a slightly opalescent solution. Water was added to give a clear solution,

which was left in an open tube at 4–7°. The solution gradually became turbid and had a silky sheen on being stirred. Fig. 7 shows the appearance of the crystalline material suspended in saturated $(\text{NH}_4)_2\text{SO}_4$.

DISCUSSION

Immunological analysis of the F and S fractions of the leucocidin preparation showed more than one antigen in each fraction (Woodin, 1959). The present work establishes that the leucocidin activity results from synergism between one constituent of each fraction, the F and S components of leucocidin. It is probable that neither component is greatly contaminated with other immunologically active substances or with material of greatly different charge or sedimentation properties. A calculation of the degree of polydispersity has not been attempted as it has been found that both components can be easily denatured and it is likely that some denatured material is present in the most highly purified preparation of each component.

SUMMARY

1. An antibody-combining-power method has been developed for the determination of the concentration of the two components of leucocidin.

2. Both the F and the S fractions of the leucocidin preparation contain deoxyribonuclease.

3. The F and the S components of leucocidin have been purified with Dowex-2 ($\times 8$) and Amberlite CG-50.

4. The F component of leucocidin crystallizes from 0.2M-phosphate buffer, pH 6.7, in the form of needles or as hexagonal plates.

5. The slow component of leucocidin has been crystallized by salting out with ammonium sul-

phate from 0.05M-phosphate buffer, pH 6.7, in the form of very fine needles.

6. Both the F and S components of leucocidin behave immunologically as single substances and on electrophoresis or ultracentrifuging show a single boundary.

I am greatly indebted to Dr W. E. van Heyningen for many discussions; to Dr D. W. Henderson, F.R.S., for providing facilities for the large-scale production of the culture filtrates and to Mr R. Elsworth and Mr A. May for carrying this out; to Miss Mollie Barr of the Wellcome Research Laboratories for supplies of staphylococcal antitoxin; to Mr L. Taylor for technical assistance.

The ultracentrifuge and electrophoresis apparatus were obtained by a grant to Dr W. E. van Heyningen from the Wellcome Trustees, and part of the work described in this paper was assisted by a further grant to Dr W. E. van Heyningen from the United States Office of Naval Research (Project 103-474).

REFERENCES

- Boman, H. G. & Westlund, L. E. (1956). *Arch. Biochem. Biophys.* **64**, 217.
 Gladstone, G. P. & van Heyningen, W. E. (1957). *Brit. J. exp. Path.* **38**, 123.
 Kay, R. M., Simmons, N. S. & Dounce, A. L. (1952). *J. Amer. chem. Soc.* **74**, 465.
 Laskowski, M. & Seidel, M. K. (1945). *Arch. Biochem.* **7**, 465.
 Ma, T. S. & Zuazaga, G. (1942). *Industr. Engng Chem. (Anal.)*, **14**, 280.
 Markham, R. (1942). *Biochem. J.* **36**, 790.
 Neisser, M. & Wechsberg, F. (1901). *Z. Hyg. InfektKr.* **36**, 299.
 Ouchterlony, Ö. (1949). *Acta path. microbiol. scand.* **26**, 507.
 Privat de Garilhe, M., Fassina, G., Pochon, F. & Pillet, J. (1958). *Bull. Soc. Chim. biol., Paris*, **40**, 1905.
 Woodin, A. M. (1959). *Biochem. J.* **73**, 225.

Biochem. J. (1960) **75**, 165

Comparative Detoxication

7. THE METABOLISM OF CHLOROBENZENE IN LOCUSTS: EXCRETION OF UNCHANGED CHLOROBENZENE AND CYSTEINE CONJUGATES*

By TERESA GESSNER AND J. N. SMITH

Department of Biochemistry, St Mary's Hospital Medical School, London, W. 2

(Received 24 September 1959)

Chlorinated cyclic hydrocarbons are among the most widely used insecticides but sometimes their value is reduced if the insects can metabolize the poison to inactive products. The importance of

detoxication mechanisms in strains of some insects resistant to 2:2-bis-(*p*-chlorophenyl)-1:1:1-trichloroethane (DDT) is now well known (cf. Metcalf, 1955), and the natural resistance of the grasshopper family probably has a similar metabolic basis (Sternburg & Kearns, 1952).

* Part 6: Kikal & Smith (1959).