of the activity was associated with the soluble fraction.

3. The activity of the thyroid gland, however, was restricted to the soluble fraction.

4. More detailed fractionation studies of cerebral cortex and kidney indicated that the particulate matter associated with enzymic activity corresponded in general to the mitochondrial fraction.

This investigation was supported by Grant B-381, National Institute of Neurological Diseases and Blindness, National Institutes of Health, U.S. Public Health Service.

REFERENCES

- Beinert, H. (1951). *J. biol. Chem.* **190**, 287.
 Booth, V. H. (1938). *J. Physiol.* **91**, 474.
 Cooper, C. & Lehninger, A. L. (1956). *J. biol. Chem.* **219**, 489.
 Datta, P. K. & Shepard, T. H., II (1959). *Arch. Biochem. Biophys.* **81**, 124.
 Greenfield, R. E. & Price, V. E. (1954). *Abstr. Amer. chem. Soc.*: September, 48 c.
 Hogeboom, G. H. (1955). *Un. int. Sci. biol. B*, **21**, 3.
 Hogeboom, G. H. & Schneider, W. C. (1948). *J. biol. Chem.* **172**, 619.
 Hogeboom, G. H. & Schneider, W. C. (1955). In *The Nucleic Acids*, vol. 2, p. 199. Ed. by Chargaff, E. & Davidson, J. N. New York: Academic Press Inc.
 Karler, R. & Woodbury, D. M. (1959). *Fed. Proc.* **18**, 409.
 Kuff, E. L. (1954). *J. biol. Chem.* **207**, 361.
 Meldrum, N. U. & Roughton, F. J. W. (1934). *J. Physiol.* **80**, 113.
 Novikoff, A. B. (1957). *Symp. Soc. exp. Biol., N.Y.*, **10**, 92.
 Paigen, K. (1956). *Biochim. biophys. Acta*, **19**, 297.
 Roughton, F. J. W. & Clark, A. M. (1951). In *The Enzymes*, vol. 1, part 2, p. 1250. Ed. by Sumner, J. B. & Myrbäck, K. New York: Academic Press Inc.
 Schneider, W. C. & Hogeboom, G. H. (1952). *J. biol. Chem.* **198**, 155.

Biochem. J. (1960) **75**, 543

The Metabolism of Phenolic Antioxidants

2. THE METABOLISM OF BUTYLATED HYDROXYANISOLE IN THE RAT*

By B. D. ASTILL, D. W. FASSETT AND R. L. ROUDABUSH

Laboratory of Industrial Medicine, Eastman Kodak Company, Rochester, New York, U.S.A.

(Received 28 September 1959)

Commercial 'butylated hydroxyanisole' consists of a mixture of 15% or less of 2-*tert*.-butyl-4-hydroxyanisole (isomer A) and 85% or more of 3-*tert*.-butyl-4-hydroxyanisole (isomer B). The introduction of butylated hydroxyanisole as a fat-stabilizer about 10 years ago, and its subsequent extensive use in edible fats, has been accompanied by studies of its acute and chronic toxicity (e.g. Graham, Teed & Grice, 1954). Wilder & Kraybill (1948), found none of the substance, other than questionable trace amounts in depot and carcass fat of rats receiving 3% of butylated hydroxyanisole in the diet over a prolonged period. Studies on the storage of the compound in dogs given daily doses of 0.3, 30 and 100 mg./kg. for more than a year (Hodge & Fassett, unpublished work) also failed to reveal any butylated hydroxyanisole in perirenal, omental or subcutaneous fat or in the brain, liver and kidney.

The metabolism of butylated hydroxyanisole has been studied in the rabbit by Dacre, Denz &

Kennedy (1956), who found that an oral 0.5 g. dose was excreted as glucuronide (60%), ethereal sulphate (12%) and free phenols (4% of the dose). This dosage is very large compared with human intake. The ingestion of phenolic antioxidants as a result of their use in fat stabilization is probably less than 0.1 mg./kg. It is known that variations in dose may alter the proportions of metabolites excreted, where more than one metabolic pathway is available (cf. phenol; Bray, Thorpe & White, 1952). To justify extrapolation from experimental doses to the usage dose, it is desirable to study the pattern of metabolism with a variety of doses, including the lowest possible so that any unsuspected changes in metabolic pattern may be detected. This paper deals in this way with the metabolism in the rat of butylated hydroxyanisole and its component isomers.

METHODS AND MATERIALS

Animals, diets and dosage. Male and female albino rats of Sprague-Dawley strain, average wt. of 128 rats 321 g. (224–443 g.), were fed with Purina Chow and kept singly or

* The paper by Astill, Fassett & Roudabush (1959) is to be regarded as Part 1.

in pairs in metabolism cages. Butylated hydroxyanisole, isomer A or isomer B was administered by oral injection in one of the media shown in Table 5. Control values for metabolites were established by analysis of the pooled urine collected for 1 day before dosing. Urines were analysed immediately or kept frozen until analysis. The total faeces in Expts. 14 and 16 were collected and analysed for unabsorbed butylated hydroxyanisole.

Materials. Butylated hydroxyanisole, m.p. 54–58°, b.p. 264–270° at 733 mm. Hg, isomer A, m.p. 62.5–64.0°, and isomer B, m.p. 64.5°, were kindly supplied by the Tennessee Eastman Co., Kingsport, Tenn., U.S.A., to whom we are indebted for the physical data. Chromatographic analysis showed that isomer A contained 3–5% of *p*-methoxyphenol. The following were purchased: 2,6-dichloroquinonechloroimide, 2,2'-dipyridyl, *tert*.-butylquinol, naphtharesorcinol, D-glucuronolactone (Eastman Organic Chemicals, Rochester, N.Y.), β -glucuronidase (ox-liver; Ketodase, Warner-Chilcott, N.Y.), Safranin blue (3-amino-7-dimethylamino-5-phenylphenazinium chloride, Allied Chemical and Dye Corp., N.Y.).

Paper chromatography. Table 1 records R_F values, colour reactions and solvent systems: strips of Whatman no. 1 paper were used in the descending method. Isomer A (2–3 μ g.) and isomer B (1 μ g.) were visually detectable with spray reagents A and C. R_F values of metabolites were determined on the purest available specimens. Chromatograms were scanned at 225 or 275 m μ with an attachment inserted in the cell-holder space of a Beckman model DU spectrophotometer (Table 2). Chromatograms were scanned quantitatively after colour development with a Photovolt Electronic Densitometer model 525 (Photovolt Corp., 95 Madison Avenue, N.Y.).

Estimation of butylated hydroxyanisole on chromatograms. From 5 to 40 μ g. of each isomer was chromatographed in solvent II for 2 hr. or solvent III for 1 hr. The dried papers were sprayed with (i) reagent A, and absorbing areas corresponding to butylated hydroxyanisole were scanned at 445 m μ , or (ii) reagent C (Gibbs's Reagent; Mahon & Chapman, 1951), dried and scanned at 595 m μ . Elution of spots located by co-chromatography gave low recoveries with reagent C spectrophotometrically. Results, recorded graphically by plotting extinction against distance along the paper and integrating, showed a linear relation between total extinction of a spot and amount chromatographed. Considerable variation from experiment to experiment was found: thus 20 μ g. of isomer B, sprayed with reagent C, gave a mean area in thirteen separate experiments of 249 (212–290) mm.², although good replicates were obtained on co-chromatography. A known amount of the isomer to be determined was co-chromatographed in each determination.

Examination of urine

Paper chromatography. Experimental and control urines (10 μ l. each) were compared as in Table 1. Only isomers A and B and a metabolite of isomer A, compound H, could be detected in experimental urines. Scanning of chromatograms in ultraviolet light revealed no metabolites because of absorption by substances of natural origin. Elution and estimation of metabolites located by co-chromatography gave low recoveries.

Ethereal sulphates. The total and inorganic sulphate content was estimated either by Folin's gravimetric or by Sperber's turbidimetric method (Bray & Thorpe, 1954).

Free isomers A and B. Urine (5 or 10 ml.) was gently shaken with ether (60 ml.) for 16–24 hr. The separated ether extract was evaporated to dryness at a temperature below 60–70°. A portion of a solution of the residue in ethanol (0.5 ml.) was chromatographed and the free isomer content was estimated chromatographically. Recoveries from control urine (5 ml.) of added isomer A (0.10–0.72 mg.) were 94 $\pm$ 4%, and of added isomer B (0.10–0.78 mg.) were 94 $\pm$ 3%. The ethereal sulphates of isomers A and B did not interfere with this estimation, as shown by separate experiments in which 0.15 μ mole of isomer A and 0.1 μ mole of isomer B were recovered from control urine (10 ml.) to which the corresponding ethereal sulphates (34 μ moles) had been added. In view of the known instability of butylated hydroxyanisole in acid solution (Dacre *et al.* 1956) the usual estimation of combined phenols by acid hydrolysis followed by ether extraction was omitted.

For glucuronides. Total glucuronides were estimated by a modified naphtharesorcinol method (Astill *et al.* 1959), and in low-dose studies by the following procedure. The glucuronide gum obtained by precipitation of the insoluble lead salts at pH 9.5 (as in the procedure of Kamil, Smith & Williams, 1951) from 10 ml. of urine was redissolved in water. A portion, giving an estimated maximum hydrolysis concentration of 0.4 mM of required glucuronide, was added to 5 ml. of 0.2 M-sodium acetate buffer, pH 4.6, and 2 ml. of β -glucuronidase ($\equiv 10^4$ Fishman units, Talalay, Fishman & Huggins, 1946). The mixture was incubated for 16–24 hr. at 37.4°, extracted with ether (2 $\times$ 50 ml.) and the extract evaporated and analysed chromatographically for the required isomers. Ether extracts of hydrolysates of similarly treated control urines gave unexpectedly high extinctions at 600 m μ with Gibbs's reagent. This was caused by substances, absent from experimental ether extracts, which gave intense colours with reagent C at R_F 0.4–0.6 (solvent II). Recoveries by the enzymic and chromatographic methods of the glucuronide of isomer B (2.3–19.0 mg., 0.14–0.36 mm in the hydrolysis) added to urine (10 ml.) were 67 $\pm$ 4% in nine experiments. Recoveries of this glucuronide (5–20 mg., 0.27 mm in the hydrolysis) from urine without a preliminary lead-salt separation were 24–96% and varied inversely as the volume of urine (2.0–0.4 ml.) in the hydrolysis.

The lead salts of the ethereal sulphates of both isomers precipitated with those of the glucuronides at pH 9.5, but interference with the glucuronide estimation was negligible. Non-enzymic hydrolysis of the ethereal sulphate of isomer A (1.1 mg.) was 1.7% and of that of isomer B (0.95 mg.) was 1.3%. β -Glucuronidase was without action on either sulphate.

Examination of faeces

Total faeces were analysed chromatographically by the ferric chloride-2,2'-dipyridyl method: the excess of reducing power over control faeces was expressed as butylated hydroxyanisole (Astill *et al.* 1959).

RESULTS

Excretion of metabolites by control animals. The average daily excretion by undosed rats was 11.9 mg. (4.8–20.4 mg.) of ethereal sulphates as butylated hydroxyanisole [ethereal sulphate/in-

Table 1. *R_F* values and colour reactions of isomers A and B and their metabolites

Whatman no. 1 filter paper was used throughout in the descending technique. Solvents were: I, organic phase of freshly mixed butanol-acetic acid-water (4:1:5, by vol.); II, organic phase of benzene-acetic acid-water (2:2:1, by vol.); III, organic phase of butanol-formic acid-light petroleum, b.p. 66-75° (1:10:10, by vol.). (1) After 10-15 cm. development; (2) after 35-40 cm. development. Spray reagents were: A, aq. 1% (w/v) silver nitrate mixed with 1 part of aq. 3*N*-NH₃ soln.; B, equal volumes, freshly mixed, of 1% (w/v) sulphuric acid in 3*N*-HCl and aq. 5% (w/v) NaNO₂; the dried paper was sprayed with aq. 2% (w/v) Na₂CO₃; C, ethanolic 0.05% (w/v) 2,6-dichloroquinonechloroimide; the dried paper was sprayed with aq. 2% (w/v) sodium borate, or, D, with aq. *N*-NH₃ soln.; E, *N*-HCl; the dried paper was heated at 70° for 3 min. and treated as in D. - indicates that no colour reaction was given.

Substance	<i>R_F</i>			Colours given by reagents				
	Solvent I	Solvent II	Solvent III	A	B	C	D	E
Isomer A	0.96	0.92	(1) 0.90 (2) 0.82	Blue-brown	Orange	Blue-grey	Dark blue	-
Isomer B	0.96	0.94	(1) 0.90 (2) 0.90	Blue-black	Orange-red	Blue-purple	Purple	-
2,5-Di- <i>tert</i> -butyl-4-hydroxyanisole	0.95	0.94	(1) 0.94	Weak brown	Weak orange	Pink	-	-
<i>p</i> -Methoxyphenol	0.95	0.80	(1) 0.32	Brown	Red	Blue	-	Blue-grey, fades to grey
Ethereal sulphate of isomer A	0.82	0.10	(1) 0.00	-	-	-	-	Bright blue, fades to pink
Ethereal sulphate of isomer B	0.79	0.07	(1) 0.00	-	-	-	-	Very weak blue
Glucuronide of isomer A	0.90	0.00	(1) 0.00	-	-	-	-	Very weak purple
Glucuronide of isomer B	0.89	0.00	(1) 0.00	-	-	-	-	Yellow
Compound H	0.83	0.00	(1) 0.00	-	Orange	Weak purple	Mauve	-
<i>tert</i> -Butylquinol	0.97	.	(1) 0.50	Black	Purple	Blue	Blue-purple	-

organic sulphate ratio 0.137 (0.068–0.201)], determined gravimetrically on fourteen groups each of eight rats, and, as determined by the procedures for low doses, on eight groups each of eight rats, 0.15 mg. (0–0.32 mg.) of glucuronides and 0.05 mg. (0–0.11 mg.) of free phenols, as isomer B.

Tables 3–5 record the excretion of metabolites in dose studies.

Metabolism of isomer A. Administration of a single dose of 0.4 g. of isomer A/kg. caused an increase in urinary ethereal sulphate output sufficient to account for 72 % of the dose, and a much smaller increase in glucuronide output. About 2 % of the dose was recovered from the urine unchanged. Chromatography revealed a minor metabolite, compound H. Repeated doses of 0.5 g./kg. depressed ethereal sulphate output and considerably elevated glucuronide output, the over-all conjugate recovery being slightly lowered. Faecal excretion in repeated-dose studies was 0.2 % of the total dose. In Expt. 15, 0.6 % of the dose was excreted as the impurity *p*-methoxyphenol and 2.6 % as its glucuronide, estimated enzymically and chromatographically.

Metabolism of isomer B. After a dose of 0.4 g./kg., increased glucuronide excretion accounted for 57–71 % and increased ethereal sulphate output for 15 % of the dose; 4 % was excreted unchanged. No minor metabolites were detected. Repeated doses of 0.5 g./kg. lowered the proportion of the dose excreted conjugated. Faecal excretion was 0.3 % of the dose.

Metabolism of butylated hydroxyanisole. A single dose of 0.4 g./kg. was metabolized in a similar manner to a single dose of 0.4 g. of isomer B/kg.; no increase in ethereal sulphate output was caused by the 15 % of isomer A in butylated hydroxyanisole. Repeated doses of 0.5 g./kg. depressed the proportion of the dose excreted conjugated. With single doses of 2–100 mg./kg., consistently high recoveries of the dose from the urine were found. Excretion of butylated hydroxyanisole was at a maximum on the first day after dosing, and the time for complete excretion of the dose decreased as the dose decreased. The glucuronide conjugation remained the principal metabolic pathway at these doses, accounting for 76 % (69–92 %) of the dose. The percentage excretion of unchanged butylated

Table 2. *Absorption spectra of some derivatives of isomers A and B*

Solvents were (a) ethanol and (b) water.

Compound	$\lambda_{\max}$	log ϵ	$\lambda_{\max}$	log ϵ
Isomer A (a)	225	3.79	288	3.56
Potassium 3- <i>tert</i> .-butyl-4-methoxyphenylsulphate (b)	225	3.85	276	3.36
Methyl(triacetyl-3- <i>tert</i> .-butyl-4-methoxyphenyl- β -glucosid)-uronate (a)	225	3.94	280	3.34
3- <i>tert</i> .-Butyl-4-methoxyphenyl- β -glucosiduronic acid (b)	223	3.89	280	3.36
Isomer B (a)	226	3.90	289	3.56
Potassium 2- <i>tert</i> .-butyl-4-methoxyphenylsulphate (b)	224	3.85	278	3.33
2- <i>tert</i> .-Butyl-4-methoxyphenyl- β -glucosiduronic acid (b)	224	3.92	280	3.39
Compound H (b)	225	—	275	—

Table 3. *Urinary excretion of butylated hydroxyanisole, isomer A and isomer B and their metabolites by rats for 5 days after a single dose of 0.4 g./kg.*

Results are expressed as % of dose fed. Compounds were injected by gastric tube as (a) a 50 % (w/v) solution in propylene glycol or (b) a 10 % (w/v) suspension in pea soup (Strained Baby Foods, Beech-Nut Co., Syracuse, N.Y.) diluted with an equal volume of water into each rat in a group of four (Expts. 1–6) or eight (Expt. 7) and urine was collected daily. Values were obtained daily on the pooled daily output of each group. Unless specified otherwise sulphates were measured gravimetrically (Bray & Thorpe, 1954) and glucuronides by the modified naphtha-resorcinol method (Astill, Fassett & Roudabush, 1959).

Compound	Expt.	Medium	Average dose (g./rat)	Free butylated hydroxyanisole	Glucuronide	Ethereal sulphate	Total accounted for
Isomer A	3	a	0.09	2	11	73*	86
	4	b	0.12	1	4†	61	66
	5	b	0.13	2	10†	87	99
Isomer B	2	b	0.11	3	71	16	90
	6	a	0.11	4	57	15*	76
Butylated hydroxyanisole	1	a	0.08	3	61	16*	80
	7	a	0.10	7	82	11	100

* Estimated by Sperber's method (Bray & Thorpe, 1954).

† Estimated enzymically.

hydroxyanisole in the urine decreased as the dose increased.

Metabolites of isomers A and B

The glucuronide of isomer A could not be isolated, but its crude methyl-triacetyl derivative, after separation on acid-washed alumina, was converted into 3-*tert*.-butyl-4-methoxyphenyl- β -glucuronamide, identical with a synthetic specimen. The ethereal sulphate of isomer A was isolated from the glucuronide gum as its Safranin bluish salt (Dodgson, Rose & Spencer, 1955) and was converted by cation exchange into potassium 3-*tert*.-butyl-4-methoxyphenyl sulphate; both salts were identical with synthetic specimens. Compound H was formulated as a 2 (or 3)-*tert*.-butyl-4-hydroxyphenyl sulphate, since mild acid hydrolysis of a chromatographically pure specimen yielded only SO_4^{2-} ions and *tert*.-butylquinol, identified spectroscopically and chromatographically. The glucuronide of isomer B was isolated via its barium salt (Dacre *et al.* 1956) from the glucuronide gum. No ethereal sulphate of isomer B was isolated, although its Safranin bluish salt was readily prepared. No other metabolites of isomers A and B were detected.

EXPERIMENTAL

Melting points are uncorrected.

Examination of urine for metabolites

Metabolism of butylated hydroxyanisole. Urine (860 ml.) from Expt. 18, total dose 15.5 g., was clarified by centrifuging and the sediment reserved; the liquid was extracted with ether (2 $\times$ 800 ml.) by gentle shaking for 2 hr., and the ether extract reserved. Glucuronide gums were obtained in the usual way by precipitation with lead acetate soln. at pH 4.5 and 9.5 (Kamil *et al.* 1951).

Sediment. The dry sediment yielded an ether-soluble fraction (0.25 g.), which two-dimensional chromatography showed to be isomer A mixed with a trace of isomer B.

Ether extract. The semi-crystalline residue yielded on distillation at 35 mm. Hg and 150–155° a crystalline solid (0.41 g.), shown chromatographically to be isomer B [R_f in solvent III (2), 0.90] with a trace of isomer A [R_f in solvent III (2), 0.85].

Glucuronide gum (pH 4.5). The aqueous solution (500 ml.) mixed with saturated barium acetate solution (8 ml.) was slowly concentrated *in vacuo*. An inorganic precipitate was discarded. The filtrate yielded a gum which was redissolved, freed from Ba^{2+} ions with CO_2 , the filtrate evaporated and the residue extracted with ethanol and the ethanol extract concentrated to a gum (4.9 g.). This was shown, by hydrolysis of 0.017 g. with β -glucuronidase, and chromatography of the ether extract of the hydrolysate, to contain 0.26 g. of mixed isomer A and B glucuronides as butylated hydroxyanisole, and no glucuronides other than those normally found in this fraction. The total SO_4^{2-} ion content of the gum, measured gravimetrically, was equivalent to 0.03 g. of ethereal sulphates as butylated hydroxyanisole.

Table 4. Urinary excretion of butylated hydroxyanisole and its metabolites by rats after a single dose

Results are expressed as % of dose fed. Each rat in a group of four (Expts. 12, 13) or eight (Expts. 8–11) received butylated hydroxyanisole by gastric tube in diluted pea soup. Analyses were performed on the pooled daily output of all rats in a group, collected until no further metabolites were detectable. Scripts indicate the number of groups in an experiment. Butylated hydroxyanisole was estimated chromatographically, and glucuronides chromatographically after enzymic hydrolysis, both as isomer B. Ethereal sulphates were estimated gravimetrically.

Expt. no.	Dose		Butylated hydroxy-anisole in medium (%)	Free butylated hydroxy-anisole	Period of excretion (days)	Glucuronide	Period of excretion (days)	Ethereal sulphate	Period of excretion (days)	Total accounted for
	(mg./rat)	(g./kg.)								
8 ¹	31.5	0.1	10	4	7–8	75	5–6	12	3–4	91
9 ¹	13.5	0.05	10	5	3–4	69	5–6	10	2–3	84
10 ¹	7.6	0.025	1	8	3–4	92	4–5	Not detectable	.	100
11 ¹	3.7	0.01	1	14	4–5	.	.	Not detectable	.	.
12 ²	3.6, 3.9	0.01	1	7, 12	2–3	70, 74	2–3	Not detectable	.	81, 82
13 ²	0.82, 0.85	0.002	0.35	9.5, 11	2–3	72*	2–3	Not detectable	.	82

* Estimated on the pooled daily output of both groups.

Table 5. *Urinary excretion of butylated hydroxyanisole and metabolites during successive doses of 0.5 g./kg.*

Results are expressed as % of dose fed. Compounds were injected by gastric tube: (a) as a 50% (w/v) solution in propylene glycol; (b) as a 10% (w/v) suspension in diluted pea soup; (c) as a 50% (w/v) solution in corn oil. Each rat of a group of rats was given successive daily doses. Urines were collected for the duration of dosing and pooled. Unless specified otherwise, sulphates were measured gravimetrically and glucuronides by the modified naphtharesorcinol method. Superscript numerals indicate the number of doses per rat.

Compound	Expt. no.	Total average dose/rat (g.)	Medium	No. of rats	Free butylated hydroxyanisole	Glucuronide	Ethereal sulphate	Total accounted for
Isomer A	14	0.81	b	4 ⁵	6	47	23*	76
	15	0.77	c	6 ⁵	5 1†	24† 10†	30 5†	59 16†
	16	0.52	a	3 ⁵	7	43	6*	56
Isomer B	17	1.05	b	6 ⁴	2	70†	3	75
Butylated hydroxyanisole	18	1.94	b	8 ¹⁰	3	46	3	52

* Estimated by Sperber's method (Bray & Thorpe, 1954).

† Estimated enzymically.

‡ Excreted in 12 days after the last dose in Expt. 15.

Glucuronide gum (pH 9.5). The aqueous solution (750 ml.), mixed with saturated barium acetate solution (10 ml.) and adjusted to pH 10.0, was slowly concentrated *in vacuo*. Needles of the barium salt of the glucuronide of isomer B (10.2 g.) separated and were converted into 2-*tert*.-butyl-4-methoxyphenyl- β -glucosiduronic acid, m.p. 168–169°, by the method of Dacre *et al.* (1956) (Found: C, 57.5; H, 6.8; Calc. for C₁₇H₂₄O₈: C, 57.3; H, 6.8%). The i.r. spectrum had a strong band at 5.75 μ (carboxyl C=O) and enhanced absorption, compared with isomer B, at 3 μ (OH).

The supernatant of the glucuronide precipitation was freed from Ba²⁺ ions with CO₂, and the filtrate gave on evaporation *in vacuo* a clear-yellow gum (19.4 g.), shown by enzymic hydrolysis to contain 0.39 g. of unprecipitated glucuronides of butylated hydroxyanisole. The gum was twice extracted with hot ethanol (100 ml.) and since the ethanol-insoluble portion contained no metabolites it was discarded. Comparison of the ethanol-soluble gum with the corresponding gum from control rat urine by enzymic hydrolysis and two-dimensional chromatography of the ether extract of the hydrolysate revealed that the only glucuronides not normally present were those of isomer A and B.

Residual urine. After the removal of lead with H₂S, the combined filtrate and washings (1.5 l.) were concentrated *in vacuo* to low volume, stirred for 1 hr. with stearic acid-deactivated charcoal (15 g.; Asatoor & Dalglish, 1956), the charcoal was collected and the filtrate, which was chromatographically free from detectable metabolites of butylated hydroxyanisole, discarded. The substances adsorbed by the washed charcoal were removed by elution with 1 l. each of (successively) aq. 7.2% (v/v) phenol, aq. 6N-NH₃ soln.-aq. 7.2% (v/v) phenol (1:5, v/v) and 6N-HCl-aq. 7.2% (v/v) phenol (1:5, v/v). Paper-chromatographic examination of the phenol-free residues from the eluates revealed no substances, other than a trace of compound H, which were not present in the corresponding fractions of control urine.

Metabolism of isomer A. (i) *Isolation of the glucuronide as its amide.* Clarified urine (260 ml.) from Expt. 14 was shown chromatographically to contain compound H and isomer A; its ether extract contained isomer A, the impurity *p*-methoxyphenol and normally occurring phenols.

The pH 4.5 glucuronide gum yielded no combined isomer A, and enzymic hydrolysis showed the pH 9.5 glucuronide gum (1.9 g.) to contain the glucuronides of isomer A and, in trace amounts, of *p*-methoxyphenol. Solvent extraction of this gum, partition chromatography, methylation and acetylation yielded intractable products. It was extracted with cold propan-2-ol (3 $\times$ 350 ml.), the slightly dextro-rotatory extract was evaporated to an amorphous solid which was methylated and acetylated, the dried product extracted with boiling hexane (200 ml.) and the extract chromatographed on a column of acid-washed alumina prepared from a hexane slurry. Fractions (100 ml.) were selected by extinction at 280 m μ ; those eluted in order by benzene-CHCl₃ (2:3, v/v), benzene-CHCl₃ (1:4, v/v) and CHCl₃ were combined and evaporated. The residue was dissolved in methanol (20 ml.), the chilled solution was saturated with anhydrous NH₃ and kept for 24 hr. at 5°. The methanol was removed and the residue crystallized from water to yield white prisms of 3-*tert*.-butyl-4-methoxyphenyl- β -glucosiduronamide (8.5 mg.), m.p. 237.0–238.5° (decomp.) not depressed on admixture with a synthetic specimen (Found: C, 57.1; H, 7.5. C₁₇H₂₂O₇N requires C, 57.4; H, 7.1%); natural and synthetic materials had identical u.v. and i.r. absorptions. We were unable to obtain the barium salt of this glucuronide prepared by Dacre *et al.* (1956).

(ii) *Isolation of the ethereal sulphate.* The copious precipitate of lead salts, obtained at pH 9.5 from clarified urine (510 ml.) from Expt. 15, was suspended in water and lead was removed as PbS; the solution was partly concentrated at 35° *in vacuo* to remove H₂S, adjusted to pH 9.5 with K₂CO₃ and evaporated *in vacuo* at 40° to a gum. This was extracted with cold ethanol (500 ml.), and the ethanol extract yielded a clear gum, from which, on addition of saturated Safranin bluish solution (10 ml.), red needles separated (1.26 g.), which crystallized from aq. ethanol. 3-Amino-7-dimethylamino-5-phenylphenazinium 3-*tert*.-butyl-4-methoxyphenyl sulphate (0.42 g.) had m.p. 230–235° (Found: C, 64.4; H, 5.9; N, 10.4. C₃₁H₃₄O₅N₄S requires C, 64.8; H, 6.0; N, 9.7%); i.r. and u.v. spectra were identical with those of synthetic material (m.p. 233–235°), also R_F (0.87) in solvent I. The Safranin salt (400 mg.) was shaken for 2 hr. with Dowex-50 (K⁺ form), and the solution was filtered and concentrated *in vacuo*; the residue, recrystal-

lized from ethanol, yielded *potassium 3-tert.-butyl-4-methoxyphenyl sulphate* (74 mg., 37%). The plates, recrystallized from propan-2-ol, had an indeterminate m.p. (Found: C, 43.8; H, 5.2; K, 13.1. $C_{11}H_{15}O_5SK$ requires C, 44.3; H, 5.1; K, 13.1%); i.r. and u.v. spectra, chemical and chromatographic properties were identical with those of synthetic material.

Properties and attempted isolation of compound H. Compound H was detected on chromatograms in solvent I of urine from all rats dosed with isomer A. Its absence from urine of rats dosed with *p*-methoxyphenol showed that it was not a metabolite of this impurity in isomer A. Compound H remained in urine after extraction with ether and after precipitation of glucuronides. Lead-free urine thus treated from Expt. 15 was concentrated to a stiff gum which was macerated under hot acetone (1 l.) and the extract evaporated to a gum; the acetone-insoluble portion was discarded. The gum was stirred in water (100 ml.) for 1 hr. with stearic acid-deactivated charcoal, when compound H was completely adsorbed. The adsorbed substances were eluted with aq. 7.2% (v/v) phenol and aq. 6N-NH₃ soln.—aq. 7.2% (v/v) phenol (1:5, v/v), and the eluates on evaporation *in vacuo* and removal of phenol yielded gummy residues which were combined (0.52 g.), streaked on eight sheets of Whatman no. 4 paper, 18.5 in. × 22 in., and chromatographed in solvent I. Compound H, located by co-chromatography, was eluted from the required cut-out zones with solvent I and the eluate was evaporated to a brown gum, which was washed with benzene and propan-2-ol. The light-brown semi-crystalline residue (10 mg.) had the reactions shown in Table 1 and the light-absorption maxima shown in Table 2. Compound H gave no Fe³⁺ ion or ninhydrin reactions on paper and no naphtharesorcinol reaction in solution. The i.r. spectrum had bands at 3.0 and 9.5 μ (S:O). Gentle warming with 3N-HCl liberated SO₄²⁻ ions and an ether-extractable phenol, which had the chromatographic properties (Table 1) and the u.v. spectrum of *tert*.-butylquinol. Both the liberated phenol and *tert*.-butylquinol gave in aq. ethanol the same pale-yellow Fe³⁺ ion colour, which deepened on addition of 1 drop of 2N-NaOH. Compound H is probably a 2 (or 3)-*tert*.-butyl-4-hydroxyphenyl sulphate, derived from isomer A by *O*-demethylation and subsequent ethereal sulphate conjugation of either hydroxyl, or by *O*-demethylation of the ethereal sulphate of isomer A.

Metabolism of isomer B. Continuous ether extraction of the urine (120 ml.) from Expt. 16 yielded isomer B, b.p. 150–155° at 35 mm. Hg, identical with the material fed. The glucuronide gum yielded 2-*tert*.-butyl-4-methoxyphenyl- β -glucosiduronic acid (0.36 g.) as before, m.p. and mixed m.p. 168–169°. Exhaustive chromatographic examination of the enzymic hydrolysates of the glucuronide gum and of the glucuronide-free urine revealed no further metabolites of isomer B. No ethereal sulphate Safranine bluish salt was obtained from the glucuronide gum.

Synthesis of ethereal sulphates

These were prepared by the method of Feigenbaum & Neuberger (1941).

Potassium 3-tert.-butyl-4-methoxyphenyl sulphate. A mixture of carbon disulphide (20 ml.), diethylaniline (5 ml.) and chlorosulphonic acid (2 g.) was slowly added with stirring to a chilled solution of isomer A (3 g.) in carbon disulphide (20 ml.) and diethylaniline (2 ml.). After 3 hr.,

an excess of aq. 50% (w/v) KOH was added slowly, followed by ether (200 ml.); the precipitate was triturated with ether (200 ml.) and extracted with hot propan-2-ol. The cooled extract yielded colourless prisms (2.4 g., 48%) of indeterminate m.p. (Found: C, 44.2; H, 5.0; S, 10.8; OCH₃, 11.0; K, 13.1. $C_{11}H_{15}O_5SK$ requires S, 10.7; OCH₃, 10.4%) having absorption at 9.42 μ (S:O). It decomposed at 65° *in vacuo* and above 120°. On gentle warming of a portion with 2N-HCl, isomer A was detected chromatographically in the ether extract of the solution and the solution also gave a strong SO₄²⁻ ion reaction. An insoluble lead salt was formed at pH 9.5. Addition of saturated aq. Safranine bluish solution to aq. 10% (w/v) potassium 3-*tert*.-butyl-4-methoxyphenyl sulphate yielded lustrous green-red needles, recrystallized from aq. ethanol. 3-*Amino-7-dimethylamino-5-phenylphenazinium 3-tert.-butyl-4-methoxyphenyl sulphate* had m.p. 233–235° (Found: C, 64.8, 64.7; H, 5.8, 6.0; N, 9.9; S, 5.3. $C_{31}H_{34}O_5N_4S$ requires S, 5.6%), *R_F* in solvent I, 0.87. It had λ_{max} . 229 (log ϵ_{max} . 4.28) and 274 m μ (log ϵ , 4.69). The Safranine salt (50 mg.) in water (10 ml.) was shaken for 1 hr. with Dowex-50 (K⁺ form) (5 g.), the solution was filtered, evaporated to dryness *in vacuo* and the residue, recrystallized from propan-2-ol, yielded potassium 3-*tert*.-butyl-4-methoxyphenyl sulphate (16.4 mg., 65%).

Potassium 2-tert.-butyl-4-methoxyphenyl sulphate. This, prepared as described above from isomer B in 50% yield, recrystallized from propan-2-ol as prisms, had an indeterminate m.p. (Found: C, 44.3; H, 5.2; S, 10.7; OCH₃, 10.1; K, 13.2. $C_{11}H_{15}O_5SK$ requires C, 44.3; H, 5.1; S, 10.7; OCH₃, 10.4; K, 13.1%) and strong absorption at 9.45 μ . Its chemical properties were similar to those of the ethereal sulphate of isomer A. 3-*Amino-7-dimethylamino-5-phenylphenazinium 2-tert.-butyl-4-methoxyphenyl sulphate*, obtained as before, gave red needles from aq. ethanol, m.p. 216–218° (Found: C, 63.8; H, 6.1. $C_{31}H_{34}O_5N_4S$, 0.5 H₂O requires C, 63.8; H, 6.0%) and had λ_{max} . 229 (log ϵ_{max} . 4.28) and 273 m μ (log ϵ_{max} . 4.69).

Synthesis of derivatives of the glucuronide of isomer A

Methyl (tri-O-acetyl-3-tert.-butyl-4-methoxyphenyl- β -glucosid)uronate. This was prepared by the procedure of Bollenback, Long, Benjamin & Lindquist (1955). Methyl (tetra-O-acetyl- β -glucopyranosid)uronate (5 g.), isomer A (5 g.) and freshly fused ZnCl₂ (1 g.), dissolved in 20 ml. of acetic anhydride-acetic acid (1:20, v/v), were heated at 100° for 20 min., excess of solvent was removed *in vacuo* at 100° for 20 min., the residue was dissolved in benzene (100 ml.) and the solution washed with 2N-NaOH (2 × 100 ml.) and water (100 ml.), dried over MgSO₄ and concentrated *in vacuo*. A semi-crystalline residue resulted after several hours at 1.5 mm. Hg, and after trituration with hexane colourless needles were collected and recrystallized from hexane. *Methyl (tri-O-acetyl 3-tert.-butyl-4-methoxyphenyl- β -glucosid)uronate* (1.6 g.) had m.p. 122–123°, $[\alpha]_D^{24}$ –25.1° in CHCl₃ (c, 16.7) (Found: C, 57.6; H, 6.7; OCH₃, 12.9. $C_{24}H_{32}O_{11}$ requires C, 58.0; H, 6.5; OCH₃, 12.5%). Chromatographic treatment of the combined mother liquors on a column of acid-washed alumina yielded a further 600 mg. of product, m.p. 118–120.5° (total yield 15%). Longer heating of the reactants (90 min. at 100°) yielded, on chromatography of the product, needles of 3-*tert*.-butyl-4-methoxyphenyl acetate (4.5 g.),

m.p. 56–58°, identical with that prepared by acetylation of isomer A in pyridine and acetic anhydride, m.p. 58.0–58.5° (Found: C, 70.2; H, 8.5; OCH₃, 13.3. C₁₃H₁₈O₃ requires C, 70.2; H, 8.2; OCH₃, 13.9%) and yellow needles (0.65 g.) of the anomeric methyl (tri-*O*-acetyl-3-*tert*.-butyl-4-methoxyphenyl- α -glucosid)uronate, m.p. 144–146°, $[\alpha]_D^{24} + 127^\circ$ in CHCl₃ (c, 9.1) (Found: C, 58.4; H, 6.5%), but no β -anomer. Anhydrous NH₃ was passed for 0.5 hr. into a chilled solution in methanol (30 ml.) of the β -anomer (300 mg.), the solution was chilled for 3 days, the solvent removed and the residue crystallized from water to yield needles (135 mg.) of 3-*tert*.-butyl-4-methoxyphenyl- β -glucosiduronamide, m.p. 238.5° (decomp.), $[\alpha]_D^{24} - 188^\circ$ in ethanol (c, 0.9) (Found: C, 57.0; H, 7.3; N, 4.3. C₁₇H₂₅O₇N requires N, 3.9%). The i.r. spectrum showed strong bands at 2.84, 2.93 and 5.98 μ and the u.v. spectrum had λ_{max} . 226 (log ϵ 3.93) and 284 m μ (log ϵ , 3.43).

Attempted preparation of the glucuronide of isomer A. Hydrolysis of the methyl tri-*O*-acetyl- β -anomer with Ba(OH)₂ solution or aq. methanolic suspensions of Ba(OH)₂ yielded amorphous products. Catalytic deacetylation of 1.34 g. with sodium (0.05 g.) in methanol (300 ml.) yielded, after 6 days, a semi-crystalline mass on evaporation *in vacuo*, which was purified through its lead salt or over Dowex-2 (H⁺ form) to give a residue which separated from propan-2-ol as a glass (270 mg.), $[\alpha]_D^{24} - 46.5^\circ$ in water (c, 3.4). The product gave a strong positive naphtharesorcinol reaction and had pH 3.7 in 0.1% solution. Hydrolysis with β -glucuronidase (10⁴ Fishman units) of 2.8 μ moles gave 2.45 μ moles (84%) of isomer A, estimated chromatographically, and u.v. (Table 2) and i.r. spectra (bands at 3.0, 5.8 μ) were closely similar to those of the glucuronide of isomer B. This was the purest specimen prepared of the glucuronide of isomer A.

DISCUSSION

The metabolism of butylated hydroxyanisole in the rat resembles that reported for the rabbit for doses of 0.13–0.55 g./kg. (Dacre *et al.* 1956). At all doses studied in the rat butylated hydroxyanisole and its component isomers, on ingestion, are absorbed and readily metabolized. The possible metabolic pathways for phenols and substances analogous to butylated hydroxyanisole (Williams, 1949; Brodie, Gillette & LaDu, 1958) suggest that 4-*O*-conjugation, *O*-demethylation, hydroxylation of the phenyl ring and side-chain oxidation ($-\text{CH}_3 \rightarrow \text{CH}_2\cdot\text{OH}$) might be involved in metabolism of butylated hydroxyanisole. However, the main metabolites are 4-*O*-conjugates, i.e. the *O*-sulphate and *O*-glucuronide of isomer A and the *O*-glucuronide of isomer B. The only other metabolic pathway is the probable *O*-demethylation of isomer A, which yields compound H. In the rabbit both isomers are metabolized largely by 4-*O*-glucuronide formation (Dacre *et al.* 1956).

The conjugation of isomer B appears to differ from that predictable from steric considerations. The 4-hydroxyl group in isomer B is clearly crowded by the 3-*tert*.-butyl group, such that conjugation

with the ethereal sulphate group would seem preferable to conjugation with the bulkier glucuronide group (Fisher–Taylor–Hirschfelder Models, Fisher Scientific Co., Pittsburgh, Pa., U.S.A.). However, isomer B forms mostly an *O*-glucuronide. Isomer A does not appear to possess steric restrictions on conjugation, and forms mainly an ethereal sulphate, *O*-glucuronide formation apparently becoming significant only when the sulphate pool is depleted by repeated large doses.

In the single-dose studies, at the lowest dose (2 mg. of butylated hydroxyanisole/kg.) the background absorption of chromatograms in the zones examined for isomer B, owing to the excretion of normal metabolites, is no longer negligible. Thus the apparent glucuronide excretion (enzymic method) of controls, expressed as isomer B, amounts to 18% (0–39%)/day of a dose of 0.82 mg. of butylated hydroxyanisole/rat. The apparent excretion of free isomer B by controls amounts to 6% (0–13.5%)/day of the same dose. In Expt. 13, however, actual control values, determined on the day before dosing, were glucuronide, 0%, and free isomer B, 3.8–4.0%, of the dose, and were assumed to be unaltered during urine collection. In other experiments at this dose, not reported, background absorption was too high to enable a reliable estimation of dose excretion to be made. This dose is clearly a lower limit for the method.

Over the range studied, the excretion of unconjugated butylated hydroxyanisole decreased from 11% (7–14%) for a 10 mg./kg. dose to 3–7% for a 400 mg./kg. dose. With 0.3 mg./kg. in dogs H. C. Hodge & D. W. Fassett (unpublished) found negligible tissue storage, and in the present study over-all recoveries of a single dose remained uniformly high (82–100%) at all doses studied. The slightly higher proportion of unconjugated butylated hydroxyanisole excreted at the lowest doses is therefore unlikely to be associated with any increase in tissue storage.

The above study approaches closer to human-usage doses than does any previous study on anti-oxidants, and, although the problem of studying the human-usage dose remains, extrapolation to that amount seems justified. No evidence has so far been presented which suggests that butylated hydroxyanisole is not largely absorbed and largely excreted at these very low doses.

SUMMARY

1. The metabolism of butylated hydroxyanisole and 2- and 3-*tert*.-butyl-4-hydroxyanisole in the rat is studied with particular emphasis on the fate of butylated hydroxyanisole at low doses.

2. Increased urinary glucuronide and ethereal sulphate excretions during 5 days after dosing with

0.4 g. of butylated hydroxyanisole/kg. account for 61–82 and 11–16 % respectively of the dose, and 5 % of the dose is excreted, unchanged.

3. For single oral doses of 0.002, 0.01, 0.025, 0.05 and 0.1 g. of butylated hydroxyanisole/kg., overall recoveries of the dose from the urine were 81–100 %. Glucuronide excretion accounts for 69–92 % of the dose, and a slight increase in excretion of unchanged butylated hydroxyanisole occurs at the lowest doses.

4. For single oral doses of 0.4 g. of 2-*tert*.-butyl-4-hydroxyanisole 72 % of the dose is excreted as ethereal sulphate in 5 days. With 3-*tert*.-butyl-4-hydroxyanisole 57–71 % of the dose is excreted as glucuronide in 5 days.

5. The administration of several successive daily doses of 0.5 g. of butylated hydroxyanisole/kg., and of 3-*tert*.-butyl-4-hydroxyanisole, decreases the percentage recovery of conjugates and the proportion of ethereal sulphate excreted during dosing; repeated doses of 0.5 g. of 2-*tert*.-butyl-4-hydroxyanisole/kg. cause a considerable increase in glucuronide and a decrease in ethereal sulphate conjugation of the dose.

6. The isolation of the major metabolites, the probable formation of a demethylation product of 2-*tert*.-butyl-4-hydroxyanisole as a minor metabolite, the synthesis of the ethereal sulphates of both isomers and the synthesis of the glucuronamide of 2-*tert*.-butyl-4-hydroxyanisole are described.

We wish to thank Dr C. J. Terhaar, and the staff of the Toxicology Laboratory, for the care and dosing of experi-

mental animals; Mr Stuart Foster of the Industrial Laboratory for infrared and ultraviolet spectra; and Mr D. F. Ketchum of the Microanalytical Laboratory for microanalyses.

REFERENCES

- Asatoor, A. & Dalgliesh, C. E. (1956). *J. chem. Soc.* p. 2291.
 Astill, B. D., Fassett, D. W. & Roudabush, R. L. (1959). *Biochem. J.* **72**, 451.
 Bollenback, G. N., Long, J. W., Benjamin, G. D. & Lindquist, J. A. (1955). *J. Amer. chem. Soc.* **77**, 3310.
 Bray, H. G. & Thorpe, W. V. (1954). *Meth. biochem. Anal.* **1**, 47.
 Bray, H. G., Thorpe, W. V. & White, K. (1952). *Biochem. J.* **52**, 423.
 Brodie, B. B., Gillette, J. R. & LaDu, B. N. (1958). *Annu. Rev. Biochem.* **27**, 427.
 Dacre, J. G., Denz, F. A. & Kennedy, T. H. (1956). *Biochem. J.* **64**, 777.
 Dodgson, K. S., Rose, F. A. & Spencer, B. (1955). *Biochem. J.* **60**, 346.
 Feigenbaum, J. & Neuberg, C. A. (1941). *J. Amer. chem. Soc.* **63**, 3529.
 Graham, W. D., Teed, H. & Grice, H. C. (1954). *J. Pharm., Lond.*, **6**, 534.
 Kamil, I. A., Smith, J. N. & Williams, R. T. (1951). *Biochem. J.* **50**, 235.
 Mahon, J. H. & Chapman, R. A. (1951). *Analyt. Chem.* **23**, 1120.
 Talalay, P., Fishman, W. H. & Huggins, C. (1946). *J. biol. Chem.* **166**, 757.
 Wilder, O. H. M. & Kraybill, H. R. (1948). *Summary of Toxicity Studies on Butylated Hydroxy Anisole*, pp. 1–5. Chicago: American Meat Institute Foundation, University of Chicago.
 Williams, R. T. (1949). *Detoxication Mechanisms*. New York: John Wiley and Sons.

Biochem. J. (1960) **75**, 551

Partial Purification and some Properties of the Blood-Clotting Factor from the Venom of *Bothrops jararaca*

By OLGA B. HENRIQUES, MINA FICHMAN AND S. B. HENRIQUES
Instituto Butantan, C.P. 65, São Paulo, Brazil

(Received 18 September 1959)

The blood-clotting activity of the venom of various species of *Bothrops* snakes has been studied by various authors, some of them attributing the clotting effect to the proteolytic activity of the venom (Jánszky, 1950; Zeller, 1951; Slotta, 1953), and others (Taborda, 1940; Deutsch & Diniz, 1955; Rosenfeld, Hampe & Kelen, 1960) giving evidence that separate enzymes are concerned with the two activities. Michl (1954) and Holtz & Raudonat (1956) were able to show that in the venom of

Bothrops jararaca the clotting activity is distinguishable from its proteolytic activity. However, Hamberg & Rocha e Silva (1956) and Henriques, Lavras & Fichman (1956) have shown that in the venom of *B. jararaca* there exist at least two proteolytic enzymes. Therefore one of them could be responsible for the clotting activity. Nevertheless, Henriques, Mandelbaum & Henriques (1959) demonstrated that the clotting activity of the venom of *Bothrops jararaca* is independent of those