sented for the existence of a bound form of ubiquinone when the heart-muscle preparation was suspended in a medium containing inorganic phosphate. It is possible that part of the oxidative-phosphorylation process is still intact in these preparations and that this bound form is a phosphate-containing derivative of ubiquinone or ubiquinol and is an intermediate in the process. Such a role would be consistent with scheme (II) in Fig. 3.

SUMMARY

1. Methods for studying the kinetics of endogenous ubiquinone in heart-muscle preparations are described.

2. The rate of reduction of ubiquinone was not equivalent to the steady-state rate of electron transfer as measured by the rate of oxidation of the substrate.

3. Studies with inhibitors indicated that ubiquinone was reduced by the flavoproteins and oxidized by way of the antimycin A-sensitive factor and cytochrome oxidase.

4. It is suggested that ubiquinone is a branch pathway of the respiratory chain, and that it may function either as an interchain carrier or as an intermediate in oxidative phosphorylation.

We wish to thank Professor R. A. Morton, F.R.S., for his interest and encouragement in this work. We are indebted to the Department of Scientific and Industrial Research for a Research Studentship to one of us (A.M.P.).

REFERENCES

- Chance, B. (1952). *Nature, Lond.*, **169**, 215.
 Chance, B. (1957). In *Proceedings of the International Symposium on Enzyme Chemistry, Tokyo and Kyoto, 1957*, p. 9. London: Pergamon Press Ltd.
 Chance, B. (1958). *J. biol. Chem.* **233**, 1223.
 Colpa-Boonstra, J. & Holton, F. A. (1959). *Biochem. J.* **72**, 4P.
 Crane, F. L., Hatefi, Y., Lester, R. L. & Widmer, C. (1957). *Biochim. biophys. Acta*, **25**, 220.
 Giuditta, A. & Singer, T. P. (1959). *J. biol. Chem.* **234**, 662.
 Green, D. E., Hatefi, Y. & Fechner, W. F. (1959). *Biochem. Biophys. Res. Commun.* **1**, 45.
 Harrison, K. (1958). *Nature, Lond.*, **181**, 1131.
 Hatefi, Y. (1959). *Biochim. biophys. Acta*, **31**, 501.
 Hatefi, Y., Lester, R. L., Crane, F. L. & Widmer, C. (1959). *Biochim. biophys. Acta*, **31**, 490.
 Joel, C. D., Karnovsky, M. L., Ball, E. G. & Cooper, O. (1958). *J. biol. Chem.* **233**, 1565.
 Keilin, D. & Hartree, E. F. (1952). *Biochem. Prep.* **2**, 1.
 Kimura, T. & Singer, T. P. (1959). *Nature, Lond.*, **184**, 791.
 Lester, R. L. & Fleischer, S. (1959). *Arch. Biochem. Biophys.* **80**, 470.
 Pumphrey, A. M. & Redfearn, E. R. (1959). *Biochem. J.* **72**, 3P.
 Pumphrey, A. M. & Redfearn, E. R. (1960). *Biochem. J.* **76**, 61.
 Pumphrey, A. M., Redfearn, E. R. & Morton, R. A. (1958). *Chem. & Ind.* p. 978.
 Redfearn, E. R. (1959). *Disc. Faraday Soc.* **27**, 260.
 Singer, T. P. & Kearney, E. B. (1957). *Meth. biochem. Anal.* **4**, 307.
 Slater, E. C. (1958). *Advanc. Enzymol.* **20**, 147.
 Tsou, C. L. (1951). *Biochem. J.* **49**, 512.
 Wu, C. Y. & Tsou, C. L. (1955). *Sci. Sinica*, **4**, 137.

Biochem. J. (1960) **76**, 71

Studies on the Biosynthesis of Proteins in the Silkworm, *Bombyx mori* (L.)

BY P. FAULKNER AND B. BHEEMESWAR*

Insect Pathology Research Institute, Sault Ste. Marie, Ontario, Canada

(Received 14 September 1959)

Although extensive studies have been made on the biosynthesis of proteins in animals, plants and micro-organisms, relatively little is known about similar mechanisms in insects. The use of labelled amino acids for studying the synthesis of silk by the giant silkworm, *Hyalophora (Platysamia) cecropia* (L.), was first reported by Zamecnik, Loftfield, Stephenson & Williams (1949), who showed that silk spun 24 hr. after injection of a mixture of carboxy-

labelled glycine and DL-alanine was radioactive. They also provided evidence for the incorporation of [1-¹⁴C]glycine *in vitro* by incubating excised silk glands aerobically in Krebs-Ringer phosphate medium. Shimura, Koide, Itabashi & Fukai (1955) demonstrated net fibroin synthesis in perfused silk glands and studied the effects of various metabolic intermediates added to the medium. In a later study, Shimura, Fukai, Suto & Hoshi (1958) showed that [1-¹⁴C]glycine injected into the silkworm was incorporated into the proteins of the silk gland, gut and dermal tissues, and that 4 hr. after injection,

* Post Doctoral Research Fellow of the National Research Council of Canada. Present address: 81 Fruit Street, Bangalore 1, India.

95% of the radioactivity in the fibroin isolated from the silk gland was in the glycine residue. Bricteux-Gregoire, Verly & Florkin (1958) have shown by injection experiments that [^{14}C]glycine serves as a precursor of the serine and alanine moieties of silk and that [^{14}C]phenylalanine (Bricteux-Gregoire, Verly & Florkin, 1956; Fukuda, 1956) is incorporated into silk protein after being converted into tyrosine. Sisakian & Kuvaeva (1957) found that [^{14}C]glycine was incorporated into the protein of the strained brei of silkworm pupae after prolonged incubation. More recently Takayama, Ito & Miura (1958) presented evidence for fibroin synthesis *in vitro* by minces of the posterior silk gland and its dependence on the high ribonucleic acid content of the tissue. These workers also found that the ability to incorporate [^{14}C]glycine into protein was almost completely lost when the silk-gland preparations were homogenized, but there was some incorporation in a preparation containing only microsomes and cell sap.

In the present paper we report on the labelling of tissue protein after injection of [^{14}C]glycine into silkworms at various stages of larval development. Some factors influencing the incorporation *in vitro* of [^{14}C]glycine into excised silk-gland protein have also been studied. Experiments in which [^{14}C]glycine has been incorporated into an ultracentrifuged cell-sap preparation are described and some of the properties of the resulting complex are given.

Some of the results have been reported briefly elsewhere (Faulkner & Bheemeswar, 1959).

EXPERIMENTAL

All the reagents were of analytical-reagent quality. Disodium salts of adenosine triphosphate (ATP) and guanosine triphosphate (GTP) were obtained from Sigma Chemical Co., St Louis, Mo., U.S.A., and adenosine 5-phosphate from Nutritional Biochemicals Corp., Cleveland, Ohio, U.S.A. Crystalline ribonuclease was obtained from Worthington Biochemicals Corp., Freehold, N.J., U.S.A. [^{14}C]Glycine (universally labelled) of specific activity 2.51 mc/m-mole, [^{14}C]glycine (carboxyl labelled) of specific activity 19.1 mc/m-mole and ammonium L-[^{14}C]glutamate of specific activity 5 mc/m-mole were purchased from Atomic Energy of Canada Ltd., Ottawa, and L-[^{14}C]phenylalanine of specific activity 0.244 mc/m-mole was purchased from Schwarz Laboratories Inc., Mt. Vernon, N.Y., U.S.A.

Insects. Silkworms, *Bombyx mori* (L.), of both sexes from the Institute stock, were used in these experiments.

Injection and isolation of tissues. Silkworm larvae were individually injected with 5 μl . of [^{14}C]glycine (2.0 μC) by means of a microsyringe (Bergold, 1941). Except in the study of the rate of incorporation of glycine at different developmental stages, the insects injected with glycine were in the middle of the fifth instar. At suitable intervals after injection insects were bled by cutting a proleg, and the blood was collected in a small test tube kept at 0°. The

protein in 0.1 ml. of blood was precipitated by adding 5 ml. of 5% trichloroacetic acid. Gut, fat body and the posterior silk gland were then dissected out and dropped into tubes containing 5% trichloroacetic acid.

Preparation of minces and method of incubation. The term 'mince' is used to describe a preparation obtained by chopping finely with scissors the posterior silk glands taken from mid-fifth instar larvae. Incubations were carried out in 5 ml. Warburg vessels and, unless otherwise stated, the vessels were incubated aerobically at 37° with shaking. Each Warburg vessel contained 5–10 mg. of the silk-gland mince. The basal incubation medium contained: potassium phosphate buffer, pH 7.3, 20 mM; MgCl_2 , 10 mM; L-malate, 10 mM; [^{14}C]glycine, 0.2 μC , 0.08 mM. Total volume was 1 ml. At the end of the incubation period 0.25 ml. of 25% trichloroacetic acid was added to stop the reaction.

Isolation of protein and radioactive assay. The acid-inactivated tissues or minces were ground in a glass tissue grinder, and the suspension, which contained about 1.0–2.5 mg. of protein (dry wt.), was washed to remove any unincorporated labelled glycine, nucleic acids and lipids, as follows: five times with 5% trichloroacetic acid at room temperature, twice with hot 5% trichloroacetic acid (15 min. at 90°), once with 95% ethanol, twice with alcohol-ether (3:1, v/v) for 3 min. at 60°, and finally with ether. Preliminary experiments indicated that the presence of unlabelled glycine in the cold trichloroacetic acid washings was not necessary. The efficiency of the washing procedure was tested in each experiment by means of a control, in which the tissue was dropped into a tube containing [^{14}C]glycine in 5% trichloroacetic acid. The specific activity of the protein in the control after following the washing schedule was not more than 15 counts/min./mg. of protein.

The washed protein was suspended in 0.2 ml. of water and a portion (0.1 ml.) was applied to a tared stainless-steel planchet and dried under a heat lamp. The planchets were dried to constant weight in a vacuum desiccator and the radioactivity was determined with a Tracerlab TGC-14 thin-window Geiger counter. Corrections were applied for background and self-absorption, and the results were expressed as specific activity (counts/min./mg. of protein). In order to compare the effect of added test substances used in different experiments, the term 'relative specific activity' has been introduced.

$$\text{Relative specific activity} = \frac{\text{Specific activity with test substances}}{\text{Specific activity without test substances}} \times 100.$$

Preparation of the pH 5 extract. The posterior portions of silk glands from mid-fifth instar larvae were used. Wet tissue (5–6 g.) was ground with 20 ml. of medium X (Zamecnik & Keller, 1954) in a glass tissue grinder. The brei was centrifuged for 10 min. at 20 000 g at 0–4° to remove the insoluble particles and the cell debris, and the opalescent supernatant was then centrifuged at 150 000 g for 50 min. in a Phywe ultracentrifuge at 20°. Protein was precipitated from the clear supernatant by slow addition of 0.1 N-HCl with constant stirring until the pH had fallen to 5.1. The protein which dissolved in about 5 ml. of 0.2 M-potassium phosphate buffer, pH 7.8, was termed the pH 5 extract. The insoluble residue was discarded. The protein concentration of the pH 5 extract was 1–3 mg./ml.

Incubation procedure with the pH 5 extract, and radioassay. Incubations were carried out in open 3.5 ml. test tubes without shaking, or in 5 ml. Warburg vessels (when anaerobic conditions were desired). At the end of the incubation period the reaction was stopped by the addition of 2 ml. of 12% trichloroacetic acid. The precipitate was washed five times with portions (3 ml.) of 5% trichloroacetic acid, and twice with ice-cold water.

The residual material was dissolved in 0.15 ml. of 0.1N-KOH and a portion (0.1 ml.) was applied to a stainless-steel planchet for radioassay in a Tracerlab TGC-14 thin-window Geiger counter. The results are expressed as specific activity (counts/min./mg. of protein). The protein content of the material was determined by the method of Ramachandran & Fraenkel-Conrat (1958).

The washing procedure was checked in each experiment by adding [^{14}C]glycine to a tube containing the pH 5 extract with 5% trichloroacetic acid and subjecting the precipitate to the same washing cycle before plating.

The glycine-activating enzyme was assayed by the method of Hoagland, Keller & Zamecnik (1956).

RESULTS

Incorporation of [^{14}C]glycine into tissue proteins of the silkworm. After injection of [^{14}C]glycine, the various tissue proteins in the larvae are rapidly labelled. Proteins of the silk gland, fat body, gut and blood are found to be radioactive within a few minutes. The data in Fig. 1 show that silk-

Fig. 1. Rate of incorporation of [^{14}C]glycine into tissue proteins. Larvae (24) taken on the fifth day of the fourth moult were each injected with $2\mu\text{C}$ of [^{14}C]glycine. At the stated times, four larvae were bled and dissected and the specific activity of the tissue proteins was determined. ○, Silk gland; ●, blood; △, gut; ▲, fat body.

gland protein is most rapidly labelled, and that the rate is roughly linear for the first 30 min. after injection but diminishes later. Proteins of the fat body appear to reach equilibrium with the label more quickly than do those of the silk gland, and there is a lag period before blood protein incorporates glycine at a linear rate. The fat body is believed to be an important site of blood-protein synthesis (Shigematsu, 1958), and the observation that incorporation into blood protein continues after incorporation into fat body reaches equilibrium is in keeping with this hypothesis.

Effect of stage of development on incorporation. Preliminary experiments indicated that there was considerable variation in the rate of glycine incorporation among a random group of fifth-instar larvae. Accordingly, we studied the rate of glycine incorporation with larvae from the same family at various ages in the fourth and fifth instars. Each day four insects were injected, beginning 1 day before the fourth moult and continuing up to the day of spinning (approximately 10 days). Fifteen minutes after injection, the proteins were isolated and the radioactivity was measured. The data given in Fig. 2 show the amount of radioactivity incorporated into the silk-gland protein at different developmental stages. During the period just before and after the fourth moult, when the insect does not feed, the rate of incorporation is low. After this period, the amount of glycine incorporated increases gradually as the insect grows rapidly. Just before spinning the rate of incorporation decreases markedly.

Factors influencing the incorporation of [^{14}C]glycine into the proteins of minces. [^{14}C]Glycine is

Fig. 2. [^{14}C]Glycine incorporation into silk-gland protein during development. Four larvae from the same family and of the same age were injected with $2\mu\text{C}$ of [^{14}C]glycine. After 15 min. the specific activity of the silk-gland protein was measured.

readily incorporated into protein when a silk-gland mince is incubated aerobically in the presence of potassium phosphate, pH 7.3. The incorporation is increased in the presence of Mg^{2+} , Mn^{2+} or Ca^{2+} ions (all at 10 mM) and by certain organic and keto acids. The data in Fig. 3 show the rate of incorporation in phosphate buffer supplemented with L-malate and Mg^{2+} ions. In this experiment, maximum incorporation of $[^{14}C]$ glycine occurred after 25 min. At this time 23% of the added glycine had been incorporated, which is equivalent to a rate of approximately 30 μ moles of glycine/g. of protein/hr. As was shown by Takayama *et al.* (1958), this rate is nearly 200 times that reported for rat-liver slices.

Data showing the effects of sugars, organic acids and keto acids on incorporation are given in Table 1. Sugars generally have little influence; however, glucose is slightly stimulatory and fructose inhibitory. All the organic acids and keto acids tested, with the exception of glyoxalate, had stimulatory effects on glycine incorporation.

Oxidative metabolism appears to be essential for the incorporation of $[^{14}C]$ glycine into silk-gland minces, since, under anaerobic condition, or aerobically in the presence of cyanide, incorporation is strongly inhibited. Considerable impairment of incorporation is observed when K^+ ions in the incubation medium are replaced by Na^+ ions

Fig. 3. Rate of $[^{14}C]$ glycine incorporation into silk-gland minces. Incubation medium contained: silk-gland mince, 5–10 mg.; $MgCl_2$, 10 mM; $[^{14}C]$ glycine (0.2 μ C), 0.08 mM; potassium phosphate, pH 7.3, 20 mM; L-malate, 10 mM. Total volume was 1 ml. Incubated at 37° for times indicated.

(Table 2). Data are also included in the table on the effect of other inhibitors. Chloramphenicol, which is an inhibitor of protein synthesis in other systems, is effective in silk-gland minces at 1 mM concentration. Hydroxylamine and pyruvic oxime at 1 mM concentration are known to inhibit a silk-gland transaminase (Bheemeswar & Faulkner, 1959). Both these substances inhibit incorporation of glycine into silk gland but the concentration required is much higher than for cell-free transaminase systems.

Silk glands are particularly rich in nucleic acids. It is reported that the ribonucleic acid content of the posterior silk gland is 2.6–4.4 mg./g. of fresh tissue (Denucé, 1952). The data in Table 2 confirm

Table 1. *Metabolites affecting the incorporation of $[^{14}C]$ glycine into protein of silk-gland mince*

Basal medium contained: potassium phosphate, pH 7.3, 20 mM; $MgCl_2$, 10 mM; silk-gland mince, 5–10 mg.; $[^{14}C]$ glycine (0.2 μ C), 0.08 mM. Total volume was 1 ml. Incubated for 45 min. at 37°.

Addition to basal medium (final concn., 10 mM)	Relative specific activity
Nil	100
D-Glucose	115
D-Fructose	74
Sucrose	99
α -Trehalose	96
L-Malate	231
Citrate	230
Succinate	161
Fumarate	146
Glyoxylate	55
α -Glycerophosphate	304
Pyruvate	218
Phenylpyruvate	305
Oxaloacetate	206
Oxoglutarate	192

Table 2. *Inhibition of $[^{14}C]$ glycine incorporation into the protein of silk-gland mince*

Basal medium contained: potassium phosphate, pH 7.3, 20 mM; $MgCl_2$, 10 mM; silk-gland mince, 5–10 mg.; $[^{14}C]$ glycine (0.2 μ C), 0.08 mM. Total volume was 1 ml. Incubated for 45 min. at 37°.

Condition tested or addition to basal medium	Inhibition (%)
Anaerobiosis	97
KCN, 10 mM	93
Arsenite, 10 mM	0
Arsenate, 10 mM	23
NaF, 10 mM	66
Sodium phosphate replacing potassium phosphate	72
Hydroxylamine { 10 mM	72
1 mM	6
Pyruvic oxime, 10 mM	48
Pyruvic oxime, 1 mM	6
Chloramphenicol, 1 mM	45
Ribonuclease, 1 mg.	99

the observation of Takayama *et al.* (1958) that pre-incubation of silk gland with ribonuclease leads to an impairment of its ability to incorporate glycine. This result suggests that intact nucleic acids are essential for incorporation.

Factors influencing [¹⁴C]glycine incorporation into the pH 5 extract. The composition of the medium for incorporation of [¹⁴C]glycine into the pH 5 extract is given in Table 3. The results show that the high incorporation rates are obtained in the presence of Mg²⁺ ions and either ATP or GTP. If no nucleotide is present, little incorporation is observed. Stimulation by ATP or GTP was detectable in concentrations as low as 1 mM, which suggests that the two nucleotides are readily inter-converted or that the incorporating enzyme in silk gland is not highly specific for them. Some source of high-energy phosphate is apparently necessary since no incorporation results when adenosine 5-phosphate is substituted for ATP or GTP. The results given in Table 4 show that incorporation proceeds under both aerobic and anaerobic conditions, but, in both cases, it is dependent on the presence of ATP.

The extent of [¹⁴C]glycine incorporation is proportional to the concentration of enzyme in the pH 5 extract (Fig. 4). Time studies have shown that incorporation is rapid for the first 5 min. of incubation at 25°, and then the rate is lower, but steady, for a further 15 min. For example, in one experiment of 20 min. duration, 50 % of the total incorporation was observed in the first 5 min. During the experimental period the amount of glycine incorporated was 0.4 μm-mole/mg. of protein in the pH 5 extract.

Incorporation of a mixture of [¹⁴C]amino acids by the pH 5 extract. Besides [¹⁴C]glycine, which is rapidly incorporated, L-[¹⁴C]phenylalanine and L-[¹⁴C]glutamate are incorporated at lower rates by the pH 5 extract in the presence of Mg²⁺ ions and ATP (Table 5). The incorporation of a mixture of labelled glycine, phenylalanine and glutamate appears to be additive. This is in keeping with the suggestion that separate enzymes are responsible

for the activation and incorporation of individual amino acids (Chantrenne, 1958).

Properties of the [1-¹⁴C]glycine-pH 5 extract complex. Experiments were performed to elucidate some of the chemical properties of the complex

Table 4. *Aerobic and anaerobic incubation of pH 5 extract*

Complete incubation medium contained: pH 5 extract (0.2 ml.) containing approx. 0.5 mg. of protein; MgCl₂, 16 mM; ATP, 3.3 mM; [¹⁻¹⁴C]glycine, (0.2 μC), 0.25 mM. Total volume was 0.3 ml. Incubations were carried out in Warburg vessels with [¹⁻¹⁴C]glycine in the side arm. Vessels were pre-incubated in air or in N₂ for 10 min.; the [¹⁻¹⁴C]glycine was then tipped in and incubation continued for a further 15 min. at 30°.

	Specific activity	
	Air	N ₂
Complete system	950	670
Complete system less ATP	165	168

Table 5. *Incorporation of radioactive glycine, phenylalanine and glutamate*

Incubation mixture contained: pH 5 extract, 0.2 ml. containing approx. 0.5 mg. of protein; MgCl₂, 10 mM; [¹⁴C]amino acid, as indicated. Total volume was 0.6 ml. and incubation was for 15 min. at 37°.

Labelled amino acid	Specific activity	
Glycine, 0.2 μC (0.16 mM)	451	643
L-Phenylalanine, 0.2 μC (0.16 mM)	107	
L-Glutamate, 0.2 μC (0.08 mM)	85	
Glycine, 0.16 mM, + phenylalanine, 0.16 mM, + glutamate, 0.08 mM	730

Fig. 4. Effect of enzyme concentration on [¹⁴C]glycine incorporation into the pH 5 extract. Each tube contained: MgCl₂, 10 mM; ATP, 2 mM; [¹⁴C]glycine, 0.2 μC, 0.16 mM. The amount of enzyme was varied as indicated and was dissolved in 0.25 ml. of potassium phosphate buffer, pH 7.8, 0.1M. Total volume was 0.5 ml. Tubes were incubated for 10 min. at 25°.

Table 3. *Cofactor requirements for incorporation into the pH 5 extract*

Incubation medium contained: pH 5 extract, 0.2 ml. containing approx. 0.5 mg. of protein; MgCl₂, 10 mM; [¹⁴C]glycine, 0.4 mM, (1 μC); ATP, 4 mM, and GTP, 4 mM, as indicated. Total volume was 0.5 ml. Incubated for 30 min. at 37°.

Additions	Specific activity
Mg ²⁺ , ATP, GTP	1095
ATP, GTP	586
Mg ²⁺ , ATP	948
Mg ²⁺ GTP	880
Mg ²⁺	147

Table 6. *Stability of the [1-¹⁴C]glycine-pH 5 extract complex*

Basic medium for both experiments contained: pH 5 extract, 0.2 ml., containing approx. 0.5 mg. of protein; MgCl₂, 10 mM; [1-¹⁴C]glycine (0.2 μ C), 0.25 mM; ATP, 3.3 mM. Total volume was 0.3 ml. Incubated for 10 min. at 25° in Expt. 1.

Treatment after incubation		Specific activity
Expt. 1.	Usual washing procedure (as described in the Experimental section)	4000
	After washing with cold trichloroacetic acid in the usual way the precipitate was washed twice with hot trichloroacetic acid (90° for 15 min.), twice with ethanol-ether, 3:1 (60° for 3 min.) and once with ether	0
	Mixture heated for 5 min. at 100° at pH 7.8 at end of incubation period before adding cold trichloroacetic acid to precipitate protein	398
Expt. 2.	Usual washing procedure	2999 after 25 min.
	Usual washing procedure	2668 after 10 min.
	Unlabelled glycine (10 m-moles) was added after 10 min. incubation and the mixture was incubated for a further 15 min. Washed by the usual procedure	1000
	Ribonuclease (50 μ g.) was added after 10 min. incubation and then incubated for a further 15 min. Washed by the usual procedure	0

Table 7. *Glycine activation and [1-¹⁴C]glycine incorporation*

The activating enzyme activity was assayed as indicated in the Experimental section. Basal medium for assaying incorporation contained: pH 5 extract, 0.2 ml., containing approx. 0.5 mg. of protein; MgCl₂, 10 mM; [1-¹⁴C]glycine (0.2 μ C), 0.16 mM; ATP, 2 mM; GTP, 0.5 mM. Total volume was 0.5 ml. Incubated for 30 min. at 37°.

Fraction	Activation (μ moles of acetoxyhydroxamic acid/mg. of protein/hr.)	Incorporation (specific activity)
pH 5 extract	0.033	851
Ammonium sulphate (0.47-0.67 saturation)	0.66	3

formed when [1-¹⁴C]glycine was incorporated into the pH 5 extract. The usual procedure at the end of an incubation was to inactivate the enzyme by the addition of trichloroacetic acid and then to wash the precipitate five times with cold 5% trichloroacetic acid. If the cold trichloroacetic acid washing is followed by washings with hot trichloroacetic acid, ethanol-ether mixture and finally with ether, the labelled amino acid is completely eliminated (Table 6, Expt. 1). The data also show that when, after the usual incubation period at pH 7.8, the mixture was heated for 5 min. at 100° and then washed with cold trichloroacetic acid in the usual way, the specific activity of the precipitate decreased by 90% from the original value.

The results given in Table 6, Expt. 2, indicate that incorporation of [1-¹⁴C]glycine was to some extent reversible, since the specific activity fell when unlabelled glycine was added during the

incubation period. Addition of ribonuclease during incubation completely split glycine from the complex.

Other experiments have shown that pre-incubation of the pH 5 extract with 1 μ g. of crystalline ribonuclease completely prevented the subsequent incorporation of added [1-¹⁴C]glycine. Incorporation was also inhibited by chloramphenicol at 2 mM concentration.

The fact that the complex formed between [1-¹⁴C]glycine and the extract is split by heating at pH 7.8, and is removed by treatment with hot trichloroacetic acid or crystalline ribonuclease, suggests that the product is a [1-¹⁴C]glycine-nucleo-protein complex. Such complexes are formed in the rat-liver pH 5.0 extract after the activation of an amino acid and subsequent condensation with the S-ribonucleic acid fraction (Hoagland, Zamecnik & Stephenson, 1957).

Relationship between glycine activation and [1-¹⁴C]glycine incorporation. It is generally believed that activation of an amino acid precedes its incorporation. Glycine is activated to a limited extent by the pH 5 extract from the silk gland, whereas tyrosine and tryptophan are activated to a greater extent (Heller, Szafranski & Sulkowski, 1959). The protein fraction precipitated between 0.47 and 0.67 saturation of ammonium sulphate contained the activating enzyme purified 20 times but the incorporating ability was almost completely removed (Table 7).

DISCUSSION

Silkworm tissues, particularly the posterior silk gland, incorporate [1-¹⁴C]glycine *in vitro* as well as *in vivo*. The rate of incorporation of [1-¹⁴C]glycine into

minces of the posterior silk gland is many times that of such mammalian tissues as rat diaphragm, rabbit bone marrow and rabbit reticulocytes (Tarver, 1954). Further, the rate of incorporation depends on the stage of development, being low at the moulting period and increasing progressively thereafter to parallel the rapid growth of the fifth-instar larvae. The rate decreases just before spinning. Demyanovskii, Vasil'ova & Konikova (1952), working with oak silkworm, *Antheraea pernyi*, observed that incorporation of [^{35}S]methionine into blood proteins depends on the stage of development, and it is possible that the observed incorporation rates in insects are under hormonal control.

Glycine incorporation *in vitro* into minces is affected by the presence of certain metabolic intermediates such as di- and tri-carboxylic acids, α -glycerophosphate and α -oxo acids, the stimulation being two- to three-fold in some cases. α -Glycerophosphate, which has been shown to be an important substrate in the oxidative pathway in the flight muscle of the housefly (Chance & Sactor, 1958), also stimulates incorporation of glycine into proteins of the silk gland. The observed reduction of incorporation by glyoxylate (Table 1) may be due to the formation of unlabelled glycine by transamination (Fukuda & Hayashi, 1958) and subsequent dilution of [^{14}C]glycine.

The results on glycine incorporation in the pH 5 extract indicate that the label is loosely bound and becomes associated with a fraction containing nucleic acid. By ammonium sulphate fractionation of the pH 5 extract it was possible to increase the specific activity of the glycine-activating enzyme many times, but the ability to incorporate [^{14}C]glycine was lost, a result which demonstrates that separate enzymes are responsible for activation and incorporation in the silk gland.

It would appear that besides the ultracentrifuge other cell fractions are necessary for complete synthesis of protein by the silk gland. In injection experiments with [^{14}C]glycine Shimura *et al.* (1958) found that the subcellular fraction which sedimented after 10 min. at 8500 *g* was, initially, the most highly labelled. It is possible that the large granules present in this fraction are required for protein synthesis.

SUMMARY

1. The incorporation of [^{14}C]glycine into tissue proteins of injected silkworm larvae has been studied, and is found to depend on the stage of development of the larvae. The rate is low during the fourth moult and increases progressively during the fifth larval instar. Just before spinning the rate of incorporation is again low.

2. Incorporation *in vitro* into silk-gland minces is an aerobic process depending upon the presence of bivalent ions. It is enhanced by the addition of intermediates of the tricarboxylic acid cycle, particularly by malate, citrate, succinate, fumarate and α -glycerophosphate.

3. A pH 5 extract was prepared by precipitating at pH 5.1 an ultracentrifugate of brei obtained from the posterior portion of the silk gland and subsequently dissolving the precipitate at pH 7.8. Incorporation of [^{14}C]glycine into this extract is rapid, and is stimulated by Mg^{2+} ions and either adenosine or guanosine triphosphate.

4. In addition to [^{14}C]glycine, labelled phenylalanine and glutamate are incorporated into the pH 5 extract but at a slower rate; in mixtures, the three amino acids are incorporated additively.

5. The [^{14}C]glycine-pH 5 extract complex is decomposed by heating at pH 7.8, by treatment with hot trichloroacetic acid, and by the action of ribonuclease.

6. A glycine-activating enzyme has been purified 20-fold by ammonium sulphate precipitation of the pH 5 extract. The purified preparation no longer incorporates [^{14}C]glycine, which indicates that separate enzymes are involved in activation and incorporation.

The technical assistance of Mrs L. Jago is gratefully acknowledged. This paper is contribution no. 3, Insect Pathology Research Institute, Sault Ste. Marie, Ont., Canada.

REFERENCES

- Bergold, G. H. (1941). *Biol. Zbl.* **61**, 158.
- Bheemeswar, B. & Faulkner, P. (1959). *J. Insect Physiol.* **3**, 349.
- Bricteux-Gregoire, S., Verly, W. G. & Florkin, M. (1956). *Nature, Lond.*, **177**, 1237.
- Bricteux-Gregoire, S., Verly, W. G. & Florkin, M. (1958). *Nature, Lond.*, **182**, 1515.
- Chance, B. & Sactor, B. (1958). *Arch. Biochem. Biophys.* **76**, 509.
- Chantrenne, H. (1958). *Annu. Rev. Biochem.* **27**, 35.
- Demyanovskii, Ja. S., Vasil'ova, N. V. & Konikova, A. S. (1952). *Biochemistry, Leningr.*, **17**, 529.
- Denucé, J. M. (1952). *Biochim. biophys. Acta*, **8**, 111.
- Faulkner, P. & Bheemeswar, B. (1959). *Fed. Proc.* **18**, 224.
- Fukuda, T. (1956). *Nature, Lond.*, **177**, 429.
- Fukuda, T. & Hayashi, T. (1958). *J. Biochem., Tokyo*, **45**, 469.
- Heller, J., Szafranski, P. & Sulkowski, W. (1959). *Nature, Lond.*, **183**, 397.
- Hoagland, M. B., Keller, E. B. & Zamecnik, P. C. (1956). *J. biol. Chem.* **218**, 345.
- Hoagland, M. B., Zamecnik, P. C. & Stephenson, M. L. (1957). *Biochim. biophys. Acta*, **24**, 215.
- Ramachandran, L. K. & Fraenkel-Conrat, H. (1958). *Arch. Biochem. Biophys.* **74**, 224.

- Shigematsu, H. (1958). *Nature, Lond.*, **182**, 880.
- Shimura, K., Fukai, H., Suto, S. & Hoshi, R. (1958). *J. Biochem., Tokyo*, **45**, 481.
- Shimura, K., Koide, F., Itabashi, H. & Fukai, H. (1955). *J. Biochem., Tokyo*, **42**, 285.
- Sisakian, A. M. & Kuvaeva, E. B. (1957). *Dokl. Akad. Nauk. S.S.S.R.: Biol. Sci. Sect. (Transl.)*, **113**, 99.
- Takayama, S., Ito, H. & Miura, Y. (1958). *Biochim. biophys. Acta*, **30**, 233.
- Tarver, H. (1954). In *The Proteins*, vol. 2, part B, p. 1199. Ed. by Neurath, H. & Bailey, K. New York: Academic Press Inc.
- Zamecnik, P. C. & Keller, E. B. (1954). *J. biol. Chem.* **209**, 337.
- Zamecnik, P. C., Loftfield, R. B., Stephenson, M. L. & Williams, C. M. (1949). *Science*, **109**, 624.

Biochem. J. (1960) **76**, 78

The Effect of the Incubation Conditions on the Respiration of Mitochondria

BY D. BELLAMY AND W. BARTLEY

Medical Research Council Unit for Research in Cell Metabolism, Department of Biochemistry, University of Oxford

(Received 12 November 1959)

The experiments described arose from the use of the polarographic method to measure the respiration of quantities of mitochondria which were too small to be handled in manometers. In the initial experiments the oxygen uptakes of mitochondria, measured polarographically and manometrically, were different. Such discrepancies have been reported by Chance & Sacktor (1958), who used the two techniques to measure the respiration of insect mitochondria.

In the present work both methods gave the same results with respiring yeast suspensions and with mitochondria after they had been shaken in the medium used for the manometric experiments. More detailed investigation showed that rapid changes in the characteristics of the mitochondrial respiration occurred during the initial period of shaking under the manometric conditions. These changes consisted of a rapid rise and subsequent fall in the oxygen uptake and the transformation of the mitochondria from a state in which the respiration was insensitive to adenosine diphosphate to a state where adenosine diphosphate stimulated the oxygen uptake.

METHODS

Yeast suspensions. Fresh baker's yeast (1 g.) was suspended in 100 ml. of 0.135 M-KCl containing 1% of glucose.

Mitochondria. Rat-liver mitochondria were prepared according to Werkheiser & Bartley (1957), except that the dissolved CO₂ was not removed from the sucrose solution. Rat-brain mitochondria were prepared by the method of Bellamy (1959) and pigeon-breast-muscle mitochondria according to the method of Chappell & Perry (1953). The mitochondria were centrifuged and finally suspended in 0.25 M-sucrose, except in some experiments on rat-liver

mitochondria, which were suspended in the saline medium of Chappell & Perry (1954).

Dry weights. Dry weights were determined as 'acid-insoluble dry matter' according to the method of Werkheiser & Bartley (1957).

Measurement of respiration

(a) **Constant-volume manometry.** The conventional Warburg technique was used with manometer flasks of volume 5.5 or 23 ml. The incubation medium contained 0.01 M-substrate, 0.01 M-K₂HPO₄-KH₂PO₄ buffer (pH 7.4, measured with a glass electrode) and 2 mM-MgCl₂. Adenosine diphosphate (0.01 M) was added where indicated. About 50 or 200 mg. (wet wt.) of mitochondria, according to the size of the manometer cups used, was added, and the solute content adjusted to 0.25 osmolar with sucrose solution. With rat-liver mitochondria suspended in saline the incubations were in 0.1 M-KCl, 0.05 M-2-amino-2-hydroxy-methylpropane-1:3-diol (tris) buffer, pH 7.4, mM-ethylenediaminetetra-acetate and 5 mM-MgSO₄ (Chappell & Perry, 1954). In all experiments the centre well contained 0.2 ml. of 2N-NaOH with filter paper. Incubations were at 20° with air as the gas phase and the oxygen uptake was measured after a 5 min. equilibration period. The initial rates of respiration were expressed as Q_{O₂} values (μl. of O₂ consumed/hr./mg. dry wt.).

(b) **Polarography.** Petering & Daniells (1938) were the first to use a dropping-mercury electrode to measure the respiration of a variety of cell suspensions and rat-liver homogenates, and the method has since been used to measure the respiration of micro-organisms, fibroblasts and mitochondria (Longmuir, 1954; Harris & Barclay, 1955; Chance & Williams, 1955).

When a potential difference of 0.6–0.7 v is applied across a platinum and a reference electrode immersed in an aqueous medium, the current flowing is proportional to the concentration of oxygen in the solution and the area of platinum exposed to the solution. The current is increased when the medium is stirred.