The Reaction of 2:2'-Diaminodiethyl Disulphide (Cystamine) with Sulphite at Low pH

By R. CECIL

Department of Biochemistry, University of Oxford

(Received 7 December 1959)

In the preceding paper (Cecil & Loening, 1960) it was shown that one of the S·S bonds of insulin reacted with sulphite in the range pH 2·5–3·5. It was of interest to know whether this reactivity at low pH could be accounted for by a positive charge in the vicinity of the bond or whether it was necessary to assume that reaction took place with the HSO_3^- ion. Previous work with simple disulphides (Cecil & McPhee, 1955; McPhee, 1956) was done in the range pH 4–11. It was established that disulphides carrying a net positive charge reacted most rapidly and that, under the conditions used, the reaction took place with the SO_3^{2-} ion only.

Accordingly, the reaction of a positively charged disulphide, cystamine, with sulphite was examined in the range pH 3–5.

Cecil & McPhee (1955) considered the reactions:

and showed that

$$k' = k_1[\text{H}^+] + k_2 \frac{K_\phi}{K_\phi + [\text{H}^+]}, \quad (1)$$

where k_1 and k_2 are the respective second-order rate constants and K_ϕ the second dissociation constant of sulphurous acid. k' is the apparent second-order rate constant calculated from the total sulphite concentration. The contribution, if any, of k_1 will be greatest at low pH. If values of k' are measured at different pH values in the acid range it is possible to solve for k_1 and k_2 .

EXPERIMENTAL

The reactions were carried out at 25° in buffered solutions in the range pH 3–5. Samples were withdrawn at measured time intervals and the reaction was stopped by decreasing the pH, with dil. HCl, to 1·8–2·0. The extent of reaction was determined by measuring the amount of thiol formed by potentiometric titration with HgCl_2 (Cecil, 1955).

The sulphite solutions were made up in water free from dissolved gases immediately before use and their concentrations determined by the usual I_2 – $\text{Na}_2\text{S}_2\text{O}_3$ method. It was shown that no appreciable oxidation took place during the time required for the reactions.

Two buffers, formate and phthalate, were used over the

same pH range in view of the sensitivity of the cystamine-sulphite system to different ionic species (McPhee, 1956).

A sample of cystamine dihydrochloride was kindly given by Horlicks Ltd., Slough, Bucks.

The reactions were complex in this pH range in that the apparent rate constants decreased with time. The values quoted for k' were obtained by extrapolation to zero time (Cecil & McPhee, 1955). K_ϕ was taken as $1\cdot6 \times 10^{-7}$ (Cecil & McPhee, 1955).

RESULTS

Reaction in formate buffer

The composition of the reaction mixture was 0·6 mM-cystamine, 18 mM- Na_2SO_3 , 30 mM-formic acid, pH 4·10. In other experiments the same mixture was used with the addition of 9 mM-HCl, giving pH 3·34. The apparent rate constant, k' , was 19 at pH 4·10 and 2·8 at pH 3·34. Substituting in equation (1) it was shown that $k_1 = 0$ and $k_2 = 8600 \text{ l. mole}^{-1} \text{ min.}^{-1}$. In the presence of formate therefore there is no reaction with HSO_3^- ion.

Reaction in phthalate buffer

The composition of the reaction mixture was 0·6 mM-cystamine, 18 mM- Na_2SO_3 and 20 mM-potassium hydrogen phthalate. HCl was added in concentrations between 22 and 10 mM to give the required pH values. The results are given in Table 1. The reaction with SO_3^{2-} ion is apparently inhibited by the phthalate, but there is a slow reaction with HSO_3^- ion.

DISCUSSION

The reaction of cystamine with sulphite is unusually sensitive to the ion species present and to the ionic strength. This is illustrated by the values

Table 1. *Reaction of cystamine with sulphite at 25° in phthalate buffer*

k' is the apparent second-order rate constant extrapolated to zero time. k_1 and k_2 are the second-order rate constants with HSO_3^- and SO_3^{2-} ions respectively. Values of k' , k_1 and k_2 are given in $\text{l. mole}^{-1} \text{ min.}^{-1}$.

pH	k'	k_1	k_2
4·91	34	3	2450
3·64	5·0		
3·26	3·5		

Table 2. *Reaction of cystamine with sulphite at 25° in various buffers*

Values quoted from McPhee (1956). k_2 is the second-order rate constant with SO_3^{2-} ion.

Buffer	I	k_2 (l. mole ⁻¹ min. ⁻¹)
Acetate	0.03	6700
Acetate + sulphate	0.10	1800
Phosphate	0.03	3900
Phosphate	0.05	2400

for k_2 taken from McPhee (1956) (Table 2). In view of these effects the value for k_2 of 8600 l. mole⁻¹ min.⁻¹ obtained in formate buffer is consistent with McPhee's value of 6700 obtained in acetate buffer. It appears that multivalent ions such as phosphate and sulphate have a marked inhibitory effect on the reaction with SO_3^{2-} ion. Phthalate is no exception to this as the value of k_2 , 2400 l. mole⁻¹ min.⁻¹, obtained in this buffer shows. The interesting thing is that, under these conditions, there is for the first time a reaction with HSO_3^- ion since k_1 is 3 l. mole⁻¹ min.⁻¹. It is possible that univalent ions such as formate inhibit the reaction with HSO_3^- ion in the same way that multivalent ions inhibit the reaction with SO_3^{2-} ion.

Although the reaction mechanisms in formate and phthalate buffers are different the apparent rate constants observed are of the same order. If therefore a disulphide reacts with sulphite in the acid range, as does one of the S·S bonds in insulin, it is reasonable to postulate that it has a positively charged environment. Without studying the reaction kinetics, however, it is not possible to say whether or not reaction is taking place with HSO_3^- ion.

SUMMARY

1. The reaction of cystamine with sulphite was studied in the range pH 3–5.
2. In the presence of formate buffer reaction took place with SO_3^{2-} ion but not with HSO_3^- ion.
3. In the presence of phthalate buffer a slow reaction took place with HSO_3^- ion. The reaction with SO_3^{2-} ion was slower than in the presence of formate.

REFERENCES

- Cecil, R. (1955). *Biochim. biophys. Acta*, **18**, 154.
 Cecil, R. & Leoning, U. E. (1960). *Biochem. J.* **76**, 146.
 Cecil, R. & McPhee, J. R. (1955). *Biochem. J.* **60**, 496.
 McPhee, J. R. (1956). *Biochem. J.* **64**, 22.

Biochem. J. (1960) **76**, 157

Synthesis of L-Ascorbic Acid in Rat-Liver Homogenates

CONVERSION OF L-GULONO- AND L-GALACTONO- γ -LACTONE AND THE RESPECTIVE ACIDS INTO L-ASCORBIC ACID

By F. A. ISHERWOOD, L. W. MAPSON AND Y. T. CHEN
Low Temperature Research Station, Cambridge

(Received 21 December 1959)

Evidence has steadily accumulated, supporting the original contention of Isherwood, Chen & Mapson (1954), that L-ascorbic acid is synthesized in animals along the following route:

This paper describes the results of an examination of cell-free extracts of rat liver prepared essentially by the method of Schneider (1948), either in iso-osmotic sucrose (0.25M) or in potassium chloride (0.15M). A detailed examination of the separate fractions of the rat-liver extract was carried out on the material prepared by homogenization in iso-osmotic sucrose. Differential centrifuging of these extracts into nuclear, mitochondrial and submicroscopic particles (microsomes) and soluble fractions showed that the

mitochondrial-microsome fractions were alone responsible for the conversion of L-gulono- and L-galactono- γ -lactone into L-ascorbic acid. The nuclear and soluble fractions not only had no

activity themselves but appeared to have no influence on the activity of the mitochondrial-microsome fractions. The microsomes were more active than the mitochondria. The free acids L-gulonic acid and L-galactonic acid were not converted by these washed particulate preparations into L-ascorbic acid. A preliminary account of this work was given at the Lind Bicentenary Symposium (Isherwood, 1953). The free acids, however, were slowly converted into L-ascorbic acid by whole homogenates of rat liver and later