whereas Sacktor (1958) has postulated a direct control of enzyme activity by means of an endogenous chelating agent. In this context the failure of *cyclopropane* to restore thoracic α -glycerophosphate of the DDT-poisoned insects, and, more especially, of the dieldrin-poisoned insects to their resting levels is of interest. *cycloPropane* quickly suppresses visible motor activity and, presumably, the increased respiration associated with motor activity (Merrill *et al.* 1946). Yet the α -glycerophosphate remains at a level normally found in the active or flying insect. If the enzymes of the α -glycerophosphate cycle are under nervous control, e.g. through the agency of a neurosecretion or hormone, it is possible that the well-established unstabilization of nervous tissue by DDT or dieldrin is also associated with an interference in the regulation of carbohydrate metabolism at the α -glycerophosphate level. There is some evidence that particulate α -glycerophosphate-oxidase activity may be under thyroid-hormone control (Lee, Takemori & Lardy, 1959).

SUMMARY

1. At the late prostrate stage there is a significant breakdown of thoracic adenosine triphosphate in DDT-poisoned houseflies. This fall could be reversed by injecting aqueous glucose.

2. There was also a significant breakdown of thoracic adenosine triphosphate in DDT-poisoned insects spared the hypermotor activity by *cyclo-*

propane anaesthesia. *cycloPropane* also failed to prevent the enhanced desiccation associated with DDT poisoning.

3. The fall in adenosine triphosphate and respiration rate of DDT-poisoned houseflies is not due to the exhaustion of endogenous reserves or to the hypermotor activity induced by DDT.

4. In both DDT- and dieldrin-poisoned houseflies there was a fall in the level of thoracic α -glycerophosphate, which could not be reversed by *cyclopropane* anaesthesia.

The authors wish to acknowledge the help of Mrs A. Brooks and Dr W. J. Roulston during some of the experiments.

REFERENCES

- Beard, R. L. (1958). *Ent. exp. appl.* **1**, 260.
 Bücher, T., Klingenberg, M. & Zebe, E. (1958). *Proc. 4th int. Congr. Biochem., Vienna*, **12**, 153.
 Heslop, J. P. & Ray, J. W. (1959). *J. insect. Physiol.* **3**, 395.
 Ingram, R. L. (1955). *Ann. ent. Soc. Amer.* **48**, 481.
 Lee, Y.-P., Takemori, A. E. & Lardy, H. (1959). *J. biol. Chem.* **234**, 3051.
 Merrill, R. S., Savit, J. & Tobias, J. M. (1946). *J. cell. comp. Physiol.* **28**, 465.
 Sacktor, B. (1958). *Proc. 4th int. Congr. Biochem., Vienna*, **12**, 138.
 Winteringham, F. P. W. (1956). *Chem. & Ind.* p. 1182.
 Winteringham, F. P. W. (1960). *Biochem. J.* **75**, 38.
 Winteringham, F. P. W. & Lewis, S. E. (1959). *Annu. Rev. Ent.* **4**, 303.
 Zebe, E., Delbrück, A. & Bücher, T. (1959). *Biochem. Z.* **331**, 254.

Biochem. J. (1960) **76**, 548

Metabolism in the New Zealand Strain of Obese Mice

By K. SUBRAHMANYAM*

Department of Biochemistry, Medical School, University of Otago, New Zealand

(Received 15 September 1959)

A mutant strain of mice discovered in the Jackson Memorial Laboratory by Ingalls, Dickie & Snell (1950) exhibits the 'hereditary obese-hyperglycaemic syndrome' which has been studied by Mayer and his collaborators (see Mayer, 1953). Guggenheim & Mayer (1952) found that the obese mice liberated $^{14}\text{CO}_2$ from $[1-^{14}\text{C}]$ acetate at a slower rate than non-obese mice and Hughes & Tolbert (1958) made a similar observation in respect of $[2-^{14}\text{C}]$ acetate. Other metabolic anomalies that have been reported in obese as compared with

normal mice include increased turnover of hepatic glycogen (Shull & Mayer, 1956) and increased activity of liver phosphorylase (Shull, Ashmore & Mayer, 1956).

The New Zealand strain of obese mice was developed in the Hugh Adam Department of Cancer Research by systematic inbreeding from a pair of mice chosen from a stock colony (Bielschowsky & Bielschowsky, 1953). The New Zealand obese mice exhibit the hyperglycaemic hereditary form of obesity but they differ from the obese strain of the Jackson Memorial Laboratory in several important respects (Bielschowsky & Bielschowsky, 1956).

* Present address: S.C.B. Medical College, Cuttack, Orissa State, India.

The work described in this paper was done with the object of comparing the metabolism of New Zealand obese mice with that of normal mice.

METHODS

Animals. The care of the mice and the composition of the stock diet are described by Bielschowsky & Bielschowsky (1956).

Measurement of respiration

Apparatus. The respirometer was a closed-circuit type of apparatus operated at constant pressure (Fig. 1) in a room which was thermostatically controlled at $24 \pm 1^\circ$ and air-conditioned. The compensator (A) was provided with a stopcock (C) opening to the atmosphere. The compensator was joined to a manometer (B), 40 cm. long (bore 3 mm.), which was half-filled with Brodie's fluid. The volume of air above the water in A was approximately equal to the total volume (400 ml.) of the enclosed air circuit on the other side of the manometer.

The manifold (DEFG) connected the manometer to the air circuit, and, by way of the three-way taps H and K, with the O_2 reservoirs. Tap H communicated with a 50 ml. burette, graduated in 0.1 ml., and tap K with calibrated gas pipettes, one of 150 ml. and the other of 250 ml. capacity, either or both of which could be put in communication with the air circuit. Two T-pieces leading to H and

K were provided for attachment of O_2 cylinders. The burette and the gas pipettes were equipped with levelling bulbs which were filled with saturated (at 24°) NaCl solution.

The animal chamber (M) was a cylindrical glass chamber (volume about 150 ml., diam. 5 cm.) consisting of two parts each with a ground-glass flange. The ground surfaces, lubricated with rubber grease to make an air-tight seal, were held together by means of a brass screw-clamp fitted with rubber washers. A stainless-steel wire gauze disk was placed at each end of the interior to prevent the animal from blocking the flow of air. The gas inlet and outlet tubes were provided with bulbs, loosely packed with glass wool, to disperse the air stream that is otherwise discomforting to the animal. The chamber and the gas-leading tubes were immersed in a glass tank of water (about 5 l.) at 24° .

The expired air was dried by passage through a train composed of the bubblers (P and R) containing sulphuric acid (sp.gr. 1.84) and a U-tube (S) three-quarters filled with lumps of dried $CaCl_2$ (analytical reagent) and one-quarter with anhydrous. The CO_2 was trapped in a detachable unit composed of a bubbler containing 10 ml. of carbonate-free 3N-NaOH solution and a guard tube three-quarters filled with soda asbestos and one-quarter with anhydrous. The soda asbestos and the anhydrous (both 14–22 mesh) were microanalytical reagents. Four of these units were connected in parallel but only one was used at a time. The CO_2 -absorbing units were followed by a bubbler (T) containing saturated $Ba(OH)_2$ solution to act as a visual check on the absorption of CO_2 .

Fig. 1. Closed-circuit apparatus for measurement of respiratory exchange in mice.
The lettering is that used in the text.

The air-circulating pump consisted of the bubbler valves (V_1 , V_2) filled with saturated (at 24°) NaCl solution, and a 25 ml. graduated glass syringe, the plunger of which was attached by a crank shaft to an eccentric pulley and driven by a geared 0.05 h.p. motor. During an experiment the ground surface of the plunger was kept well lubricated with machine oil. The pump operated at about 60 strokes/min., circulating the air at approximately 1 l./min.

When the apparatus was tested for leaks, under a positive or a negative pressure equal to 7 mm. Hg, with the pump working, the change in volume did not exceed 0.6 ml. in 6 hr. Since no experiment lasted more than 100 min. and the O_2 uptake was not less than 60 ml., the error due to leakage was negligible.

Procedure. The 150 ml. and 250 ml. pipettes were filled to the lower calibration marks with O_2 at atmospheric pressure and the tap K was closed. The burette was filled similarly and tap H closed. In these operations the T-pieces were used to flush the connexions with gas.

The mouse was placed in the animal chamber and the two parts were sealed together. A small mat of filter paper was placed beneath the animal to absorb any urine passed. The chamber was immersed in the tank with the gas inlet joined to the circuit but the gas outlet unattached. The tank was covered with a black cloth and open ventilation started. The mouse was usually active for about 30 min. After it had been at rest continuously for about 10 min. the circuit was closed, the O_2 burette opened and the mouse allowed to become accustomed to the chamber for a further 10 min. The compensator was closed and the barometer read.

The pump was stopped, a tared CO_2 -absorbing unit was substituted for the unit used in the preliminary operations, and the O_2 burette read after the manometric levels had been equalized. The position of the head of the plunger in the syringe was read and the pump started. These operations did not take more than 0.5 min. and were always carried out in the same order. O_2 was slowly admitted to the system from the appropriate O_2 pipette (depending on the size of the animal) by manipulating tap K in such a way as to maintain the levels approximately equal on both sides of the manometer. Thus the rate of admission of O_2 was approximately equal to the rate of consumption. When the levelling fluid reached the upper calibration mark, the screw clip on the rubber connexion to the levelling bulb was closed and for a few minutes O_2 was added to the circuit from the burette. The pump was stopped and the plunger adjusted to the initial reading. The manometric levels were equalized by adjusting the levelling-bulb attached to the burette, and the final burette reading was taken. The O_2 consumption was the sum of the calibrated volume of the O_2 pipette and the volume contraction in the burette. The volume of O_2 was corrected to dry gas at s.t.p. Production of CO_2 was determined as the increase in weight of the whole absorbing unit (bubbler and guard tube). Rates of CO_2 formation were determined by using several of the readily interchangeable units.

The respiration of the normal mouse was measured for 60–100 min. and that of the obese mouse for 30–60 min. Duplicate measurements in the resting mouse did not deviate by more than $\pm 1\%$. Water did not condense on the inner walls of the chamber at any time during the experiments. Dissipation of body heat appears to have been satisfactory since there was no detectable change in the internal temperature of the chamber.

Studies with ^{14}C -labelled compounds. In experiments involving the incorporation of ^{14}C into expired CO_2 , the mouse was injected intraperitoneally with 0.2 ± 0.002 ml. of a solution of the ^{14}C -labelled compound (0.1 m-mole containing approximately $1.0 \mu C$) immediately before it was placed in the animal chamber. In these experiments the preliminary open ventilation period was omitted and the collection of expired CO_2 began within 2 min. The compounds used ($[1-^{14}C]$ glucose, [generally labelled (G)- ^{14}C]glucose, $[1-^{14}C]$ - and $[2-^{14}C]$ -sodium acetate) were obtained from The Radiochemical Centre, Amersham, Bucks, England. The solutions of $[^{14}C]$ acetate were sterilized by steaming in sealed tubes and those of $[^{14}C]$ glucose by filtration.

Analytical methods

Determination of glycogen and fatty acids in the carcass and liver. After removal of the liver the weighed carcass was quickly chopped into pieces about the size of a pea and the whole transferred to 60% (w/v) KOH solution (2 ml./g. of carcass) which was heated under reflux for 5 hr. The digest was diluted with 2 vol. of water and decanted from the inorganic residue of the skeleton. Glycogen was precipitated by addition of 1.5 vol. of ethanol, allowed to stand for 4 hr., separated by centrifuging and dissolved in 50 ml. of water. A small portion of the solution was reserved for estimation of total glycogen. The remainder was treated with 2 vol. of ethanol to reprecipitate the glycogen, which was separated by centrifuging and redissolved in 10 ml. of 10% (w/v) trichloroacetic acid. The solution was decolorized with activated charcoal and filtered. The glycogen in the clear filtrate was precipitated by addition of 4 vol. of ethanol, separated by centrifuging and dissolved in 15 ml. of water. Any insoluble material was removed by centrifuging. Precipitation with ethanol was repeated, the product washed successively with ethanol and acetone and dried *in vacuo* over P_2O_5 . It was free from N_2 and gave about 95% of reducing substance (calculated as glucose) on acid hydrolysis.

The supernatant liquid obtained from the first precipitation of glycogen contained the potassium salts of fatty acids and the unsaponifiable lipids. It was brought to pH 4 with $10N-H_2SO_4$, extracted repeatedly with light petroleum (b.p. $40-60^\circ$) and the extract washed with water. The extract was treated with anhydrous Na_2SO_4 , the light petroleum removed and the residue dried to constant weight *in vacuo* over P_2O_5 to give the weight of the fatty acids and unsaponifiable lipids.

With liver the digestion was performed in 30% (w/v) KOH solution (3 ml./g. of liver) heated under reflux for 3 hr. The same procedure was adopted in purifying the glycogen except that the use of decolorizing charcoal was unnecessary. The fatty acids were isolated and determined in the same way as the carcass fatty acids.

Glycogen. This was estimated by the anthrone method of Seifter, Dayton, Novic & Muntwyler (1950).

Body nitrogen. This was determined by a macro-Kjeldahl incineration of mice approximately of the same age and weight as those used for the experiments involving radioactive compounds.

Combustions. Combustions of radioactive glycogen (15 mg.) and fatty acids (about 10 mg.) were carried out with the reagents of Van Slyke & Folch (1940) in the apparatus described by Sakami (1955) and the $^{14}CO_2$ was trapped in carbonate-free $3N-NaOH$ solution.

Table 1. *Respiration of normal (NZC) and obese mice (NZO) after 3, 6 and 16 hr. of starving*

NZC mice: eight males, six females; mean weight ($\pm$ S.D.), 30.8 ± 5.0 g.; range 23.4–39.1 g. NZO mice: seven males, eight females; mean weight ($\pm$ S.D.), 50.0 ± 10.3 g.; range 35.7–66.0 g. The age of animals was 5.0–8.5 months. Temp. of respirometer was 24° . Values given are the means $\pm$ S.E.M. (range in parentheses) for single observations on 14 NZC and 15 NZO mice.

Observations on 14 NZC and 15 NZO mice.					Significance of difference between means (<i>t</i> test) <i>P</i>
Basis of comparison	Period of starvation (hr.)	NZC	NZO		
		Oxygen consumption (m-moles/g./hr.)			
Weight	3	0.151 ± 0.0022 (0.130–0.167)	0.162 ± 0.0030 (0.144–0.185)		<0.02 ≈ 0.01
	6	0.162 ± 0.0021 (0.149–0.172)	0.172 ± 0.0030 (0.160–0.194)		<0.02 > 0.01
	16	0.143 ± 0.0023 (0.124–0.155)	0.172 ± 0.0021 (0.159–0.188)		<0.001
		Oxygen consumption (m-moles/g. ^{0.7} /hr.)			
(Weight) ^{0.7}	3	0.420 ± 0.011 (0.349–0.485)	0.522 ± 0.009 (0.476–0.604)		<0.001
	6	0.445 ± 0.011 (0.338–0.490)	0.554 ± 0.014 (0.466–0.641)		<0.001
	16	0.399 ± 0.010 (0.333–0.454)	0.557 ± 0.009 (0.507–0.633)		<0.001
		Respiratory quotient			
	3	0.73 ± 0.006 (0.70–0.77)	0.73 ± 0.005 (0.71–0.78)		—
	6	0.72 ± 0.006 (0.69–0.78)	0.72 ± 0.004 (0.70–0.75)		—
	16	0.73 ± 0.006 (0.69–0.78)	0.72 ± 0.006 (0.69–0.77)		—

Measurement of radioactivity. All samples of $^{14}\text{CO}_2$ were plated in triplicate on filter paper as $\text{Ba}^{14}\text{CO}_3$ and counted under a thin end-window Geiger-Müller counter. At least 5000 counts were recorded for each plate. The counts were corrected for background and self-absorption (to infinite thinness). The specific activities of the plates in the triplicate set differed from the mean by less than $\pm 5\%$.

RESULTS

Respiration of normal and obese mice

The oxygen consumption and carbon dioxide production of fourteen normal strain (NZC) and fifteen mice of New Zealand obese strain (NZO) were determined after withdrawal of food, but not water, for 3, 6 and 16 hr. Each mouse was returned to the stock diet for 3–7 days between the determinations. Table 1 shows a comparison of respiratory rates on the basis of body weight and body surface [(weight)^{0.7}]. This has been done because the problems of correlating respiratory activity with body size are not fully resolved (Brody, 1945). These problems include the assessment of fat-free body weight and its use as a basis for comparison. In this connexion it is to be noted that the total lipid content of the NZO mouse is about 38 % of the body weight (Bielschowsky & Bielschowsky, 1956) as against 13 % for the normal mouse; and the nitrogen content of the NZO mouse is about 1.7 % of the body weight as against 2.3 % for the NZC mouse. The nitrogen figures represent a small number of determinations. NZC mice, weighing 35.6 and 32.8 g., contained 22.8 and 24.1 mg. of nitrogen/g. respectively; the corresponding figures for NZO mice, weighing 60.4 and 55.8 g., were 17.1 and 17.9 mg. of nitrogen/g. In view of these find-

Table 2. *Oxygen consumption of mice from the Jackson Memorial Laboratory*

The oxygen consumption of mice starved for 16 hr. has been re-calculated from the data of Mayer, Russell, Bates & Dickie (1952). Compare Table 1. Values are given as means $\pm$ S.D.

	Normal mice	Obese mice
Oxygen consumption		
(m-moles/g./hr.)	0.165 $\pm$ 0.032	0.074 $\pm$ 0.003
(m-moles/g. ^{0.7} /hr.)	0.469 $\pm$ 0.083	0.263 $\pm$ 0.008
Weight (g.)	28.2 $\pm$ 1.9	53.7 $\pm$ 9.1

ings and of the known activity of adipose tissue (Wertheimer & Shapiro, 1948) no attempt has been made to relate respiration to fat-free body weight. Irrespective of the basis of comparison used, the mean oxygen consumption of NZO mice is significantly greater than that of NZC mice after all periods of fasting (Table 1).

Table 2 gives the mean oxygen consumption of normal and obese mice from the Jackson Memorial Laboratory for comparison with that of NZC and NZO mice (Table 1, fasted 16 hr.). The strains of normal mice from both Laboratories consume oxygen at approximately the same rate, but the difference between the two strains of obese mice is striking. Whereas the obese mice of the Jackson Memorial Laboratory use oxygen at a slower rate than their normal controls (56 % less, computed on the basis of body weight), the NZO mice consume oxygen at a faster rate than NZC mice (20 % greater on the basis of body weight).

The R.Q. values for NZO and NZC mice are not significantly different and they are not affected significantly by the period of starvation; but the

mean rate of oxygen consumption of NZC mice is significantly lower ($P < 0.001$) after 16 hr. than after 6 hr. of starvation, whereas that of NZO mice is unchanged (Table 1). In both strains of mice the mean oxygen consumption/g. is somewhat greater after 6 hr. than after 3 hr. of starvation (NZC strain, $P < 0.001$; NZO strain, $P < 0.05 > 0.02$). This trend, which was consistent in all the mice examined, suggests that some adjustment of metabolism may occur between the third and sixth hours of starving.

None of the obese mice used in this work showed glycosuria or ketonuria. The absence of these conditions is characteristic of NZO mice in the age group used.

Oxidation of [^{14}C]acetate and [^{14}C]glucose

The oxygen consumption and carbon dioxide output of NZO and NZC mice, starved for 16 hr., were measured immediately after intraperitoneal injection of 0.1 m-mole of [^{14}C]acetate or [^{14}C]glucose. The carbon dioxide absorber was renewed every

0.5 hr. during the first 2 hr. and later at hourly intervals. The collection of carbon dioxide began a few minutes after the injection, the period of adjustment of the animal to its surroundings being omitted (see Methods section). Results of typical experiments are shown in Table 3, and the numbers of animals used in the whole series of experiments are given in Table 4.

The specific activity of $^{14}\text{CO}_2$ formed by both strains of mice after injection of [$1\text{-}^{14}\text{C}$]- or [$2\text{-}^{14}\text{C}$]-acetate is greatest in the first sample (0–0.5 hr.) and diminishes rapidly in the course of 3 hr. On the other hand, the maximum specific activity is obtained in the second sample (0.5–1.0 hr.) after injection of [$1\text{-}^{14}\text{C}$]- or [$\text{G-}^{14}\text{C}$]-glucose. The same effect is seen in fed mice after injection of [$\text{G-}^{14}\text{C}$]-glucose. The rates of change in the specific activity of $^{14}\text{CO}_2$ expired by NZC and NZO mice after administration of all these substances conform to patterns which have been observed in other normal mice (Siegel & Hughes, 1956) and rats (Gould, Sinex, Rosenberg, Solomon & Hastings, 1949) and

Table 3. *Specific activity of $^{14}\text{CO}_2$ formed by obese (NZO) and normal (NZC) mice after administration of ^{14}C -labelled compounds*

The ^{14}C -labelled compounds (0.2 ml. of solution containing 0.1 m-mole) were injected intraperitoneally. Temp. of respirometer was 24° . The labelled compounds used (with specific activities in 10^6 counts/min./m-mole) were: Expt. 1, [$1\text{-}^{14}\text{C}$]acetate, 1.06; Expt. 2, [$2\text{-}^{14}\text{C}$]acetate, 0.9; Expt. 3, [$1\text{-}^{14}\text{C}$]glucose, 1.02; Expt. 4, [$\text{G-}^{14}\text{C}$]glucose, 1.73. The mice were starved for 16 hr. in Expts. 1–4 but were fed in Expt. 5.

Weight of mouse (g.)		Time (hr.)	CO_2 produced			
			Amount (m-moles)		Specific activity (counts/min./m-mole)	
NZC	NZO		NZC	NZO	NZC	NZO
33.0	59.5	0–0.5	2.38	3.36	30 300	20 900
		0.5–1.0	2.32	2.94	8 070	7 160
		1.0–1.5	2.34	3.25	1 350	1 880
		1.5–2.0	2.61	3.30	570	730
		2.0–3.0	5.46	6.51	270	430
33.0	63.0	0–0.5	2.27	4.02	20 170	11 000
		0.5–1.0	2.10	3.72	10 840	5 290
		1.0–1.5	1.94	3.91	3 500	1 890
		1.5–2.0	1.95	3.93	1 660	950
		2.0–3.0	3.84	7.68	560	390
30.6	58.2	0–0.5	2.16	3.41	7 010	2 410
		0.5–1.0	2.01	3.40	12 200	4 290
		1.0–1.5	1.99	3.18	7 350	3 060
		1.5–2.0	2.04	3.16	4 610	2 610
		2.0–3.0	4.01	6.28	2 110	1 520
29.0	53.0	0–0.5	2.61	3.91	4 510	3 020
		0.5–1.0	2.16	3.00	9 110	5 750
		1.0–1.5	1.88	2.88	6 970	4 450
		1.5–2.0	1.70	2.54	4 420	2 500
		2.0–3.0	3.53	6.10	2 780	1 350
		3.0–4.0	3.58	5.78	1 790	690
		4.0–5.0	3.91	5.10	1 090	410
31.0	58.0	0–0.5	2.05	3.68	8 290	4 760
		0.5–1.0	1.75	3.31	12 280	5 470
		1.0–1.5	1.95	3.24	7 440	4 170
		1.5–2.0	1.77	2.97	6 030	2 510
		2.0–3.0	3.54	6.04	2 530	1 970

in the normal and obese mice from the Jackson Memorial Laboratory (Guggenheim & Mayer, 1952). Since NZO mice have a substantially larger $^{12}\text{CO}_2$ output than normal mice, they should expire $^{14}\text{CO}_2$ with a specific activity lower than that of $^{14}\text{CO}_2$ expired by NZC mice. This expectation is realized (Table 3), but other factors, e.g. differences in substrate pool size, may affect the degree of dilution of $^{14}\text{CO}_2$.

Table 4 shows the mean cumulative excretion of $^{14}\text{CO}_2$ after the administration of [^{14}C]acetate and [^{14}C]glucose to starved (16 hr.) NZC and NZO mice. In 3 hr. about 90% of the ^{14}C administered as [^{14}C] or [^{14}C]acetate is eliminated as $^{14}\text{CO}_2$ and there is no significant difference between the responses of the normal and the obese mice. Fig. 2 shows that the NZC and NZO mice expire $^{14}\text{CO}_2$ derived from [^{14}C]acetate at rates which are not significantly different. Essentially the same result was obtained with [^{14}C]acetate; but, in accor-

dance with the usual finding in [^{14}C]acetate metabolism (Weinman, Strisower & Chaikoff, 1957), the initial rate of $^{14}\text{CO}_2$ formation was somewhat faster than with [^{14}C]glucose.

Similar experiments with [^{14}C]glucose revealed a different situation (Table 4). With NZC mice the mean recoveries of ^{14}C as $^{14}\text{CO}_2$ formed during a 3 hr. period from [^{14}C]glucose and [^{14}C]glucose were about 68 and 66% respectively. These figures agree closely with those found by Siegel & Hughes (1956) in their normal mice. Under the same conditions the NZO mice gave recoveries which were about 56 and 52% of the ^{14}C administered as [^{14}C] and [^{14}C]glucose respectively. This difference between the normal and obese strains is highly significant (Table 4). The difference in the rate of elimination of $^{14}\text{CO}_2$ formed from [^{14}C]glucose persists throughout a 5 hr. period (Fig. 3) and occurs in fed as well as in starved mice (Table 4). Almost identical curves were obtained after injection

Fig. 2. Cumulative excretion of ^{14}C in expired CO_2 after injection of [^{14}C]acetate (0.1 m-mole containing 90 000 counts/min.). The curves represent the mean percentage of ^{14}C recovered in single experiments with eight NZO mice (s.d. $\pm 3.4\%$) and eight NZC mice (s.d. $\pm 3.6\%$). O, NZO; ●, NZC.

Fig. 3. Cumulative excretion of ^{14}C in expired CO_2 after injection of [^{14}C]glucose (0.1 m-mole containing 103 700 counts/min.). The curves represent the mean percentage of ^{14}C recovered in single experiments with eight NZO mice (s.d. $\pm 4.6\%$) and eight NZC mice (s.d. $\pm 4.4\%$). O, NZO; ●, NZC.

Table 4. Cumulative excretion of $^{14}\text{CO}_2$ by obese (NZO) and normal (NZC) mice after administration of ^{14}C -labelled compounds

The ^{14}C -labelled compounds (0.2 ml. of solution containing 0.1 m-mole) were injected intraperitoneally. The mice were 5-9 months old; there were approximately equal numbers of males and females in each group.

No. of mice of each strain	Condition	Substance injected	Total radioactivity (counts/min.)	Mean percentage ($\pm$ S.E.M.) of ^{14}C recovered as $^{14}\text{CO}_2$ in 3 hr.		Significance of difference of the means (<i>t</i> test) <i>P</i>
				NZC	NZO	
8	Starved for 16 hr.	[^{14}C]Acetate	106 000	91.6 $\pm$ 0.9	94.1 $\pm$ 1.1	> 0.1
8	Starved for 16 hr.	[^{14}C]Acetate	90 000	87.8 $\pm$ 1.2	90.4 $\pm$ 0.8	> 0.1
10	Starved for 16 hr.	[^{14}C]Glucose	102 000	68.1 $\pm$ 1.5	56.4 $\pm$ 1.5	< 0.001
8	Starved for 16 hr.	[^{14}C]Glucose	104 000	65.8 $\pm$ 2.0	51.7 $\pm$ 1.3	< 0.001
5	Fed	[^{14}C]Glucose	104 000	72.0 $\pm$ 2.8	61.9 $\pm$ 1.6	< 0.05 > 0.02

of [$1\text{-}^{14}\text{C}$]glucose, showing that there are no significant differences in the rates of $^{14}\text{CO}_2$ formation from the two isotopic forms of glucose.

Incorporation of ^{14}C into glycogen and fatty acids

In 2 hr. NZC and NZO mice in the fed state oxidize about 62 and 54% respectively of the [$\text{G-}^{14}\text{C}$]glucose carbon atoms administered. Other transformations of [^{14}C]glucose were studied by isolating glycogen and fatty acids from both liver and carcass. The material designated 'fatty acids' was not treated to remove non-saponifiable lipids. The animals were allowed free access to food and were given an intraperitoneal injection of non-isotopic glucose (50 mg. in 0.2 ml. of solution) 45 min. before [$\text{G-}^{14}\text{C}$]glucose was injected by the same route. The animals were killed 2 hr. after administration of the [$\text{G-}^{14}\text{C}$]glucose. The results of experiments on five normal and five obese mice are shown in Tables 5 and 6.

It will be seen that the mean weight of the livers of NZO mice is approximately twice that of NZC mice. The hepatic glycogen is distinctly lower in NZO (mean $1.9 \pm 0.2\%$ of the liver weight) than in

NZC mice (mean $3.6 \pm 0.7\%$ of the liver weight), whereas the fatty acid content of the liver is distinctly greater in the obese strain (mean $7.2 \pm 2.0\%$ as against $4.7 \pm 1.0\%$). The mean weight of the carcass fatty acids in the obese strain is about four times that in the normal strain; and the mean weight of the carcass glycogen, though variable, is about the same in both strains of mice.

Glycogenesis. The mean recovery of ^{14}C as glycogen was about 5.7% of the dose in the controls and about 2.5% in the obese mice. This difference is largely due to the higher specific activity of the hepatic glycogen formed by the normal strain (Table 5). When the percentages of ^{14}C incorporated/100 mg. of glycogen are compared in the two strains, it will be seen that the liver glycogen of the NZC mice contains nearly five times the amount of ^{14}C found in that of the obese strain (Table 6). With the carcass glycogen the difference between the means is barely significant ($P < 0.05 > 0.02$).

Lipogenesis from glucose. Table 5 shows that the incorporation of ^{14}C derived from glucose into the fatty acids of liver and carcass is substantially greater in the obese (about 9.7%) than in the

Table 5. *Incorporation of ^{14}C into the glycogen and fatty acids of liver and carcass in normal and obese mice*

Five NZC mice and five NZO mice, all males aged 5–6 months, were used. Mean weights ($\pm$ S.D.) were: NZC mice, 28.0 ± 3.3 g.; NZO mice, 51.6 ± 2.4 g. Mean weights ($\pm$ S.D.) of livers were: NZC, 0.89 ± 0.11 g.; NZO, 1.99 ± 0.13 g. The animals were given 50 mg. of inert glucose and 0.25 m-mole of [$\text{G-}^{14}\text{C}$]glucose (specific activity, 344 000 counts/min./m-mole) by intraperitoneal injection (0.2 ml. of solution) 2 hr. 45 min. and 2 hr. respectively before they were killed. Total radioactivity injected was 516 000 counts/min. No attempt was made to separate non-saponifiable lipids from the material designated 'fatty acids' (see Methods section).

Strain of mice	Material isolated	Mean wt. ($\pm$ S.D.) (mg.)	Mean specific activity ($\pm$ S.D.) (counts/min./m-mole)	^{14}C incorporated (mean percentage $\pm$ S.D. of the dose)
NZC	Liver glycogen	32.3 ± 8.1	$11\,750 \pm 1650$	2.5 ± 0.7
	Carcass glycogen	80.1 ± 29.6	$6\,770 \pm 3200$	3.2 ± 1.4
	Liver fatty acids	41.7 ± 7.4	610 ± 240	0.28 ± 0.08
	Carcass fatty acids	3680 ± 1070	86 ± 39	3.4 ± 0.7
NZO	Liver glycogen	35.6 ± 4.7	$2\,440 \pm 990$	0.55 ± 0.14
	Carcass glycogen	76.5 ± 21.1	$3\,590 \pm 320$	1.9 ± 0.7
	Liver fatty acids	141.7 ± 34.7	390 ± 160	0.62 ± 0.05
	Carcass fatty acids	$14\,690 \pm 1920$	55 ± 18	9.1 ± 2.2

Table 6. *Distribution of ^{14}C between the fatty acids and glycogen of liver and carcass in normal and obese mice*

Derived from the data of Table 5.

Mean percentage ($\pm$ S.D.) of the dose of ^{14}C incorporated/100 mg. of material isolated

Strain of mice	Liver		Carcass	
	Fatty acids	Glycogen	Fatty acids	Glycogen
NZC	0.71 ± 0.27	7.8 ± 1.4	0.11 ± 0.06	4.3 ± 2.1
NZO	0.47 ± 0.16	1.6 ± 0.6	0.06 ± 0.03	2.3 ± 0.3
Significance of difference of the means (P)	$<0.05 > 0.02$	<0.001	$<0.05 > 0.02$	$<0.05 > 0.02$

normal mice (about 3.7 %). This appears to be due mainly to the difference in body-fat content. Owing to greater dilution, the fatty acid ^{14}C of the carcass has a lower specific activity than that of the liver in both NZC and NZO mice; but comparisons between the mean specific activities of the fatty acid ^{14}C obtained from the liver, or from the carcass, show that the differences between the two strains of mice are barely significant ($P < 0.05 > 0.02$). Similarly, the differences between the means are barely significant when the incorporation of ^{14}C is expressed as a percentage of the dose/100 mg. of fatty acids (Table 6).

DISCUSSION

The NZO mouse presents the picture of a 'hereditary obese-hyperglycaemic syndrome'. Various facts described by Bielschowsky & Bielschowsky (1956) point to defective hormonal regulation of carbohydrate and fat metabolism. For example, adult animals show evidence of hyperinsulinism and are resistant to exogenous insulin; and stilboestrol lowers the elevated blood sugar and prevents excessive deposition of body fat in young animals.

The metabolic changes responsible for obesity in NZO mice are likely to be quantitative rather than qualitative. On the basis of comparison used (Table 1), it has been found that the obese strain has a greater respiratory exchange after starving than a control strain of normal mice (NZC). This observation, which is the opposite of the finding in the hyperglycaemic-obese mice of the Jackson Memorial Laboratory (Mayer, Russell, Bates & Dickie, 1952), may be related to the apparently greater activity of the New Zealand strain, which is also distinguished by its fertility and other characteristics (Bielschowsky & Bielschowsky, 1956).

The suggestion that a depressed rate of acetate oxidation may signify abnormal lipogenesis and be a factor in the causation of the obese syndrome (Guggenheim & Mayer, 1952) is not applicable to NZO mice since these animals oxidize [^{14}C]acetate at approximately the same rate as NZC mice (Fig. 2). On the other hand, NZO mice, either fed or starved, eliminate $^{14}\text{CO}_2$ derived from [^{14}C]glucose at a significantly lower rate (about 12 %) than the normal controls. This suggests the possibility that increased lipogenesis from carbohydrate may contribute to the development of the obese condition.

The experiments on the metabolism of [G- ^{14}C]glucose are consistent with this possibility (Table 5). The incorporation of ^{14}C into the liver glycogen of NZO mice (about 0.6 %) is substantially less than that observed in NZC mice (about 2.5 %). The dif-

ference, which depends on the higher specific activity of the glycogen formed by the liver of the NZC mouse, could be due to a slower rate of glycogenesis in the obese strain.

The total recovery of ^{14}C given as [G- ^{14}C]glucose in the glycogen and fatty acids of the liver and carcass was about 9.5 % in the normal strain and about 12.2 % in the obese strain. The fraction of the dose of ^{14}C that is incorporated into the 'fatty acids' is considerably greater in the obese mice and is roughly proportional to the mass of body fat (Table 5); but when the lipogenic activity of the tissues is expressed as a percentage incorporation of the dose of ^{14}C /100 mg. of fatty acids, the difference between the two strains of mice is merely on the verge of statistical significance (Table 6). If real, this difference might provide a mechanism for the genesis of the obese state and an explanation for a slower rate of glucose oxidation in the NZO mice. An investigation of isolated tissues is desirable.

The comparisons between the NZO and NZC mice were made after injection of the same amount of a ^{14}C -labelled compound into all animals irrespective of body weight. If the rate of equilibration of ^{14}C with body carbon is fast compared with the rate of oxidation of the injected ^{14}C -labelled compound, it would be expected that a lower percentage of ^{14}C would appear in the carbon dioxide expired by the animal of larger body weight. Thus it is possible that difference in body weight alone could explain the fact that NZO mice eliminate $^{14}\text{CO}_2$ derived from [^{14}C]glucose at a slower rate than NZC mice. On the other hand, the big difference in body composition may be sufficient to invalidate this argument. The NZO mouse contains only about 27 % more nitrogen (assumed to represent 'active tissue') than the NZC mouse although its body weight is about 70 % greater. For this reason an appropriate fixed dose of ^{14}C (cf. Hughes & Tolbert, 1958) was chosen in preference to a dose proportional to body weight. Differences in body size and composition alone would be expected to affect the specific activities of glycogen and lipids by dilution of ^{14}C ; and since ^{14}C is eliminated continuously from the body, the specific activities of these body constituents will also be affected by unknown turnover rates. The extent to which the observed specific activities (Table 5) are influenced by difference in body size or by difference in metabolic behaviour is uncertain.

The differences between the metabolism of the New Zealand strain of obese mice and that of the obese mice from the Jackson Memorial Laboratory serve to emphasize the opinion of Bielschowsky & Bielschowsky (1956) that the 'hereditary hyperglycaemic-obese syndrome' in rodents is not a single entity.

SUMMARY

1. A closed-circuit respirometer has been devised for measurement of the respiratory exchange in mice. It is particularly suitable for ^{14}C -tracer experiments.

2. In the post-absorptive state the New Zealand strain of obese mice has a greater oxygen consumption and carbon dioxide production/g. body weight than a control strain of normal mice.

3. New Zealand obese mice oxidize $[1-^{14}\text{C}]$ - or $[2-^{14}\text{C}]$ -acetate at approximately the same rate as the control mice. The rates of oxidation of both $[G-^{14}\text{C}]$ - and $[1-^{14}\text{C}]$ -glucose are about 12 % lower in the obese than in the normal strain.

4. The incorporation of ^{14}C derived from $[G-^{14}\text{C}]$ -glucose into the glycogen and fatty acids of liver and carcass was determined in New Zealand obese and control mice. Incorporation of ^{14}C into the liver glycogen, but not the carcass glycogen, was considerably less in obese than in normal mice.

5. In several respects the metabolism of the New Zealand strain of obese mice differs from that of the obese strain of the Jackson Memorial Laboratory.

The author is greatly indebted to the Government of New Zealand for financial assistance under the Colombo Plan; to Professor N. L. Edson for his interest and encouragement, to Dr R. D. Batt for introducing him to the isotope technique and to Drs M. and F. Bielschowsky for a generous gift of the NZO and NZC mice.

REFERENCES

- Bielschowsky, M. & Bielschowsky, F. (1953). *Proc. Univ. Otago med. Sch.* **31**, 29.
 Bielschowsky, M. & Bielschowsky, F. (1956). *Aust. J. exp. Biol. med. Sci.* **34**, 181.
 Brody, S. (1945). *Bioenergetics and Growth*, p. 352. New York: Reinhold Publishing Corp.
 Gould, R. G., Sinex, F. M., Rosenberg, I. N., Solomon, A. K. & Hastings, A. B. (1949). *J. biol. Chem.* **177**, 295.
 Guggenheim, J. & Mayer, J. (1952). *J. biol. Chem.* **198**, 259.
 Hughes, A. M. & Tolbert, B. M. (1958). *J. biol. Chem.* **231**, 339.
 Ingalls, A. M., Dickie, M. M. & Snell, G. D. (1950). *J. Hered.* **41**, 317.
 Mayer, J. (1953). *Physiol. Rev.* **33**, 472.
 Mayer, J., Russell, R. E., Bates, M. W. & Dickie, M. M. (1952). *Endocrinology*, **50**, 318.
 Sakami, W. (1955). *Handbook of Isotope Tracer Methods*. Ed. by Sakami, W. Cleveland, Ohio: Western Reserve University School of Medicine.
 Seifter, S., Dayton, S., Novic, B. & Muntwyler, E. (1950). *Arch. Biochem.* **25**, 191.
 Shull, K. H., Ashmore, J. & Mayer, J. (1956). *Arch. Biochem. Biophys.* **62**, 210.
 Shull, K. H. & Mayer, J. (1956). *J. biol. Chem.* **218**, 885.
 Siegel, B. V. & Hughes, A. M. (1956). *Amer. J. Physiol.* **184**, 356.
 Van Slyke, D. D. & Folch, J. (1940). *J. biol. Chem.* **136**, 509.
 Weinman, E. O., Strisower, E. H. & Chaikoff, I. L. (1957). *Physiol. Rev.* **37**, 252.
 Wertheimer, E. & Shapiro, B. (1948). *Physiol. Rev.* **28**, 451.

Biochem. J. (1960) **76**, 556

A Protein Disulphide Reductase from Pea Seeds

BY M. D. HATCH AND J. F. TURNER

Plant Physiology Unit, Division of Food Preservation and Transport, C.S.I.R.O., Botany School, University of Sydney, Australia

(Received 28 December 1959)

Sulphydryl groups play an important role in the reactions and properties of many proteins. The activity of many enzymes depends on the presence of sulphydryl groups; in addition to such functions as the binding of substrates or cofactors and participation as acyl acceptors, Swenson & Boyer (1957) and Boyer (1959) have suggested that sulphydryl groups may have a general role in maintaining proteins in a suitable configuration for enzyme activity. Oxidation of sulphydryl groups in proteins may explain oxygen-toxicity effects in whole tissues and extracts (Gerschman, Gilbert, Nye, Dyer & Fenn, 1954; Haugaard, Hess & Itskovitz, 1957; Turner & Mapson, 1958; Turner, Turner,

Shortman & King, 1958; Hatch & Turner, 1959, 1960) and may be responsible in part for damage by ionizing radiations (Barron & Johnson, 1954; Ginzburg, Pandre & Binus, 1957; Hope, 1958; Clubb & Wills, 1959). There is evidence suggesting that the formation of protein-sulphydryl groups may precede cell division (Mazia, 1954; Nickerson & Falcone, 1956*b*; Kawamura & Dan, 1958; Sakai & Dan, 1959).

Nickerson & Falcone (1956*a*) reported that a particulate fraction from yeast reduced disulphide groups in a yeast cell-wall protein in the presence of a liver-coenzyme concentrate. Succinate and ethanol acted as hydrogen donors but the exact