- Leeds, N. S., Fukushima, D. K. & Gallagher, T. F. (1954). *J. Amer. chem. Soc.* **76**, 2943.
- Lemieux, R. U. & Charanduk, R. (1951). *Canad. J. Chem.* **29**, 759.
- Levitz, M., Spitzer, J. R. & Twombly, G. H. (1956). *J. biol. Chem.* **222**, 981.
- Marrian, G. F. (1957). *Report on the Conference to Evaluate Methods for Determining Oestrogens in Body Fluids and Tissues*, p. 24. Shrewsbury, Mass.: Worcester Foundation for Experimental Biology.
- Marrian, G. F., Loke, K. H., Watson, E. J. D. & Panattoni, M. (1957). *Biochem. J.* **66**, 60.
- Marrian, G. F., Watson, E. J. D. & Panattoni, M. (1957). *Biochem. J.* **65**, 12.
- Martin, A. J. P. & Synge, R. L. M. (1941). *Biochem. J.* **35**, 1358.
- Pincus, G. & Pearlman, W. H. (1942). *Endocrinology*, **31**, 507.
- Pope, G. S., McNaughton, M. J. & Jones, H. E. H. (1957). *Biochem. J.* **66**, 206.
- Vogel, A. I. (1951). *A Textbook of Practical Organic Chemistry*. London: Longmans, Green and Co. Ltd.
- Westerfeld, W. W., Thayer, S. A., MacCorquodale, D. W. & Doisy, E. A. (1938). *J. biol. Chem.* **126**, 181.

Biochem. J. (1960) **77**, 406

A Chemical Method for the Quantitative Determination of 2-Methoxy-oestrone, Oestrone, Ring D α -Ketolic Oestrogens, Oestradiol-17 β , 16-*epi*Oestriol and Oestriol in Human Urine

By M. L. GIVNER,* W. S. BAULD† AND KITTY VAGI

Department of Metabolism and McGill University Clinic, Montreal General Hospital, Montreal, Canada

(Received 1 February 1960)

The object of this investigation was to develop a method for the quantitative determination of 2-methoxyoestrone, oestrone, ring D α -ketolic oestrogens, oestradiol-17 β , 16-*epi*oestriol and oestriol in human urine. The development of the method began with a detailed study of the experimental conditions required for quantitative separation of pure ketonic and non-ketonic oestrogens by means of the Girard reaction and the development of partition columns for the separation of the oestrogens in each of the Girard fractions (Givner, 1959; Givner, Bauld & Vagi, 1960).

The technique described in this report involves enzymic hydrolysis of the urine, ether extraction, separation of the ketonic from the non-ketonic oestrogens by the Girard reaction, the separation of the individual oestrogens by column-partition chromatography on Celite and colorimetric estimation. It differs from the methods of Brown (1955) and of Bauld (1956), which measure three urinary oestrogens, and from the method of Stimmel (1958), which separates five oestrogens by counter-current distribution.

Extensive specificity studies on the urinary-oestrogen fractions are also described. In addition, the method is evaluated for accuracy, adaptability and sources of error.

* Fellow of the Salmon Foundation for Medical Research. Present address: Department of Endocrinology, Southwest Foundation for Research and Education, San Antonio, Texas, U.S.A.

† Deceased.

METHODS

Purification of materials. Diethyl ether, benzene, hexane, ethanol, methanol, Kober reagents, Girard reagent P, Celite and asbestos were purified as described by Givner *et al.* (1960). Glusulase, a *Helix pomatia* extract (Endo Laboratories, New York) and bacterial β -glucuronidase (Sigma Chemical Co., St Louis, Mo., U.S.A.) were used as received. Blue tetrazolium reagent (dianisole-bis-4:4'-(3:5-diphenyl)tetrazolium chloride was recrystallized from ethanol (96 %, by wt.) by the addition of diethyl ether. pH papers (ranges pH 2-10 and pH 9-11) were purchased from Fisher Scientific Co., Montreal. Glassware was cleaned as previously described (Bauld, 1956).

Collection of urine. Urine samples were collected for 24 hr. without preservative and stored at 5° if necessary.

Fractionation of urine

The purification procedure for urine is described below; letter references are to the flow sheet of Fig. 1.

Hydrolysis of late-pregnancy urine (A). A 50 ml. (approx.) portion of a 24 hr. urine sample was adjusted to pH 5.2 with 0.1 N-acetic acid or 0.1 N-NaOH and then hydrolysed with Glusulase (500 Fishman units of β -glucuronidase/ml. of urine) for 24 hr. at 39 $\pm$ 0.5°. The units were as defined by the supplier and were based on the assay procedure of Talalay, Fishman & Huggins (1946). The urine was cooled to 5° and two 20 ml. portions of the hydrolysed urine were transferred to separating funnels, each containing 80 ml. of water.

Hydrolysis of male urine (A). A 24 hr. urine sample was diluted to 2.5 l. with water; a 250 ml. portion (approx.) was adjusted to pH 6.5, and hydrolysed with bacterial β -glucuronidase (Sigma Chemical Co.), 20 units/ml. of urine for 48 hr. at 37 $\pm$ 0.5°, and then cooled to 5°. The

Fig. 1. Flow sheet of the separation of 2-methoxyoestrone, oestrone, ring D α -ketolic oestrogens, oestradiol-17 β , 16-*epi*oestrone and oestrone from urine.

units were as defined by the supplier and were essentially Fishman units except for small changes in the assay procedure (Sigma Chemical Co. 1951). Two 100 ml. portions of the hydrolysed urine were transferred to separating funnels and extracted. For certain male urines, a larger portion of hydrolysed urine was necessary for analysis and the quantities of reagents used in the extraction procedure were adjusted accordingly.

Extraction (B). The 100 ml. portion of dilute urine was extracted with ether (4×30 ml.). The ether extract was washed with 12 ml. of m-NaHCO_3 , washed and rinsed with water (3 ml. for each procedure) to neutrality (pH paper, range pH 2–10), dried over anhydrous Na_2SO_4 and distilled to dryness on a boiling-water bath. The term 'wash' is used here to describe the procedure that involved inversions of the separating funnel for the equilibration of the aqueous and organic phases, whereas a rinsing describes the addition of water for the removal of alkali from the inside walls of the separating funnel. The flask containing the extract was connected to a water-suction pump (1–25 mm. Hg) for 5 min. at room temp. ($21 \pm 3^\circ$) to bring about partial removal of volatile impurities. The residue was transferred to a 100 ml. round-bottomed flask with ethanol (3×3 ml.), which was then removed *in vacuo*.

Girard reaction (C) and (H). The extract was dried over anhydrous CaCl_2 for 4 hr. in a vacuum desiccator and the reaction was carried out at room temperature. A second Girard reaction was carried out at an elevated temperature on the non-ketonic fraction. The experimental conditions for both procedures have been described previously (Givner *et al.* 1960).

Purification of non-ketonic fraction (I) and (J). This fraction was boiled on an electric heater under reflux for 30 min. with 82 ml. of n-NaOH . The solution was cooled, transferred to a separating funnel and brought to pH 9.3–9.5 (pH paper, range pH 9–11) by passage of CO_2 . The solution was extracted with ether (4×40 ml.), the ether extracts were combined (J) and shaken with 2N-NaOH (7.6 ml.). The aqueous phase, without removal from the separating funnel, was partially neutralized with m-NaHCO_3 (30.4 ml.); the ether and aqueous phases were again shaken. If the pH of the lower phase was 10 or less it was discarded; if the pH was higher than 10, more m-NaHCO_3 was added and the phases were again shaken. The ether phase was washed with m-NaHCO_3 (7.6 ml.), water (5 ml.) and distilled to dryness.

Partition chromatography of the non-ketonic fraction (K) and (L). The non-ketonic fraction was transferred to a Celite pad and the chromatogram was prepared as previously described (Givner *et al.* 1960). The oestradiol-17 β , 16-epioestril and oestril fractions were eluted and the solvents were removed *in vacuo*.

Partition chromatography of the ketonic fractions (D) and (E). The ketonic fraction was transferred to the Celite pad and the chromatogram was prepared as previously described (Givner *et al.* 1960). The 2-methoxyoestrone, oestrone and ring D α -ketolic oestrogen fractions were eluted and the solvents were removed *in vacuo*.

Purification of the oestradiol-17 β fraction (M). This fraction was boiled under reflux for 30 min. with n-NaOH (10 ml.), acidified with $10\text{N-H}_2\text{SO}_4$ (1.1 ml.) and extracted with benzene (25 ml.). The extract was washed with $0.5\text{M-Na}_2\text{CO}_3$ (4 ml.) and water (4 ml.). This was followed by a rinsing with water (4 ml.). The benzene extract was

transferred to a test tube containing quinol (25 ± 5 mg.) and the solvent removed by heating in a boiling-water bath in a stream of filtered air.

Purification of the 2-methoxyoestrone fraction (F). This was purified by the procedure described above for the oestradiol-17 β fraction with one difference, namely the substitution of m-NaHCO_3 (4 ml.) for $0.5\text{M-Na}_2\text{CO}_3$ (4 ml.).

Purification of the oestrone fraction (G). The extract was transferred to a separating funnel with ether (4×40 ml.) and the ether extract was shaken with 2N-NaOH (7.6 ml.). The aqueous phase, without removal from the separating funnel, was partially neutralized with m-NaHCO_3 (30.4 ml.) and the phases were shaken again. If the pH of the lower phase was 10 or less, it was discarded; however, if the pH was higher than 10, more m-NaHCO_3 was added and the phases were again shaken. The ether extract was washed with m-NaHCO_3 (7.6 ml.) and water (5 ml.). The ether solution was distilled to dryness.

Colour development on purified extracts. The oestrogen extracts (with the exception of the oestradiol-17 β and 2-methoxyoestrone extracts, which were already in test tubes) were dissolved in ethanol and portions were transferred to test tubes; quinol (25 ± 5 mg.) was added and the solvent evaporated by heating in a boiling-water bath. The Kober reaction was carried out by the procedure described by Givner *et al.* (1960). Standard curves were prepared with six pure oestrogens (2-methoxyoestrone, 0–10 $\mu\text{g.}$; oestrone, 0–10 $\mu\text{g.}$; 16 α -hydroxyoestrone, 0–12 $\mu\text{g.}$; oestradiol-17 β , 0–10 $\mu\text{g.}$; 16-epioestril, 0–9 $\mu\text{g.}$; oestril, 0–10 $\mu\text{g.}$). However, the amount of each oestrogen in each fraction was calculated from its corrected extinction and that of the appropriate standard by simple proportion. The 16 α -hydroxyoestrone standard was used as the reference compound for the ring D α -ketolic oestrogen fraction.

Special procedures for specificity studies

In the identification of the column-chromatography eluates, paper chromatography was used. All systems were maintained at a constant temperature of $32 \pm 1^\circ$ as described by Mitchell & Davies (1954). The lid was sealed with starch paste (Zaffaroni, Burton & Keutman, 1951). The solvent systems were equilibrated overnight in the cabinet and the tank was equilibrated for 5 hr. Chromatography was performed on Whatman no. 1 filter paper washed as described by Axelrod (1954). Samples, standards of pure oestrogens and a solvent blank were applied to parallel strips (2 cm. $\times$ 46 cm.) by means of ethanol (3×0.05 ml.) and the solvent was evaporated by a stream of air. The diameters of the starting spots were 0.5 cm. (approx.). The strips were equilibrated for 1 hr. before the run was started. Detection of spots was by means of Folin & Ciocalteu's (1927) phenol reagent; elution was with ethanol by an apparatus described by Axelrod (1954). The solvent was evaporated *in vacuo*. For paper chromatography, methanol–water (70:30, v/v) was employed as the stationary phase with the following mobile phases: hexane–benzene (90:10, v/v) for 2-methoxyoestrone and oestrone fractions; hexane–benzene (80:20, v/v) for ring D α -ketolic fraction; hexane–benzene (95:5, v/v) for the oestradiol-17 β fraction; hexane–benzene (50:50, v/v) for the 16-epioestril fraction; hexane–benzene (20:80, v/v) for the oestril fraction.

For additional specificity studies, the six urinary-oestrogen fractions were prepared as described previously

and the absorption and fluorescence spectra measured (Bauld, Givner, Engel & Goldzieher, 1960). For ultra-violet-absorption studies, the oestrogen extracts were dissolved in methanol (3 ml.). In addition, the absorption spectra of the oestrogen extracts after the Kober reaction were determined. All the spectra were recorded on a Beckman DK-2 ratio-recording spectrophotometer.

RESULTS

Recovery of oestrogens added at different stages of the fractionation procedure

Table 1 shows that the purification procedures cause little or no significant loss of oestrogen. The recoveries of oestrogens after the Girard reaction and the Celite transfer are not shown in this table but have been described previously (Givner *et al.* 1960).

Specificity studies on each of the six urinary-oestrogen fractions

Late-pregnancy urine (500 ml.) was taken through the procedure as described in the Methods section and the six fractions were submitted to the different tests.

Retention volume. The retention volumes (V_r) of the six oestrogen fractions were in agreement with the values of pure standards (16 α -hydroxyoestrone and 16-oxo-oestradiol-17 β were used as reference standards for the ring D α -ketolic fraction). In some unpublished experiments, pure samples of the six oestrogens were added to male urine and the fractionation procedure was carried out. The Kober reaction was carried out on each of a large number of small eluate fractions from the Celite columns and the results showed no evidence that the characteristic retention volumes of the added oestrogens were affected by contaminants from the urinary extracts. Since the position of a compound in a chromatogram depends on its partition coefficient, the retention volume is of some value in its identification.

Paper chromatography of the six oestrogen fractions. The 2-methoxyoestrone, 16-epioestrinol and oestrinol fractions showed single spots which had the same chromatographic mobilities as the pure reference compounds.

Oestrone fraction. Chromatography revealed three spots, two of which had mobilities identical with reference oestrone and 16-oxo-oestrone. When saponification was applied to the column eluate, the number of spots was reduced to two. Alkali washing (stage G, Methods section) proved superior, for chromatography after this procedure yielded a single spot with the mobility of oestrone and on elution gave a Kober reaction.

Ring D α -ketolic fraction. This fraction separated into a deep-blue spot, with mobility like that of 16 α -hydroxyoestrone and 16-oxo-oestradiol-17 β standards, and a pale-blue spot between the 16-epioestrinol and 16-oxo-oestradiol-17 β standards. These areas were eluted and subjected to further studies (see below).

Oestradiol-17 β fraction. Chromatography revealed three spots. The spot with the mobility of oestradiol-17 β gave a positive Kober reaction, whereas the two contaminants did not. No ketonic (oestrone) material was present in this fraction.

Kober-reaction spectra of the six oestrogen fractions. Kober reactions were carried out on pure compounds and on each of the six fractions and the absorption spectra recorded. The wavelengths of peak absorption of the fractions were in good

Fig. 2. Kober-reaction absorption spectrum of a 2-methoxyoestrone urinary fraction (A) with a modified reagent. B, 2-Methoxyoestrone (10 μ g.).

Table 1. Recovery of oestrogens added at different stages of fractionation procedure

The recovery of oestrogens after the Girard reaction and the 'Celite transfer' have been reported elsewhere (Givner, Bauld & Vagi, 1960).

Oestrogen (10 μ g.) added	No. of expts.	Stage investigated	Recovery (%)	
			Mean	Range
2-Methoxyoestrone	8	F*	96	91-100
Oestradiol-17 β	11	M*	96	89-102
Oestrinol	7	I	95	90-100
Oestrone	5	G	99	98-101

* The use of electric heaters (e.g. Electrothermal Bunsen) for the saponification stages gave higher recoveries of added oestrogen and decreased etching of glassware.

agreement with the appropriate reference standard (e.g. Fig. 2).

Ultraviolet-absorption spectra of the six oestrogen fractions. All fractions except 16-epioestriol showed curves comparable with that of the reference compound (e.g. Fig. 3); 16-epioestriol did not because of the presence of impurities.

Absorption and fluorescence spectra of the six oestrogen fractions in 75 % (v/v) sulphuric acid

Oestrone, 16-epioestriol and oestriol fractions. These fractions and the pure reference steroids showed maximum absorption and fluorescence at similar wavelengths (e.g. Fig. 4).

Ring D α -ketolic fraction. Fig. 5 illustrates that 16-oxo-oestradiol-17 β fluoresces at a slightly shorter wavelength than does 16 α -hydroxy-oestrone. A solution containing 16-oxo-oestradiol-17 β (10 μ g.) and 16 α -hydroxyoestrone (12 μ g.)

Fig. 3. Ultraviolet-absorption spectrum of an oestriol urinary fraction (A). B, Oestriol (50 μ g.).

Fig. 4. Absorption (A) and fluorescence (F) spectra of 10 μ g. of oestriol (S) and oestriol urinary fraction (F) in 75 % (v/v) H_2SO_4 .

showed maximum fluorescence at a wavelength characteristic of 16 α -hydroxyoestrone (Fig. 6). The '16 α -hydroxyoestrone spot' showed maximum fluorescence at the wavelength associated with 16 α -hydroxyoestrone (Fig. 6) or at an intermediate wavelength between the 16 α -hydroxyoestrone and 16-oxo-oestradiol-17 β peaks (Fig. 5). The unidentified 'x spot' showed maximum fluorescence at a wavelength which was slightly longer than the typical fluorescence peak of 16-oxo-oestradiol-17 β (Fig. 5).

If it is assumed that the fluorescence emitted/unit wt. was the same for the '16 α -hydroxyoestrone' and 'x spots', then approximately 10 % of the ring D α -ketolic fraction was made up of 'x'

Fig. 5. Fluorescence spectra of (1) the eluted '16 α -hydroxy-oestrone spot' (from ketolic fraction), (2) 16 α -hydroxy-oestrone (12 μ g.), (3) 16-oxo-oestradiol-17 β (10 μ g.), (4) eluted 'x spot' and (5) eluted paper blank in 75 % (v/v) H_2SO_4 .

Fig. 6. Fluorescence spectra of (1) 16-oxo-oestradiol-17 β (10 μ g.), (2) 16-oxo-oestradiol-17 β (10 μ g.), plus 16 α -hydroxyoestrone (12 μ g.), (3) eluted '16 α -hydroxyoestrone spot' (from ketolic fraction) and (4) 16 α -hydroxyoestrone (12 μ g.) in 75 % (v/v) H_2SO_4 .

compound or compounds. Moreover, the corrected extinction of 'x spot' in the Kober reaction (with Oestriol reagent) was approximately 11 % of the corrected extinction of the '16 α -hydroxyoestrone spot' under similar conditions.

Fig. 7 illustrates that the absorption spectra of 16 α -hydroxyoestrone and 16-oxo-oestradiol-17 β are different in aqueous sulphuric acid and that a solution containing the two oestrogens had an absorption spectrum characteristic of 16 α -hydroxyoestrone. The '16 α -hydroxyoestrone spot' had a wavelength of maximum extinction which was slightly shorter than the peak wavelength of 16 α -hydroxyoestrone.

2-Methoxyoestrone and oestradiol-17 β fractions. These fractions did not show well-defined spectra because of the presence of impurities.

Colour reactions on the six oestrogen fractions

'Blue tetrazolium' reaction. The method of Izzo, Keutman & Burton (1957) proved to be satisfactory when the ethanol was distilled from a 'blue tetrazolium' solution. Pure 16 α -hydroxyoestrone, '16 α -hydroxyoestrone spot' and 'x spot' gave positive reactions and were estimated by readings against

Fig. 7. Absorption spectra of (1) 16-oxo-oestradiol-17 β (20 μ g.), (2) 16 α -hydroxyoestrone (24 μ g.), (3) 16-oxo-oestradiol-17 β (20 μ g.) plus 16 α -hydroxyoestrone (24 μ g.) and (4) eluted '16 α -hydroxyoestrone spot' (from ketolic fraction) in 75 % (v/v) H_2SO_4 .

an appropriate paper-blank solution. Tetrazolium reactions are given by compounds with an α -ketol or dihydroxyacetone side chain (Axelrod, 1955).

Zimmermann reaction. The 2-methoxyoestrone, oestrone and ring D α -ketolic oestrogen fractions gave positive reactions by the procedure of Sobel, Golub, Henry, Jacobs & Basu (1958). This reaction is given by 17-oxo steroids (Klyne, 1957).

David reaction. The oestriol fraction gave a positive David (1934) reaction. The test was inconclusive with the 16-epioestriol fraction because of the presence of impurities. The reaction appears to be specific for oestriol, 16-epioestriol and their 3-methyl ethers; 17-epioestriol, 16-oxo-oestradiol-17 β - and 16 α -hydroxyoestrone do not give the reaction (Diczfalusy, Tillinger & Westman, 1957).

Standard calibration curves prepared with pure 2-methoxyoestrone, 16 α -hydroxyoestrone and 16-epioestriol. Since curves have not been reported for these oestrogens in the Kober reaction, it was considered essential to investigate this before recovery experiments were attempted. The results showed a satisfactory linear relationship between the concentrations and extinctions of 2-methoxyoestrone (0–10 μ g.), 16 α -hydroxyoestrone (0–12 μ g.) and 16-epioestriol (0–9 μ g.) solutions.

Recoveries of oestrogens added to enzymically hydrolysed urine. Known amounts of the six pure oestrogens were added to portions of male urine and the oestrogen content of each fraction was determined after the fractionation procedure. At the same time urine blank values (in duplicate) were determined and used to correct the recovery experiment for endogenous oestrogen. Table 2 shows that the recovery was, on the average, from 65 to 95 % depending on the oestrogen added. The lowest recovery was obtained with 16 α -hydroxyoestrone.

Sources of error

The precautions reported by Bauld (1956) are of equal importance in this method. Heavy metals, dust, traces of cleaning solution, pitted glassware and solvent contamination are particularly destructive. In addition, care is required for the temperature control in the evaporation of methanol from the Celite pad containing the oestrogen extract.

Samples analyses

In preliminary experiments oestriol and the ring D α -ketolic oestrogens were measured in one-twentyfifth of a 24 hr. collection of male urine. In one subject, 28 μ g. of oestriol and 40 μ g. of ring D α -ketolic oestrogens per 24 hr. were excreted. These values were corrected for known experimental losses, which were determined by the recoveries of added oestrogen to a portion of the analysed urine. Recently, by the use of this

Table 2. Recoveries of oestrogens added to normal male urine

No. of urines analysed	No. of determinations	Oestrogen added (μ g.)	Recovery (%)	
			Mean	Range
6	12	2-Methoxyoestrone (9.8)	95	92-97
6	12	Oestrone (5.1)	90	88-94
6	12	16 α -Hydroxyoestrone (7.9)	65	57-71
6	12	Oestradiol-17 β (5.0)	66	62-67
6	12	16- <i>epi</i> Oestriol (4.6)	87	83-92
6	12	Oestriol (5.0)	76	74-81

oestrogen method, Dr R. Hobkirk (personal communication) measured the 24 hr. urinary-oestrogen excretion of a normal pregnant woman (35 weeks) and found 315 μ g. of 2-methoxyoestrone, 1890 μ g. of oestrone and 4020 μ g. of ring D α -ketolic oestrogen. In another pregnant woman (30 weeks), Dr Hobkirk found no 2-methoxyoestrone, 1775 μ g. of oestrone, 3960 μ g. of ring D α -ketolic oestrogen, 578 μ g. of oestradiol-17 β , 578 μ g. of 16-*epi*oestriol and 12 620 μ g. of oestriol in a 24 hr. urine sample.

DISCUSSION

The instability of 16-oxo-oestradiol-17 β to hot-acid hydrolysis in urine (Watson & Marrian, 1955) and the known losses of oestrone, oestradiol-17 β and oestriol during acid hydrolysis (Marrian, 1956) emphasized the need for a hydrolytic technique other than by acid treatment.

Katzman, Straw, Beuhler & Doisy (1954) found evidence that the major portion of oestrone, oestradiol-17 β and oestriol is excreted in conjugation with glucuronic acid and that hydrolysis with bacterial β -glucuronidase resulted in urinary-oestrogen estimations equal to or higher than the values obtained by acid hydrolysis. Gallagher *et al.* (1954) reported that ox-liver β -glucuronidase is an effective procedure to hydrolyse urinary oestrogens. Brown (1957) found that in non-pregnancy urines, enzymic hydrolysis with a preparation of β -glucuronidase and phenol sulphatase from the limpet *Patella vulgata* resulted in higher values for urinary oestrone, oestradiol-17 β and oestriol than those obtained with acid hydrolysis. Engel (1957) administered radioactive oestradiol-17 β to human subjects and found that, for the most part, the urinary radioactive fraction was hydrolysed with 2 units of bacterial β -glucuronidase/ml. of urine on incubation for 2-48 hr. Recently, Dr R. Hobkirk (personal communication) found that Glusulase was more effective than

bacterial β -glucuronidase in the hydrolysis of oestrogen metabolites in late-pregnancy urine. Thus in our method for urinary-oestrogen analysis, Glusulase was used for the hydrolysis of oestrogen metabolites in pregnancy urine and bacterial β -glucuronidase for male urine.

An important step in this chemical method to measure six urinary-oestrogen fractions included the use of the Girard reaction for the separation and purification of the ketonic oestrogens. Recent experiments on this reaction (Givner *et al.* 1960) indicated that under certain conditions a quantitative and non-destructive separation is obtainable.

For the specificity studies on the complete method, late-pregnancy urine (500 ml.) was used. The object here was to provide a searching test for the purification and separation procedures. The behaviour of the six oestrogen fractions on the partition columns, the paper chromatograms and in the multiple-extraction procedures supported the specificity of the method. The fluorescence and absorption spectra of the fractions in aqueous sulphuric acid, the ultraviolet spectra in methanol, the different colour reactions (Kober, Folin & Ciocalteu, 'blue tetrazolium', Zimmermann, David) and the Girard reaction provided additional evidence in support of the method's specificity. Recently, a 25-tube countercurrent distribution in the system benzene-water of a 16-*epi*oestriol fraction obtained from human-pregnancy urine (by the method described above) exhibited a single peak with the same *K* value as authentic 16-*epi*oestriol (Dr R. Hobkirk, personal communication). In these latter experiments, [14 C]oestriol (2000-5000 counts/min.) added to pregnancy urine and carried through the method did not contaminate the 16-*epi*oestriol fraction to a significant extent; [14 C]oestrone (2000-4000 counts/min.) did not contaminate the oestradiol-17 β fraction nor was it detectable in the first 10 ml. of eluate from the

non-ketonic columns; ^{14}C -labelled 16-oxo-oestradiol-17 β (2000 counts/min.) did not contaminate the oestradiol-17 β fraction. Thus the results indicated the satisfactory separation of urinary 2-methoxyoestrone, oestrone, ring D α -ketolic oestrogens, oestradiol-17 β , 16-*epio*estrinol and oestrinol.

On the basis of the fluorescence and absorption spectra of the eluted ketolic zones from the paper chromatograms, the ring D α -ketolic fraction was composed chiefly of 16 α -hydroxyoestrone and 16-oxo-oestradiol-17 β . An unidentified compound (or compounds) was also detected in this fraction. The properties of the unidentified compound were similar to those of 16 α -hydroxyoestrone. First, the fluorescence spectra in aqueous sulphuric acid were alike. This property may be significant because of the different spectral characteristics of the ring D α -ketolic oestrogens when compared with the other oestrogens (Bauld *et al.* 1960). Secondly, both compounds gave the 'blue tetrazolium' reaction, suggesting that the unknown compound was an α -ketol. Marrian, Watson & Panattoni (1957) have reported that 16 α -hydroxyoestrone has marked reducing power in the 'blue tetrazolium' reaction. Finally both compounds had similar spectra in the Kober reaction. However, the unidentified compound had lower mobility on a paper chromatogram than 16 α -hydroxyoestrone (in this system 16 α -hydroxyoestrone and 16-oxo-oestradiol-17 β showed the same chromatographic mobility). Recently, Layne & Marrian (1958) isolated 16 α -hydroxyoestrone, 16 β -hydroxyoestrone and 16-oxo-oestradiol-17 β from pregnancy urine. Thus the unidentified compound in the ring D α -ketolic fraction might have been 16 β -hydroxyoestrone. It is improbable that the unidentified compound was 18-hydroxyoestrone (Loke, Marrian & Watson, 1959) since the fluorescence spectrum of the latter is unlikely to be similar to that of 16 α -hydroxyoestrone.

It seems likely that the low concentration of 2-methoxyoestrinol (Fishman & Gallagher, 1958) relative to oestrinol did not permit its detection in the non-ketonic fraction during the specificity studies.

The accuracy of the method was estimated by the percentage recovery of six oestrogens added to hydrolysed male urine. The Kober reaction (Bauld, 1954) was used for the final determinations because of its established reliability and specificity. The recoveries were 65–95% and the individual values were dependent on the nature of the added oestrogen. The relatively low recoveries of 16 α -hydroxyoestrone (65%) were to be expected on the basis of known losses in the Girard reaction and in the Celite-transfer technique. The recovery experiments were carried out on low concentrations of

oestrogens in order to simulate the oestrogen content in male urine. Recently, Dr R. Hobkirk (personal communication) obtained the following mean recoveries by this method: oestrone (unlabelled and ^{14}C -labelled compounds added to non-pregnancy and pregnancy urines) $86 \pm 4.6\%$; 16 α -hydroxyoestrone (unlabelled compound added to non-pregnancy urine) and 16-oxo-oestradiol-17 β (^{14}C -labelled compound added to pregnancy urine) $61 \pm 3.9\%$; oestrinol (labelled and unlabelled compounds added to non-pregnancy and pregnancy urines respectively) $80 \pm 4.2\%$; oestradiol-17 β (unlabelled compound added to non-pregnancy urine) $80 \pm 6.6\%$. In no case, up to the present time, has there been any difference in the recovery between small amounts of unlabelled oestrogens added to non-pregnancy urine and ^{14}C -labelled oestrogens added to pregnancy urine.

A larger number of analyses are required before the precision can be evaluated adequately by statistics. The available experimental data indicate very good agreement between the duplicate estimations and therefore satisfactory reproducibility. Recently, Dr R. Hobkirk (personal communication) found that the precision of the newer method (Table 3) compared favourably with Bauld's method (Bauld, 1956). However, the following points should be noted: first, the values in Table 3 are for oestrogen analyses on pregnancy urine and not for recovery experiments; secondly, the procedure was shortened by the omission of the second Girard reaction and the refluxing with alkali procedure for the oestradiol-17 β and 2-methoxyoestrone fractions. The omission of refluxing with alkali procedure is possible when the oestrogen to chromogen ratio is high as in late-pregnancy urine.

The sensitivity of the method has not been determined. An important limiting factor is the sensitivity of the Kober reaction. Under appropriate conditions, pure 2-methoxyoestrone (0.7 $\mu\text{g.}$), oestrone (0.3 $\mu\text{g.}$), 16 α -hydroxyoestrone (0.6 $\mu\text{g.}$), oestradiol-17 β (0.4 $\mu\text{g.}$), 16-*epio*estrinol (0.4 $\mu\text{g.}$) and oestrinol (0.4 $\mu\text{g.}$) can be measured accurately. The experiments showed that six oestrogen fractions can be measured in as little as 20 ml. of late-pregnancy urine and that the ring D α -ketolic oestrogens and oestrinol can be measured in 100 ml. of male urine. Greater sensitivity would be achieved by the use of fluorimetric methods, which remain to be applied to this new method.

The method is satisfactory for the determination of 2-methoxyoestrone, oestrone, ring D α -ketolic oestrogens, oestradiol-17 β , 16-*epio*estrinol and oestrinol in pregnancy urine. This is equally true for male urine with the possible exception of the 2-methoxyoestrone fraction, where the high ratio of non-specific chromogens to oestrogens may create

Table 3. *Standard deviation between duplicates of consecutive analyses of oestrogens ($\mu\text{g./24 hr.}$) from pregnancy urines*

Data from Dr R. Hobkirk (personal communication).

Oestrogen	Method	Urinary oestrogen range (mg./24 hr.)	N*	s*
Oestrone	Bauld (1956)	0.3-1.0	206	0.043
	Bauld (1956)	1.0-3.0	76	0.067
	Givner <i>et al.</i>	0.4-2.4	44	0.074
Oestradiol-17 β	Bauld (1956)	0.1-0.4	224	0.019
	Bauld (1956)	0.4-0.8	38	0.031
	Givner <i>et al.</i>	0.2-1.0	34	0.053
Oestriol	Bauld (1956)	1.0-10.0	157	0.295
	Bauld (1956)	10.0-30.0	149	0.607
	Givner <i>et al.</i>	3.0-23.0	30	0.510
16- <i>epi</i> Oestriol	Givner <i>et al.</i>	0.2-1.0	34	0.045
Ring D α -ketolic	Givner <i>et al.</i>	1.0-10.0	40	0.314
2-Methoxyoestrone	Givner <i>et al.</i>	Insufficient data	—	—

* $s = \sqrt{(ed^2/2N)}$, where s = estimate of standard deviation, d = difference between duplicates and N = number of duplicate analyses (Snedecor, 1952).

difficulties in the application of the Allen colour-correction formula. However, the presumed low concentration of 2-methoxyoestrone in male urine make this difficulty of doubtful importance.

In 8 working days one person can measure the oestrogen content of four different urines (in duplicate) and thereby measure 48 fractions. The method is easily modified for the measurement of only 2-methoxyoestrone, or oestrone or ketolic oestrogens.

SUMMARY

1. A new chemical method which separates and quantitatively measures 2-methoxyoestrone, oestrone, ring D α -ketolic oestrogens, oestradiol-17 β , 16-*epi*oestriol and oestriol in human urine is described.

2. The method involves: enzymic hydrolysis; ether extraction; separation of the ketonic from the non-ketonic oestrogens by the Girard reaction; purification of the non-ketonic fraction by refluxing with alkali and alkali washes; separation and purification of the oestradiol-17 β , 16-*epi*oestriol and oestriol fractions by column-partition chromatography; purification of the oestradiol-17 β fraction by refluxing with alkali; separation and purification of the 2-methoxyoestrone, oestrone and ring D α -ketolic oestrogen fractions by column-partition chromatography; purification of the 2-methoxyoestrone fraction by refluxing with alkali and the oestrone fraction by alkali washes. The purified fractions are measured by the Kober-colour reaction with a spectrophotometric correction for non-oestrogen chromogenic substances.

3. The efficiency of the different procedures is described.

4. The method is evaluated for accuracy, specificity, adaptability and sources of error.

The authors are deeply indebted to Dr J. W. Goldzieher for reading the entire manuscript and for his helpful criticisms and suggestions. Also, we wish to thank Professor G. F. Marrian, F.R.S., Dr R. Hobkirk, Dr L. L. Engel, Dr E. H. Bensley and Professor J. S. L. Browne for their interest and valuable suggestions. Thanks are also due to Dr R. Hobkirk, who sent us advance data based on his experience with the method. Grateful acknowledgement is made to Dr R. M. Greenway and Dr D. Douglas for their contribution in the synthesis of 16 α -hydroxyoestrone. Generous gifts of 16-*epi*oestriol were received from Professor G. F. Marrian, 16-oxo-oestradiol-17 β from Professor Marrian and Dr M. Levitz and of 2-methoxyoestrone from Dr T. F. Gallagher and Dr L. L. Engel. One of us (M. L. G.) wishes to thank the Salmon Foundation for Medical Research for a Fellowship during this study. The investigation was supported by generous grants from the National Research Council of Canada, Maurice Pollack Foundation, Parke, Davis and Co. Ltd., Abbott Laboratories Ltd. and Mr David L. Harrington. This work is taken in part from the thesis of Morris L. Givner submitted to McGill University, Montreal, Canada, in partial fulfilment of the requirements for the degree of Doctor of Philosophy (1959).

REFERENCES

- Axelrod, L. R. (1954). *Recent Progr. Hormone Res.* **9**, 69.
- Axelrod, L. R. (1955). *Analyt. Chem.* **27**, 1308.
- Bauld, W. S. (1954). *Biochem. J.* **56**, 426.
- Bauld, W. S. (1956). *Biochem. J.* **63**, 488.
- Bauld, W. S., Givner, M. L., Engel, L. L. & Goldzieher, J. W. (1960). *Canad. J. Biochem. Physiol.* **38**, 213.
- Brown, J. B. (1955). *Biochem. J.* **60**, 185.
- Brown, J. B. (1957). *Report on the Conference to Evaluate Methods for Determining Oestrogens in Body Fluids and Tissues*, p. 127. Shrewsbury, Mass.: Worcester Foundation for Experimental Biology.

- David, K. (1934). *Acta brev. Neerl. Physiol.* **4**, 64.
- Diczfalusy, E., Tillinger, K. G. & Westman, A. (1957). *Acta endocr., Copenhagen*, **26**, 303.
- Engel, L. L. (1957). *Report on the Conference to Evaluate Methods for Determining Oestrogens in Body Fluids and Tissues* p. 142. Shrewsbury, Mass.: Worcester Foundation for Experimental Biology.
- Fishman, J. & Gallagher, T. F. (1958). *Arch. Biochem. Biophys.* **77**, 511.
- Folin, O. & Ciocalteu, V. (1927). *J. biol. Chem.* **73**, 627.
- Gallagher, T. F., Bradlow, H. L., Fukushima, D. K., Beer, C. T., Kritchevsky, T. H., Stokem, M., Eidinoff, M. L., Hellman, L. & Dobriner, K. (1954). *Recent Progr. Hormone Res.* **9**, 411.
- Givner, M. L. (1959). *Proc. Canad. Fed. Biol. Soc., 2nd Meeting, Toronto*, p. 25.
- Givner, M. L., Bauld, W. S. & Vagi, K. (1960). *Biochem. J.* **77**, 400.
- Izzo, A. J., Keutman, H. & Burton, R. B. (1957). *J. clin. Endocrin. Metab.* **17**, 889.
- Katzman, P. A., Straw, R. F., Buehler, H. J. & Doisy, E. A. (1954). *Recent Progr. Hormone Res.* **9**, 45.
- Klyne, W. (1957). *The Chemistry of the Steroids*. London: Methuen and Co. Ltd.
- Layne, D. S. & Marrian, G. F. (1958). *Biochem. J.* **70**, 244.
- Loke, K. H., Marrian, G. F. & Watson, E. J. D. (1959). *Biochem. J.* **71**, 43.
- Marrian, G. F. (1956). *Rep. Brit. Emp. Cancer Campgn.: Oestrogen Methodology Conference, London*.
- Marrian, G. F., Watson, E. J. D. & Panattoni, M. (1957). *Biochem. J.* **65**, 12.
- Mitchell, F. L. & Davies, R. E. (1954). *Biochem. J.* **56**, 690.
- Sigma Chemical Co. (1951). *Bulletin*, **105**, 1.
- Snedecor, G. W. (1952). *Biometrics*, **8**, 85.
- Sobel, C., Golub, O. J., Henry, R. J., Jacobs, S. L. & Basu, G. K. (1958). *J. clin. Endocrin. Metab.* **18**, 208.
- Stimmel, B. F. (1958). *Proc. 40th Meeting Endocr. Soc., San Francisco*, p. 45.
- Talalay, P., Fishman, W. H. & Huggins, C. (1946). *J. biol. Chem.* **166**, 757.
- Watson, E. J. D. & Marrian, G. F. (1955). *Biochem. J.* **61**, xxiv.
- Zaffaroni, A., Burton, R. B. & Keutman, E. H. (1951). *J. biol. Chem.* **188**, 763.