Apart from these technical difficulties there is also the problem that reaction (1) occurs under the experimental conditions simultaneously with reaction (2), so that the spectrophotometric measurements reflect both reactions (1) and (2). For the purpose of the determination of K_2 it was assumed that reaction (1) did not significantly disturb the equilibrium in reaction (2). The ground for this assumption was the calculation from the data of Alexander & Greenberg (1956) that the concentrations of serine and glycine were not seriously changed due to reaction (1). A mean value of 0.098 was obtained for K_2 , and since K_1 , the equilibrium constant for reaction (1), is 3.2×10^4 M (Blakley, 1960), this gives a value for K_3 (the equilibrium constant for reaction 3) of 3.1×10^3 M ($K_1 \times K_2$). This is in good agreement with the value of 2.76×10^3 reported by Alexander & Greenberg (1956).

SUMMARY

1. Serine transhydroxymethylase has been further purified five- to nine-fold by the use of ion-exchange columns. By this additional step the overall purification of the enzyme from rabbit-liver extracts is about 250-fold. The behaviour of the enzyme on ion-exchange resins indicates that it is a basic protein with isoelectric point above pH 7.7.

2. The spectrophotometric evidence indicates that one isomer of DL-methylenetetrahydropteroylglutamic acid, produced by the action of formaldehyde on DL-tetrahydropteroylglutamic acid, is a substrate for serine transhydroxymethylase.

3. Similar evidence has been obtained which indicates that only one isomer of tetrahydropteroylglutamic acid participates in the serine-transhydroxymethylase reaction. This isomer is formed by dihydropteroylglutamic reductase from dihydropteroylglutamic acid and in the presence of serine is converted by serine transhydroxymethylase into a product with an absorption spectrum identical with that of methylenetetrahydropteroylglutamic acid. Like methylenetetrahydropteroylglutamic acid the product is instantly converted into tetrahydropteroylglutamic acid in the presence of bisulphite.

4. The equilibrium constants have been obtained for the serine-transhydroxymethylase reactions from spectrophotometric data.

I am greatly indebted to Mr J. McKeough for all preparations of serine transhydroxymethylase up to the stage preceding ion-exchange treatment, for enzyme assays and for other technical assistance.

REFERENCES

- Alexander, N. & Greenberg, D. M. (1956). *J. biol. Chem.* **220**, 775.
 Blakley, R. L. (1954). *Biochem. J.* **58**, 448.
 Blakley, R. L. (1957). *Biochem. J.* **65**, 331.
 Blakley, R. L. (1958). *Nature, Lond.*, **182**, 1719.
 Blakley, R. L. (1959). *Biochem. J.* **72**, 707.
 Blakley, R. L. (1960). *Biochem. J.* **74**, 71.
 Futterman, S. (1957). *J. biol. Chem.* **228**, 1031.
 Huennekens, F. M., Osborn, M. J. & Whiteley, H. R. (1958). *Science*, **128**, 120.
 Osborn, M. J. & Huennekens, F. M. (1958). *J. biol. Chem.* **233**, 969.
 Porath, J. (1957). *Ark. Kemi*, **11**, 97.

Biochem. J. (1960) **77**, 465

Purification and Properties of a Protease from the Gut of *Etropolis suratensis*

By SAROJA SUNDARAM AND P. S. SARMA*
University Biochemical Laboratory, Madras 25, India

(Received 26 April 1960)

The proteases of fishes form a group in the field of proteolytic enzymes where relatively little work has been undertaken. Generally, investigations have been carried out hitherto on the distribution of proteolytic enzymes in various fishes, particularly in their digestive tracts (Buchs, 1954; Hamoir, 1955). However, purification of these

enzymes appears to have been done only to a limited extent. The main objective of these investigations has been the possible use of these enzymes as a cheap substitute for pancreatin, which is used industrially as a leather bate.

The crystallization of pepsin from the gut of salmon by Norris & Elam (1939) and the isolation of halibut, shark and tuna pepsins by Eriksen (1943), Sprissler (1942) and Norris & Mathies (1953)

* Present address: Department of Biochemistry, Indian Institute of Science, Bangalore 12, India.

respectively, constitute a few of the isolated cases where a high degree of purification of a fish protease has been achieved. A study of trypsin-like fish proteases acting at alkaline pH therefore appeared logical. A preliminary survey conducted in this laboratory (Saroja Sundaram, 1959) showed that the gut of the freshwater fish *Etroplus suratensis* constituted a convenient source of such an enzyme. This paper embodies the results of a detailed study on the purification and properties of the *Etroplus* enzyme.

EXPERIMENTAL

Materials

Fish gut. Freshly caught fish (*Etroplus suratensis*) were obtained from the Fisheries Department, Government of Madras. After washing them well in running tap water their guts were removed and the food particles in them were gently squeezed out.

Adsorbant. The adsorbant used in this study was tricalcium phosphate gel prepared by the method of Tsuboi & Hudson (1954).

Soya-bean trypsin inhibitor. A solution of soya-bean trypsin inhibitor (Worthington Biochemical Corp., Freehold, N.J., U.S.A.) of the required concentration was prepared in 2.5 mM-HCl.

Buffers. In most of the experiments, the buffer system was chosen from one of the following two series. Buffer series 1 consisted of three buffers (Britton, 1942): pH 2.0-6.0, 0.2N-citric acid-0.2N-sodium hydroxide; pH 6.9-11.82, 0.2N-dipotassium hydrogen phosphate-0.2N-sodium hydroxide; pH 7.46-10.03, 0.2N-boric acid-0.2N-sodium hydroxide. Buffer series 2 also comprised three buffers (Colowick & Kaplan, 1955): pH 2.2-6.0, 0.2M-potassium hydrogen phthalate-0.2M-hydrochloric acid; pH 7.2-8.4, 0.2M-tris-0.2M-hydrochloric acid; pH 8.6-10.6, 0.2M-glycine-0.2M-sodium hydroxide.

Protein substrates

Casein. Hammarsten casein was dissolved in 0.05N-sodium hydroxide to give a solution of the required concentration.

Haemoglobin. Haemoglobin (Eastman Kodak Co.) was dissolved in 0.05N-sodium hydroxide to give a solution of the required concentration.

Egg albumin. A 10% (w/v) solution of egg albumin (Merck) in 0.05N-sodium hydroxide was prepared.

Bovine-serum albumin. A 10% (w/v) solution of bovine-serum albumin (The Armour Laboratories, Chicago, Ill., U.S.A.) was prepared in 0.05N-sodium hydroxide.

Heat-denatured egg albumin and heat-denatured bovine-serum albumin. These were prepared by keeping the respective protein solutions at 100° for 5 min. and by subsequent cooling.

Synthetic substrates

The following synthetic peptides, kindly supplied by Dr E. L. Smith, Salt Lake City, Utah, U.S.A., were used: α -benzoylargininamide-HCl, benzoxycarbonylglycyl-L-phenylalaninamide and benzoxycarbonyl-L-glutamyl-L-

tyrosine. L-Leucylglycylglycine and L-leucylglycine were products of A. F. Hoffman La Roche Co. Ltd., Basle, Switzerland. Solutions of these peptide substrates were prepared in boric acid (0.2N)-sodium hydroxide (0.2N) buffer, pH 9.5 (Britton, 1942).

Methods

Determination of enzyme activity. The incubation system for the estimation of proteolytic activity of the enzyme during the purification procedure consisted of 1.8 ml. of boric acid-sodium hydroxide buffer (Britton, 1942), pH 9.5, 1 ml. of 10% (w/v) casein solution and 0.2 ml. of enzyme solution. The mixture of buffer and substrate solution was first incubated at 37° for 20 min. and then the enzyme extract was added and the reaction was allowed to proceed for 30 min. at 37°. For the blank, water was added instead of the enzyme extract. At the end of incubation, the reaction was stopped by adding 1 ml. of 10% (w/v) trichloroacetic acid and, after standing for 30 min., the mixture was made up to 10 ml. with water and filtered. The extent of proteolysis during the incubation period was determined on suitable portions of the filtrate as follows: to the portion, water was added to bring the volume to 3.0 ml., followed by 4 ml. of 0.5N-NaOH and 1 ml. of the thrice-diluted phenol reagent (Folin & Ciocalteu, 1927). After thorough mixing of the solution and standing for 10 min., the blue colour formed was measured at 650 m μ in a Lumetron Colorimeter (model 400A, Photovolt Corporation, New York City). A standard curve was constructed with pure tyrosine.

The enzyme activity is expressed in units, a unit being defined as the amount of enzyme that liberates (in 1 min.) products not precipitated by trichloroacetic acid which give the same colour with phenol reagent as 1 μ g. of tyrosine.

Determination of protein nitrogen in the enzyme extracts. In the initial stages, the nitrogen content of the enzyme extracts was determined by the micro-Kjeldahl method (Hawk, Oser & Summerson, 1947) and, in the later stages of the enzyme purification, where the concentration of the protein was low the nesslerization method was adopted for the estimation of protein nitrogen (Hawk *et al.* 1947).

RESULTS

Purification of the fish protease. The details of the various stages of the purification procedure and the results obtained in a typical fractionation experiment are summarized in Table 1. Nearly 200-fold purification of the enzyme could be obtained by the method adopted, with a yield of 21%. The supernatant after step 5 was used throughout for the study of the properties of the enzyme.

Optimum pH. In studies on the pH optimum of the enzyme, both buffer series 1 and 2 were employed and the results are presented in Fig. 1. The fish protease showed maximum activity between pH 9.0 and 10.5. The same results are obtained irrespective of whether buffer series 1 or 2 was employed and whether casein or haemoglobin was the substrate.

Table 1. *Purification of the protease of Etroplus gut*

Fresh gut (160 g.) was added to 700 ml. of chilled 2.5% NaCl solution and mixed in a Waring Blender for 2 min. Celite (10 g.) was added and the mixture filtered under suction through a bed of Celite. The clear yellowish red filtrate was the crude extract. All procedures were carried out at 5°, and, except where otherwise stated, all precipitates were centrifuged at 13 500 g for 10 min.

Step no.	Details of purification	Total vol.	Total activity	Protein nitrogen (mg.)	Specificity activity (units/mg. of protein)	Yield (%)
1	Crude extract	700	280 000	1242.5	230	95
2	Supernatant after extract had been dialysed against 1% NaCl for 60 hr. and then centrifuged	705	257 325	629.7	822	88
3	Supernatant was treated with (NH ₄) ₂ SO ₄ to 0.8 saturation and kept 24 hr. The ppt. in 125 ml. of 2.5% NaCl was dialysed against 1% NaCl	127	125 000	21.47	5 821	42
4	Ppt. formed from (3) by 0.3–0.6 saturation with (NH ₄) ₂ SO ₄ was dissolved in 30 ml. of 2.5% NaCl and dialysed	31	60 000	5.5	11 270	21
5	An equal volume of calcium phosphate gel was added to solution (4). After 24 hr. the mixture was centrifuged at 1800 g for 10 min. and the supernatant was examined	62	60 000	1.27	48 540	21

Fig. 1. Influence of hydrogen-ion concentration on the activity of *Etroplus* protease. The incubation systems comprised 2 ml. of the appropriate buffer, 1 ml. of 5% (w/v) substrate solution and 0.1 ml. of the purified enzyme. Incubation was at 37° for 30 min. Substrates: A, casein, buffer series 1; B, casein, buffer series 2; C, haemoglobin, buffer series 2.

Fig. 2. Stability of *Etroplus* protease at different pH values. A portion (0.3 ml.) of the purified enzyme was incubated with 1.2 ml. of the appropriate buffer (series 2) for 60 min. at 37°. Part (0.5 ml.) of the mixture was then incubated with 1 ml. of 5% casein solution and 2 ml. of boric acid-sodium hydroxide buffer, pH 9.5.

The stability of the enzyme preparation at different pH values was studied with buffer series 1 and the result obtained is presented in Fig. 2. As can be seen from the figure, the enzyme was found to be most stable over the range pH 6.0–8.5.

Kinetics of enzymic action

Time course and order of the reaction. The time course of proteolysis with 500, 250, 125 and 62.5 mg. of casein in 15 ml. of the incubation mixture is represented in Fig. 3. With 500, 250 and 125 mg. of the substrate, the reaction appeared to follow almost zero-order kinetics over a considerable period of time (at least for the first hour), whereas when the amount of substrate is 62.5 mg. the reaction tends to obey more the requirements of a first-order reaction. At higher concentrations of the substrate, the reaction conformed to zero-order kinetics for longer periods than at the lower concentration.

Effect of enzyme concentration on reaction rate. The effect of enzyme concentration on the rate of proteolysis was determined by varying the enzyme concentration from 0 to 0.0058 mg. of protein nitrogen/2.5 ml. The enzyme concentration-reaction velocity plot exhibited linearity over the range of concentration studied.

Michaelis constant. The dependence of enzymic activity upon substrate concentration was studied at two enzyme concentrations, and from the Lineweaver & Burk plot (Fig. 4) the Michaelis constant was evaluated, the mean value being 22.5 mg.atoms

Fig. 3. Time course of casein hydrolysis by *Etroplus* protease. The incubation mixture comprised 5 ml. of casein solution of the appropriate concentration, 9 ml. of boric acid-sodium hydroxide buffer, pH 9.5, and 1 ml. of the enzyme (purified enzyme diluted eight times with buffer). Incubation was at 37° and at different intervals 1 ml. portions of the reaction mixture were withdrawn. Substrate concentration (mg. of casein/15 ml. of the incubation mixture): A, 500; B, 250; C, 125; D, 62.5.

of casein nitrogen/l. The Michaelis constants for proteases acting on proteins have not been evaluated with much accuracy, although in recent years this constant has been studied in detail with

Fig. 4. Lineweaver & Burk plot for *Etroplus* protease. The incubation mixture comprised 0.5 ml. of casein solution containing the desired amount of the protease in 2.5 ml. of boric acid-sodium hydroxide buffer, pH 9.5, and 0.2 ml. of enzyme. Incubation was for 15 min. at 37°. A, Purified enzyme diluted four times; B, purified enzyme diluted eight times. [S], Concentration of substrate expressed as mg.atoms of casein nitrogen/l. (mm). V, Initial velocity of reaction expressed as tyrosine (μ moles) liberated/min.

Fig. 5. Influence of temperature on the velocity constant of *Etroplus* protease. The incubation mixture comprised 0.5 ml. of 10% (w/v) casein solution, 2.3 ml. of boric acid-sodium hydroxide buffer, pH 9.5, and 0.2 ml. of enzyme. Incubation was for 15 min. at the desired temperature. A, Purified enzyme diluted four times; B, purified enzyme twice diluted. ○ and ● in curve A represent values obtained from two separate but identical experiments. T, Temperature of incubation. k_0 , Zero-order velocity constant (μ g. of tyrosine liberated/min.).

synthetic substrates (Hofmann & Bergmann, 1941; Kaufman, Neurath & Schwert, 1949; Kaufman, Neurath, Schwert & Snoke, 1948; MacAllister, Harmon & Niemann, 1949).

Influence of temperature on reaction rate and energy of activation. The proteolysis of casein by the purified enzyme was studied at 25°, 30°, 35°, 40°, 45° and 50° and the results are presented in Fig. 5. The experimental conditions in this case were so chosen that the reaction followed zero-order kinetics and the period of incubation was as short as possible to prevent enzyme inactivation. The straight-line nature of the plot of $1/T$ against $\log k_0$ (Fig. 5) satisfies the requirements of the Arrhenius equation.

The close agreement of the several values calculated from the Arrhenius equation for activation energy is also indicative of the validity of the results obtained. The mean value of the activation energy for *Etroplus* protease was found to be 18 149 cal./mole.

Substrate specificity of the enzyme. The proteolytic activity of the enzyme on various proteins (casein, haemoglobin, egg albumin, bovine-serum albumin, heat-denatured egg albumin and heat-denatured bovine-serum albumin) was estimated during a 2 hr. incubation period. The results are presented in Fig. 6. Casein was the protein most susceptible to the protease; among the other substrates, egg albumin and bovine-serum albumin were not attacked even after 2 hr. of incubation. However, heat-denaturation of these proteins before incubation made them subsequently susceptible to the enzyme.

The results presented in Table 2 show that the enzyme attacked α -benzoyl-L-argininamide, which is a specific substrate for trypsin, whereas benzoxycarbonylglycyl-L-phenylalaninamide, which is a typical chymotrypsin substrate, was completely resistant.

Further experiments at pH 9.5 showed that benzoxycarbonyl-L-glutamyl-L-tyrosine and L-leucylglycine, which are substrates for pepsin and peptidase respectively, were not hydrolysed. The

paper-chromatographic method of Giri (Block, Durrum & Zweig, 1958) was used to test for hydrolysis products.

Effect of activators and inhibitors on the activity of the fish protease. The effect of various metal ions and other miscellaneous substances on the proteolytic activity of the fish protease was studied by incubating the test compounds at the required concentration with the enzyme for 15 min. before the addition of the substrate (casein).

Fig. 6. Action of *Etroplus* protease on protein substrates. The incubation mixture comprised 3 ml. of 10% (w/v) protein solution, 5.8 ml. of boric acid-sodium hydroxide buffer, pH 9.5, and 0.2 ml. of purified enzyme. Incubation was at 37°. Portions (1 ml.) were withdrawn at different time intervals. A, Casein; B, haemoglobin; C, heat-denatured bovine-serum albumin; D, heat-denatured egg albumin.

Table 2. Action of *Etroplus* protease on α -benzoyl-L-argininamide and benzoxycarbonylglycyl-L-phenylalaninamide

The incubation system comprised 1.8 ml. of boric acid-NaOH buffer, pH 9.5, containing the substrate and 0.2 ml. of enzyme. Incubation was at 37° for the time indicated, at the end of which 1 ml. of 10% trichloroacetic acid was added and the ammonia estimated by Folin's aeration method modified by Van Slyke & Cullen (1914).

Period of incubation (min.)	Substrate	NH ₃ liberated (as µg.atoms of nitrogen)	Substrate split (%)
30	α-Benzoyl-L-argininamide (8 µmoles)	2.05	25.6
60		3.97	49.6
30	Benzoxycarbonylglycyl-L-phenylalaninamide (10 µmoles)	0.07	Negligible
60		0.01	Negligible

The results obtained for the effect of various metal ions on the activity of protease are presented in Table 3. The enzyme was unaffected by the metal ions tested even at the relatively high concentration employed (0.01 M), except for Mn^{2+} ions, which stimulated the activity, and Fe^{3+} and Zn^{2+} ions, which slightly inhibited the enzyme.

Among the other miscellaneous activators and inhibitors tested, sulphanilamide, *p*-aminobenzoic acid, urea and potassium thiocyanate, at 0.01 M concentration, exerted little or no effect on the enzyme. *O*-Iodosobenzoate and *p*-chloromercuribenzoate at a concentration of mM were virtually without any effect. This, as well as the fact that none of the reducing agents and sulphydryl compounds had any marked stimulating effect and that the oxidizing agent ferricyanide, even at 0.01 M concentration, lowered the activity by only

14%, shows that the protease is not a sulphydryl enzyme. However, glutathione at 0.01 M concentration activated the enzyme by about 30%. Both dipyridyl and ethylenediaminetetra-acetate had no effect on the enzyme. The inference is warranted therefore that no metal activator is essential for the *Etropilus* protease.

The results of studies with the natural soya-bean trypsin inhibitor presented in Table 4 show that at a concentration of 20 μ g. of the inhibitor/3 ml., the *Etropilus* enzyme was inhibited to the extent of 84%.

DISCUSSION

The *Etropilus* protease differs from mammalian trypsin in having a slightly higher pH optimum. However, several instances are known of fish proteases having pH optima around 9 (Labarre, Tremblay & Dussalt, 1951; Akiya, Teshima & Tshikawa, 1949; Utjino & Nishiwaki, 1951), and a proteolytic enzyme exhibiting maximum activity at as high as pH 12.5 has been reported.

The susceptibility of both egg albumin and bovine-serum albumin to the enzyme only after heat-denaturation recalls a parallel observation made in respect of trypsin (Sri Ram & Maurer, 1957) and is indicative of the similarity between the two enzymes. This is further supported by the fact that the *Etropilus* protease attacks only the trypsin substrate. Activation and inhibition studies show that it does not require a metal ion for its activity and is not a sulphydryl enzyme. The inhibition by soya-bean trypsin inhibitor corroborates the conclusion that the enzyme is a protease of the trypsin type.

SUMMARY

1. From the gut of *Etropilus suratensis* a protease acting at alkaline pH has been purified 207-fold with a yield of about 20%.

2. The enzyme exhibits maximum activity at pH 9.5–10.5 and is most stable at pH 6.0–8.5. The time course of the proteolysis of casein by the enzyme and the enzyme concentration–reaction velocity relationship have also been investigated.

3. The Michaelis constant for the protease evaluated from the Lineweaver & Burk plot is 22.5 mg.atoms of casein nitrogen/l.

4. The relationship between the temperature and velocity reaction follows the requirements of the Arrhenius equation over the range 25–50°. The value for the apparent energy of activation calculated from these data is 18 149 cal./mole.

5. Casein is the most susceptible protein substrate; haemoglobin is less susceptible and bovine-serum albumin and egg albumin are attacked only after heat-denaturation.

Table 3. Influence of metal ions on the protease of the gut of *Etropilus suratensis*

The incubation system comprised 2 ml. of boric acid–NaOH buffer, pH 9.5, 0.3 ml. containing the test substance (0.1 M), 0.5 ml. of 10% casein solution (0.5 ml. of water in the blank) and 0.2 ml. of purified enzyme. The volume (0.3 ml.) of the test substance was replaced by 0.3 ml. of water in the control. Incubation was for 60 min. at 37°. The proteolysis was determined as usual.

Metal ion (at 0.01 M concentration)	Percentage of control activity
Hg ²⁺	96
Mg ²⁺	100
Mn ²⁺	140
Co ²⁺	97
Cu ²⁺	107
Fe ³⁺	86
Zn ²⁺	90
Ca ²⁺	97

Table 4. Effect of soya-bean trypsin inhibitor on *Etropilus* protease

The incubation system comprised 2.3 ml. of boric acid–NaOH buffer, pH 9.5, containing the desired amount of trypsin inhibitor, 0.5 ml. of 10% casein solution and 0.2 ml. of purified enzyme (0.2 ml. of water in the blank). Incubation was for 30 min. at 37°. The extent of proteolysis was estimated as usual in the trichloroacetic acid filtrate.

Concn. of inhibitor (μ g./3 ml.)	Percentage inhibition
2	32
4	46
6	71
8	71
10	76
15	79
20	84
30	84

6. The protease attacks only α -benzoyl-L-argininamide among the several peptides tested, which shows that it is of the trypsin type.

7. The enzyme is practically unaffected by any of a variety of compounds tested. It is not a sulphhydryl enzyme and does not require metal ions for activity and is inhibited by soya-bean trypsin inhibitor.

The generosity of the Fisheries Department, Government of Madras, in supplying the fish material used in this investigation is gratefully acknowledged. Also, we are deeply indebted to Professor E. L. Smith, College of Medicine, Salt Lake City, Utah, U.S.A., for the gift samples of the synthetic protease substrates used in this study and to Dr E. Raghupathy for going through the manuscript. The authors wish to thank the University of Madras for the award of a research studentship to one of them (S.S.) and for kind permission to publish the results, which form part of a thesis approved for the degree of Master of Science.

REFERENCES

- Akiya, S., Teshima, S. & Tshikawa, Y. (1949). *Symp. enzyme Chem. Japan*, **3**, 43.
- Block, R. J., Durum, E. L. & Zweig, G. (1958). *A Manual of Paper Chromatography and Paper Electrophoresis*. New York: Academic Press Inc.
- Britton, H. T. S. (1942). *Hydrogen Ions*, vol. 1, p. 300. London: Chapman and Hall Ltd.
- Buchs, S. (1954). *Z. vergl. Physiol.* **36**, 165.
- Colowick, S. P. & Kaplan, N. O. (1955). *Methods in Enzymology*, vol. 1, p. 138. New York: Academic Press Inc.
- Eriksen, N. (1943). Thesis: University of Washington.
- Folin, O. & Ciocalteu, V. (1927). *J. biol. Chem.* **73**, 627.
- Hamoir, G. (1955). *Advanc. Protein Chem.* **10**, 227.
- Hawk, P. B., Oser, B. L. & Summerson, W. H. (1947). *Practical Physiological Chemistry*. New York: Blackiston Co.
- Hofmann, K. & Bergmann, M. (1941). *J. biol. Chem.* **138**, 231, 243.
- Kaufman, S., Neurath, H. & Schwert, G. W. (1949). *J. biol. Chem.* **177**, 793.
- Kaufman, S., Neurath, H., Schwert, G. W. & Snoke, J. E. (1948). *J. biol. Chem.* **172**, 221.
- Labarre, J., Tremblay, J. L. & Dussalt, H. P. (1951). *Rev. Canad. Biol.* **10**, 140.
- MacAllister, R. V., Harmon, K. M. & Niemann, C. (1949). *J. biol. Chem.* **177**, 767.
- Norris, E. R. & Elam, D. W. (1939). *Science*, **90**, 399.
- Norris, E. R. & Mathies, J. C. (1953). *J. biol. Chem.* **204**, 673.
- Saroja Sundaram (1959). *J. sci. industr. Res.* **18C**, 182.
- Sprissler, G. P. (1942). Dissertation: Catholic University of America, Washington, D.C.
- Sri Ram, J. & Maurer, P. H. (1957). *Arch. Biochem. Biophys.* **70**, 185.
- Tsuboi, K. K. & Hudson, P. B. (1954). *Arch. Biochem. Biophys.* **53**, 341.
- Utjino, S. & Nishiwaki, I. (1951). *Acta Sch. med. Univ. Kyoto*, **29**, 1.
- Van Slyke, D. D. & Cullen, G. E. (1914). *J. biol. Chem.* **19**, 211.

Biochem. J. (1960) **77**, 471

A Protein from Leaves Catalysing the Reduction of Metmyoglobin and Triphosphopyridine Nucleotide by Illuminated Chloroplasts

By H. E. DAVENPORT

University of Bristol, Research Station, Long Ashton, Bristol

(Received 5 May 1960)

In his early experiments with isolated chloroplasts Hill (1939) was able to show that when these are illuminated in the presence of an extract of acetone-dried leaf, oxygen is evolved. The extracts therefore contained soluble material in substrate amounts capable of accepting hydrogen in the photochemical system. Although the pathway of hydrogen transport in this reaction has not yet been determined, two leaf-protein fractions have since been shown to be active catalysts of hydrogen transport from illuminated chloroplasts.

One of these, occurring in acetone-dried leaf powders, was first observed to catalyse the reduction of metmyoglobin (Davenport, Hill & Whatley, 1952). The active protein has since been obtained

from pea leaves in a highly purified state, homogeneous both electrophoretically and in the ultracentrifuge (Davenport & Hill, 1960) and was shown also to promote the reduction of other haem-protein compounds including cytochrome *c* and cytochrome *b₅*.

Meanwhile San Pietro & Lang (1958) isolated a spinach-leaf-protein fraction active in catalysing the photochemical reduction of triphosphopyridine nucleotide. This also has since been purified (San Pietro, 1958) and named 'photosynthetic pyridine nucleotide reductase'.

During some experiments aimed at confirming the observations of San Pietro & Lang (1958) it was noticed that the capacity of leaf-protein fractions to