body but rather that both ethyl trichloro- and tri-fluoro-acetate are effective as lipotropic agents because of the CCl_3 or CF_3 group. These groups are larger than the CH_3 group and may exert their effects through a steric blockage of an enzyme system.

Ethyl dichloroacetate was not only without lipotropic action but was highly toxic when given to rats in an established choline-deficient state. Rats fed on a ration which supported the breeding colony tolerated the same dosage with no noticeable impairment of function. It is probable that it is more toxic when the quality of protein in the diet is poor since it was also toxic when choline was added to the experimental diet (Kratzing, 1960). Laug (1946) stated that dichloroacetate was a more effective inhibitor of mouse-liver respiration than corresponding concentrations of trichloroacetate, whereas acetate, at the same concentration range, was not inhibitory. But Woodard, Lange, Nelson & Calvery (1941) found that rats tolerated larger oral doses of dichloroacetic acid than they did of trichloroacetic acid or acetic acid before showing signs of toxicity. In general the non-toxicity of the poly-halogen derivatives of acetic acid is in marked contrast to the toxicity of the monohalogen derivatives (Kratzing, 1960).

SUMMARY

1. Ethyl trifluoroacetate appeared as effective as ethyl trichloroacetate in decreasing excess of

liver-lipid content of choline-deficient rats; whereas ethyl acetate and ethyl dichloroacetate were without effect.

2. Because of failure of the rats to grow, no lipotropic properties could be attributed to ethyl tribromoacetate.

We would like to express our appreciation for the technical assistance of Mrs G. Dabelstein and Miss M. Muir in this investigation. This work was supported by grants from the Commonwealth Research Fund and the N.H. and M.R.C. of Australia.

REFERENCES

- Best, C. H., Hartroft, W. S. & Sellers, E. A. (1952). *Gastroenterology*, **20**, 375.
 Best, C. H., Lucas, C. C. & Ridout, J. H. (1954). *Ann. N.Y. Acad. Sci.* **57**, 646.
 Best, C. H. & Ridout, J. H. (1938). *J. Physiol.* **94**, 47.
 Griffith, W. H. (1940). *J. Nutr.* **19**, 437.
 Hartroft, W. S. (1954). *Ann. N.Y. Acad. Sci.* **57**, 633.
 Kratzing, C. C. (1960). *Biochem. Pharmacol.* **3**, 245.
 Kratzing, C. C. & Windrum, G. M. (1957). *Nature, Lond.*, **180**, 859.
 Kratzing, C. C. & Windrum, G. M. (1959). *Aust. J. exp. Biol. med. Sci.* **37**, 321.
 Laug, E. P. (1946). *Proc. Soc. exp. Biol., N.Y.*, **61**, 178.
 Moore, H. C. (1957). *J. Path. Bact.* **74**, 171.
 Wheelon, L. W. & Collins, F. D. (1957). *Biochem. J.* **66**, 435.
 Windrum, G. M. & Kratzing, C. C. (1960). *Arch. Path.* **70**, 156.
 Woodard, G., Lange, S. W., Nelson, K. W. & Calvery, H. O. (1941). *J. industr. Hyg.* **23**, 78.

Biochem. J. (1961) **78**, 748

Studies in the Biosynthesis of Fungal Metabolites

2. THE BIOSYNTHESIS OF ALTERNARIOL AND ITS RELATION TO OTHER FUNGAL PHENOLS*

By R. THOMAS†

Department of Biochemistry, London School of Hygiene and Tropical Medicine, University of London, Keppel Street, London, W.C. 1

(Received 20 July 1960)

The mode of biosynthesis of certain types of fungal phenols involves the condensation of acetate units (for a review see Birch, 1957). However, the biosynthesis of alternariol, $\text{C}_{14}\text{H}_{10}\text{O}_5$ (I), a metabolite of *Alternaria tenuis* auct. (Raistrick, Stickings & Thomas, 1953), was of particular

interest since, in addition to its rarity among known fungal products as a diphenyl derivative, it appeared to constitute a classical example of the acetate hypothesis. Thus it can in theory be derived solely by the head-to-tail condensation of seven acetate units without requiring additional stages involving oxidation, reduction, decarboxylation, etc. Another example of a fungal phenol, exclusively derived from acetate, is the key substance orsellinic acid, $\text{C}_8\text{H}_8\text{O}_4$ (II), isolated from *Chaeto-*

* Part 1: Chaplen & Thomas (1960).

† Present address: Department of Microbiology, Squibb Institute for Medical Research, New Brunswick, New Jersey, U.S.A.

Fig. 2. Growth curves for *A. tenuis*. Average results from four combined flasks. ■, $10^{-1} \times$ glucose (%); □, $10^{-1} \times$ mycelial weight (g.); ○, other extract (g.) (crude alternariol); ●, petroleum extract (g.).

Table 1 lists the degradation products whose activities were determined in order to calculate the labelling of the individual alternariol carbon atoms.

A Kuhn-Roth oxidation yielded acetic acid containing the methyl carbon and carbon atom 6 of alternariol (I). This was partially converted into its *p*-bromophenacyl ester, and the remainder was further degraded by pyrolysis of the lithium salt to lithium carbonate and acetone, which was then converted into iodoform with alkaline iodine (Cornforth, Hunter & Popják, 1953). The iodoform in turn was oxidized to carbon dioxide and assayed as barium carbonate.

Further methylation of alternariol trimethyl ether (III) under aqueous alkaline conditions produced the ester $C_{19}H_{22}O_6$ (IV). This was demethylated and decarboxylated by refluxing with a mixture of red phosphorus and hydriodic acid, yielding the tetrahydroxymethyldiphenyl (V); the effluent carbon dioxide corresponded to the lactone carbonyl of alternariol.

An unsuccessful attempt to prepare a pentanitro derivative of (V) by the action of a mixture of sulphuric acid and nitric acid gave rise instead to a product $C_{13}H_6O_{14}N_4S$, which was interpreted as resulting from the introduction of four nitro groups and the formation of a δ -sultone as shown in the structure (VI). Cullinane, Morgan & Plummer (1937) obtained the corresponding sultone of 2-hydroxydiphenyl-2'-sulphonic acid on refluxing *o*-phenylphenol with 70% sulphuric acid.

It was considered that the stability of the δ -sultone ring was responsible for the failure of a nitro group to enter position 2'. The product (VI) was then oxidized with barium hypobromite, and the resulting bromopicerin separated by steam distillation, converted by wet combustion into carbon dioxide and assayed as barium carbonate. The absence of any activity was consistent with the derivation of the four nitro C atoms 3, 5, 4' and 6' from the methyl C atom of acetate. However, only 58% of the theoretical yield could be obtained, whereas a parallel oxidation of trinitro-oreinol yielded bromopicerin equivalent to 85% of theory. Consequently it is possible that the bromopicerin was derived from only three of the four nitro C atoms present in the sultone (VI), although in view of the assigned structure it seems improbable that any one group would be completely resistant to attack. Assuming that all four nitro C atoms do in fact contribute to the bromopicerin and are consequently devoid of radioactivity, then it follows that C-2' in alternariol is also inactive, since it is indistinguishable from C-6' in the tetrahydroxymethyldiphenyl (V) from which the sultone was derived.

Alkaline permanganate oxidation of alternariol trimethyl ether (III) yielded a mixture of 3:5-dimethoxyphthalic acid and 4:6-dimethoxyphthalonic acid (VII) (Raistrick *et al.* 1953) containing ring B of alternariol. The latter product was refluxed with hydriodic acid in the presence of red

Fig. 3. Alternariol degradation scheme.

Table 1. Activities of alternariol degradation products

Degradation product	Alternariol derived C atoms	Activity ($\mu\text{C}/\text{mole}$)	
		Found	Theory*
Alternariol trimethyl ether	1-6, 1'-6', CO, Me	460	7×65.7
<i>p</i> -Bromophenacyl acetate	6, Me	59.1	65.7
<i>p</i> -Bromophenacyl acetate	1, 1'	59.4	65.7
BaCO_3 (via IV)	CO	74.7	65.7
BaCO_3 (via VII)	CO	74.7	65.7
BaCO_3	Me	0	0
BaCO_3	2	69.1	65.7
BaCO_3 (ex bromopicrin)	3, 5, 2', 4', 6'	0	0
BaCO_3	1	0	0
3:5-Dimethoxyphthalic anhydride	1, 1'-6', CO	259	263
Orcinol	1, 1'-6'	184	197
2:3':4:5'-Tetramethoxy-6-methyldiphenyl- 2'-carboxylic ester	1-6, 1'-6', CO, Me	454	460
2:3':4:5'-Tetrahydroxy-6-methyldiphenyl	1-6, 1'-6', Me	385	394
4:6-Dimethoxyphthalonic acid	1, 2, 1'-6', CO	328	328
3:5-Dihydroxyphenylacetic acid	1, 2, 1'-6'	247	263

* Based on a theoretical incorporation of seven acetate units.

phosphorus when demethylation, reduction and decarboxylation took place giving rise to 3:5-dihydroxyphenylacetic acid (VIII). The carbon dioxide evolved again corresponded to the lactone carbonyl of alternariol and its activity when assayed as barium carbonate was in good agreement with that previously obtained from the ester (IV). Decarboxylation of the acid (VIII) yielded orcinol and carbon dioxide derived from C-2. Finally, a Kuhn-Roth oxidation of orcinol produced acetic acid which was degraded as before to determine the labelling of carbon atoms 1 and 1'.

In conjunction with the previous results, a knowledge of the activity of the phthalonic acid permitted the calculation by difference of the activity of C-4. However, the hydration of the phthalonic acid (which analysed as $C_{11}H_{10}O_6 \cdot H_2O$ even after vacuum drying at 100°) made it an equivocal standard. A more reliable assessment of the combined activities of its constituent carbon atoms was computed from the sum of the specific activities of 3:5-dimethoxyphthalic anhydride, prepared by dehydration of the free acid and barium carbonate, derived from C-2 in the course of decarboxylation of 3:5-dihydroxyphenylacetic acid. However, this alternative evaluation proved to be almost identical with the observed activity of the phthalonic acid monohydrate. The average activity of carbon atoms 3' and 5' was calculated from the known values of carbons 1, 1', 2', 4', 6' and orcinol.

The observed distribution of labelling (Table 2) clearly substantiated the postulated pathway involving acetate, thereby justifying the classification of alternariol with a number of other fungal phenols of similar biosynthetic origin. Some slight variation in the activities of alternate carbon atoms was apparent which may be significant. In particular the contrast between the lactone carbonyl and C-6 was outside the normal limits of experimental error; however, at present no obvious explanation of these results has been found.

The nature of the post-acetate intermediates is unknown, but if a sequential condensation of

acetate units is assumed, then a consideration of the various alternative pathways leads to some interesting possible interrelationships. One mechanism of biosynthesis which has been suggested involves the oxidative coupling of orsellinic acid and 3:5-dihydroxyphthalic acid (Aghoramurthy & Seshadri, 1954). However, the orientation of hydroxyl substituents in alternariol is such that this would necessitate an abnormal type of phenol coupling (Barton & Cohen, 1957). The simpler direct pathway involving the condensation of seven acetate units (as in Fig. 1) requires carbon atoms 1 and 1' to be derived from a single C_2 unit, thus precluding the independent origin of the two benzenoid rings. On this basis, orsellinic acid (arising from four acetate units, for example via IX) could in theory give rise to either but not both of rings A and B of alternariol (Fig. 4).

In order to test this possibility [*carboxy*- ^{14}C]-orsellinic acid was synthesized from orcinol and sodium [^{14}C]cyanide and fed to *A. tenuis* under the conditions previously used for the uptake of acetate. However, the resulting alternariol was inactive, as it was also when orsellinic acid was added in the form of its ethyl ester with a view to overcoming possible disadvantages of the free acid, such as ease of decarboxylation or cell-wall-impermeability factors. This finding does not exclude the possibility of there being a C_{14} intermediate containing an intact six-membered ring, e.g. ' α -orsellinyl triacetic acid' (X), in which the orsellinic moiety corresponds to ring A of alternariol (Fig. 5). The particular theoretical advantage of this intermediate is that it can also give rise to a number of other natural phenols, as it is capable of various alternative modes of ring closure, including five (routes A-E) in which a second benzenoid ring is formed.

Thus biochemically feasible oxidation and reduction steps could lead to a product of structure (XI) which is a probable precursor of citromycin (XII), a metabolite of *Penicillium frequentans* and other species, whose origin from acetate has been established by Birch *et al.* (1958a). It has been previously pointed out that the product (XI) could serve as a common precursor of citromycin and fulvic acid (XIII) (involving oxygen-bridge formation), or alternatively it could yield the naphthalene derivatives fusarubin, purpurogenone and javanicin (probable structure XIV) following a carbocyclic condensation analogous to the route A folding of the intermediate (X) (Dean, Eade, Moubasher & Robertson, 1957; Whalley, 1958) as indicated in Fig. 5.

The complete derivation of javanicin from acetate in this manner is perhaps less certain than that of fusarubin and purpurogenone, as this would involve the reduction of the aromatic naphthalene

Table 2. *Distribution of labelling in alternariol derived from [^{14}C]acetic acid*

The relative activity is based on overall activity of seven units. Activities in parentheses were obtained by difference: activities of carbon atoms 3' and 5' represent the mean of their combined activities.

C atom	Relative activity	C atom	Relative activity
1	0	1'	0.90
2	1.05	2'	0
3	0	3'	(0.95)
4	(1.11)	4'	0
5	0	5'	(0.95)
6	0.90	6'	0
Me	0	CO	1.14

Fig. 4. Orsellinic acid as a possible alternariol precursor.

carboxyl to a methyl group, a process for which there does not at present appear to be a known analogy. Furthermore, as Birch *et al.* (1958*b*) showed that one of the methyl side chains of the dimethyl-substituted benzoquinone aurantioglucladin arises by the replacement of an acetate carboxyl group by a C_1 unit (formate), one would expect that the corresponding methyl group of javanicin is derived from formate or some other C_1 donor.

This raises the interesting possibility that an analogous substitution of a polyisoprenoid side chain into an acetate-derived 3-carboxy-2-methyl-1:4-naphthaquinone by displacement of the carboxyl group could account for the biosynthesis of vitamins K_1 and K_2 , in just the same way as Birch and co-workers have postulated a corresponding biosynthetic relationship between aurantioglucladin and coenzyme Q.

An alternative carbocyclic condensation of the intermediate (X) is between C-atoms 1 and 6 (route B), yielding a benzophenone derivative, which could precede the formation of both the lichen product lichexanthone (XV) (Asahina & Nogami, 1942) and griseofulvin (XVI), another mould product of proven acetate origin (Birch *et al.* 1958*c*). On the other hand, a 6:7 condensation

(route C) followed by δ -lactone formation would yield alternariol (I).

Although the product (XVII) of a 1:8 condensation (route D) has at present no known C_{14} analogue, it is closely related to the C_{15} plant phenol eleutherinol (XVIII) which occurs together with eleutherin (XIX) in *Eleutherine bulbosa* (Schmid, 1952). The carbon skeletons of both of these products could arise by the addition of a further acetate to the carboxyl of the acid (X) followed by decarboxylation and subsequent condensations analogous to routes D and A respectively. Furthermore, a route B type condensation of this C_{15} intermediate would yield a potential precursor of the fungal *p*-methylbenzophenone derivatives sulochrin (Nishikawa, 1939), geodin, erdin (Barton & Scott, 1958) and pinselin (Munekata, 1953), whereas in theory all three modes of folding (routes A, B and D) could precede formation of the polyhydroxy-2-methylanthraquinone group of mould pigments.

No examples of natural products related to the acid (XX) resulting from a 4:2' condensation (route E) of (X) have been described. This may simply reflect the reluctance of a phenolic hydroxyl to undergo condensation, although the acid (X) need not represent the actual common precursor

Fig. 5

but only its carbon skeleton; thus the six-membered ring could still be present at a pre-aromatic stage.

In addition to the support given to this mechanism through the co-occurrence of eleutherin and eleutherinol, a further example of the possible use of alternative condensations of a common precursor exists in *Carpenteles brefeldianum*, which produces fulvic acid (XIII) on Raulin-Thom medium and griseofulvin (XVI) on Czapek-Dox (Dean *et al.* 1957).

If α -orsellinyl triacetic acid (X) does act as a common precursor of various bicyclic C_{14} fungal phenols, it follows that the branched side chain can itself serve as an aromatic precursor. Consequently if both rings of alternariol are formed in a similar manner, then the original ring [corresponding to orsellinic acid (II)] would arise from α -acetyltri-acetic acid (IX) as illustrated in Fig. 4. This latter acid is the open-chain form of dehydroacetic acid (XXI) (Fig. 6) and it is of interest that alternaric acid (XXII), a metabolite of *Alternaria solani* (Bartels-Keith & Grove, 1959) may be regarded as a dihydro-dehydroacetic acid derivative.

Dehydroacetic acid has been shown in *Penicillium urticae* to inhibit strongly the production of 6-methylsalicylic acid and gentisyl alcohol (Ehrens-värd, 1955). However, this could be explained as a consequence of competitive inhibition, since 6-methylsalicylic acid (being a deoxyorsellinic acid) would require in the above pathway the mediation of a dihydro derivative of dehydroacetic acid. It is perhaps significant that the particular dihydro-dehydroacetic acid required for 6-methylsalicylate synthesis is precisely that shown to be present in alternaric acid. Dehydroacetic acid could also ring-close to form acetophenone derivatives similar to phloracetophenone and clavatul (Hassall & Todd, 1947).

An alternative biosynthetic scheme involving the folding of linear poly- β -keto acid precursors has been elaborated by Birch (1957) and also by Woodward (1957), who in addition suggested the

possible mediation of macrocyclic lactones of the type known to be present in the 'macrolide' class of antibiotics. On this basis the mould metabolite curvularin could be considered as an advanced intermediate, since it contains both a benzene and a macrocyclic lactone ring and has been shown to arise from acetate (Birch, Musgrave, Rickards & Smith, 1959).

However, although the linear polyacetate hypothesis could also explain the biosynthesis of alternariol, it is inadequate in cases such as citromyccetin, fulvic acid, sulochrin, pinselin and geodin, which clearly require a branched-chain precursor.

EXPERIMENTAL

All melting points are uncorrected. Analyses for C, H, N and S were carried out by Weiler and Strauss of Oxford and by Mr F. H. Oliver of Parke, Davis and Co. Ltd., Hounslow, Middlesex.

Culture and medium. The culture of *Alternaria tenuis* was the London School of Hygiene and Tropical Medicine strain number S.M. 108, which was grown in surface culture on a Czapek-Dox medium as described by Raistrick *et al.* (1953).

Radioactive assay. The ^{14}C content of samples mounted on 1.5 cm. diameter planchets was determined at infinite thickness with a thin end-window Geiger-Müller counter, as outlined by Chaplen & Thomas (1960).

Determination of alternariol production. Czapek-Dox medium (350 ml./flask) was inoculated with a mycelial suspension of *A. tenuis* in sterile water and incubated in the dark at 24°. Over 10 weeks, sample batches of four flasks were removed at regular intervals (4 days), and the combined culture filtrates examined for pH, residual glucose (determined polarimetrically) and ferric reaction. The combined mycelium was dried, weighed and extracted in a Soxhlet apparatus, initially with light petroleum (b.p. 40–60°) for 24 hr. and subsequently with dry ether for a further 24 hr. These extracts were evaporated to dryness; the ether extract was washed with boiling light petroleum before ultimately drying and weighing.

The pH of the culture filtrate rose from 4.40 to 7.65 in the first 2 weeks and increased only a further 0.2 unit in the following 5 weeks. The filtrate at 2 weeks and thereafter yielded a green colour on addition of a few drops of $FeCl_3$, which changed to a dark-brown solution accompanied by

Fig. 6

a precipitate in the presence of an excess of the reagent. The other results are shown graphically in Fig. 2.

Incorporation of sodium [^{14}C]acetate. Autoclaved solutions of sodium [^{14}C]acetate (60 $\mu\text{C}/5\text{ ml.}$) were added to the metabolism liquors of 19-day cultures by careful pouring down the wall of each flask so as to pass under the mycelial pad. In this way, labelled acetate was added to ten flasks which were subsequently harvested after a total of 29 days' growth and extracted as above. The crude light-petroleum extract (2.60 g.) and the dry ether extract (2.72 g.) of the mycelium (36.06 g.) possessed closely similar activities (13 $\mu\text{C}/\text{g.}$). Thus the incorporation of added acetate into the light-petroleum extract (lipid fraction) and the ether extract (consisting largely of alternariol and its methyl ether) was in each case approx. 6%.

Demethylation of the ether extract. The ether extract (1.582 g.) was demethylated by refluxing with constant-b.p. HI (10 ml.) in the presence of phenol (4 g.) which facilitated solution of the alternariol methyl ether. The effluent methyl iodide was absorbed in pure dimethyl-aniline; the crystalline precipitate of phenyltrimethyl-ammonium iodide which began to form after about 30 min., was filtered at 2 hr. and washed with chloroform. The dried crystals, m.p. 218°, when assayed for ^{14}C content were found to be devoid of radioactivity.

Demethylation of the ether extract was continued for a further 10 hr., after which time no further methyl iodide was evolved. The crude alternariol (1.452 g.) was recovered by addition of water and filtration of the resulting precipitate, which was washed thoroughly with hot water before drying.

Preparation and dilution of alternariol trimethyl ether. Crude alternariol (1.36 g.) was refluxed with dimethyl sulphate (6 ml.) and anhydrous K_2CO_3 (20 g.) in dry acetone (150 ml.) until the solution no longer gave a purple ferric reaction (8 hr.). It was then filtered and concentrated to yield alternariol trimethyl ether, $\text{C}_{17}\text{H}_{16}\text{O}_5$ (1.43 g.), which after crystallization from acetone melted at 161°. This product (1.003 g.) (activity, 4127 counts/min.) was diluted with pure inactive trimethyl ether (9.027 g.) and recrystallized from acetone (400 ml.), yielding the source material for the degradation studies whose activity (410 counts/min. $\equiv$ 460 $\mu\text{C}/\text{mole}$) exceeded the average background count rate approximately 50-fold.

Degradation of alternariol trimethyl ether. The preparation of many of the products used in these experiments has been described in detail by Raistrick *et al.* (1953) and where possible the same procedures were adopted.

Kuhn-Roth oxidation of alternariol trimethyl ether. Alternariol trimethyl ether (608 mg.) when oxidized by refluxing with a mixture of 5*N*-chromic acid (120 ml.) and conc. H_2SO_4 (30 ml.) produced a quantitative yield of acetic acid, which was separated by steam-distillation and neutralized with 0.1*N*-LiOH. The aqueous solution was concentrated by distillation and a portion converted into *p*-bromophenacyl acetate, which was purified by crystallization from light petroleum (b.p. 80–100°) and vacuum sublimation, yielding colourless needles, m.p. 84–85° (Found: activity, 59.1 $\mu\text{C}/\text{mole}$). The remaining lithium acetate was evaporated to dryness and pyrolysed in a current of N_2 at 390°; the effluent acetone was absorbed in water and converted into iodoform by treatment with aqueous alkaline iodine. The iodoform was oxidized to CO_2

and assayed as BaCO_3 , which was found to be free from radioactivity.

Preparation of 2:3':4:5'-tetrahydroxy-6-methyldiphenyl. Alternariol trimethyl ether (1.036 g.) was hydrolysed by gently refluxing with *N*-NaOH (20 ml.) for 15 min. The resulting solution was cooled, dimethyl sulphate (2 ml.) added and the mixture boiled for about a minute. This procedure gave rise to a crystalline deposit (1.166 g.) which was purified by crystallization from methanol, yielding the methyl ester of 2:3':4:5'-tetramethoxy-6-methyldiphenyl-2'-carboxylic acid, m.p. 124° (Raistrick *et al.* 1953) (Found: activity, 454 $\mu\text{C}/\text{mole}$). A mixture of this ester (0.499 g.), red P (2 g.) and constant-b.p. HI (15 ml.) was heated in an oil bath at 140°, and the effluent CO_2 assayed as BaCO_3 (Found: activity, 74.7 $\mu\text{C}/\text{mole}$). The hydrolysis was continued for 2 hr. in order to obtain complete demethylation, after which the solution was filtered and cooled. The resulting crystalline precipitate (0.271 g.) was purified by sublimation in a high vacuum at 200°, yielding 2:3':4:5'-tetrahydroxy-6-methyldiphenyl, m.p. 246–248° (Found: activity, 385 $\mu\text{C}/\text{mole}$).

Nitration of 2:3':4:5'-tetrahydroxy-6-methyldiphenyl. 2:3':4:5'-Tetrahydroxy-6-methyldiphenyl (0.59 g.) was heated at 100° with conc. H_2SO_4 (1.5 ml.) for 30 min. The mixture was cooled in an ice bath and conc. HNO_3 (4 ml.) slowly added, the temperature being maintained below 5° until the initial exothermic reaction was complete. The temperature was then raised to 70°, when a vigorous reaction took place giving rise to a yellow crystalline precipitate. When the reaction had subsided, the mixture was diluted with water (10 ml.) and cooled. After filtration and washing with ice-cold water, an initial crop of yellow microneedles was obtained (0.502 g.), m.p. 243°. A small second crop of needles (0.023 g.) separated from the filtrate on standing overnight. Ether extraction of the mother liquor from these two crops produced further material (0.240 g.).

The nitration product, after repeated crystallization from 0.1*N*-HCl, melted at 246° with vigorous decomposition. The analysis corresponded to that required for the δ -sultone of 2:3':4:5'-tetrahydroxy-6-methyl-3:4':5:6'-tetranitrodiphenyl-2'-sulphonic acid (Found: C, 32.6; H, 1.3; N, 11.4, 11.6; S, 7.0. $\text{C}_{13}\text{H}_6\text{O}_{14}\text{N}_4\text{S}$ requires C, 32.9; H, 1.3; N, 11.8; S, 6.8%).

Hypobromite degradation of the tetranitrosultone. The sultone (200 mg.) was dissolved in boiling water (10 ml.) and hot $\text{Ba}(\text{OH})_2$ solution (20 ml.) added, giving rise to a reddish-brown precipitate. The mixture was then cooled to 4° and treated with a solution of Br_2 (0.5 ml.) in saturated $\text{Ba}(\text{OH})_2$ (50 ml.) at the same temperature. The characteristic pungent odour of bromopierin, $\text{CBr}_3\cdot\text{NO}_2$, was rapidly generated. The mixture was kept overnight and the product was collected by steam-distillation into a centrifuge tube. After repeated centrifuging and intermediate washing with water, the bromopierin droplet was frozen and transferred to a flask for wet combustion to CO_2 under van Slyke-Folch conditions with chromic acid solution and subsequently precipitated as BaCO_3 (195 mg. $\equiv$ 58% of theoretical yield). The washed and dried product was devoid of any radioactivity.

Potassium permanganate oxidation of alternariol trimethyl ether. Alternariol trimethyl ether (3.90 g.) was refluxed with *N*-NaOH (65 ml.) for 15 min., after which the resulting solution was cooled and oxidized at room temperature with excess of aq. KMnO_4 (260 ml. of a 10% solution)

and kept overnight. The heavy MnO_2 precipitate was removed by filtration. The filtrate was adjusted to pH 7 with conc. HCl, and the excess of permanganate reduced by the addition of sodium dithionite. It was then acidified to pH 2 before extracting with ether (3 x 1 vol.). Evaporation of this extract yielded crystals of 3:5-dimethoxyphthalic acid (0.558 g.), m.p. 153–156°, and a partially crystalline residue (0.161 g.).

The ether-extracted oxidation solution was neutralized with 2N-NaOH and partially evaporated *in vacuo* to about 200 ml.; it was then reacidified and extracted continuously with ether for 24 hr. Concentration of this second ether extract yielded a largely crystalline pale-yellow gum (2.089 g.) which on repeated crystallization from water yielded 4:6-dimethoxyphthalonic acid, m.p. 173° (decomp.). Drying to constant weight at 100° in a high vacuum produced the acid monohydrate (Found: C, 48.1; H, 4.5; activity, 328 μ C/mole. Calc. for $C_{11}H_{10}O_7 \cdot H_2O$; C, 48.5; H, 4.4%).

3:5-Dimethoxyphthalic anhydride. The $KMnO_4$ oxidation product, 3:5-dimethoxyphthalic acid, was purified by crystallization from water and dehydrated at 190° for 5 min., by which time the evolution of water was complete. On cooling, the melt crystallized and pure 3:5-dimethoxyphthalic anhydride was prepared by crystallization from *o*-xylene, followed by sublimation in a high vacuum, giving colourless needles, m.p. 148–149° (Found: activity, 259 μ C/mole).

Reductive decarboxylation of 4:6-dimethoxyphthalonic acid. 4:6-Dimethoxyphthalonic acid (0.473 g.) was heated with a mixture of red P (1.0 g.) and constant-b.p. HI at 140° for a total of 8 hr. The effluent CO_2 was trapped as $BaCO_3$ which was recovered by filtration, washed with CO_2 -free water, dried and assayed (Found: activity, 74.7 μ C/mole). The reaction mixture was filtered while hot, cooled, and extracted with ether. Successive washing of the extract (150 ml.) with water (5 ml.) then conc. $Na_2S_2O_3$ solution (5 ml.) removed HI and I_2 respectively and on evaporation to dryness a partly crystalline colourless gum (0.231 g.) was obtained which completely crystallized on trituration with ether. The product (3:5-dihydroxyphenylacetic acid) was purified by vacuum sublimation at 130°, giving a crystalline sublimate, m.p. 130° (Found: activity 247 μ C/mole).

Decarboxylation of 3:5-dihydroxyphenylacetic acid. 3:5-Dihydroxyphenylacetic acid (0.165 g.) was decarboxylated by heating the solid at 270° in a stream of N_2 gas; the effluent CO_2 was trapped as $BaCO_3$ (0.130 g.) (Found: activity, 69.1 μ C/mole). The decarboxylation product was collected by sublimation on a 'cold finger' and then purified by further sublimation in a high vacuum at 70°. This yielded a crystalline sublimate (0.091 g.) of m.p. 107–108°, which was unchanged on admixture with orcinol purified by a similar sublimation procedure (Found: activity, 184 μ C/mole).

Oxidation of the orcinol with CrO_3 under Kuhn–Roth conditions gave a quantitative yield of acetic acid, part of which was converted into its *p*-bromophenacyl ester as before (Found: activity, 59.4 μ C/mole). The remainder was pyrolysed as lithium acetate to give acetone, subsequent alkaline-iodine treatment of which yielded iodoform as described above. When oxidized and converted into $BaCO_3$, this product proved to be inactive.

Synthesis of [carboxy- ^{14}C]orsellinic acid. [formyl- ^{14}C].

Orsellinic aldehyde was synthesized by a modification of the method of Adams & Levine (1923) in which 1 mc of $Na^{14}CN$ (9.8 mg.) was added to $Zn(CN)_2$ (3 g.) before reacting with orcinol (2.0 g.) in the appropriate manner. The resulting crude orsellinic aldehyde (2.16 g., m.p. 181°) was then converted into orsellinic acid as described by Hoesch (1913). This further sequence of reactions involved the initial formation of the diethoxycarbonyl derivative, m.p. 58°, followed by oxidation of the aldehyde with $KMnO_4$ in aqueous acetone to the acid, m.p. 115°, which was finally hydrolysed with dilute NaOH to orsellinic acid (1.50 g.). The free acid was purified by crystallization from water and sublimation in a high vacuum, yielding colourless needles, m.p. 184–185° (Found: activity, 72 μ C/g.). The pure orsellinic acid, which melted at a higher temperature than the normally recorded value (176°), produced the expected purple ferric reaction in ethanolic solution. The identity of the product was confirmed by preparation of its methyl ester, m.p. 137° (recorded m.p. 138–140°), and also its ethyl ester, m.p. 130° (recorded m.p. 131–132°), by brief treatments of the acid with diazomethane and diazoethane respectively.

Attempted incorporation of orsellinic acid and ethyl orsellinate into alternariol. Approx. 7 μ C of the above orsellinic acid (100 mg.) in ethanolic solution (2 ml.) was added to each of two flasks containing 18-day cultures of *A. tenuis*. One flask was harvested at 21 days and the other at 25 days. The mycelium was only slightly active (24 counts/min.) and the activities of both the light-petroleum and ether extracts of the mycelium were relatively negligible. A corresponding experiment with labelled ethyl orsellinate (3 μ C in 50 mg.) in ethanol (2 ml.) also failed to produce significant incorporation of activity.

SUMMARY

1. The biosynthesis of alternariol, $C_{14}H_{10}O_6$, a phenolic product of *Alternaria tenuis*, has been studied.
2. Chemical degradation of labelled alternariol derived from sodium [^{14}C]acetate clearly demonstrated a biosynthetic mechanism involving the head-to-tail condensation of acetate units.
3. A possible common precursor with other fungal phenols containing C_{14} skeletons is discussed.

I am grateful to Imperial Chemical Industries Ltd. for the purchase of some of the tracer equipment and ^{14}C -labelled compounds used in this work, which was carried out during the tenure of an I.C.I. Research Fellowship. I also wish to thank Professor J. H. Birkinshaw for his continued interest and encouragement.

REFERENCES

- Adams, R. & Levine, I. (1923). *J. Amer. chem. Soc.* **45**, 2373.
Aghoramurthy, K. & Seshadri, T. R. (1954). *J. sci. industr. Res.* **13A**, 114.
Asahina, Y. & Nogami, H. (1942). *Bull. chem. Soc. Japan*, **17**, 202.

- Bartels-Keith, J. R. & Grove, J. F. (1959). *Proc. chem. Soc., Lond.*, p. 398.
- Barton, D. H. R. & Cohen, T. (1957). In *Festschrift Arthur Stoll*, p. 117. Ed. by Stoll, A. Basle: Birkhäuser.
- Barton, D. H. R. & Scott, A. I. (1958). *J. chem. Soc.* p. 1767.
- Birch, A. J. (1957). *Fortschr. Chem. org. Naturst.* **14**, 186.
- Birch, A. J., Fitton, P., Pride, E., Ryan, A. J., Smith, H. & Whalley, W. B. (1958a). *J. chem. Soc.* p. 4576.
- Birch, A. J., Fryer, R. I. & Smith, H. (1958b). *Proc. chem. Soc., Lond.*, p. 343.
- Birch, A. J., Massey-Westropp, R. A., Rickards, R. W. & Smith, H. (1958c). *J. chem. Soc.* p. 360.
- Birch, A. J., Musgrave, O. C., Rickards, R. W. & Smith, H. (1959). *J. chem. Soc.* p. 3146.
- Chaplen, P. & Thomas, R. (1960). *Biochem. J.* **77**, 91.
- Cornforth, J. W., Hunter, G. D. & Popják, G. (1953). *Biochem. J.* **54**, 597.
- Cullinane, N. M., Morgan, N. M. E. & Plummer, C. A. J. (1937). *Rec. Trav. chim. Pays-Bas*, **56**, 627.
- Dean, F. M., Eade, R. A., Moubasher, R. A. & Robertson, A. (1957). *Nature, Lond.*, **179**, 366.
- Ehrensward, G. (1955). *Exp. Cell Res. Suppl.* **3**, 102.
- Gatenbeck, S. & Mosbach, K. (1959). *Acta chem. scand.* **13**, 1561.
- Hassall, C. H. & Todd, A. R. (1947). *J. chem. Soc.* p. 611.
- Hoesch, K. (1913). *Ber. dtsh. chem. Ges.* **46**, 886.
- Munekata, H. (1953). *J. Biochem., Tokyo*, **40**, 451.
- Nishikawa, H. (1939). *Acta phytochim., Tokyo*, **11**, 187.
- Raistrick, H., Stickings, C. E. & Thomas, R. (1953). *Biochem. J.* **55**, 421.
- Rosett, T., Sankhala, R. H., Stickings, C. E., Taylor, M. E. U. & Thomas, R. (1957). *Biochem. J.* **67**, 390.
- Schmid, H. (1952). *Fortschr. Chem. org. Naturst.* **11**, 124.
- Thomas, R. (1959). *Proc. chem. Soc., Lond.*, p. 88.
- Whalley, W. B. (1958). *Chem. & Ind.* p. 131.
- Woodward, R. B. (1957). In *Festschrift Arthur Stoll*, p. 524. Ed. by Stoll, A. Basle: Birkhäuser.

Biochem. J. (1961) **78**, 758

The Metabolism of 3:5-Di-*tert.*-butyl-4-hydroxytoluene and 3:5-Di-*tert.*-butyl-4-hydroxybenzoic Acid in the Rabbit

By J. C. DACRE

Toxicology Research Unit of the Medical Research Council of New Zealand, Medical School, University of Otago, Dunedin, New Zealand

(Received 27 April 1960)

The compound known commercially as butylated hydroxytoluene (3:5-di-*tert.*-butyl-4-hydroxytoluene) is an antioxidant developed initially for use with petrol, petroleum products and rubber. Dugan, Marx, Weir & Kraybill (1954) found butylated hydroxytoluene to be an effective antioxidant for animal fats at a concentration of 0.01%, and legislative approval for its addition to fats and foods containing fats intended for human consumption has already been given in eight countries including U.S.A., Denmark, Sweden and the United Kingdom. The toxicity of the compound to rabbits, rats, guinea pigs, cats and dogs was investigated by Deichmann, Clemmer, Rakoczy & Bianchine (1955), who found that when it was given orally to rabbits, there was an increased excretion of both glucuronide and ethereal sulphate in the urine. The causes of the toxic and lethal effects in rabbits given large doses of butylated hydroxytoluene were investigated by Denz & Llauro (1957) and some aspects of its chronic toxicity in rats were studied by Brown, Johnson & O'Halloran (1959).

It will be shown in this paper that 3:5-di-*tert.*-butyl-4-hydroxybenzoic acid is a metabolite of

butylated hydroxytoluene in both the free and conjugated forms. The metabolism of this acid and of the antioxidant butylated hydroxytoluene has been investigated in the rabbit and the major metabolites excreted in the urine have been isolated and characterized. A summary of these results has been published (Dacre, 1960).

EXPERIMENTAL

Animals, diet and dosage. The rabbits used were does of the New Zealand white strain, weighing 1.33–2.0 kg. They were maintained on a standard pellet diet (wheat, 40%; pollard, 33%; bran, 27%) and during experiments were kept singly in metabolism cages designed to permit the separate collection of urine and faeces. For the isolation of metabolites in quantity, rabbits were maintained on a modified diet (wheat, 35%; pollard, 20%; meat meal, 15%; bran, 10%; maize meal, 10%; milk powder, 7.5%; bone meal, 1.7%; cod-liver oil, 0.8%). Water and food were supplied *ad lib.* and the urine was collected daily and stored at -10° . In the initial experiments, single doses of 0.8 g. of butylated hydroxytoluene (BHT)/rabbit were dissolved in 7.2 ml. of olive oil and administered by stomach tube. In later experiments, doses of 0.5 and 1.0 g. in 5 ml. of olive oil/rabbit or 0.8 g. of BHT in 4 ml. of olive oil/kg. body wt. of the