however, the catechins are present as minor components associated with high-molecular-weight tannins, whereas in tea leaves they form the major phenolic components.

SUMMARY

1. The distribution of flavonoid compounds in the heartwoods and barks of four wattles (*Acacia* spp.) has been studied.

2. (-)-Fisetinidol, (+)-fustin, (+)-7:3':4'-trihydroxyflavan-3:4-diol, fisetin and polymeric leuco-fisetinidin tannins are common components to the heartwoods of *A. mearnsii* (*A. mollissima*) (black wattle), *A. decurrens* (green wattle), *A. dealbata* (silver wattle) and *A. pycnantha* (golden wattle).

3. The barks of *A. mearnsii* and *A. decurrens* contain (-)-robinetinidol, (+)-catechin and (+)-galocatechin; *A. dealbata* contains these catechins together with (-)-epicatechin and (-)-epigallocatechin, and *A. pycnantha* contains these together with (-)-epicatechin, (-)-epicatechin gallate, (-)-epigallocatechin, (-)-epigallocatechin gallate and gallic acid.

4. The possible significance of the relative distribution of heartwood and bark components to the taxonomy of these wattles is discussed.

Thanks are due to Miss M. Tindale, Royal Botanic Gardens, Sydney, Australia, for confirming the relationships between these *Acacia* spp.; to Mr S. P. Sherry and Professor H. Shaw, Wattle Research Institute, Pietermaritzburg, Natal, for authentic bark and heartwoods of *A. mearnsii*, *A. decurrens* and *A. dealbata*, and for black x green wattle hybrids; to the Director, Union Department of Forestry, Pretoria, for the wood and bark of *A. pycnantha*; and to Dr R. Child, Tea Research Institute of East Africa,

Kericho, Kenya, for dried green tea. This work was financed by the annual grant of the South African Wattle Growers' Union to the Leather Industries Research Institute.

REFERENCES

- Bradfield, A. E. & Bate-Smith, E. C. (1950). *Biochim. biophys. Acta*, **4**, 441.
 Bradfield, A. E. & Penney, M. (1948). *J. chem. Soc.* p. 2249.
 Bradfield, A. E., Penney, M. & Wright, W. B. (1947). *J. chem. Soc.* p. 32.
 Brennan, J. P. M. & Melville, R. (1960). *Kew Bull.* **14**, 37.
 Clark-Lewis, J. W. & Roux, D. G. (1959). *J. chem. Soc.* p. 1402.
 Erdtman, H. (1959). *Proc. 4th int. Congr. Biochem., Vienna*, **2**, 1.
 Evelyn, S. R. (1956). *J. Soc. Leath. Tr. Chem.* **40**, 335.
 Forsyth, W. G. C. (1955). *Biochem. J.* **60**, 108.
 Kalbe, H. (1954). *Hoppe-Seyl. Z.* **297**, 19.
 Keppler, H. H. (1957). *J. chem. Soc.* p. 2721.
 Mayer, W. & Bauni, G. (1957). *Liebigs Ann.* **611**, 264.
 Nordström, C. G. & Swain, T. (1953). *J. chem. Soc.* p. 2764.
 Roberts, E. A. H. & Wood, D. J. (1951). *Biochem. J.* **49**, 414.
 Roberts, E. A. H. & Wood, D. J. (1953). *Biochem. J.* **53**, 332.
 Roux, D. G. (1953). *J. Soc. Leath. Tr. Chem.* **37**, 229.
 Roux, D. G. (1957). *Nature, Lond.*, **179**, 158.
 Roux, D. G. & Evelyn, S. R. (1958a). *Biochem. J.* **69**, 530.
 Roux, D. G. & Evelyn, S. R. (1958b). *Biochem. J.* **70**, 344.
 Roux, D. G. & Evelyn, S. R. (1960). *Biochem. J.* **76**, 17.
 Roux, D. G. & Maihs, E. A. (1958). *Nature, Lond.*, **182**, 1798.
 Roux, D. G. & Maihs, E. A. (1960a). *Biochem. J.* **74**, 44.
 Roux, D. G. & Maihs, E. A. (1960b). *J. Chromat.* **4**, 65.
 Roux, D. G. & Paulus, E. (1961a). *Biochem. J.* **78**, 120.
 Roux, D. G. & Paulus, E. (1961b). *Biochem. J.* **78**, 785.
 Vuataz, L., Brandenberger, H. & Egli, R. H. (1959). *J. Chromat.* **2**, 173.
 Weinges, K. (1958). *Liebigs Ann.* **615**, 203.

Biochem. J. (1961) **78**, 839

A Method for the Identification of the Immediate Product of Decarboxylation Reactions

By R. F. PALMER*

Department of Pharmacology and Therapeutics, University of Florida, Gainesville, Florida, U.S.A.

(Received 12 May 1960)

Under most physiological conditions the state of an enzymic decarboxylation product as bicarbonate ion, carbonic acid or carbon dioxide is determined mainly by local conditions of pH and gas concentration. The fact that the immediate end product of a decarboxylation reaction may actually

* Present address: Department of Medicine, The Johns Hopkins Hospital, Baltimore 5, Maryland, U.S.A.

be any one of the three forms would be masked by its rapid conversion into a form demanded by the buffer and gas composition of the medium. This conversion is rapid in systems where carbonic anhydrase is present. It has therefore been difficult with conventional methods to determine the immediate reaction product in carbon dioxide-producing systems.

Carbonic anhydrase can be used as a tool in studying such reactions, as was first suggested by Roughton and co-workers (Meldrum & Roughton, 1933; Ferguson & Roughton, 1934; Roughton, 1935). Its use in the Warburg manometer established that carbon dioxide was the end product of the urea-urease system. In this system the relation of urease to urea was such that total urea-splitting occurred in a very short time, producing an initial abrupt rise of carbon dioxide pressure and a gradual return to equilibrium. In the presence of carbonic anhydrase no such rise occurred. Krebs & Roughton (1948) used this technique with yeast pyruvic decarboxylase and demonstrated that carbon dioxide was the immediate end product. Conway & O'Malley (1953), by rapid addition of carbonic anhydrase from the side arm of a Warburg flask to a mixture of yeast carboxylase and pyruvate producing carbon dioxide at a steady rate, altered the rate of production of the gas in a mathematically predictable way depending on whether the product were carbonic acid or carbon dioxide. They concluded that the product obtained depended to a certain extent on the source and purity of the yeast carboxylase.

It has been presumed (Woodbury & Esplin, 1959; Koch & Woodbury, 1960) that carbonic acid was the primary product of brain-tissue decarboxylations and that the role of carbonic anhydrase in brain cells was catalysis of bicarbonate to carbon dioxide. To test this hypothesis and to study further the physiological effect of carbonic anhydrase and its inhibitors in the central nervous system (Maren & Robinson, 1960), we utilized Krebs & Roughton's and Conway & O'Malley's methods on brain tissue of mice to determine the dominant end product of brain decarboxylations. It became apparent that inherent in both these methods was the necessity of a more powerful decarboxylase preparation than could be obtained with concentrated brain homogenates or minces in conventional Warburg vessels. In addition, the fact that the tissue was consuming oxygen at the same time as it was producing carbon dioxide obscured the small deflections of pressure necessary to differentiate the end products. A method was needed which was not influenced by oxygen consumption and which was relatively independent of the magnitude of the rate of carbon dioxide production.

In this paper a new technique is described that permits differentiation of these end products in physiological or other complex systems producing carbon dioxide at slow rates. The technique introduces a means of removing carbon dioxide from the medium by bubbling carbon dioxide-free air through the reactants. Carbon dioxide is thus removed as it is formed either primarily from the decarboxylation of substrate, or secondarily from

the dehydration of carbonic acid as it is formed from substrate. The rate of collection of carbon dioxide is measured. Conditions, particularly rate of air flow and buffer pH, are chosen in such a way that the products can be distinguished kinetically. The distinction will ultimately depend on the velocity constants of several simultaneous reactions which are described in the theoretical section.

If carbon dioxide were the primary product, increase of flow rate will increase the rate of liberation of gas. Addition of carbonic anhydrase will decrease the rate of liberation of carbon dioxide. If carbonic acid or bicarbonate were the primary product, the rate of liberation of the gas will be relatively unaffected by flow rate but strongly increased by the addition of carbonic anhydrase. In this paper the method will be described and its application to the yeast pyruvic-decarboxylase system presented as a model.

MATERIALS AND METHODS

Apparatus. (See Fig. 1.) The reaction chamber consisted of a Toribara Ultra Filtration tube (obtained from Will Corp., Atlanta, Georgia) through which air previously passed through a soda-lime cannister and moistened by bubbling through distilled water was allowed to flow at a constant rate. Air was dispersed by the sintered-glass filter into minute bubbles, flowed through the reactants (usually enzyme, substrate and buffer) and was collected in a gas washing bottle (collecting vessel) containing a measured amount of NaOH in CO_2 -free water. Phenolphthalein was added to the collecting solution, and the time for the indicator to reach a faint-pink end point was measured. Solutions could be added rapidly to the reaction vessel during constant air flow through a 15-gauge blunt syringe

Fig. 1. Reaction chamber and collecting vessel.
For explanation see text.

needle inserted through the rubber stopper. The needle was attached to a two-way metal stopcock, and syringes containing the solutions were connected to the cock. A typical experiment began with an accurately-measured constant flow of air passing through the empty reaction chamber and collected in the vessel containing 100 ml. of CO_2 -free water and a few drops of phenolphthalein. The collecting vessel containing the base and indicator was found to be 99–100% efficient in collecting all CO_2 under the most rapid flow rates employed. Two syringes were attached to the stopcock, one containing 1 ml. of 0.2M-sodium pyruvate and 1.0 ml. of 0.5M-phosphate buffer, pH 7.8, and the other containing 0.5 ml. of yeast carboxylase of known activity. A few seconds before addition of the reactants to the reaction chamber, 0.5 ml. of 0.01N-NaOH was added to the collecting vessel, thereby turning the phenolphthalein pink. At zero time the stopcock was opened, the syringes were emptied into the reaction vessel, and the time to the phenolphthalein end point in the collecting vessel was recorded. When carbonic anhydrase was used, it was included in the syringe containing the carboxylase preparation.

Yeast. Freshly pressed brewer's yeast, obtained from Schwarz Laboratories, Mount Vernon, New York, was spread in thin layers and allowed to dry at room temperature with the aid of a fan and then stored in the refrigerator until use. A portion (20 g.) of this dried preparation was incubated with 60 ml. of water for 2 hr. at 25° and then centrifuged, and the clear supernatant used as a source of

under a vacuum and to perform control experiments on all solutions to ensure that the reactants were not contaminated with bicarbonate or other volatile acids.

Air flow. At flows greater than 60 ml./min., a calibrated flow meter (Flowmizer S/N 23976, tube size 2-15-3; Brooks Rotometer Co., Lansdale, Pa., U.S.A.) was used, accurate to approximately 2%. At lower flows, where meters are less accurate, flow was determined by counting the number of bubbles of air per minute flowing through the water bottle before entering the reaction vessel. By counting the number of bubbles and simultaneously collecting the gas in an inverted water-filled graduated cylinder it was possible to measure flow quite reproducibly and accurately.

Warburg manometer. For estimation of yeast carboxylase activity, Warburg shaking and constant-temperature apparatus manufactured by Gilson Medical Electronics, Middleton, Wis., U.S.A., was used. Calibrated conical flasks of approximately 20 ml. capacity with double side arms were employed.

Corrections were made for buffer effects by addition of HCl from the second side arm as described by Umbreit, Burris & Stauffer (1957).

THEORETICAL

The following scheme describes the events which may take place in the reaction and collecting vessels:

carboxylase. The supernatant was then further diluted with water to give an activity as measured in the Warburg at 25° of 1–2 μmoles of CO_2 /min. with a standard pyruvate solution.

Pyruvate. Sodium pyruvate, supplied by Schwarz Laboratories, Mount Vernon, New York, was prepared in an 0.2M-concentration with CO_2 -free distilled water. A portion (1 ml.) of this solution added to 0.5 ml. of diluted yeast carboxylase in 2 ml. of phosphate buffer yielded a steady rate of CO_2 production over a 10 min. period.

Carbonic anhydrase. A freeze-dried preparation of the chloroform-ethanol extract prepared from human blood by the method of Meldrum & Roughton (1933) containing 37 enzyme units (e.u.)/mg. was used. Enzyme activity was determined by the method of Maren, Ash & Bailey (1954). One unit by this method is roughly equivalent to 5 Meldrum-Roughton units. In control experiments with carbonic anhydrase, acetazolamide was added to the enzyme preparation in a concentration sufficient to totally inhibit the enzyme. Acetazolamide had no effect on the action of yeast carboxylase in the standard manometric procedure.

Results would be seriously affected if any of the solutions were contaminated with CO_2 or HCO_3^- . It was therefore extremely important to shake and then store all reactants

where X and Z are the rates of formation of CO_2 and HCO_3^- (or H_2CO_3) respectively in $\mu\text{moles/min.}$

If CO_2 is the immediate product it can conceivably follow two pathways. It can be blown off by the air which is flowing through the reactants at a rate Y measured in ml./min. Such a rate of removal is given by

$$\frac{d[\text{CO}_2]}{dt} = -k_{\text{app.}} Y [\text{CO}_2] \quad (2)$$

where $k_{\text{app.}}$ is a constant for the apparatus, including the diffusion constant of CO_2 , a constant for bubble size and distribution, turbulence of liquid and air and volume of the reaction vessel and collecting tubing. This overall constant can be obtained by the addition of 10.0 $\mu\text{equiv.}$ of NaHCO_3 to strong acid in the reaction vessel and measuring the length of time (t) required to neutralize 5.0 $\mu\text{equiv.}$ of NaOH in the collecting bottle as various rates of flow. Then:

$$k_{\text{app.}} = \frac{2.3 \log 2}{t_1 \cdot Y} = 0.03 \text{ ml.}^{-1} \quad (3)$$

for the flow rates used.

The second pathway CO_2 may take is that of hydration and secondary dehydration. From the scheme we may write the rate of change of CO_2 as:

$$\frac{d[\text{CO}_2]}{dt} = \frac{X}{V} - k_{\text{app.}} \cdot Y[\text{CO}_2] - k_{\text{CO}_2}[\text{CO}_2] + k_{\text{H}_2\text{CO}_3}[\text{H}_2\text{CO}_3] \quad (4)$$

where V is the volume of the solution in the reaction vessel.

The hydration and dehydration reactions have been extensively and accurately studied. Rate constants are given and methods reviewed by Roughton (1943) and Kern (1960). At 25° k_{CO_2} is 0.037 sec.^{-1} and $k_{\text{H}_2\text{CO}_3}$ is 20 sec.^{-1} (from the review by Kern, 1960).

Since H_2CO_3 or HCO_3^- is instantly interconverted (Roughton, 1941) we may write the sum of the rates of change of these two species as:

$$\begin{aligned} \frac{d[\text{HCO}_3^-]}{dt} + \frac{d[\text{H}_2\text{CO}_3]}{dt} \\ = \frac{Z}{V} + k_{\text{CO}_2}[\text{CO}_2] - k_{\text{H}_2\text{CO}_3}[\text{H}_2\text{CO}_3]. \end{aligned} \quad (5)$$

The equilibrium constant of the ionization is written as:

$$K_{\text{H}_2\text{CO}_3} = \frac{[\text{H}^+][\text{HCO}_3^-]}{[\text{H}_2\text{CO}_3]}. \quad (6)$$

The values for the equilibrium constant $K_{\text{H}_2\text{CO}_3}$ are reviewed by Kern (1960). The value of 1.8×10^{-4} is taken from the work of Scheurer, Brownell & LuValle (1958).

Since the hydrogen-ion concentration is maintained by the buffer, the rate of change of H_2CO_3 is proportional to the rate of change of HCO_3^- . From equation (6):

$$\frac{d[\text{HCO}_3^-]}{dt} = \frac{K_{\text{H}_2\text{CO}_3}}{[\text{H}^+]} \cdot \frac{d[\text{H}_2\text{CO}_3]}{dt}. \quad (7)$$

Inserting this into equation (5), we obtain:

$$\begin{aligned} \left(1 + \frac{K_{\text{H}_2\text{CO}_3}}{[\text{H}^+]}\right) \cdot \frac{d[\text{H}_2\text{CO}_3]}{dt} \\ = \frac{Z}{V} + k_{\text{CO}_2}[\text{CO}_2] - k_{\text{H}_2\text{CO}_3}[\text{H}_2\text{CO}_3]. \end{aligned} \quad (8)$$

A solution to the two simultaneous first-order differential equations (4) and (8) is given in the Appendix.

Inspection of equations (4) and (8) shows that if CO_2 is the primary product, at high flows ($k_{\text{app.}} \cdot Y$ very large) CO_2 will be quantitatively removed; the rate-limiting factor is its production (X). If, on the other hand, H_2CO_3 or HCO_3^- is the primary product, at high flows the removal of formed CO_2 is limited by the rate of the dehydration reaction. Approximate equations applicable to the conditions employed here are:

(i) For the hypothetical case of CO_2 production in terms of total CO_2 liberated (R). Let

$$\begin{aligned} m = 1 + \frac{K_{\text{H}_2\text{CO}_3}}{[\text{H}^+]}. \\ R = X \left[\frac{k_{\text{app.}} \cdot Y}{k_{\text{CO}_2} + \frac{k_{\text{H}_2\text{CO}_3}}{m}} \right] \left[\frac{-1}{\frac{k_{\text{H}_2\text{CO}_3}}{m} + k_{\text{CO}_2} + k_{\text{app.}} \cdot Y} + t \right] \end{aligned} \quad (9)$$

(ii) For the hypothetical production of H_2CO_3 or HCO_3^- .

$$\begin{aligned} R = Z \left[t - \left\{ 1 + \frac{k_{\text{CO}_2}}{\left(\frac{k_{\text{H}_2\text{CO}_3}}{k_{\text{app.}} \cdot Y} \right) \cdot m} \right\} \right. \\ \left. \times \left\{ 1 - \exp \left(\frac{-k_{\text{app.}} \cdot Y \cdot t}{1 + \frac{k_{\text{CO}_2}}{k_{\text{H}_2\text{CO}_3} \cdot m}} \right) \right\} \right] \end{aligned} \quad (10)$$

RESULTS

Experiments with yeast carboxylase

Two syringes were attached to the stopcock, one containing 0.5 ml. of diluted yeast carboxylase and 2 ml. of 0.5M-phosphate buffer, pH 7.8, and the other containing 1 ml. of 0.2M-sodium pyruvate. For each series of experiments the activity of the yeast extract was measured in the Warburg. Production of CO_2 (X) varied from 0.8 to 2.6 $\mu\text{moles/min.}$ The collecting vessel contained 5 μmoles of NaOH in 100 ml. of water. The term R is then equivalent to the amount of NaOH added minus the amount required to bring the CO_2 -free water to a faint-pink end point (usually 1.5 $\mu\text{moles/100 ml.}$). The flow rate was varied from 10 to 60 ml./min. The solutions were injected and the time to the phenolphthalein end point was measured. The results of several experiments are given on Table 1, and compared with the predicted values for primary CO_2 formation obtained by substitution in equation (9.) A representative experiment is plotted in Fig. 2.

Experiments with bicarbonate

The hypothetical situation in which a substrate-enzyme system produces HCO_3^- or H_2CO_3 was simulated by the slow constant addition of NaHCO_3 solution to phosphate buffer under conditions of constant air flow. A syringe filled with 0.03M- NaHCO_3 , whose plunger was slowly depressed at a constant rate by a micrometer device attached to a low-speed motor, was attached to the stopcock on the reaction flask. The rate of addition was adjusted by a rheostat which controlled the motor. With a constant flow of air through 3.5 ml. of 0.5M-phosphate buffer in the reaction chamber, the

Table 1. *Reaction times at various rates of air flow and various rates of yeast pyruvic-decarboxylase activity*

The collecting vessel contains, in all cases, 3.5 μ moles of phenolphthalein-titratable NaOH (*R*). Predicted values are obtained from substitution in equation (9).

Carboxylase activity CO ₂ production ($\bar{X}$) (μ moles/min.)	Rate of air flow (<i>Y</i>) (ml./min.)	Experimental reaction times (<i>t</i>) (min.)		Predicted reaction time (min.)	<i>R/X</i> (min.)
0.8	22	10.7	11.0	12.8	4.35
	49	6.2	6.0	5.8	—
	61	4.8	4.9	4.7	—
1.06	24	7.6	7.4	8.9	3.3
	37	4.1	—	3.4	—
	51	3.5	3.4	3.3	—
1.87	10	10.8	11.0	12.2	1.9
	15	6.9	7.4	8.3	—
	27	3.9	4.2	4.7	—
	40	3.0	2.9	3.2	—
	48	2.5	2.7	2.7	—
	60	2.0	2.1	2.1	—
2.85	12	6.1	6.3	6.9	1.2
	26	3.0	3.2	3.3	—
	42	2.0	2.2	2.1	—

Table 2. *Reaction times at various rates of air flow and various rates of addition of 0.03 M-sodium hydrogen carbonate to phosphate buffer, pH 7.8, in the reaction chamber*

The collecting bottle contains 3.5 μ moles of phenolphthalein-titratable NaOH (*R*). Theoretical times are calculated by substitution in equation 10 of the text.

Rate of addition of 0.03 M-NaHCO ₃ (<i>Z</i>) (μ moles/min.)	Rate of air flow (<i>Y</i>) (ml./min.)	Experimental reaction times (<i>t</i>) (min.)		Theoretical reaction times (min.)
1.20	23	13.8	13.9	14.3
	30	13.9	13.8	13.8
1.87	18	12.0	12.1	13.0
	23	12.1	12.2	11.1

motor was started and the time required to reach the phenolphthalein end point in the collecting vessel was measured. Flow was varied from 10 to 60 ml./min. Results at different rates of addition are given in Table 2 and compared with the predicted values for primary HCO₃⁻ formation from equation (10). A representative experiment is plotted in Fig. 2.

Experiments with carbonic anhydrase

In these experiments the flow rate was held constant at 30 ml./min.; 0.2 ml. of carbonic anhydrase solution containing 370 e.u./ml. was mixed with the yeast extract; with this exception the technique was the same as that previously described for the carboxylase system. In experiments with constant flow of NaHCO₃, carbonic anhydrase was first added to the phosphate buffer. As a control the same volume of enzyme solution plus 0.1 ml. of 1 mg./ml. acetazolamide was added. Results are given in Table 3. It is evident that carbonic anhydrase prolongs the reaction time in the yeast carboxylase system and shortens it in the experiments with NaHCO₃.

DISCUSSION

Results show that it is possible to distinguish between the two end products, as predicted by the theoretical treatment. Comparison of the experimental curves in Fig. 2 shows that it is principally the rate of air flow that enables the differences in the reaction velocities to become apparent. Velocities principally concerned are the terms

$k_{app.} \cdot Y$ contrasted with $k_{H_2CO_3} \frac{[H^+]}{K_{H_2CO_3}}$. If $k_{app.} \cdot Y$ is sufficiently large, the amount of CO₂(*R*) which is blown off is equal to *X* $\cdot t$. On the other hand, in the HCO₃⁻ experiments the term $k_{app.} \cdot Y$ plays no role at flow rates beyond a point where it is permissible to neglect the reverse reaction. The rate of liberation of CO₂ ($\frac{R}{t}$) is determined principally by

$$k_{H_2CO_3} \cdot \frac{[H^+]}{K_{H_2CO_3}}.$$

The two model systems of Fig. 2 are arranged so that the rates of HCO₃⁻ addition on the one hand and CO₂ formation on the other are constant and equal. Regardless of the magnitude of these rates,

Fig. 2. Representative curves of reaction times versus rate of air flow in the yeast pyruvic-decarboxylase system (O) and in the NaHCO_3 addition to buffer experiments (●). X and $Z = 1.87 \mu\text{moles/min.}$ For details see text.

the differences between the two species will now be apparent at high flow. Confirmation is given by the use of carbonic anhydrase, which by increasing the magnitude of both k_{CO_2} and $k_{\text{H}_2\text{CO}_3}$ decreases the rate of liberation of CO_2 if CO_2 is the end product and increases it if HCO_3^- or H_2CO_3 is the end product (Table 3).

The effect of carbonic anhydrase on the reaction time in the NaHCO_3 experiments is much greater than in the CO_2 (yeast carboxylase) system. Equations (4) and (8) show that the effect on NaHCO_3 dehydration is exponential, and on CO_2 hydration is inversely proportional, accounting for the differences observed in Table 3.

Experiments with yeast pyruvic decarboxylase show unequivocally that CO_2 is the dominant, if not the sole, end product. The response to various rates of flow and to the addition of carbonic anhydrase is entirely consistent with this. This finding is somewhat in contrast with the results of Conway & O'Malley (1953), who concluded that, in the decarboxylation of keto acids by yeast carboxylase, CO_2 and H_2CO_3 (or HCO_3^-) are both primary

products but that one or the other may predominate, depending on the total conditions. It is possible with the present method to make an analysis concerning the dominance of CO_2 as an end product, as follows: from Fig. 2 it is evident that at flow rates where the curve is asymptotic $t = R/X$. If a significant amount of HCO_3^- or H_2CO_3 were being formed in addition to CO_2 (Conway & O'Malley, 1953) this would not be the case, t at asymptote being greater than R/X . From these data, it appears that if any HCO_3^- or H_2CO_3 is formed in addition to CO_2 , it is undetectable.

APPENDIX

Solutions to equations (4) and (8) in the text may be obtained for $[\text{CO}_2]$, $[\text{H}_2\text{CO}_3]$ and $[\text{HCO}_3^-]$ by the method of operators. These solutions are in the form $a + be^{-r_1t} + ce^{-r_2t}$, where a , b and c are different constants for the concentrations of the three substances. The rate constants of the two exponential terms are the two roots, $D = -r_1$ and $D = -r_2$, of the equation

$$m \cdot D^2 + (k_{\text{H}_2\text{CO}_3} + m \cdot k_{\text{app}} \cdot Y + m \cdot k_{\text{CO}_2}) D + k_{\text{app}} \cdot Y \cdot k_{\text{H}_2\text{CO}_3} = 0 \quad (1)$$

$$\text{where } m = 1 + \frac{K_{\text{H}_2\text{CO}_3}}{[\text{H}^+]}$$

Equation (1) holds whether CO_2 , H_2CO_3 or HCO_3^- is the primary product. Designating C as the sum of the concentrations of the three components, and C_∞ as the value of C at infinite time, the solution for C , if CO_2 is the primary product, may be written

$$C = C_\infty - \frac{C_\infty r_2 - \frac{X}{V}}{r_2 - r_1} \cdot e^{-r_1t} - \frac{\frac{X}{V} - C_\infty r_1}{r_2 - r_1} \cdot e^{-r_2t} \quad (2)$$

where C_∞ is given by

$$C_\infty = \frac{X}{V} \left(\frac{1}{k_{\text{app}} \cdot Y} \right) \left(1 + \frac{k_{\text{CO}_2}}{k_{\text{H}_2\text{CO}_3}} \cdot m \right) \quad (3)$$

Table 3. Effect of carbonic anhydrase on the reaction times of the yeast-carboxylase system and the simulated bicarbonate- or carbonic acid-producing system at constant flow rate

Carbonic anhydrase (74 e.u.) is added to the syringe containing yeast carboxylase and buffer. In experiments with NaHCO_3 , carbonic anhydrase is added to the phosphate buffer in the reaction vessel. As a control, 0.1 ml. of 1.0 mg./ml. acetazolamide is added to the carbonic anhydrase solution. Air flow is constant at 30 ml./min. The collecting bottle contains 3.5 μmoles of phenolphthalein-titratable NaOH (R). Figures are means of five experiments $\pm$ S.E.M.

System	Rate of production of CO_2 or HCO_3^- (X or Z) ($\mu\text{moles/min.}$)	Reaction times (min.)		
		No carbonic anhydrase	Carbonic anhydrase plus acetazolamide	Carbonic anhydrase alone
Yeast carboxylase	1.87 (X)	4.0 $\pm$ 0.07	4.0 $\pm$ 0.06	4.6 $\pm$ 0.09
NaHCO_3	1.87 (Z)	12.0 $\pm$ 0.03	12.1 $\pm$ 0.04	4.8 $\pm$ 0.15

P values < 0.01 between experiments with and without carbonic anhydrase.

If HCO_3^- or H_2CO_3 is the primary product, Z replaces X in equation (2) above and C_∞ is given by

$$C_\infty = \frac{Z}{V} \left(\frac{1}{k_{\text{app}} \cdot Y} \right) \left(1 + \frac{k_{\text{app}} \cdot Y + k_{\text{CO}_2} \cdot m}{k_{\text{H}_2\text{CO}_3}} \right). \quad (4)$$

The amount of CO_2 removed, R , is given by

$$R = Xt - CV \quad (5)$$

when CO_2 is the primary product. When H_2CO_3 or HCO_3^- is the primary product

$$R = Zt - CV. \quad (6)$$

It is then possible to calculate precisely the amount of CO_2 blown off from either system in any circumstances. If the values of the hydration and dehydration velocity constants at 25° are substituted and $Y = 27.5$ ml./min. is chosen, equation (1) above yields

$$r_1 = 0.000462 \text{ sec.}^{-1} \text{ and } r_2 = 0.52 \text{ sec.}^{-1}$$

at a pH of 7.80. Therefore the first exponential term in equation (2) above becomes relatively small only after 10^4 sec., whereas the second exponential term is negligible after 10 sec. For the times actually employed in these experiments the first exponential term of equation (2) may be expanded with retention of the first term of the series. The second exponential term of equation (2) may be dropped.

Certain limiting cases are of interest. If $k_{\text{app}} \cdot Y$ is large, we may neglect the first term in equation (1) above. If $k_{\text{app}} \cdot Y$ is small we may neglect the third term in equation (1). r_1 and r_2 are then given by simple expressions. Substitution of the values for r_1 and r_2 in equation (2) and solving equations (3) and (5) yield, for the case of CO_2 as the primary product,

$$R = X \left[1 - \frac{k_{\text{H}_2\text{CO}_3} + k_{\text{CO}_2}}{m} \right] \left[\frac{-1}{k_{\text{app}} \cdot Y} + t \right] \quad (7)$$

when $k_{\text{app}} \cdot Y$ is large, and text equation (9) when $k_{\text{app}} \cdot Y$ is small. For the case of HCO_3^- or H_2CO_3 production, retention of only the first term of the expansion of the first exponential term in equation (2) is not sufficient. If the first exponential term of equation (2) is not expanded and the second exponential term is neglected we obtain

$$R = Z \left[t - \frac{m}{k_{\text{H}_2\text{CO}_3}} \left\{ 1 - \exp \left(\frac{-k_{\text{H}_2\text{CO}_3} \cdot t}{m} \right) \right\} \right] \quad (8)$$

for H_2CO_3 or HCO_3^- production when $k_{\text{app}} \cdot Y$ is large. Expansion of equation (8) and retention of only two terms results in

$$R = Z \frac{k_{\text{H}_2\text{CO}_3}}{m} \cdot t^2 \quad (9)$$

which is applicable for short reaction times. If $k_{\text{app}} \cdot Y$ is small, text equation (10) is obtained, which yields, on expansion to two terms,

$$R = Z \frac{k_{\text{app}} \cdot Y \cdot t^2}{1 + \frac{k_{\text{CO}_2}}{k_{\text{H}_2\text{CO}_3}} \cdot m}. \quad (10)$$

SUMMARY

1. A method is proposed to distinguish between carbon dioxide and HCO_3^- ion as end products of decarboxylation reactions.

2. The method is dependent upon the removal of carbon dioxide by flowing air at various rates through reactants and then quantitatively collecting and measuring the liberated carbon dioxide. Theoretical and experimental evidence are presented, demonstrating that the distinction is dependent on the velocity constant of the dehydration of carbonic acid.

3. Experiments with yeast pyruvic carboxylase and the constant addition of sodium hydrogen carbonate, both with and without carbonic anhydrase, confirmed the validity of the assumptions.

4. Yeast pyruvic decarboxylase produces carbon dioxide as its dominant, if not sole, end-product.

The author wishes to thank Dr T. H. Maren for his support, enthusiasm and helpful suggestions. He is indebted to Dr A. L. Koch for his interest and advice, particularly in the development of the theoretical portion.

REFERENCES

- Conway, E. J. & O'Malley, E. (1953). *Biochem. J.* **54**, 154.
 Ferguson, J. J. W. & Roughton, F. J. W. (1934). *J. Physiol.* **83**, 68.
 Kern, D. M. (1960). *J. chem. Educ.* **37**, 14.
 Koch, A. & Woodbury, D. M. (1960). *Amer. J. Physiol.* **198**, 434.
 Krebs, H. A. & Roughton, F. J. W. (1948). *Biochem. J.* **43**, 550.
 Maren, T. H., Ash, V. I. & Bailey, E. M., jun. (1954). *Johns Hopk. Hosp. Bull.* **95**, 244.
 Maren, T. H. & Robinson, B. (1960). *Johns Hopk. Hosp. Bull.* **106**, 1.
 Meldrum, N. V. & Roughton, F. J. W. (1933). *J. Physiol.* **80**, 143.
 Roughton, F. J. W. (1935). *Physiol. Rev.* **15**, 263.
 Roughton, F. J. W. (1941). *J. Amer. chem. Soc.* **63**, 2930.
 Roughton, F. J. W. (1943). *Harvey Lect.* **39**, 96.
 Scheurer, P., Brownell, R. & LuValle, J. (1958). *J. phys. Chem.* **62**, 809.
 Umbreit, W. W., Burris, R. H. & Stauffer, J. F. (1957). *Manometric Techniques*, chapt. 2, p. 18. Minneapolis: Burgess Publishing Co.
 Woodbury, D. M. & Esplin, D. W. (1959). *Res. Publ. Ass. nerv. ment. Dis.* **37**, 24.