stituent amino acids (glutamic acid, α -amino-n-butyric acid, and glycine); transpeptidation also occurred in the nuclear fraction, and γ -glutamyl transpeptidase is the enzyme which brings about hydrolysis and transpeptidation in the nuclear fraction. The breakdown of ophthalmic acid in the supernatant fraction gives 5-oxopyrrolidine-2-carboxylic acid rather than glutamic acid, and is ascribed to the action of γ -glutamyl lactamase. The localization of γ -glutamyl lactamase in the supernatant fraction was shown when γ -glutamylglycine was used as substrate.

5. In the subcellular fractionation of calf lens, nearly all the protein is in the supernatant fraction. The enzymes that are most readily detected in lens extracts are those which are present in the supernatant fraction of liver.

We should like to thank our colleagues at this Laboratory for advice and encouragement, Miss Stewart for technical assistance, The Staffs of the Oxford Co-operative and Eastwyck Farm Slaughterhouses for their co-operation, the National Council to Combat Blindness Inc., New York, for a Fight for Sight Grant, and The Distillers Co. Ltd. (Biochemicals) for a gift of glutathione. One of us (E.E.C.) was in receipt of a Thornton Bursary.

REFERENCES

- Allfrey, V. (1959). In *The Cell*, vol. 1, p. 193. Ed. by Brachet, J. & Mirsky, A. E. New York: Academic Press Inc.
- Bailie, M. J. & Morton, R. K. (1958). *Biochem. J.* **69**, 35.
- Bates, H. M. & Lipmann, F. (1960). *J. biol. Chem.* **235**, PC22.
- Binkley, F., Davenport, J. & Eastall, F. (1959). *Biochem. biophys. Res. Commun.* **1**, 206.
- Cliffe, E. E. & Waley, S. G. (1958a). *Biochem. J.* **69**, 649.
- Cliffe, E. E. & Waley, S. G. (1958b). *Nature, Lond.*, **182**, 804.
- Cliffe, E. E. & Waley, S. G. (1959a). *Biochem. J.* **73**, 25p.
- Cliffe, E. E. & Waley, S. G. (1959b). *Biochem. J.* **72**, 12p.
- Cohn, W. E. & Carter, C. E. (1950). *J. Amer. chem. Soc.* **72**, 4273.
- Connell, G. E. & Hanes, C. S. (1956). *Nature, Lond.*, **177**, 377.
- Fodor, P. J., Miller, A. & Waelsch, H. (1953). *J. biol. Chem.* **202**, 551.
- Hanes, C. S., Hird, F. J. R. & Isherwood, F. A. (1952). *Biochem. J.* **51**, 25.
- Harrison, M. F. (1953). *Nature, Lond.*, **171**, 611.
- Hird, F. J. R. & Springell, P. H. (1954a). *Biochem. J.* **56**, 417.
- Hird, F. J. R. & Springell, P. H. (1954b). *Biochim. biophys. Acta*, **15**, 31.
- Hogeboom, G. H. & Schneider, W. C. (1952). *J. biol. Chem.* **197**, 611.
- Hogeboom, G. H., Schneider, W. C. & Striebich, M. J. (1952). *J. biol. Chem.* **196**, 111.
- Holleman, J. W. (1954). *C.R. Acad. Sci., Paris*, **238**, 1360.
- Hunter, G. (1957). *J. clin. Path.* **10**, 161.
- Lang, K., Siebert, G. & Müller, L. (1951). *Naturwissenschaften*, **38**, 528.
- Lofffield, R. B., Grover, J. W. & Stephenson, M. L. (1953). *Nature, Lond.*, **171**, 1024.
- Morton, R. K. (1958). *Nature, Lond.*, **181**, 540.
- Neubeck, C. E. & Smythe, C. V. (1944). *Arch. Biochem.* **4**, 443.
- Rademaker, W. & Soons, J. B. J. (1957). *Biochim. biophys. Acta*, **24**, 451.
- Reindel, F. & Hoppe, W. (1954). *Ber. dtsh. chem. Ges.* **87**, 1103.
- Revel, J. P. & Ball, E. G. (1959). *J. biol. Chem.* **234**, 577.
- Robinson, H. W. & Hogden, C. G. (1940). *J. biol. Chem.* **135**, 727.
- Roodyn, D. B. (1959). *Int. Rev. Cytol.* **8**, 279.
- Rosenthal, O., Gottlieb, B., Gorry, J. D. & Vars, H. M. (1956). *J. biol. Chem.* **223**, 469.
- Schneider, W. C. (1948). *J. biol. Chem.* **176**, 259.
- Snoke, J. E. & Bloch, K. (1955). In *Methods in Enzymology*, vol. 2, p. 342. Ed. by Colowick, S. P. & Kaplan, N. O. New York: Academic Press Inc.
- Wade, H. E. & Morgan, D. M. (1955). *Biochem. J.* **60**, 264.
- Waley, S. G. (1958). *Biochem. J.* **68**, 189.
- Waley, S. G. (1959). *Biochem. J.* **71**, 132.
- Wanko, T. & Gavin, M. A. (1959). *J. biophys. biochem. Cytol.* **6**, 97.
- Webster, G. C. (1953). *Arch. Biochem. Biophys.* **47**, 241.

Biochem. J. (1961) **79**, 128

Studies on the Flash Photolysis of Visual Pigments

1. PIGMENTS PRESENT IN FROG-RHODOPSIN SOLUTIONS AFTER FLASH-IRRADIATION

By C. D. B. BRIDGES

Visual Research Division, M.R.C. Ophthalmological Research Unit, Institute of Ophthalmology, Judd Street, London, W.C. 1

(Received 27 April 1960)

Irradiation of visual-pigment solutions with brief intense bursts of light makes it possible to detect products of bleaching which have only a transient existence or which are themselves photo-reactive. Ideally, the requirement is for an in-

stantaneous flash of intensity sufficient to excite all the visual-pigment molecules: in practice, electronically-operated discharge tubes of the type used in photography provide the best available light sources. Such tubes are capable of producing

flashes of high luminous intensity with durations ranging from a few microseconds to several milliseconds.

With a 64J argon-filled flash-tube assembly delivering a 2–6 μ sec. flash, Wulff, Adams, Linschitz & Abrahamson (1958) irradiated solutions of cattle rhodopsin. Specially designed equipment, which measured absorption spectra more or less immediately after the flash, revealed the presence of rapidly fading photoproducts with maxima displaced 12 $m\mu$ towards the blue. After about 1 hr. all thermal changes had ceased, and the solutions were found to contain a stable photosensitive substance with $\lambda_{\max}$ at 490 $m\mu$: 8 $m\mu$ below that of the parent rhodopsin.

Solutions of frog rhodopsin were used in the present work and were exposed to the flash from a 400J xenon-filled discharge tube. These flashes were of higher intensity but longer duration than those used by Wulff and his colleagues, yet the results are essentially similar. Initially produced thermally unstable pigments faded rapidly, and the material which remained after thermal changes were complete was stable and photosensitive. When this stable material was tested for homogeneity by the method of partial bleaching (Dartnall, 1952) two photosensitive pigments were detected.

METHODS

Preparation of frog-rhodopsin solutions

Frogs (*Rana temporaria*) were dark-adapted for about 16 hr. and killed by decapitation, and the retinæ removed in dim red light (Lythgoe, 1937). The retinæ were then washed three times with McIlvaine's buffer, pH 4.6, and extracted twice with aqueous 2% (w/v) digitonin solution (powdered digitonin obtained from L. Light and Co. Ltd.). Sufficient saturated sodium borate solution was added to the resultant extract to bring the pH to 8.2–8.7. Extracts prepared in this manner had $E_{\min}/E_{\max}$ ranging from 0.30 to 0.59.

One extract was prepared by floating off the rod outer segments in isopycnotic sucrose solution (Saito, 1938) and then proceeding as for whole retinæ.

Hydroxylamine solutions

Aqueous hydroxylamine (0.1M) was prepared from the hydrochloride and neutralized (to pH 7.0) by the addition of a few drops of concentrated sodium hydroxide solution.

Absorption spectra

In the experiments designed to measure thermally unstable pigments a Beckman DK-2 automatically recording spectrophotometer was used (1 cm. silica cells). With this apparatus spectra from 620 to 380 $m\mu$ were measured in about 1–1½ min. Thermally stable material was examined in a Unicam SP. 500 spectrophotometer (1 cm. glass cells) or, when partial-bleaching investigations were carried out, in a spectrophotometer (0.5 cm. glass cells) similar in

principle to that described by Dartnall (1952). With the DK-2 and SP. 500 measurements were made at $25 \pm 3^\circ$. With the author's spectrophotometer the temperature was maintained at $25 \pm 0.5^\circ$.

Bleaching apparatus

Flash equipment. The flash tube was a xenon-filled Siemens SF E.22, operated from a Majorette power pack supplied for this purpose by Clive Courtenay and Co. Ltd. The energy dissipated through the tube in all experiments was 400J. Light-output was found to be reproducible within 1%, and the duration of the flash was about 4 msec. (see Fig. 1).

The tube was in the form of a tight helix of 1½ turns and about 6 cm. diam. It was surrounded by a ground-glass cylinder surmounted by a disk carrying an optical-cell holder. During a flash the silica or glass cell filled with rhodopsin solution was suspended in the centre of the helix. Thus light from the discharge tube irradiated the solution through both optical faces. The temperature was $25 \pm 3^\circ$. Although of such high intensity, the flash produced no detectable change in the temperature of the solution.

Partial-bleaching equipment. The author's spectrophotometer was designed as a dual-purpose instrument. By widening the monochromator entry and exit slits, changing to an intense ribbon-filament light-source and using the appropriate stray-light filters, solutions could be exposed to monochromatic radiation. A mirror was placed behind the glass optical cells during all bleaching operations, thus effectively doubling the light-intensity. The temperature was maintained at $25 \pm 0.5^\circ$.

Normal white bleaching. Normal white bleaching was used to convert all photosensitive pigments into indicator yellow. This was carried out by exposing solutions for 15 min. to the radiation from a 15W lamp at a distance of about 1 ft. (illumination ~ 25 ft.-candles).

RESULTS

In Fig. 2 are shown the changes in absorption spectrum after flash-irradiation of frog rhodopsin. Curve A is the pre-flash spectrum. Measurement of curve B was commenced within 30 sec. of exposure to the flash and took about 1 min. to complete. The $\lambda_{\max}$ of this curve is at 478 $m\mu$. Allowing 8 $m\mu$ correction for $E_{\min}/E_{\max}$. (Collins & Morton, 1950; Crescitelli & Dartnall, 1954), the

Fig. 1. Light-output curve of xenon-filled discharge tube.

true $\lambda_{\max}$ is estimated to be at $486 \text{ m}\mu$. Curve *C* ($\lambda_{\max}$ after $10 \text{ m}\mu E_{\min}/E_{\max}$ correction = $493 \text{ m}\mu$) was obtained after $1\frac{1}{2}$ hr. in darkness. A considerable loss in extinction had occurred at wavelengths longer than about $405 \text{ m}\mu$, the difference spectrum for this process having $\lambda_{\max}$ at approx. $477 \text{ m}\mu$. Subsequently, the extinction of the solution remained constant, but what remained was photosensitive, as is shown by the further fall in extinction to curve *D* after exposure to ordinary white light. The difference spectrum of this material showed a maximum at about $494 \text{ m}\mu$.

The difference spectra of the fading process (henceforward called Fraction 1), the residual thermally stable photosensitive material (henceforward called Fraction 2) and the parent frog rhodopsin are compared in the lower half of Fig. 2. The rhodopsin difference spectrum has been obtained by subtracting the final fully bleached curve *D* from the pre-flash curve *A*; it is, of course, independent of any intervening processes.

In another similar experiment, the light of the flash was passed through an orange Ilford 201 filter before irradiating the solution. Fraction 1

was spectroscopically identical with that produced by an unfiltered flash. Fraction 2, however, although produced in substantially the same quantity as by the unfiltered flash, had $\lambda_{\max}$ as high as $500 \text{ m}\mu$. This observation is considered further in the next paper (Bridges, 1961).

Effect of hydroxylamine. The quality and quantity of Fraction 2 are not altered by the presence of 0.01 M - or 0.03 M -hydroxylamine. Thus in experiments on solutions without hydroxylamine Fraction 2 had $\Delta E_{\max}$ $32.45 \pm 2.07\%$ of that of the parent rhodopsin; in the presence of hydroxylamine this percentage was 32.22 ± 0.99 . Again, the difference spectra of Fraction 2 produced in the presence and the absence of hydroxylamine were identical from 620 to about $480 \text{ m}\mu$. Divergence of the curves below this point (Fig. 3) was attributed to the differing absorptions of the photoproducts retinene oxime and alkaline indicator yellow (*N*-retinylidene opsin).

Is Fraction 2 a mixture of photosensitive pigments?

Fig. 4 illustrates a typical experiment designed to test the homogeneity of Fraction 2. Curve *B* is the absorption spectrum of a solution of frog rhodopsin measured $1\frac{1}{2}$ hr. after a flash. All thermal changes were completed, as shown by the fact that the return measurements from 610 to $390 \text{ m}\mu$ follow the curve of the outward measurements from 380 to $620 \text{ m}\mu$ (cf. Dartnall, 1952). Spectra *C*, *D* and *F* are respectively after irradiation for 1, 5 and 3 hr. with $\lambda 600 \text{ m}\mu$ light (irradiance of the order of 10^{13} quanta $\text{sec}^{-1} \text{ cm}^{-2}$). The spectrum *E* was measured after the solution had been left

Fig. 2. Effect of flash-irradiating a solution of frog rhodopsin. Curve *A*, initial absorption spectrum; curve *B*, absorption spectrum of initial photoproduct: this curve was measured from 600 to $400 \text{ m}\mu$ (the reading at $500 \text{ m}\mu$ was 30 sec. after the flash); curve *C*, after $1\frac{1}{2}$ hr. in darkness; curve *D*, after exposure to normal white light (all spectra measured on automatic recorder). The lower half of the Figure shows the following difference spectra: parent pigment (*D* minus *A*); Fraction 1 (*C* minus *B*); Fraction 2 (*D* minus *C*).

Fig. 3. A comparison of difference spectra produced when Fraction 2 is bleached with white light: +, in the presence of 0.01 M -hydroxylamine; x, no hydroxylamine added.

overnight in darkness and was due to an alteration in irrelevant absorption. Since this experiment was carried out in the presence of 0.01 M-hydroxylamine, no regeneration could occur. Finally, curve *G* is the absorption spectrum of the solution after exposure to a 15 w lamp for 15 min.

In the lower half of Fig. 4 are shown the difference spectra of each bleaching operation, namely *C* - *B*, *D* - *C*, *F* - *E* and *G* - *F*. Their maxima shift from 502 m μ for the first λ 600 m μ irradiation to 486 m μ for the final white bleach. This experiment, and the others listed in Table 1, clearly show that two photosensitive pigments are present in Fraction 2. The three difference spectra shown in Fig. 5 have been obtained by subtracting from the fully bleached absorption spectrum of a solution the corresponding absorption spectrum measured after

Fig. 4. The non-homogeneity of Fraction 2. Upper half of Figure: curve *B*, absorption spectrum of a solution of frog rhodopsin 1½ hr. after a flash; curves *C*, *D* and *F*, after 1, 5 and 3 hr. irradiation respectively with λ 600 m μ light; curve *G*, after irradiation with normal white light; curve *E*, after leaving the solution overnight in darkness. Lower half of Figure: the difference spectra for each bleaching operation (the solution contained 0.01 M-hydroxylamine).

exhaustive irradiation with long-wavelength light (curve *G* - *F* of Fig. 4 is such a difference spectrum). These difference spectra are identical within experimental error and show that one of the photosensitive pigments has λ_{max} at 486 ± 2 m μ . Since this pigment cannot be extracted from frog retina, and therefore has no claim to be a visual pigment, it is proposed to call it photosensitive pigment 486 or P486. In this way it can be distinguished from a truly visual pigment, e.g. frog rhodopsin or VP 502.

The other pigment of Fraction 2 seems to be identical with the parent frog rhodopsin, for, as Fig. 6 shows, the difference spectrum resulting from an initial short red irradiation and that produced when frog rhodopsin is bleached are identical. The difference spectrum produced when Fraction 2 is bleached by exposure to white light (as in Fig. 2) is the sum of the difference spectrum produced by bleaching the component pigments. In Fig. 7 the difference spectra produced by bleaching pure P486 and frog rhodopsin respectively are shown scaled according to the proportions in which they occur in Fraction 2. The figure represents the mean of nine results, in which it was found that P486 accounted for $52.3 \pm 3.2\%$ of the total photosensitive material. In terms of the ΔE_{max} values, the total photosensitive pigment in Fraction 2 amounted to $32.87 \pm 1.59\%$ of the pre-flash rhodopsin. (It should be noted that the overall ΔE_{max} for Fraction 2 amounts to only 98.4% of the summed ΔE_{max} values of the component difference spectra.)

Fig. 5. Difference spectra of terminal white bleachings after the exhaustive irradiation of Fraction 2 with long-wavelength light. Corresponding experiments are listed in Table 1. O, Expt. no. 3; +, expt. no. 4; x, expt. no. 5.

Table 1. *Summary of experiments showing the presence of two photosensitive pigments in Fraction 2*

Stray light filters were used in conjunction with the monochromator. Ilford filters 203 and 201 were used for $\lambda 600 \text{ m}\mu$ and $\lambda 580 \text{ m}\mu$ bleaches respectively.

	Expt. no.	Wavelength of bleaching light ($\text{m}\mu$)	Duration (hr.)	ΔE_{max} (0.5 cm.)	λ_{max} ($\text{m}\mu$)
No hydroxylamine	1	600	1	0.0295	500
		580	$\frac{1}{2}$	0.0375	498
		White	$\frac{1}{2}$	0.0460	491
	3	600	4	0.0733	502
		600	2	About 0.0014	About 500
		580	1	0.0114	490
		White	$\frac{1}{2}$	0.0292	485
Hydroxylamine (0.01 M)	4	600	2	0.0620	502
		580	1	0.0287	498
		580	2	0.0220	494
		White	$\frac{1}{2}$	0.0194	488
	5	600	1	0.0497	502
		600	5	0.0560	497
		600	3	0.0185	495
		White	$\frac{1}{2}$	0.0204	486

Fig. 6. Correspondence between the difference spectrum produced when frog rhodopsin is bleached in the presence of 0.01 M-hydroxylamine (+) and that produced by an initial red bleaching of Fraction 2 (x) i.e. curve C minus B of Fig. 4.

Effect of the flash on a freshly bleached solution of frog rhodopsin

A solution of frog rhodopsin was exposed to white light until all the visual pigment had been converted into alkaline indicator yellow. This material was then flash-irradiated, left in darkness for 1 hr. and its absorption spectrum measured.

Fig. 7. Composition of Fraction 2. ●, Difference spectrum produced when Fraction 2 is bleached with white light; ◐, difference spectrum produced when pure pigment 486 is bleached; ○, difference spectrum produced when frog rhodopsin is bleached (0.01 M-hydroxylamine present).

The flash produced a fall in extinction at short wavelengths, but a very small rise in extinction beyond about 450 $\text{m}\mu$. This rise was maximal at about 500 $\text{m}\mu$ and amounted to only about 3% of the original rhodopsin. Exposure of this solution to white light caused bleaching, showing that a photosensitive pigment had been generated by the flash. Its difference spectrum (obtained by subtracting the post-flash absorption spectrum from that of the final fully bleached solution) was identical with that of frog rhodopsin (Fig. 8).

Fig. 8. Nature of the photosensitive pigment produced by flash-irradiation of indicator yellow. A solution of frog rhodopsin ($E_{500\text{ m}\mu}$ 1.4685) was completely bleached by exposure to normal white light. The indicator yellow produced was then flash-irradiated and left for 1 hr. +, Extinction changes (differences spectrum) which took place on further exposure to white light; ○, difference spectrum of frog rhodopsin when bleached by a similar exposure to white light.

DISCUSSION

In the experiments described here, frog-rhodopsin solutions were exposed to intense flashes of light lasting about 4 msec. About $\frac{1}{2}$ min. later, when measurements were begun, the extinction at all wavelengths above 405 mμ was found to be decreasing. After an hour in darkness no further changes were observed, and the solution contained roughly equal amounts of frog rhodopsin and a new photosensitive pigment with λ_{max} at 486 mμ.

Wulff and his colleagues suggested that the initial photoproduct of flash-irradiation of cattle rhodopsin consisted of two fractions. One of these (Fraction 1) decomposed in darkness at room temperature; the other (Fraction 2) was photosensitive and thermally stable. The present observations can be interpreted along similar lines, although another possibility should be considered. Low-temperature studies (Bridges, 1960) have shown that a photoproduct of frog rhodopsin (with λ_{max} at about 480 mμ at -5°) can be thermally converted into the parent photosensitive pigment. It could be therefore that part of Fraction 2 is generated during the thermal-decay process. In the low-temperature work conversions of this nature were characterized by a bodily shifting of the absorption spectrum towards longer wavelengths as the photoproduct changed to a stable photosensitive pigment. This effect would be scarcely noticeable, provided that only a small proportion of Fraction 1 was converted into a Fraction 2 with little change of λ_{max} . However, careful scrutiny of the present results and also those of Wulff *et al.* has revealed no evidence to support this notion.

The situation seems more akin to the low-temperature studies of Wald, Durell & St George (1950) and of Kropf & Hubbard (1958), where it is likely that the product of low-temperature irradiation of cattle rhodopsin is a mixture of thermolabile and thermostable photosensitive pigments (the latter resembling Fraction 2).

In all probability therefore the constituents of Fraction 2 are present throughout the post-flash period, according to Scheme 1.

Unpublished observations by the author (from experiments similar to that of Fig. 2) show that other visual pigments react similarly to flash-irradiation. Rat rhodopsin (λ_{max} 498 mμ) yields a Fraction 1 with λ_{max} at about 473 mμ and Fraction 2 with λ_{max} at 490 mμ. The vitamin A₂ porphyropsin of perch (*Perca fluviatilis*; λ_{max} about 541 mμ) also yields an unstable Fraction 1 and a

Scheme 1

stable photosensitive Fraction 2: both have maxima at about 525 m μ .

Part of the frog rhodopsin in Fraction 2 may be unbleached parent pigment: the remainder, together with P 486, is generated during the flash period or very shortly after it (within a few microseconds in Wulff's experiments). Regeneration from irradiated indicator yellow cannot account for Fraction 2 since its quality and quantity are unaffected by hydroxylamine, which binds the retinene of indicator yellow and renders it incapable of regeneration. In any case, flash-irradiation of indicator yellow yields only a very small quantity of rhodopsin. The mechanism by which Fraction 2 is produced is considered further in the next paper (Bridges, 1961).

Pigment 486

It was suggested by Wulff *et al.* (1958) that their Fraction 2 was a mixture of cattle isorhodopsin (based on isoretinene_{1a}) and cattle rhodopsin (based on neoretinene_{1b}). Cattle rhodopsin has $\lambda_{\max}$ at 498 m μ , cattle isorhodopsin at 487 m μ (Hubbard & Wald, 1952). Thus on changing the steric configuration of the retinene isomer in cattle rhodopsin one obtains a new pigment with $\lambda_{\max}$ shifted by 11 m μ towards the blue. The $\lambda_{\max}$ of P 486 is 16 m μ below that of frog rhodopsin, and it is quite possible that it bears a similar relationship to frog rhodopsin as does cattle rhodopsin to isorhodopsin, i.e. the retinene isomer has been altered. Isorhodopsin is understood to refer to a synthetic photosensitive pigment based on isoretinene_{1a}. Since, however, the relevant retinene isomers have not yet been characterized for the frog pigments, it seems advisable to retain the term P 486 for the flash-generated photosensitive derivative described here.

SUMMARY

1. Alkaline (pH 8.2–8.7) aqueous digitonin solutions of frog rhodopsin at 25° were exposed to

intense 4 msec.-duration flashes from a 400 J xenon-filled discharge tube.

2. Two types of pigment were found in the irradiated solutions. One, with $\lambda_{\max}$ at 477 m μ , was thermally unstable (Fraction 1) and, when left in darkness at 25°, decayed to alkaline indicator yellow over the course of about 1 hr. The other, which was thermally stable and photosensitive (Fraction 2), had $\lambda_{\max}$ at 494 m μ .

3. When Fraction 2 was tested for homogeneity by the method of partial bleaching, it was found to consist of two pigments. One of these, which accounted for 52.3 $\pm$ 3.2 % of the total photosensitive material, had $\lambda_{\max}$ at 486 $\pm$ 2 m μ (photosensitive pigment 486). The other was identical with the parent frog rhodopsin.

4. Fraction 2 was not altered in composition or amount when produced in solutions containing hydroxylamine (0.01 or 0.03 M).

5. Flash-irradiation of alkaline indicator yellow (i.e. an alkaline solution of frog rhodopsin bleached to completion with normal white light) generated only a very small quantity (about 3 %) of the parent rhodopsin.

The author gratefully acknowledges the assistance of Mrs P. M. Rudkin.

REFERENCES

- Bridges, C. D. B. (1960). *Nature, Lond.*, **186**, 292.
- Bridges, C. D. B. (1961). *Biochem. J.* **79**, 135.
- Collins, F. D. & Morton, R. A. (1950). *Biochem. J.* **47**, 3.
- Crescitelli, F. & Dartnall, H. J. A. (1954). *J. Physiol.* **125**, 607.
- Dartnall, H. J. A. (1952). *J. Physiol.* **116**, 257.
- Hubbard, R. & Wald, G. (1952). *J. gen. Physiol.* **36**, 269.
- Kropf, A. & Hubbard, R. (1958). *Ann. N.Y. Acad. Sci.* **74**, 266.
- Lythgoe, R. J. (1937). *J. Physiol.* **89**, 331.
- Saito, Z. (1938). *Tohoku J. exp. Med.* **32**, 432.
- Wald, G., Durell, J. & St George, R. C. C. (1950). *Science*, **111**, 179.
- Wulff, V. J., Adams, R. G., Linschitz, H. & Abrahamson, E. W. (1958). *Ann. N.Y. Acad. Sci.* **74**, 281.