- Milas, N. A. & Sussman, S. (1937). *J. Amer. chem. Soc.* **59**, 2345.
- Nakada, H. I., Friedmann, B. & Weinhouse, S. (1955). *J. biol. Chem.* **216**, 583.
- Nakazawa, Y. (1950). *Folia pharm. jap.* **46**, 6.
- Oettingen, W. F. von. (1943). *Publ. Hlth Bull., Wash.*, no. 281. p. 166.
- Parke, D. V. (1956). *Biochem. J.* **62**, 339.
- Parke, D. V. (1961). *Biochem. J.* **78**, 262.
- Parke, D. V. & Williams, R. T. (1953). *Biochem. J.* **54**, 231.
- Pohl, J. (1923*a*). *Handb. exp. Pharm.* **1**, 860.
- Pohl, J. (1923*b*). *Handb. exp. Pharm.* **1**, 867.
- Pons, C. A. & Custer, R. P. (1946). *Amer. J. med. Sci.* **211**, 544.
- Ratner, S., Nocito, V. & Green, D. E. (1944). *J. biol. Chem.* **152**, 119.
- Riker, W. F. & Gold, H. (1942). *J. Amer. pharm. Ass. (sci. ed.)* **31**, 306.
- Scholz, J. (1956). *Arzneim. Forsch.* **6**, 227.
- Smyth, H. F., jun., Seaton, J. & Fischer, L. (1941). *J. industr. Hyg.* **23**, 259.
- Spector, W. S. (1956*a*). *Handbook of Toxicology*, vol. 1, p. 134. Philadelphia and London: W. B. Saunders Co. Ltd.
- Spector, W. S. (1956*b*). *Handbook of Toxicology*, vol. 1, p. 152. Philadelphia and London: W. B. Saunders Co. Ltd.
- Umbreit, W. W., Burris, R. H. & Stauffer, J. F. (1949). *Manometric Techniques and Tissue Metabolism*, p. 110. Minneapolis: Burgess Publishing Co.
- Weinhouse, S. (1955). In *Amino Acid Metabolism*, p. 637. Ed. by McElroy, W. D. & Glass, H. B. Baltimore: The Johns Hopkins Press.
- Weinhouse, S. & Friedmann, B. (1951). *J. biol. Chem.* **191**, 707.
- Weinhouse, S. & Friedmann, B. (1952). *J. biol. Chem.* **197**, 733.

Biochem. J. (1961) **79**, 489

Induction and Repression of β -Galactosidase in Non-growing *Escherichia coli*

By J. MANDELSTAM

National Institute for Medical Research, Mill Hill, London, N.W. 7

(Received 18 November 1960)

It has been known since the earliest studies of enzyme induction that glucose will often prevent the induced synthesis of enzymes concerned with the metabolism of other carbohydrates. The glucose effect and similar phenomena have been summarized in a number of reviews (Monod & Cohn, 1952; Mandelstam, 1956; Pollock, 1959). It appears to be an instance of a more general type of reaction, now termed repression (Vogel, 1957), in which the rate of synthesis of an enzyme is specifically decreased by a repressor, which is generally, but not necessarily, a normal metabolite of the cells.

The inhibitory effect of glucose upon the induced synthesis of β -galactosidase in growing cultures of *Escherichia coli* has been examined by Cohn & Horibata (1959). The present experiments show that, in non-growing cells (i.e. in the absence of net protein synthesis), the repressor action of glucose is not specific, and that the synthesis of β -galactosidase is effectively inhibited by all substances tested which the cells can utilize as sources of carbon and energy.

MATERIALS AND METHODS

Organisms. The following strains of *E. coli* were used: ML 30 and K₁₂ (wild type); ML 328c (Lac⁺) (requiring

leucine and able to form β -galactosidase); 160-37 (requiring arginine); 26-26 (requiring lysine); N.C.T.C. 8009 (requiring methionine); 15⁻ (requiring thymine); W 3301 (requiring uracil and biotin). All experiments were done at 35°. The wild-type strains were grown with shaking in a synthetic medium (pH 7.2) containing phosphate and inorganic salts (Mandelstam, 1958*a*) and glucose or another carbon source (1%, w/v). All organic acids were neutralized to pH 7.0 with NaOH; concentrations given in the text refer to the acid and not to its salt. For the auxotrophic mutants, the medium was supplemented with the appropriate growth factor as follows: DL-leucine (300 μ g./ml.), L-arginine-HCl (100 μ g./ml.), L-lysine-HCl (150 μ g./ml.), L-methionine (100 μ g./ml.), thymine (10 μ g./ml.), uracil (100 μ g./ml.), biotin (0.05 μ g./ml.).

Induction of β -galactosidase. The cells were harvested when the culture contained about 0.8 mg. dry wt. of bacteria/ml., and were washed with phosphate buffer (having the same composition and pH as the phosphate in the growth medium). In the earlier experiments the bacteria were first starved of their specific requirement or, with the wild-type strains, of ammonium salts (see below). In later experiments this was omitted, and the cells, after washing, were suspended in the same phosphate buffer (at 0.2–0.4 mg. dry wt./ml.) with methyl β -D-thiogalactoside (MTG; 0.1 mg./ml.) as inducer. MTG was synthesized by the method of Helferich & Türk (1956).

Incorporation of [¹⁴C]leucine. Washed bacteria were incubated in buffer as already described, with the addition of DL-[¹⁴C]leucine (200 μ g./ml.; specific activity about 30

counts/min./ $\mu\text{g.}$). Samples (40 ml.) were taken at intervals. The precipitate obtained with trichloroacetic acid was prepared for counting at infinite thickness as described by Mandelstam (1958*b*). Carbon sources, when present, were used at 1% (w/v) unless otherwise stated. The suspensions were shaken in a water bath and samples (5 ml.) taken periodically and centrifuged immediately. This was found to stop the formation of β -galactosidase as effectively as the addition of chloramphenicol. The cells were treated with toluene in sodium phosphate buffer and β -galactosidase activity was measured as described by Rickenberg & Lester (1955), with *o*-nitrophenyl β -D-galactoside as substrate. A unit is defined as the amount of enzyme which liberates 1 μmole of *o*-nitrophenol/hr.

RESULTS

Effect of amino acid- or nitrogen-starvation upon the synthesis of β -galactosidase. Experiments were first carried out to determine to what extent cells starved of nitrogen or of an amino acid were capable of producing β -galactosidase.

Arginine-requiring cells were grown to 0.70 mg./ml. in glucose-synthetic medium, and divided into

Fig. 1. Effect of arginine-starvation upon the subsequent production of β -galactosidase in washed suspensions. Arginine-requiring cells were starved for 30 min. (see the Materials and Methods section), washed and shaken in buffer with MTG. O, Starved cells; ●, unstarved cells supplemented with L-arginine-HCl (100 $\mu\text{g.}/\text{ml.}$).

two equal portions. One was left in the flask at room temperature. The other was centrifuged, washed once with phosphate buffer (50 ml.), suspended in the same volume of medium with arginine omitted and shaken for 30 min. in a water bath at 35°. This period of starvation was previously shown to be sufficient to exhaust arginine and to cause the cells to accumulate and discharge into the medium considerable amounts of amino acids (Mandelstam, 1958*a*). Both the starved and unstarved bacteria were then centrifuged, washed once and suspended in 100 ml. of buffer containing MTG. In addition, the suspension of unstarved cells was supplemented with L-arginine-HCl (100 $\mu\text{g.}/\text{ml.}$).

Production of β -galactosidase in the shaken cultures is shown in Fig. 1. The starved cells formed the enzyme nearly as rapidly as the unstarved cells, the specific activities being 22.5 and 26 units/mg. dry wt. of bacteria at the end of 4 hr. It may be noted for comparison that cultures grown with lactose as carbon source contained about 100 units/mg. whereas cultures grown with succinate and MTG, which is a more efficient inducer, contained about 400 units/mg.

Similar experiments were carried out with wild-type *E. coli* (K₁₂ and ML 30) and these were subjected to starvation for 30 min. by omitting ammonium salts from the medium. Again there was little or no impairment of the ability to produce β -galactosidase subsequently in washed suspensions, and this was also the case with cells which had been starved of leucine (see below), methionine or lysine.

Effect of thymine- or uracil-starvation upon the synthesis of β -galactosidase. The effect of uracil-starvation upon the ability of *E. coli* to form induced enzymes has been the subject of conflicting reports (see Pollock, 1959), and in thymine deficiency it has been reported (Cohen & Barner, 1954, 1955) that xylose-isomerase formation is not impaired. The effect of depriving *E. coli* of these pyrimidines has been examined.

Thymine-less *E. coli* (15_r-) was washed and suspended in glucose-synthetic medium (200 ml.) at 0.36 mg. dry wt./ml. and the suspension divided into two 100 ml. lots. Thymine (final concentration 10 $\mu\text{g.}/\text{ml.}$) was added to one culture, and both were shaken in a water bath at 35°. For the first 60 min. growth in the two flasks was equal, but after this there was a progressive fall in the growth rate of the thymine-starved cells and growth had ceased after 100 min., when the bacterial density was 0.97 mg./ml. The unstarved culture had reached 1.3 mg./ml. at this stage. Portions of both cultures were centrifuged, washed and suspended in buffer (50 ml.) at the same bacterial density (0.35 mg./ml.). Thymine (final concentration 10 $\mu\text{g.}/\text{ml.}$)

ml.) was added to the unstarved cells and MTG to both.

With the uracil-requiring strain (W 3301), washing was sufficient to deplete the cells and no growth occurred after resuspending the cells in glucose medium without uracil. Starvation was therefore continued for only 30 min., during which the culture containing uracil increased from 0.34 to 0.50 mg./ml. while the starved culture remained at 0.35 mg./ml. Both cultures were washed and resuspended at the same density (0.3 mg./ml.) in buffer (50 ml.) containing MTG. The unstarved cells were also given uracil (20 μ g./ml.).

The results with uracil (see Fig. 2) are consistent with those of Pardee (1955), and show that uracil deprivation, under these conditions, does not significantly affect the ability to form β -galactosidase. The effect of longer periods of starvation was not investigated because it was found that starvation for 60 min. seriously damaged the cells. Oxygen uptake with glucose as substrate was decreased by over 50 % (see also Cohen & Barner, 1955), and the cells failed to grow when subsequently transferred to a complete medium.

Thymine deficiency reduced the rate of formation of β -galactosidase by about 30 % (Fig. 2).

Fig. 2. Effect of pyrimidine-starvation upon the subsequent production of β -galactosidase in washed suspensions. Bacteria requiring thymine or uracil were starved (see the Materials and Methods section), washed and shaken in buffer with MTG. O, Starved cells; ●, unstarved cells supplemented with pyrimidine (see text); —, uracil-requiring; - - -, thymine-requiring.

Effect of a carbon source upon the synthesis of β -galactosidase. Leucine-requiring cells [ML 328c (Lac^+)] grown in succinate medium were washed and resuspended at 0.1 mg. dry wt./ml. in buffer containing MTG and Fe^{2+} and Mg^{2+} ions and the following additions: (a) none; (b) NH_4^+ ions; (c) succinate; (d) NH_4^+ ions and succinate; (e) NH_4^+ ions, succinate and leucine (i.e. complete medium). All additions were as in the growth medium. The cultures were shaken and samples taken for measurement of β -galactosidase and extinction.

The cells in complete medium grew almost without lag and the bacterial mass increased almost fourfold (0.1–0.38 mg./ml.) in 2 hr. This should be borne in mind when comparing enzyme production in this culture (see Fig. 3) with that of the four suspensions in which no growth occurred. The graph makes it apparent that omission from the complete medium of leucine alone or of leucine and ammonium salts led to complete repression of β -galactosidase. The further omission of succinate restored the ability to produce the enzyme although the rate was, of course, greatly below that observed in the growing culture.

The finding, that a carbon source which allowed a high rate of enzyme synthesis under normal growth conditions became totally inhibitory as soon as

Fig. 3. Effect of succinate, ammonium salts and leucine upon production of β -galactosidase in leucine-requiring cells. Washed bacteria were shaken in buffer with MTG, Mg^{2+} ions, Fe^{2+} ions and further additions (see text) as follows: O, none; ●, NH_4^+ ions; Δ, succinate; ■, NH_4^+ ions and succinate; □, NH_4^+ ions, succinate and leucine. The last-named is a complete medium, and the cells increased fourfold in 2 hr.; there was no growth in the other suspensions.

growth was prevented, appeared to hold generally for all the strains examined, and for all carbon sources tested (succinate, pyruvate, lactate, gluconate, glycerol, malate, fructose, maltose). The specific cause for the stoppage of growth was immaterial; that is omission of ammonium salts or of a specific amino acid or of a pyrimidine produced total inhibition of β -galactosidase synthesis, which was relieved by removal of the carbon source.

Comparison of β -galactosidase synthesis and the incorporation of DL-[^{14} C]leucine into protein in the presence or the absence of succinate. The possibility had to be examined that the inhibition of β -galactosidase synthesis was non-specific, i.e. in non-growing cells the presence of a carbon source might, for some reason, decrease protein synthesis generally.

For these experiments leucine-requiring cells grown in succinate-synthetic medium were washed and suspended (at 0.5 mg. dry wt./ml.) in buffer (200 ml.) containing MTG and DL-[^{14} C]leucine. A similar flask contained succinate (1%, w/v) in addition. The suspensions were shaken at 35°. Samples (40 ml.) were taken every 40 min. and added to 1 ml. of chloramphenicol solution (1 mg./ml.) to stop further incorporation and enzyme synthesis. A sample (2.0 ml.) was taken for esti-

mation of β -galactosidase and the rest of the suspension used to prepare protein for radioactivity measurement.

From Fig. 4 it is apparent that the formation of β -galactosidase bore little relation to the protein synthesis as measured by incorporation of leucine. The effect of succinate was to increase incorporation while decreasing the synthesis of β -galactosidase. The uptake of radioactivity into protein was similar to that found previously in several strains of *E. coli* and indicated a rate of protein turnover of about 5%/hr. (see Mandelstam, 1958b).

Reversal of repression by malonate or 2:4-dinitrophenol. These results suggested that stopping growth caused the accumulation of metabolites which acted as repressors of β -galactosidase. It therefore seemed likely that suitable inhibitors might reverse the repression caused by carbon sources and that this might help to indicate what sort of metabolites the active repressor was derived from. Most of the experiments were carried out with W 3301 (requiring uracil), mainly because

Fig. 4. Comparison of effect of succinate upon incorporation of DL-[^{14} C]leucine and upon synthesis of β -galactosidase. Leucine-requiring cells were washed and incubated in buffer with MTG and DL-[^{14} C]leucine. ○, With succinate; ●, no succinate. - - -, β -Galactosidase; —, radioactivity incorporated into material insoluble in hot trichloroacetic acid (see the Materials and Methods section).

Fig. 5. Repression of β -galactosidase by succinate and reversal of repression by malonate. Washed cells were incubated in buffer with MTG and various concentrations of malonate, and β -galactosidase formation was measured after 2 hr. ○, No succinate; ●, with succinate (0.25%, w/v).

this strain produced about twice as much enzyme in washed suspensions as most other strains of *E. coli*. However, some of the experiments have been repeated with ML 30 (wild-type) and M 328c (Lac⁺) and similar results have been obtained.

(i) Malonate. It is well known that malonate competes with succinate for succinic dehydrogenase (Quastel & Wooldridge, 1928). Its effect upon the repression produced by succinate was examined as follows. Cells grown in glucose medium were washed and incubated in buffer (10 ml.) with MTG and various concentrations of malonate in the presence or the absence of succinate (0.25 %, w/v). The β -galactosidase formed after 2 hr. was measured. In the absence of succinate, malonate enhanced enzyme production (48–64 units) at a concentration of 0.5 % (w/v) whereas higher concentrations were inhibitory (Fig. 5). In the presence of succinate, and with no malonate, enzyme production was repressed (16 units). Malonate not only overcame the repression but at a concentration of 1.5 % allowed the production of about twice as much enzyme (106 units) as was found in the control suspension without either succinate or malonate (Fig. 5). Concentrations above 1.5 % produced some inhibition.

When these experiments were done with cells originally grown in succinate medium instead of glucose, the repression was not reversed by malonate at any concentration up to 2.5 %.

(ii) 2:4-Dinitrophenol. The bacteria were grown in synthetic medium with the compound to be tested as the sole source of carbon and energy. After washing, the cells were suspended as before in phosphate buffer, and various dilutions of 2:4-dinitrophenol in the same buffer were added. The suspensions (total volume 10 ml.) were induced with MTG as usual by shaking at 35° for 2 hr. and β -galactosidase was then measured. A similar set of flasks contained the carbon compound (final concentration 1 %, w/v).

Fig. 6 (a) shows the enzyme produced in 2 hr. as a function of 2:4-dinitrophenol concentration with and without succinate. In the absence of succinate there was a continuous fall in the rate of enzyme synthesis with increasing amounts of 2:4-dinitrophenol. In the presence of succinate, the repression (about 85 %) produced by succinate in the absence of 2:4-dinitrophenol was largely overcome at 0.125 mM-2:4-dinitrophenol. In these experiments the extent of the repression was somewhat variable but was never less than 80 %, and more often almost complete. Both the extent of reversal achieved and the concentration of 2:4-dinitrophenol producing the greatest effect also varied from one experiment to another. Thus the amount of enzyme synthesized in the presence of the optimum concentration of 2:4-dinitrophenol lay

between 80 and 150 % of that produced in the control suspension without succinate (see Table 1), and the optimum concentration of 2:4-dinitrophenol varied from 0.125 to 0.25 mM.

Fig. 6. (a) Repression of β -galactosidase by succinate and reversal of repression by 2:4-dinitrophenol. Washed cells were incubated in buffer with MTG and various concentrations of 2:4-dinitrophenol, and β -galactosidase formation was measured after 2 hr. $\circ$, No succinate; $\bullet$, with succinate (1 %, w/v). (b) Repression of β -galactosidase by gluconate and partial reversal of repression by 2:4-dinitrophenol. Details were as given for (a). $\circ$, No gluconate; $\bullet$, with gluconate (1 %, w/v).

The repression by succinate and the reversal with 2:4-dinitrophenol also depended upon the previous history of the cells. If the cells had originally been grown in glucose medium, the inhibition subsequently produced in washed suspension by succinate was less (about 50 % instead of 85–100 %) and it was reversed at a lower concentration of 2:4-dinitrophenol.

A number of other carbon compounds were tested in the same way. They could be divided into three groups. (a) Compounds producing repression which was often not complete and which was appreciably reversed by 2:4-dinitrophenol, i.e. 2:4-dinitrophenol restored 50 % or more of the ability to synthesize β -galactosidase. Succinate was typical of this group. (b) Compounds producing complete repression and exhibiting a low degree of reversal with 2:4-dinitrophenol, i.e. less than 10 % restoration of the ability to form the enzyme. Gluconic acid acted in this way (see Fig. 6b). (c) Compounds producing complete repression and not significantly reversible by 2:4-dinitrophenol. Glycerol and glucose are examples of this group.

The compounds tested are listed in Table 1. As might be expected, those compounds which are metabolized aerobically showed a high degree of reversibility with 2:4-dinitrophenol whereas those which can be utilized by anaerobic pathways were not reversible.

Repressor action of lactose. It has been shown by

Table 1. *Repression of β -galactosidase synthesis by carbon sources and the effect of 2:4-dinitrophenol*

Washed cells (W 3301) were incubated for 2 hr. with the compound (1%, w/v) to be tested and MTG (0.1 mg./ml.) as inducer. 2:4-Dinitrophenol was added to give the following final concentrations (mM); 0, 0.125, 0.25, 0.5 and 1.0. The second column gives the concentration of 2:4-dinitrophenol at which the highest enzyme activity was obtained. The last column expresses the highest enzyme value in the series as a percentage of the control value obtained in the absence both of carbon source and of 2:4-dinitrophenol. The control value varied somewhat from one experiment to another but was in the range 40–60 enzyme units/mg. dry wt. of bacteria. 'Irreversible' indicates that enzyme formation was less than 2 % of the control at all concentrations of 2:4-dinitrophenol (DNP).

Compound	Optimum concn. of DNP (mM)	Reversibility (%)
Succinate	0.125–0.25	80–150
Lactate	0.5	50–80
Pyruvate	0.5	90–100
Malate	0.25	90–150
Gluconate	0.25–0.5	5–10
Glycerol	—	Irreversible
Glucose	—	Irreversible
Fructose	—	Irreversible
Maltose	—	Irreversible

Rickenberg, Cohen, Buttin & Monod (1956) that galactosides are concentrated in *E. coli* by an active transport system which they called galactoside permease. The permease is induced in parallel with β -galactosidase and requires energy for its functioning (Kepes, 1960). The possibility had to be considered that the repressors produced their effect not by acting upon the enzyme-forming system but by preventing the entry of the inducer into the cells.

In an attempt to rule this out, the following experiment was done with lactose, which as a galactoside is transported into the cells by galactoside permease, and which as a carbon source should produce a repressor effect. Cells, grown in succinate medium, were induced in washed suspension in the usual way. At the end of an hour, when an appreciable amount of enzyme had been formed, the suspension was divided into three portions and the following additions were made: (a) none; (b) succinate (final concentration 1%, w/v); (c) lactose (final concentration 1%, w/v). Enzyme production was followed for the next 2 hr. The results (see Fig. 7) showed that in partially induced cells lactose could act as an efficient repressor in much the same way as succinate. The fact that it had this effect indicates that it was entering the cell and that the galactoside permease was therefore functioning.

Fig. 7. Repression of β -galactosidase by succinate and by lactose. Washed cells were incubated in buffer with MTG. After 1 hr. the suspension was divided into three and additions were made as follows: O, none; ●, succinate (1%); Δ, lactose (1%).

Fig. 8. Induction of β -galactosidase by lactose and MTG. Washed cells were incubated with phosphate buffer with additions as follows: $\circ$, lactose; $\bullet$, MTG; $\blacktriangle$, MTG and lactose (1%).

If, however, lactose was added at the beginning of the incubation period together with MTG, the repressor effect was observed only after some enzyme had been formed, so that the lactose could be metabolized.

In Fig. 8 a comparison has been made of the induction of β -galactosidase by lactose and by MTG separately and together. It can be seen that lactose first acts as an inducer which is about as efficient as MTG, and then when sufficient enzyme has been formed, the repressor effect becomes manifest not only in the cells treated with lactose but in those treated with both MTG and lactose. These results make it reasonable to conclude that the repressor action of lactose is an effect upon the enzyme-forming system and imply that other carbon sources also act in this way and not upon the galactoside permease.

DISCUSSION

It was shown by Monod, Pappenheimer & Cohen-Bazire (1952) that in a growing culture of *E. coli* the amount of β -galactosidase formed was proportional to the net protein synthesis, i.e. the enzyme was a constant fraction of the total new protein being synthesized by the cells. In the present experiments, when the cells were starved of nitrogen or of an amino acid, there was no net synthesis of protein but there was nevertheless considerable production of β -galactosidase. The

amount of enzyme formed in 1 hr. in the non-growing condition is about 10–30 units/mg. dry wt., depending upon the strain of *E. coli*. This rate of enzyme production should be considered in relation to the actual protein synthesis. It has been shown (Mandelstam, 1957, 1958b) that in non-growing *E. coli* protein is degraded and resynthesized at about 5%/hr. Thus for every 1% of protein 'turned over' in the presence of MTG, the bacteria acquire a specific enzyme activity of 2–6 units/mg. If this relationship could be maintained constant until all the protein had been degraded and resynthesized the activity of the cells would be 200–600 units/mg. This is the same order of activity as that of cells induced under conditions of normal growth. For instance, after growth for ten generations in lactose the activity is about 100 units/mg. and after growth in the presence of succinate and MTG it is about 400 units/mg. It is therefore apparent that β -galactosidase forms a more or less constant fraction of the total protein synthesized whether the cells are growing or not, provided, of course, that, in the latter condition, no carbon source is present. The kinetics of formation of β -galactosidase and its relationship to turnover during the period when cells are adapting to growth on a new carbon source have been discussed by Mandelstam (1960).

The inhibitory effect of a carbon source on the synthesis of β -galactosidase in a pyrimidine-less strain of *E. coli* had been noted by Pardee (1955). The present results show that the effect is general and that a carbon source becomes inhibitory as soon as growth is prevented by omitting any essential ingredient from the medium. This explains why Monod *et al.* (1952) obtained no β -galactosidase in *E. coli* auxotrophs deprived of their specific requirements.

Although β -galactosidase formation is inhibited by a carbon source, the synthesis of protein as measured by the incorporation of [^{14} C]leucine is, if anything, slightly enhanced. It follows that it may be misleading to use formation of β -galactosidase as an indication of the general ability of the cells to synthesize protein.

The results also show that the 'glucose effect' is not specific to glucose but can be produced by a variety of carbon sources, provided that growth of the cells is prevented. The fact that the repressor effect of these substances is manifest only under such conditions suggests that when growth is blocked, metabolites accumulate, one (perhaps more) of which functions as a repressor of the β -galactosidase-induction system. These conclusions support the suggestion made by Neidhardt & Magasanik (1956, 1957) that glucose acted as a repressor because it was more rapidly metabolized than other carbon sources, so that, even under

growing conditions, intermediary metabolites accumulated which repressed the inducible enzymes concerned with the utilization of alternative carbon sources.

From Table 1 it is apparent that the repression produced by substances which are metabolized entirely aerobically (e.g. succinate) is reversed by DNP, whereas that produced by compounds like glucose which can also be metabolized anaerobically is not reversed. This could mean that the repressor is a product, e.g. adenosine triphosphate, common both to aerobic and anaerobic metabolism, but there is no direct evidence to support this idea.

The suggestion has been made, on the basis of genetic experiments (Pardee, Jacob & Monod, 1958), that in wild-type strains of *E. coli* β -galactosidase is inducible because the cytoplasm of the cells contains a specific repressor, whose effect upon the β -galactosidase system has to be reversed by adding an inducer if enzyme synthesis is to be obtained. In constitutive strains this repressor is assumed to be absent.

The present experiments pose the question whether the condition of blocked growth leads to the formation, from the carbon source, of more of a repressor which is assumed to be present in any case, or whether it leads to the accumulation of some other substance with the properties of a repressor. Again there is no evidence that will provide a definite answer, but the fact that repressor can apparently be produced from all carbon sources and by aerobic and anaerobic metabolism suggests that it is not as specific a substance as the repressor postulated by Pardee *et al.* (1958).

SUMMARY

1. Various strains of *Escherichia coli* deprived of ammonium salts, or of an essential amino acid or pyrimidine, are nevertheless able to produce considerable amounts of induced β -galactosidase in washed suspension.

2. Under non-growing conditions, compounds which the cells can use as sources of carbon and energy repress enzyme formation. The repressors appear to act upon the enzyme-forming system and not upon the galactoside permease.

3. Incorporation of [^{14}C]leucine into protein is unaffected under conditions which prevent β -

galactosidase synthesis. Formation of this enzyme thus appears to be unreliable as a guide to the protein-forming ability of the cells.

4. The repression produced by compounds which are metabolized aerobically is reversed by 2:4-dinitrophenol; that produced by compounds metabolized anaerobically is not reversible.

5. The repressor of β -galactosidase appears to be a substance common to aerobic and anaerobic metabolism.

6. Consideration of the kinetics of enzyme synthesis shows that, in the absence of a carbon source, β -galactosidase constitutes about the same fraction of the total protein being synthesized as it does in the growing culture.

Auxotrophic mutants of *E. coli* were kindly supplied by Dr T. Alper, Dr B. D. Davis and Dr S. Spiegelman. I am much indebted to Miss R. Coyle for technical assistance.

REFERENCES

- Cohen, S. S. & Barner, H. (1954). *Proc. nat. Acad. Sci., Wash.*, **40**, 885.
- Cohen, S. S. & Barner, H. (1955). *J. Bact.* **69**, 59.
- Cohn, M. & Horibata, K. (1959). *J. Bact.* **78**, 601, 613, 624.
- Helferich, B. & Türk, D. (1956). *Chem. Ber.* **89**, 2215.
- Kepes, A. (1960). *Biochim. biophys. Acta*, **40**, 70.
- Mandelstam, J. (1956). *Int. Rev. Cytol.* **5**, 51.
- Mandelstam, J. (1957). *Nature, Lond.*, **179**, 1179.
- Mandelstam, J. (1958a). *Biochem. J.* **69**, 103.
- Mandelstam, J. (1958b). *Biochem. J.* **69**, 110.
- Mandelstam, J. (1960). *Bact. Rev.* **24**, 289.
- Monod, J. & Cohn, M. (1952). *Advanc. Enzymol.* **13**, 67.
- Monod, J., Pappenheimer, A. M. & Cohen-Bazire, G. (1952). *Biochim. biophys. Acta*, **9**, 648.
- Neidhardt, F. C. & Magasanik, B. (1956). *Nature, Lond.*, **178**, 801.
- Neidhardt, F. C. & Magasanik, B. (1957). *J. Bact.* **73**, 253.
- Pardee, A. B. (1955). *J. Bact.* **69**, 233.
- Pardee, A. B., Jacob, F. & Monod, J. (1958). *C.R. Acad. Sci., Paris*, **246**, 3125.
- Pollock, M. R. (1959). In *The Enzymes*, 2nd ed. p. 619. Ed. by Boyer, P. D., Lardy, H. & Myrback, K. New York: Academic Press Inc.
- Quastel, J. H. & Wooldridge, W. R. (1928). *Biochem. J.* **22**, 689.
- Rickenberg, H. V., Cohen, G. N., Buttin, G. & Monod, J. (1956). *Ann. Inst. Pasteur*, **91**, 829.
- Rickenberg, H. V. & Lester, G. (1955). *J. gen. Microbiol.* **13**, 279.
- Vogel, H. J. (1957). *Proc. nat. Acad. Sci., Wash.*, **43**, 491.