of normal male rats in this way, and Fulco & Mead (1959) have shown how eicosa-5:8:11-trienoic acid arises in fat-deficient rats.

SUMMARY

1. The liver vitamin A stores of fat-deficient male rats fed on a diet containing cholesterol (1%) did not differ significantly from those of controls which received linoleic acid.

2. The liver vitamin A stores of fat-deficient female rats fed on a 1% cholesterol diet were 48% lower than those of their controls which had been given linoleic acid.

3. The addition of cholesterol to a fat-free diet aggravated fat deficiency as judged by more rapid cessation of growth at a lower body weight. Both effects were appreciably more pronounced in males than in females.

4. Cholesterol-supplementation of the diet caused much larger accumulations of cholesteryl esters in livers than are caused by fat deficiency alone. The effect was much greater in males. Linoleic acid (100 mg./day) did not alter the severity of this phenomenon in males; a smaller dose (25 mg./day) reduced the effect considerably in females.

5. The rise in liver triglyceride concentration caused by fat deficiency was not altered by adding cholesterol to the diet. Cholesterol caused an increase in the concentrations of polyunsaturated acids in triglycerides, both in deficient animals and their controls. This effect was more pronounced in males.

6. Dietary cholesterol decreased the concentrations of dienoic and tetraenoic acids, particularly the latter, in the liver phospholipids of control animals. Both effects were appreciably greater in males than in females.

7. Dietary cholesterol consistently raised the triene:tetraene ratio of liver phospholipids.

8. Dietary cholesterol invariably raised the diene:tetraene ratio in total liver lipids, phospholipids and triglycerides, and lowered this value in the triglycerides. These effects occurred both in deficient animals and in controls.

REFERENCES

- Bromer, W. W. (1953). Ph.D. Thesis, Part II: Indiana University. University Microfilms Doctoral Dissertation Series 7429, University of Michigan, Ann Arbor.
- Dittmer, J. C. & Hanahan, D. J. (1959). *J. biol. Chem.* **234**, 1983.
- Fulco, A. J. & Mead, J. F. (1959). *J. biol. Chem.* **234**, 1411.
- Green, B., Horner, A. A., Lowe, J. S. & Morton, R. A. (1957). *Biochem. J.* **67**, 235.
- Greenberg, S. M., Calbert, C. E., Savage, E. E. & Deuel, H. J., jun. (1950). *J. Nutr.* **41**, 473.
- Holman, R. T. (1957). *Meth. biochem. Anal.* **4**, 112.
- Holman, R. T. (1960). *J. Nutr.* **70**, 405.
- Holman, R. T. & Burr, G. O. (1948). *Arch. Biochem.* **19**, 474.
- Holman, R. T. & Peifer, J. J. (1960). *J. Nutr.* **70**, 411.
- Horner, A. A. (1958). Ph.D. Thesis: University of Liverpool.
- Horner, A. A. & Morton, R. A. (1960). *Biochem. J.* **74**, 301.
- Klein, P. D. (1959). *Arch. Biochem. Biophys.* **81**, 382.
- Lynn, W. S. & Brown, R. H. (1959). *Arch. Biochem. Biophys.* **81**, 353.
- Mead, J. F., Slaton, W. H., jun. & Decker, A. B. (1956). *J. biol. Chem.* **218**, 401.
- Morton, R. A. & Horner, A. A. (1961). *Biochem. J.* **79**, 631.
- Okey, R. & Lyman, M. M. (1957). *J. Nutr.* **61**, 523.
- Patil, V. S. & Magar, N. G. (1960). *Biochem. J.* **74**, 441.
- Peifer, J. J. & Holman, R. T. (1955). *Arch. Biochem. Biophys.* **57**, 520.
- Peifer, J. J., Janssen, F., Ahn, P., Cox, W. & Lundberg, W. O. (1960). *Arch. Biochem. Biophys.* **86**, 302.
- Popják, G. (1946). *Biochem. J.* **40**, 608.
- Rieckehoff, I. G., Holman, R. T. & Burr, G. O. (1949). *Arch. Biochem.* **20**, 331.

Biochem. J. (1961) **79**, 642

Ester-Linked Long-Chain Fatty Acids of Nervous Tissue

By R. W. R. BAKER

Department of Chemical Pathology, Guy's Hospital Medical School, London, S.E. 1

(Received 26 September 1960)

It has long been recognized that nervous tissue contains a high proportion of lipid which on hydrolysis gives long-chain fatty acids in substantial amounts (Thudichum, 1901). Extracts of this lipid comprise a number of compounds that are now well characterized; the quantities in which these are present, and their chemical structures, have in many cases been elucidated (see LeBaron &

Folch, 1957). Although the results of early qualitative work and of later quantitative studies on the acids from isolated fractions of different lipids extracted from nervous tissue have appeared in the literature, no survey of the nature and distribution of these acids in total lipid extracts of nervous tissues had been reported before the account by Johnston & Kummerow (1960) of the acids in two

samples of whole-brain lipid. Much of the acid is combined in ester linkage, as in the glyceryl-phosphatidyl compounds, the balance occurring largely in the form of amides, as in sphingomyelin. Small amounts of apparently free acid (Laser, 1949) were isolated from brain and characterized as octadec-11-enoic acid by Morton & Todd (1950).

Lecithins and cephalins from many sources contain C_{16} and C_{18} acyl residues, extending in the even-numbered series to C_{24} and even C_{26} (Klenk & Montag, 1958*a*, *b*). Acyl groups in the plasmalogens of ox-heart lipid are of similar chain length (Klenk & Krickau, 1957). Lipids nearly always contain unsaturated acids, often with more than one double bond in the molecule. Oleic acid and arachidonic acid were the only unsaturated acids that Levene & Rolf (1922*a*, *b*) could isolate from lecithin and cephalin of brain; they were unable to detect the corresponding dienoid acids. Arachidonic acid is an important component of lecithins from tissues other than the nervous system; with suprarenal gland or liver as the source, it accounts for over a fifth of the acids obtained by hydrolysis of the lecithin (Ault & Brown, 1940). Klenk & Böhm (1951) showed that hydrolysis of brain phospholipids yielded also C_{20} and C_{22} acids with up to four double bonds in the shorter-chain and six in the longer-chain acids. Further work in Klenk's laboratory clarified the structure of still more of the acids from brain phospholipid (Klenk & Bongard, 1952; Klenk & Lindlar, 1955*a*, *b*).

Fatty acids and their α -hydroxy derivatives occur as amides in the phospholipid fraction of nervous tissue in sphingomyelin, and in the cerebrosides, sulpholipids and ganglioside fractions of brain (Deuel, 1951). Acids liberated by hydrolysis of these are generally of higher molecular weight than those from the esters, and chains of 24 and 26 carbon atoms were found in the acids from cerebrosides (Klenk, 1928).

Knowledge of the acids incorporated into lipids of the nervous system has been limited in the past by the difficulties of their quantitative separation. Although countercurrent partition and paper chromatography gave new impetus, it was not until the methods of vapour-phase chromatography had been developed that semi-microanalytical work could be undertaken without recourse to laborious and imperfect quantitative techniques. In the work described below, a preliminary survey of the nature and quantities of ester-linked acids in some different regions of the human nervous system was undertaken, without attempting to relate the acids to chemically distinct lipid fractions. Striking differences between the acids in sciatic nerve and those in spinal cord and brain have been found. Within the brain, white and grey regions also showed points of contrast.

MATERIALS AND METHODS

Normal unfixed human brain and other nervous tissue from adults of either sex and from 38 to 81 years old was obtained at post-mortem examinations. Fatty acids for the preparation of methyl esters as marker materials were of laboratory-reagent grade (British Drug Houses Ltd. or L. Light and Co. Ltd.). Solvents and other substances were of A.R. quality wherever possible.

Extraction of lipids from brain and spinal cord was carried out by the method of Folch, Lees & Sloane-Stanley (1957); the weighed sample, chopped with scissors, was homogenized in 19 ml. of chloroform-methanol 2:1 (v/v) mixture/g. and the suspension filtered. The filtrate was washed with 0.2 vol. of water, the phases were separated by centrifuging and the upper (aqueous) layer was rejected. Solvent was removed from the remainder in a rotary evaporator at 50° under reduced pressure, and residual lipid was dried to constant weight over $CaCl_2$ and paraffin wax in an evacuated desiccator. Sciatic nerve was freeze-dried and extracted with chloroform-methanol mixture, followed by washing etc. as above.

The fatty esters were hydrolysed by homogenizing the lipid (about 100 mg.) in 22.5 ml. of ethanol, mixing with 2.5 ml. of aq. 10*N*-KOH and agitating for 30 min. in a water bath at 38° by passing a stream of nitrogen through the suspension.

To separate the fatty acids formed, the hydrolysates were diluted with 30 ml. of water and acidified (Congo-red paper) with approx. 20 ml. of 2*N*-HCl; the organic acids were extracted by shaking gently first with $\frac{1}{2}$ vol. and then with $\frac{1}{4}$ vol. of light petroleum (b.p. 40–60°). The combined light-petroleum solutions were washed with 0.5 vol. of 0.1*N*-HCl, separated, and extracted with 0.5 vol. of aq. 0.05*N*-NaOH by gentle inversion. The light-petroleum layer was stratified centrifugally and rejected, and the aqueous solution acidified (Congo-red paper) with an equal volume of 0.1*N*-HCl and twice extracted as before with light petroleum. The extracts were combined, washed with 0.5 vol. of 0.1*N*-HCl, separated, dried over Na_2SO_4 , filtered, and evaporated under nitrogen; the acids remaining were dried *in vacuo* over $CaCl_2$ and paraffin wax to constant weight.

The mixtures of acids so obtained were converted into methyl esters by interaction in diethyl ether with diazomethane; this was prepared (De Boer, 1954) in a mixture of ether (2 ml.) and ethanol (3 ml.) in Quickfit B10 apparatus from 200 mg. of *N*-methyl-*N*-nitrosotoluene-*p*-sulphonamide (Diazald, Aldrich Chemical Co.) by reaction with 1 ml. of aq. 40% (w/v) KOH. The diazomethane was distilled with the ether (James & Webb, 1957*a*) into a solution of about 100 mg. of fatty acid in 1.5 ml. of ether. A trap consisting of a drying tube loosely packed with paper tissue damped with crude oleic acid absorbed escaping excess of diazomethane. Centrifuge tubes containing the reaction mixtures were placed in 150 ml. of ice-cold water in a fume chamber, until they had reached room temperature, when the ether was evaporated under nitrogen. Methyl esters required as reference substances in gas chromatography were prepared on a larger scale by leaving overnight at room temperature a solution of the acid in 25 vol. of methanol containing 2% (v/v) of H_2SO_4 . Most of the methanol was distilled off, the residue was diluted with 5 vol. of water, and the ester was extracted with light petroleum. The extract was dried over Na_2SO_4 , the solvent was

evaporated off and the esters were distilled under reduced pressure (approx. 0.4 mm. Hg) from an oil bath. Final purification was achieved by gas chromatography on a preparative scale with a column 2.5 cm. in diameter and 50 cm. long, packed with Apiezon L-coated Celite. With this column a full-flow catharometer detector designed for the purpose with directly heated Thermistor beads (Stantel Type P15, Standard Telephones and Cables Ltd., Footscray, Kent) was used. Currents in detector and reference Thermistors (the latter in a stream of pure carrier gas at the temperature of the column) were controlled at the same value by transistor stabilizers, and a balanced transistor amplifier was made to compare the potentials across the Thermistor elements. The carrier gas was nitrogen. Samples were collected by passing the effluent stream through drying tubes loosely packed with cotton wool washed with diethyl ether and moistened with methanol. Tubes were changed as indicated by the detecting system to collect predetermined peaks. The cotton wool was then extracted with diethyl ether, which was finally evaporated to leave pure ester. Methyl esters of lauric, myristic, palmitic, stearic, oleic, linoleic, arachidonic and clupanodonic acids were prepared. In the treatment of unsaturated acids, temperatures were not allowed to exceed 50° except under nitrogen or *in vacuo*.

Analytical methods. A colorimetric method (Stern & Shapiro, 1953) was used to estimate acyl-ester groups in the hydrolysis experiments described below, with tristearin as the standard.

Vapour-phase chromatography (see James, 1960; Rose 1959; Farquhar, Insull, Rosen, Stoffel & Ahrens, 1959) in packed columns 130 cm. high and of bore 3–4 mm. was applied for analysis of 0.02–1.0 μ l. samples of mixtures of methyl esters. The apparatus was essentially as described by James, Lovelock & Piper (1959), with argon as the carrier gas and ^{90}Sr as the radioactive source in an ionization detector (Lovelock, 1958) in which a modified sparking plug was used as collector electrode. Lovelock's amplifier and high-tension supply unit were used in conjunction with a recording milliammeter. Chromatograms were also recorded with a simple battery-operated circuit in which current was limited by a resistance of $5 \times 10^9 \Omega$; signals were derived from a load resistor of $10^7 \Omega$ in series with the detector cell, battery and limiter, and were fed directly to a high-impedance recording millivoltmeter (Kent, Multelec Mark 3).

Commercial Celite 545 was sieved into fractions passing

85–100 or 100–120 mesh screens, calcined at 300° for 3 hr., washed with HCl, water and 1% KOH in methanol for use as support for the stationary phase. This was Apiezon L hydrocarbon grease, Reoplex 400 (Stoffel, Insull & Ahrens, 1958), or a polyester prepared in the laboratory, e.g. polydiglycol succinate and adipate (Lipski, Landowne & Godet, 1959). Working temperatures were in the range 165–205°, with gas-flow rates of 25–55 ml./min.

The identities of eluted esters were provisionally established from retention-volume data obtained with different stationary phases, with pure methyl palmitate or stearate as reference substance. Confirmation was obtained by direct comparison with the methyl esters prepared as above. Relative quantities in the mixtures of esters examined were assessed by planimetry of the chromatograms. Linearity of response within the limits of load and polarizing potential used was within 5% when tested with mixtures of palmitic and stearic esters of known compositions.

RESULTS

The yields of total lipid extracted from whole brain and from the seven regions of the nervous system are shown in Table 1. Considerable variation in content of fat from one region to another is evident. In the same Table are recorded the quantities of fatty acid obtained after hydrolysis of the esters.

The efficiency of the procedure chosen for hydrolysis (see Dawson, 1954) was investigated and confirmed in experiments in which the decrease in ester equivalent of the biologically derived and of other fatty esters was followed during the course of reaction, with results indicated in Table 2. The acyl-ester content of the reaction mixture was estimated by applying the hydroxamic acid method to acidified samples. It was necessary to estimate the ester as rapidly as possible after acidifying, since in the acid solution re-esterification occurred at a steady rate of 2.6%/hr. at room temperature. At 0°, no measurable change occurred in 2 hr. Notable lability of the acyl groups and uniformity of the speeds of hydrolysis were revealed. The standard deviation in replicate determinations of

Table 1. *Yields of total lipid extracted from nervous tissues and of fatty acids obtained by mild alkaline hydrolysis of the lipid*

S.D. = Standard deviation (no. of samples in parentheses).

		Brain						Spinal cord (thoracic)	Sciatic nerve
		Whole	White (corpus callosum)	Grey					
				Cortex	Thalamus	Caudate nucleus	Putamen		
Lipid content (g./100 g. of fresh tissue)	Mean value	10.2	21.4	8.3	16.6	12.3	9.6	19.9	24.5
	S.D. (6)	1.5	3.4	1.9	1.1	4.0	2.8	6.4	8.1
Fatty acid content (g./100 g. of fresh tissue)	Mean value	2.32	2.04	1.64	2.85	2.46	1.74	3.59	8.07
	S.D. (6)	0.27	0.62	0.56	0.59	0.65	0.23	2.24	2.76

acyl ester was 4% with tristearin and 6% with a typical sample of hydrolysis mixture. Slightly higher extinctions (e.g. 0.209, compared with 0.207) were consistently given by tristearin added to hydrolysis mixtures, and observations were corrected in this ratio.

To check the method described above for extracting the acids formed by hydrolysis, diluted and acidified hydrolysis reaction mixtures were shaken serially with $\frac{1}{2}$ vol. quantities of light petroleum, and each extract was washed with 0.1N-HCl and evaporated; the weights of the fatty acid residues were determined. Of the total acid obtained from three serial extractions, 92.8% (w/w) was present in the first, 5.6% in the second and 1.6% in the third extract.

A classification of the fatty acids obtained from nervous tissues and the relative quantities in which they were present are summarized in Table 3. Lower acids (C_8 - C_{12}) were found only in

traces in all the regions of brain studied, but were quite prominent in spinal cord and still more so in sciatic nerve. Myristic acid occurred in all the samples, the quantities in corpus callosum and in cord being 1.6% of total acid; sciatic nerve contained over 4%. An insignificantly small proportion of these lower acids was unsaturated. From 15 to 30% of the acids were of the C_{16} straight-chain type in all the lipids analysed, palmitic acid preponderating. Palmitoleic acid accounted for an eighth or less of this group except in the nerve (where it constituted nearly a third) and in spinal cord.

The C_{18} acids comprised approximately one- to two-thirds of the total in all cases, but stearic acid itself was unevenly distributed, its amounts ranging from 5% in sciatic nerve to 23% in whole brain. The largest class seen in every region was that of the octadecenoic acids; these were not resolved into position isomers and are referred to simply as oleic

Table 2. *Time-courses of hydrolysis of lipids*

The lipid (0.5-1.2 g.) was treated with 25 ml. of ethanolic 90% (v/v) N-KOH at 38°. The results are expressed as percentages of acyl ester unhydrolysed (value at zero time = 100%).

Time (min.) ...	2.5	5	10	15	20	25	30
Whole-brain lipid	48	27	12	6.0	5.0	4.0	4.0
White-matter lipid	42	17	5.0	2.0	4.0	3.0	3.0
Grey-matter lipid	40	20	6.0	4.0	3.0	3.0	3.0
Spinal-cord lipid	35	17	6.0	5.0	4.0	3.0	3.0
Sciatic-nerve lipid	57	25	10	7.0	5.0	4.0	4.0
Human-liver lipid	37	15	9.0	8.0	7.0	6.0	6.0
Triolein	64	24	12	8.0	6.0	—	—
Tristearin	42	13	3.0	1.0	1.0	—	—
Cholesterol oleate	42	23	19	2.0	2.0	—	—

Table 3. *Mean values, as percentage of total acid, for ester-bound acids in total lipid from nervous tissue*

+, < 0.05; ++, < 0.50, > 0.05.

Fatty acid	No. of carbon atoms	No. of double bonds	Brain						Spinal cord (thoracic)	Peripheral nerve (sciatic)
			Whole	Corpus callosum	Cortex	Caudate nucleus	Thalamus	Putamen		
Caprylic	8	0	+	+	+	+	+	+	+	+
Capric	10	0	+	+	+	+	+	+	0.1	1.7
Lauric	12	0	+	0.1	+	+	+	+	0.2	1.0
Myristic	14	0	0.1	1.6	0.7	+	0.2	+	1.6	4.2
Palmitic	16	0	24.0	15.0	28.7	22.5	20.5	25.3	14.3	19.2
Palmitoleic	16	1	3.2	++	++	++	++	++	3.9	8.3
	16	2	++	++	++	++	++	++	0.6	1.5
Stearic	18	0	23.0	16.2	15.5	21.0	19.1	20.7	7.7	5.0
Oleic	18	1	32.6	39.4	22.0	25.1	33.9	30.4	51.6	49.7
Linoleic,	18	2, 3	++	3.5	3.1	5.0	3.7	0.1	1.1	8.6
linolenic	20	1-3	3.9	10.7	5.9	13.7	6.7	5.8	6.6	0.7
	20	4-6	6.5	9.2	6.0		8.5	9.6	7.4	0.2
	22	1-4	3.6	1.1	6.4	8.9	1.4	0.7	4.0	0.2
	22	5, 6	2.0	3.2	11.8		3.1	4.3	1.0	0.2
	24	1-5	++	++	++	3.9	3.3	2.0	0.2	0
Total			98.9	100.0	100.1	100.1	100.4	98.9	100.3	100.5

acid. It formed about a third of the brain acids and half of those from the cord and nerve. The C_{18} group included *n*-octadecadienoic and traces of the trienoic acids.

With increasing chain length, unsaturation increased. Only traces of arachidic acid and behenic acid were detected, the latter only in a few brain samples; yet the C_{20} class accounted for 10–20 %, the C_{22} class for 4–18 % and the C_{24} class for up to 4 % of the esterified fatty acids of brain and cord lipids. Arachidonic acid was the chief representative of the C_{20} group, constituting up to 8 % of the total acid. Among the C_{22} acids, members with up to three double bonds were scarce, but the more unsaturated acids formed about 6 % of the whole, with clupanodonic and docosahexenoic acids as the major representative. The C_{24} acids were lower in proportion and polyenoid in character.

Although spinal cord and brain showed substantial and similar quantities of the C_{20} acids, peripheral nerve yielded less than 1 %. The C_{22} acids formed 18 % of the yield from cortex, 9 % of that from the caudate nucleus, but only 5 % in other samples, except sciatic nerve, in which only traces were seen. Although C_{24} acids formed 4 % of the acid in the caudate nucleus and thalamus, they occurred to a less extent in other regions of the brain, were barely detectable in spinal cord, and were not seen at all among the acids derived from sciatic nerve. C_{15} , C_{17} and C_{19} acids were found as minor components in most of the samples.

DISCUSSION

The yields of lipid found for whole brain and for corpus callosum are close to the values reported by Folch & LeBaron (1957) for whole brain and white matter, whereas cortex showed a higher content than that quoted by these authors. Their range for the lipid content of peripheral nerve (16.7–21.5 g./100 g.) is also lower than the mean (24.5 g./100 g.) recorded above for human sciatic nerve, which is difficult to separate from connective tissue, as discussed below.

The comparatively mild conditions used for hydrolysis, for extraction of acids and for the formation of their methyl esters were shown to be quantitative, and do not expose the labile polyenoid acids to serious risk of change. Neither was such change found when substances of similar character were submitted to gas chromatography under the conditions obtaining in the present work (James & Webb, 1957*b*; Stoffel *et al.* 1958). It is considered, therefore, that the acids found are true constituents of the original tissue. The small quantities of residual and apparently stable acyl ester appearing after 30 min. hydrolysis (see Table 2) may be, at least in part, artifacts, for the

quantities involved are comparable with the error of the method.

Although essentially quantitative recovery of the acids formed by hydrolysis was demonstrated, the product was still accompanied by impurities, e.g. cholesterol. Esters made by methylation of the acids from brain and spinal cord, but not from sciatic nerve, were yellow solids with melting points sometimes as high as 60°. After distillation, the ester mixtures were almost colourless and melted below 30°. The weights of acid recorded in Table 1 include the weights of contaminants, and are about 16 % in excess of the theoretical values calculated from the ester equivalents of the lipids and the mean molecular weights for the acid mixtures, assessed from their compositions. Nevertheless, crude and distilled esters gave identical gas chromatograms when somewhat larger samples of the crude ester were taken. The contaminants therefore did not emerge from the columns; this is consistent with the view that the impurities are polar substances of high molecular weight.

Evaluation of the purity of the acids extracted is difficult. No reliance can be put on titrimetric or gravimetric methods, since the absence of acidic groups from the impurities cannot be assumed. Similarly determination of ester equivalent after methylation is not admissible. Molecular distillation could be used only with caution, because highly unsaturated acids easily form resins when heated. Comparison of peak areas on chromatograms with those given by pure substances is not accurate because of the difficulty of applying samples of the necessary size reproducibly to the column.

Johnston & Kummerow (1960) treated whole-brain lipid by prolonged acid methanolysis, and obtained values for the fatty acids combined both in amide and ester linkage. There is general agreement between their values and those shown above for whole brain, but comparison in detail is precluded by the inclusion in their values of the amide acids, known to be of longer average chain-length.

The results in Table 3 reveal a sharp difference in several respects of peripheral nerve from the other nervous tissue. Peripheral nerve contains appreciable amounts of myristic and of lower acids, which occur to a minor extent at the other sites. Palmitoleic acid is a prominent component in the nerve, but is relatively insignificant in brain. Equally sharp in contradistinction from brain is the low quantity of stearic acid in sciatic nerve, offset by a greater abundance of oleic acid and of linoleic acid. The paucity of higher acids is another notable characteristic of the nerve lipid. The results for sciatic nerve resemble those for adipose tissue (Table 4), which also is clearly different in its acid composition from the lipids of brain and spinal

Table 4. *Percentage composition of esterified fatty acids from human adipose tissue and sciatic nerve*

Fatty acid	Adipose tissue		Sciatic nerve
	(Cramer & Brown, 1943)	(Moore & Cook, 1959)	
Lauric	1	1-2	2-5
Myristic	2-6	3-5	3-7
Myristoleic	1	—	0-5
Palmitic	25	22-31	13-26
Palmitoleic	5-8	5-14	4-11
Stearic	5-9	2-15	2-9
Oleic	45-47	42-51	43-55
Linoleic	10-11	1-6	8-10
Total of means	99.5	100	96

cord. This supports the hypothesis that much of the lipid extracted from the peripheral nerve derives from associated connective tissue, and is not properly of nervous origin.

Spinal cord approximates to the nerve in containing lower acids, although in far smaller quantities. Its poverty in palmitic acid gives it a place with corpus callosum, but in its contents of palmitoleic acid, stearic acid and the highly unsaturated C_{22} acids it is intermediate between the nerve and brain. The C_{20} acids in spinal cord do not differ significantly from those in brain.

The cerebral cortex differs from corpus callosum in containing nearly twice as much palmitic acid but only half as much oleic acid. There is also a differing distribution of the longer-chain acids, cortex showing a smaller total of C_{20} but a relative excess of C_{22} acids.

The three nuclear areas taken seem not to differ markedly among themselves in acid contents; they resemble the grey rather than the white portion of brain, reflecting the anatomical relationships. A common finding for the brain fractions was the low quantity of palmitoleic acid and the proportion, approximately 25 %, of C_{20} - C_{24} unsaturated acids.

The high proportion of sphingomyelin in nerve lipid (Carter, Haines, Ledyard & Norris, 1947; Johnson, McNabb & Rossiter, 1948), and the predominance of higher (C_{20} - C_{24}) acids in this compound (Thannhauser & Boncoddio, 1948) give reason for believing that only ester-bound acids were assayed in the present work, since these acids were barely detected in lipid from sciatic nerve. Such higher acids found in the brain and spinal cord thus appear to be truly of ester origin. Further, fatty acid amide linkages are known to be stable to alkali at 37°, and the conditions described for the hydrolysis of amide linkages are very different from those used here for the fission of ester groups. Prolonged heating in strongly acidic methanol was employed by Klenk (1927) and by Thannhauser & Boncoddio (1948). It is therefore

probable that no significant splitting of amides occurs in the gentle hydrolysis of the esters described above.

SUMMARY

1. The esterified fatty acids in lipids extracted from normal human whole brain, from five regions of brain and from spinal cord and sciatic nerve were liberated by hydrolysis and examined as methyl esters by gas chromatography. Yields of lipid and of fatty acid were measured, and the composition of the fatty acid fraction was determined.

2. The fatty acid pattern of sciatic nerve was distinct from that of the other samples, and resembled that of body fat.

3. The compositions of fatty acid mixtures from brain varied little from site to site. The approximate values were palmitic acid 20 %, stearic acid 20 %, C_{20} acids 16 %, C_{22} acids 5 % and C_{24} acids 4 %, the remaining 35 % being mostly oleic acid. Lower acids (up to C_{14}) were barely detectable.

4. Fatty acid values for spinal-cord lipid lay between those of brain and of the nerve.

For most helpful discussions on the gas-chromatographic method the author is greatly indebted to Dr A. T. James, Dr J. E. Lovelock and Miss J. Webb. For the suggestion that this method should be applied to problems in neurochemistry the author thanks Professor R. H. S. Thompson. The Multiple Sclerosis Society gave a grant with which apparatus used in this work was constructed in the laboratory. Miss Pamela Fowler assisted throughout in the chemical and analytical work.

REFERENCES

- Ault, W. C. & Brown, J. B. (1940). *J. biol. Chem.* **134**, 331.
 Carter, H. E., Haines, W. J., Ledyard, W. E. & Norris, W. P. (1947). *J. biol. Chem.* **169**, 77.
 Cramer, D. L. & Brown, J. B. (1943). *J. biol. Chem.* **151**, 427.
 Dawson, R. M. C. (1954). *Biochim. biophys. Acta*, **14**, 374.
 De Boer, Th. J. (1954). *Rec. Trav. chim. Pays-Bas*, **73**, 229.
 Deuel, H. J. (1951). *The Lipids, Their Chemistry and Biochemistry*, vol. 1, p. 496. New York: Interscience Publisher Inc.
 Farquhar, J. W., Insull, W., Rosen, P., Stoffel, W. & Ahrens, E. H. (1959). *Nutr. Rev. (Suppl.)*, **17**, 1.
 Folch, J. & LeBaron, F. N. (1957). In *Metabolism of the Nervous System*, p. 67. Ed. by Richter, D. London: Pergamon Press Ltd.
 Folch, J., Lees, M. & Sloane-Stanley, G. H. (1957). *J. biol. Chem.* **226**, 497.
 James, A. T. (1960). *Meth. biochem. Anal.* **8**, 1.
 James, A. T., Lovelock, J. E. & Piper, E. A. (1959). *Ann. N.Y. Acad. Sci.* **72**, 720.
 James, A. T. & Webb, J. P. W. (1957a). *Biochem. J.* **66**, 515.
 James, A. T. & Webb, J. P. W. (1957b). *Proc. 4th int. Conf. Biochem. Prob. Lipids, Oxford*, p. 3. Ed. by Sinclair, H. M. London: Butterworths Scientific Publications.

- Johnson, A. C., McNabb, A. R. & Rossiter, R. J. (1948). *Biochem. J.* **43**, xxii.
- Johnston, P. V. & Kummerow, F. A. (1960). *Proc. Soc. exp. Biol., N.Y.*, **104**, 201.
- Klenk, E. (1927). *Hoppe-Seyl. Z.* **166**, 268.
- Klenk, E. (1928). *Hoppe-Seyl. Z.* **174**, 214.
- Klenk, E. & Böhm, P. (1951). *Hoppe-Seyl. Z.* **288**, 98.
- Klenk, E. & Bongard, W. (1952). *Hoppe-Seyl. Z.* **291**, 104.
- Klenk, E. & Krickau, G. (1957). *Hoppe-Seyl. Z.* **308**, 98.
- Klenk, E. & Lindlar, F. (1955a). *Hoppe-Seyl. Z.* **299**, 74.
- Klenk, E. & Lindlar, F. (1955b). *Hoppe-Seyl. Z.* **301**, 156.
- Klenk, E. & Montag, W. (1958a). *J. Neurochem.* **2**, 226.
- Klenk, E. & Montag, W. (1958b). *J. Neurochem.* **2**, 232.
- Laser, H. (1949). *J. Physiol.* **110**, 338.
- LeBaron, F. N. & Folch, J. (1957). *Physiol. Rev.* **37**, 539.
- Levene, P. A. & Rolf, P. (1922a). *J. biol. Chem.* **54**, 91.
- Levene, P. A. & Rolf, P. (1922b). *J. biol. Chem.* **54**, 99.
- Lipski, S. R., Landowne, R. A. & Godet, M. R. (1959). *Biochim. biophys. Acta*, **31**, 336.
- Lovelock, J. E. (1958). *J. Chromat.* **1**, 35.
- Moore, C. H. & Cook, R. P. (1959). *Biochem. J.* **73**, 43P.
- Morton, I. D. & Todd, A. R. (1950). *Biochem. J.* **47**, 327.
- Rose, B. A. (1959). *Analyst*, **84**, 574.
- Stern, I. & Shapiro, B. (1953). *J. clin. Path.* **6**, 158.
- Stoffel, W., Insull, W. & Ahrens, E. H. (1958). *Proc. Soc. exp. Biol., N.Y.*, **99**, 238.
- Thannhauser, S. J. & Boncoddio, N. F. (1948). *J. biol. Chem.* **172**, 141.
- Thudichum, J. L. W. (1901). *Die Chemische Konstitution des Gehirns des Menschen und der Tiere*. Tübingen: Granz Peitzske.

Biochem. J. (1961) **79**, 648

The Composition of Collagen from Subcellular Fractions of Guinea-Pig Granuloma Tissue

By J. E. EASTOE*

Department of Dental Science, Royal College of Surgeons of England, London, W.C. 2

(Received 31 October 1960)

Investigations of the amino acid composition of vertebrate collagens have been largely confined to preparations directly derived from whole tissues of high collagen content (Eastoe & Leach, 1958). Most studies have been made on the insoluble fibrous protein remaining after treatment of the tissue with solvents which could remove other tissue components with minimum degradation of the collagen (Bowes & Kenten, 1948). Such preparations would be expected to consist largely of extracellular collagen fibres with characteristic 640 Å striations. Recently analyses have been made of collagen fractions extracted from tissues by neutral sodium chloride and slightly acid sodium citrate solutions (Bowes, Elliott & Moss, 1955; Jackson, Leach & Jacobs, 1958). These soluble collagens represent only a small percentage of the total collagen, and much evidence, based on the incorporation of radioactive amino acids, has accumulated which shows that the neutral-salt-soluble fraction is synthesized before the insoluble fibres, which are probably formed by aggregation of the smaller particles in the soluble fraction (Harkness, Marko, Muir & Neuberger, 1954; Jackson, 1957; Green & Lowther, 1959).

The object of the work described here was to determine the composition of the protein material extracted by sodium chloride solution from three

subcellular fractions of rapidly developing mesenchymal tissue from guinea-pig granuloma. It forms part of a more extensive investigation involving radioactive tracer studies of proline hydroxylation in the subcellular fractions and their examination by electron microscopy (Lowther, Green & Chapman, 1961; Chapman, 1961). Since the protein fractions had already been shown to contain hydroxyproline and to have solubility properties similar to neutral-salt-soluble collagen extracted from whole tissues, confirmation of the characteristic overall amino acid pattern would be important evidence of the intracellular synthesis of the collagen molecule. The very small quantities of protein, obtainable from a large bulk of granuloma tissue, required the adoption of a modified semi-micro method of amino acid analysis, based on resin chromatography, which is described in an Appendix to this paper.

MATERIAL AND METHODS

Subcellular collagen fractions. The collagen fractions were isolated and purified by Dr N. M. Green and Dr D. A. Lowther (see also Lowther *et al.* 1961) by the following methods. Granulomata, produced by subcutaneous injection of carrageenin solution into guinea pigs, were removed 7 days later, immediately after killing the animals. The tissue was chopped finely with two 6 in. razor blades and serum proteins were largely removed by centrifuging a

* Senior Leverhulme Fellow in Biochemistry.