tions above 2.5 mM and of cationic cetyltrimethylammonium bromide at concentrations above 0.7 mM.

3. Digitonin, a non-ionic detergent, at a concentration of 2.0 mM solubilized about one-third of the microsomal protein without releasing lipid and adenosine-triphosphatase activity. At 2.8 mM, lipid and adenosine triphosphatase were released. Digitonin at concentrations 0.01–2.0 mM stimulated the activity of this enzyme in the presence of Na^+ ions; this effect was diminished at 2.8 mM, when the enzyme and lipid were solubilized.

4. Digitonin at all concentrations examined failed to release appreciable quantities of cholesterol into solution, owing presumably to the formation of insoluble complexes which were sedimented. The relative proportions of different lipids solubilized by all other agents were, however, similar to those present in the original microsomes.

5. About one-half of the microsomal constituents was solubilized after ultrasonic irradiation. Three-quarters of the Na^+ ion-stimulated adenosine-triphosphatase activity was recovered after 20 min. treatment, and only one-half was recovered after 40 min. treatment. Recoveries of the activity in the absence of Na^+ ions were one-half of those measured in the presence of Na^+ ions.

6. The choice of solubilizing agent in relation to these properties and to the maintenance of the activity of the adenosine triphosphatase is discussed.

We are grateful to Professor H. McIlwain for many helpful discussions. We also acknowledge with gratitude the assistance provided by the Medical Research Council, in the form of a grant for special research to D. B. G. and by the

provision of the analytical ultracentrifuge. P.D.S. was supported by a Special Fellowship BT-942 from the National Institute of Neurological Diseases and Blindness, U.S. Public Health Service.

REFERENCES

- Doty, P. & Schulman, J. H. (1949). *Disc. Faraday Soc.* **6**, 25.
 Eley, D. D. & Hedge, D. G. (1956). *Disc. Faraday Soc.* **21**, 221.
 Gammack, D. B. (1963). *Biochem. J.* **88**, 373.
 Hanzon, V. & Toschi, G. (1959). *Exp. Cell Res.* **16**, 256.
 Hughes, D. E. & Nyborg, W. L. (1962). *Science*, **138**, 108.
 Lowry, O. H., Rosebrough, N. J., Farr, A. L. & Randall, R. J. (1951). *J. biol. Chem.* **193**, 265.
 McBain, M. E. L. & Hutchinson, E. (1955). *Solubilization and Related Phenomena*, p. 61. New York: Academic Press Inc.
 Martin, J. B. & Doty, D. M. (1949). *Analyt. Chem.* **21**, 965.
 Rideal, E. K. & Taylor, F. H. (1957). *Proc. Roy. Soc. B*, **146**, 225.
 Rideal, E. K. & Taylor, F. H. (1958). *Proc. Roy. Soc. B*, **148**, 450.
 Ruysen, R. H. & Loos, R. (1947). *J. Colloid Sci.* **2**, 429.
 Salem, L. (1962). *Canad. J. Biochem. Physiol.* **140**, 1287.
 Saunders, L., Perrin, J. & Gammack, D. (1962). *J. Pharm., Lond.*, **14**, 567.
 Schwartz, A., Bachelard, H. S. & McIlwain, H. (1962). *Biochem. J.* **84**, 626.
 Skou, J. C. (1963). *Biochem. biophys. Res. Commun.* **10**, 79.
 Swanson, P. D., Bradford, H. F. & McIlwain, H. (1964). *Biochem. J.* **92**, 235.
 Toschi, G. (1959). *Exp. Cell Res.* **16**, 232.
 Wagner, H., Hörhammer, L. & Wolff, P. (1961). *Biochem. Z.* **334**, 175.
 Weissler, A. (1959). *J. Amer. chem. Soc.* **81**, 1077.
 Wherrett, J. R. & McIlwain, H. (1962). *Biochem. J.* **84**, 232.

Biochem. J. (1964), **92**, 254

Glycolytic Enzymes in Human Brain

By N. ROBINSON AND B. M. PHILLIPS

Department of Chemical Pathology, Institute of Neurology, The National Hospital, Queen Square, London, W.C. 1

(Received 2 September 1963)

Histochemical studies of animal brain have elucidated the sites of activity of enzymes associated with the different metabolic routes, and reliable data are now available for the comparison of glycolytic enzyme activities in the different regions (Breckenridge & Crawford, 1961; Naidoo, 1962; Pope, 1960; Robins & Lowe, 1961; Adams, Davison & Gregson, 1963). Few reports have

appeared on the activity of a particular enzyme in human brain or of a specific enzyme associated with a disease (Friede, 1961; Friede & Fleming, 1962; Lazarus, Wallace & Volk, 1962; Hess & Pope, 1961; Aronson, Saifer, Perle & Volk, 1961). In the present investigation, enzymes concerned with glucose metabolism in the human brain and cord have been studied to determine the rate at which

the enzymes can utilize glucose under the given experimental conditions.

The main glycolytic pathways studied were: (1) glycogen synthesis catalysed by phosphorylase; (2) Embden-Meyerhof oxidation of glucose (phosphoglucose isomerase, aldolase and lactate dehydrogenase); (3) direct oxidation of glucose (glucose 6-phosphate dehydrogenase and 6-phosphogluconate dehydrogenase). Regions investigated were the frontal and temporal lobes, white and grey matter; cerebellar white and grey matter; dorsal and lumbar cord. Enzyme activities were determined spectrophotometrically. Wherever possible comparisons were made with results from animal brains; these were, however, invariably killed, decapitated and instantly immersed in liquid nitrogen and were not post-mortem in the usual sense. Preservation of human material was maintained under the conditions required for valid estimations on post-mortem brain reported by Robins, Smith, Daesch & Payne (1958) and Naidoo & Pratt (1954).

MATERIALS AND METHODS

Two normal human adult male post-mortem brains and cords of persons known to be unaffected by neurological disorders and also unaffected biopsy specimens from three lobes (two frontal and one temporal) were used. The epineural connective tissue, meninges, major blood vessels and adjacent nervous tissue were eliminated in dissection, and the remainder was rapidly frozen and stored at -28° within 4-5 hr. after death; three lobes from biopsies were dissected within 10 min. after the operation and stored at -28° . During storage, while the experimental work was being conducted, no change was found in enzymic activity; activities were determined as soon as possible according to enzyme stability.

Concentrations of homogenates varied for different enzymes; generally 40 mg. wet wt. of white matter and 20 mg. wet wt. of grey matter/ml. were satisfactory stock suspensions. Solutions were made up in ice-cold distilled and ion-free water (for dehydrogenases), homogenized in a motor-driven Potter-Elvehjem Teflon-pestle homogenizer for 2 min. and then frozen. Homogenates were kept frozen when not being sampled; they were discarded after 24 hr.

Trial experiments were performed to establish the optimum enzyme concentration to give a linear relationship with reaction velocity throughout the period of incubation. The methods used, with reasonable care, were reproducible. Controls and blanks were prepared simultaneously with test solutions and the test results subsequently corrected. Up to six determinations were made to obtain reproducibility within 10%. Optical measurements were made on a Unicam SP. 500 spectrophotometer. In reactions where inorganic phosphate was released, phosphorus was determined by a micro-method (Robinson & Phillips, 1963).

Incubation mixtures

Phosphoglucose isomerase. The method was similar to that of Schwartz & Bodansky (1961). To 250 ml. of sodium di-

ethylbarbiturate-acetate buffer, 0.143 M with respect to each salt, was added 250 ml. of 0.1 N-HCl, followed by the slow addition of 83.3 ml. of 0.03 M-sodium glucose 6-phosphate previously adjusted to pH 7.4, and the mixture was diluted to 1 l.; the final concentration of the buffered substrate was 2.5 mM. After the incubation of 2 ml. of buffered substrate with 0.5 ml. of homogenate (12.5-50 μ g. wet wt. of tissue) at 38° for 30 min. the reaction was stopped with 2.5 ml. of 5% (w/v) trichloroacetic acid and the mixture centrifuged. Colour development of fructose was by the Seliwanoff reaction, with 10 N-HCl and ethanol-resorcinol (0.1%).

Phosphorylase. This was measured by the conversion of glucose 1-phosphate into glycogen with the release of phosphate. The method was adapted from Buell, Lowry, Roberts, Chang & Kapphahn (1958). The substrate was 30 mM-glucose 1-phosphate with added glycogen (0.5%), AMP (1 mM), NaF (50 mM) and cysteine (15 mM) (the hydrochloride neutralized to pH 6.7, this solution being added just before use). Incubation with 4-8 mg. wet wt. of tissue/ml. of reaction mixture was at 38° for 30 min. The reaction mixtures were arrested by the addition of an equal volume of 5% (w/v) trichloroacetic acid, and the mixtures were instantly cooled in ice-water and determinations of the released phosphorus carried out immediately. To diminish hydrolysis of labile glucose 1-phosphate and at the same time obtain maximum sensitivity for colour development, the ammonium molybdate solution was acidified with H_2SO_4 to give a final concentration of 0.2 N- H_2SO_4 .

Glucose 6-phosphate dehydrogenase and 6-phosphogluconate dehydrogenase. The method was adapted from those used by Glock & McLean (1953) and Marks (1961). The homogenate was prepared in ice-cold ion-free water; the suspension was then centrifuged at 3000g for 30 min. to give a supernatant liquid optically suitable for measurement. Final reaction mixtures contained 1.5-3.0 mg. wet wt. of tissue/ml., 0.5 ml. of 0.1 M- $MgCl_2$, 0.5 ml. of 0.25 M-glycylglycine buffer (pH 7.6), 0.1 ml. of 2.5 mM-NADP, and, in turn, substrates of 0.5 ml. of 0.01 M-glucose 6-phosphate, 0.01 M-6-phosphogluconate and finally combined substrate of 0.5 ml. of each of the two last named salts. Volumes were made up to a total of 2.5 ml. with ion-free water at 20° ; for extinction measurements blank cells were devoid of NADP. The effects of Ca^{2+} and Mg^{2+} ions, each at a concentration of 0.01 M, were studied in 0.01 M-tris buffer. For the determination of optimum pH, 0.25 M-glycylglycine buffer at pH intervals of 0.4 between 7.0 and 8.6 was used. The increase in extinction was taken at 0.5 min. intervals for 4 min. at 340 m μ .

Aldolase. The micro-procedure of Friedmann & Lapan (1958) was used; the reaction mixture contained 0.4-0.5 mg. wet wt. of tissue.

Lactate dehydrogenase. The method utilized the extinction changes at 340 m μ due to the NAD-NADH₂ interconversion and followed the details of Wróblewski & La Due (1955). The brain suspension was centrifuged at 3000g for 30 min. and the supernatant removed for assay. The final reaction mixtures consisted of 0.1 ml. of brain supernatant (2.0-4.0 mg. wet wt. of tissue/ml.), 0.1 ml. of 0.02 M-sodium pyruvate adjusted to pH 7.90 with NaOH, 0.1 ml. of NADH₂ (2 mg. of NADH₂/ml. of ion-free water) and 2.7 ml. of Sørensen phosphate buffer, pH 7.90. Readings were taken at 0.5 min. intervals over a period of 0-4 min. and the rate was determined graphically.

RESULTS

Enzyme activities were all measured under the optimum conditions that are believed to avoid significant destructive and inhibitory factors: modifications and adaptations to the methods have only

been made to suit the physicochemical measurement of the amount of substrate transformed; otherwise the methods used have been recognized for some time.

Preliminary experiments on the activation of cerebral glucose 6-phosphate dehydrogenase and

Table 1. *Glycolytic enzyme activities in human brain and cord*

A 1 and A 2 refer to the two autopsy specimens, B 1 and B 2 refer to the two frontal-lobe biopsy specimens and B 3 to the temporal-lobe biopsy specimen. For details of measurement of enzyme activities see the Materials and Methods section. Enzyme activities are expressed as μ moles of substrate converted/min./g. wet wt. of tissue, except for lactate dehydrogenase where the activities are in m-moles of substrate converted/min./g. wet wt. of tissue. The results are given as means $\pm$ S.D.

	Phosphoglucose isomerase					Phosphorylase				
	A 1	A 2	B 1	B 2	B 3	A 1	A 2	B 1	B 2	B 3
Cerebrum										
White matter	28.6	31.6	35.6	28.8	22.3	4.50	4.95	4.32	5.33	5.33
	± 2.1	± 2.2	± 2.2	± 1.9	± 2.0	± 0.3	± 0.3	± 0.3	± 0.4	± 0.3
Cortex	29.8	41.0	52.0	53.3	61.5	6.32	6.80	7.15	8.40	8.00
	± 0.8	± 0.4	± 0.4	± 0.4	± 0.5	± 0.4	± 0.4	± 0.4	± 0.5	± 0.5
Cerebellum										
White matter	22.5	26.7	—	—	—	2.80	3.06	—	—	—
	± 2.2	± 2.5	—	—	—	± 0.2	± 0.2	—	—	—
Cortex	44.5	46.7	—	—	—	2.92	3.56	—	—	—
	± 1.7	± 1.1	—	—	—	± 0.2	± 0.3	—	—	—
Cord										
Thoracic	—	9.7	—	—	—	—	2.86	—	—	—
	—	± 0.5	—	—	—	—	± 0.2	—	—	—
Lumbar	13.1	14.3	—	—	—	2.70	3.80	—	—	—
	± 0.8	± 0.7	—	—	—	± 0.2	± 0.3	—	—	—
	Aldolase					Lactate dehydrogenase				
	A 1	A 2	B 1	B 2	B 3	A 1	A 2	B 1	B 2	B 3
Cerebrum										
White matter	11.5	11.6	10.0	11.1	15.0	6.10	6.20	5.80	6.50	7.30
	± 0.3	± 0.4	± 0.5	± 0.4	± 0.6	± 0.58	± 0.60	± 0.54	± 0.59	± 0.61
Cortex	13.0	17.6	21.5	23.7	27.0	8.30	8.90	9.60	12.7	13.7
	± 0.7	± 1.0	± 1.1	± 1.2	± 1.3	± 0.78	± 0.80	± 0.91	± 1.1	± 1.2
Cerebellum										
White matter	8.3	10.9	—	—	—	10.1	10.7	—	—	—
	± 0.4	± 0.5	—	—	—	± 1.0	± 0.9	—	—	—
Cortex	17.8	26.5	—	—	—	10.9	12.3	—	—	—
	± 1.2	± 1.8	—	—	—	± 1.1	± 1.2	—	—	—
Cord										
Thoracic	—	8.15	—	—	—	—	6.40	—	—	—
	—	± 0.4	—	—	—	—	± 0.5	—	—	—
Lumbar	3.56	4.30	—	—	—	6.30	7.10	—	—	—
	± 0.1	± 0.1	—	—	—	± 0.6	± 0.6	—	—	—
	Glucose 6-phosphate dehydrogenase					6-Phosphogluconate dehydrogenase				
	A 1	A 2	B 1	B 2	B 3	A 1	A 2	B 1	B 2	B 3
Cerebrum										
White matter	1.10	1.15	2.35	2.56	2.90	1.35	1.58	1.68	1.77	2.33
	± 0.10	± 0.12	± 0.21	± 0.24	± 0.28	± 0.12	± 0.14	± 0.16	± 0.15	± 0.24
Cortex	0.85	0.93	1.43	1.48	1.83	1.13	1.27	1.38	1.32	1.48
	± 0.09	± 0.08	± 0.13	± 0.14	± 0.17	± 0.12	± 0.12	± 0.13	± 0.12	± 0.16
Cerebellum										
White matter	2.42	2.55	—	—	—	2.17	2.40	—	—	—
	± 0.25	± 0.23	—	—	—	± 0.20	± 0.21	—	—	—
Cortex	0.82	1.00	—	—	—	0.75	1.00	—	—	—
	± 0.09	± 0.09	—	—	—	± 0.06	± 0.09	—	—	—
Cord										
Thoracic	—	1.10	—	—	—	—	1.07	—	—	—
	—	± 0.10	—	—	—	—	± 0.10	—	—	—
Lumbar	1.45	2.36	—	—	—	1.33	2.55	—	—	—
	± 0.14	± 0.21	—	—	—	± 0.11	± 0.21	—	—	—

6-phosphogluconate dehydrogenase by Mg^{2+} and Ca^{2+} ions at a concentration of 10 mM in 10 mM-tris buffer showed that both enzymes were stimulated to different extents by these ions; at this concentration the activity of glucose 6-phosphate dehydrogenase increased by 56 % with Mg^{2+} ions and 48 % with Ca^{2+} ions, whereas 6-phosphogluconate dehydrogenase was stimulated to a smaller extent: Mg^{2+} ions increased activity by 31 % and Ca^{2+} ions by 26 %. Further work showed that Mg^{2+} ions activated both enzymes to a greater extent than Ca^{2+} ions over the concentration range of 1.0 mM–0.1 M. In the presence of the optimum concentration, 20 mM- Mg^{2+} ion, the activities of glucose 6-phosphate dehydrogenase and 6-phosphogluconate dehydrogenase were increased by 59 and 32 % respectively. Final estimations of the enzyme activities were made in glycylglycine buffer to remove traces of inhibitory metal ions (Glock & McLean, 1953). In this buffer the corresponding activities were less than 5 % higher than in the tris buffer, and, since this value was within the standard deviation for the enzyme preparations, the effect of the two different buffers on activities was not regarded as significant. The optimum pH for the reaction was 7.6, the activity falling to 79 % of the maximum at pH 7.0 and 82 % at pH 8.6. The K_m values were 21 μ M for glucose 6-phosphate dehydrogenase and 24 μ M for 6-phosphogluconate dehydrogenase.

Enzyme determinations, expressed in μ moles of substrate converted/min./g. wet wt. of tissue, are shown in Table 1. Values quoted by other investigators have been converted into similar units for comparison. Previous investigators have reported activities in terms of wet weight, dry weight and lipid-free weight. Although the lipid content differs widely in various parts of the brain, the lipid-free dry weight appears to be a nearly constant fraction (10–12 %) of the wet weight of tissue (Buell *et al.* 1958); where comparisons were made with results expressed on a lipid-free basis the factor 0.1 was used.

DISCUSSION

Most of the work reported on brain has been carried out with animals, mainly the rabbit; a few scattered studies on human tissue, mostly in relation to disease, have been reported.

A human adult brain having a respiratory rate of 2 μ moles of oxygen/min./g. of tissue could consume 0.3 μ mole of glucose/min./g., of which about 0.05 μ mole of glucose/min. would be utilized for the formation of 0.10 μ mole of lactate/min. From this all the glycolytic enzyme activities measured indicate that they would normally be functioning at no more than 10 % of their potential activity. In the cerebrum, where the potential activity of

enzymes in the Embden–Meyerhof route is much greater, they would function at an even lower rate, especially in white matter where the respiratory rate is lowered to 1 μ mole of oxygen/min./g.

In the absence of oxygen, however, lactate formation can increase in the cerebral cortex to 1.66 μ moles/min./g., requiring a threefold increase in normal enzyme working capacity. Anaerobic glycolysis may operate at very high rates for short periods of high cerebral activity and during convulsions (McIlwain, 1959), when 6.6 μ moles of lactate/min./g. may be formed; lactate dehydrogenase and aldolase will then be working at least half capacity.

Phosphorylase. This enzyme catalyses the condensation of glucose 1-phosphate to form the (1 $\rightarrow$ 4)-glucosidic links of glycogen. In the uridine-linked glycogen cycle phosphorylase has been reported as a reversible enzyme (Greville, 1962) and also as a degrading enzyme only (Sidman, 1960), but it appears to operate less in brain than in other tissues. Phosphorylase activity has recently been measured in the opposite direction by the catalytic phosphorolysis of glycogen, the rate obtained being 37 % of that by the more usual method (Breckenridge & Norman, 1962). 'Active' and 'inactive' forms of phosphorylase have been found in the brain, the inactive form that requires AMP for activation predominating. Activation is almost instantaneous. The activity found in the present study is less than that found in whole brain of mice (Breckenridge & Norman, 1962) and rabbit (Buell *et al.* 1958), but comparison with cerebellar cortex from the monkey (Robins, Smith & Jen, 1957) showed a similar value.

Significant differences have been obtained in different regions of the brain. If the maximum rate of glycogen accumulation in the cerebral cortex does not exceed 0.03 μ mole/min./g., phosphorylase as a synthesizing enzyme is adequate for this rate of glycogenesis. The activity of glycogen phosphorylase in the human cerebral cortex was just over 60 % of the values found for rabbit (Breckenridge & Crawford, 1961). Human cerebral and cerebellar cortex both had higher activities than the white matter; this and similar areas of lower activity may utilize glycogen for a special role depending on the efficiency of hexokinase for the direct utilization of glucose. Cerebellar cortex showed the highest activity of all sections examined; Buell *et al.* (1958) also found high phosphorylase activity in the molecular layer of the rabbit cerebellum. Robins (1960) suggested that the lightly myelinated tracts favoured direct utilization of glucose via hexokinase, whereas utilization of glycogen via phosphorylase and phosphoglucomutase would be relatively important in heavily myelinated tracts. This does not appear to be borne out by values for

the human frontal lobe; the dorsal column of the rabbit and dorsal and lumbar regions of the human cord also showed a lower activity than other more lightly myelinated regions. Glycogen disappears from animal brain at a rate of at least $1.5 \mu\text{moles}$ of hexose/min./g. (Breckenridge & Crawford, 1961); this would suggest that, if glycogenolysis were the consequence of phosphorylase activity, the enzyme (presumably debranching) would be working at maximum capacity in parts of the brain and cord, in which case this enzyme is likely to be rate-controlling in brain glycogenesis.

Phosphoglucose isomerase. This enzyme was first demonstrated in extracts of brain by Lohmann (1933). Buell *et al.* (1958) investigated phosphoglucose isomerase in rabbit brain by the same method used in the present work and found that the K_m of the reaction was small since the velocity was as rapid with a 2.5 M substrate as with 90 mM for the conversion of the same small fraction of substrate. The rate was found to decrease with substrate conversion as predicted for a reversible reaction, and hence the fractions of substrate converted in the experiments were kept small.

The enzyme showed a high absolute activity in all regions, the greatest activity appearing in the cerebral and cerebellar cortex. In these regions the enzyme was at least 35-fold as active as glucose 6-phosphate dehydrogenase, suggesting the greater importance of anaerobic glucose utilization. McIlwain (1959) has suggested that a high activity is related to securing the isomerization and subsequent glycolysis of glucose phosphate in competition with the direct oxidative pathway. The affinity of fructokinase for its substrate necessitates a rate of isomerization of glucose 6-phosphate of $10 \mu\text{moles/min.}/\text{g.}$ (calculated for rat brain); human cerebral and cerebellar cortex would be functioning at 15–25 % capacity to meet this demand.

Aldolase. The activity of aldolase varied considerably in different regions of the brain examined; similar findings were reported for the rabbit by Buell *et al.* (1958). In human cerebellar cortex aldolase activity was only 90 % of that reported by Robins *et al.* (1957) for the monkey, but in both the activities were more than 50 % higher than the average for whole rat and rabbit brain. Human cerebellar white matter showed similar aldolase activity to that in the rabbit. Low activity in the human cord corresponded to values for the rabbit dorsal column; both were only 30 % of the highest activity found in brain.

Hexose monophosphate oxidation. The oxidation of glucose 6-phosphate and the oxidative decarboxylation of 6-phosphogluconate are both NADP-dependent, and measurement of the conversion of NADP with glucose 6-phosphate as substrate is a summation of the two successive stages

of dehydrogenase activities. The methods of Glock & McLean (1953) and Marks (1961) enabled the rates of both reactions to be determined.

Two cations are known to stimulate the activities of these dehydrogenases. Previous investigations showed that yeast (Kornberg & Horecker, 1955), *Leuconostoc mesenteroides* (De Moss, 1955) and liver (Glock & McLean, 1953) dehydrogenases were strongly activated by Mg^{2+} ions; cerebral preparations showed a similar effect; Ca^{2+} ions provoked more activation in brain than that reported in liver. Heavy-metal ions inhibit the reaction of liver dehydrogenases (Glock & McLean, 1953), but traces of the heavy-metal ions were effectively removed by a binding mechanism when glycylglycine was used as the buffer. By using this buffer to observe the effect of pH on the reaction, it was found that the activity in cerebral preparations was suppressed to a greater extent below than above the optimum of pH 7.6.

The two successive dehydrogenations in the direct oxidative route took place at comparable rates in the different regions of human brain and cord but showed the lowest absolute activities of the enzymes investigated. The low K_m for this reaction indicates that this stage of glycolysis could compete favourably with the alternative anaerobic route for glucose 6-phosphate. Cerebral and cerebellar white matter was more active than cortex; the heavily myelinated tracts of the cord also showed higher activity, suggesting that the direct oxidative pathway could play a greater part in the utilization of glucose 6-phosphate in these regions. Greville (1962), however, has shown that the metabolism of glucose in the brain of several animals is mainly by the Embden-Meyerhof route. It is possible that the direct oxidative pathway may have a special function in white matter possibly related to myelin or oligodendroglia.

Lactate dehydrogenase. This enzyme, catalysing the last stage of the Embden-Meyerhof route, possessed the lowest activity of the enzymes examined in anaerobic glycolysis. The cerebellum was a relatively active region. Enzymic activities in the myelinated region of the cerebrum and the heavily myelinated lumbar and dorsal cords were within 10 % of each other. Aronson *et al.* (1961) gave slightly higher values for brain, but no recent comparisons are available for human cord. In studies on the developing rat brain Robins & Lowe (1961) tentatively suggested that, during growth, lactate dehydrogenase was related to differentiation of glia and deposition of myelin in addition to a general glycolytic metabolism, but in the adult brain the special functions disappeared.

In conclusion, the present study shows that the relative magnitudes of most enzyme activities in the glycolytic pathways of the human brain and

cord are similar to those of lower animals; the overall level, however, is generally lower in human tissue.

Myelinated regions of the human brain are invariably lower in glycolytic enzyme activity than other areas, exemplified particularly by the heavily myelinated dorsal and lumbar regions of the cord. In these areas of lower activity the absolute values of phosphorylase suggest the likelihood of an increase in glycogenesis. The high activity of the enzyme of the Embden-Meyerhof route in grey matter may, in part, be due to the greater blood supply and a greater concentration of lightly myelinated nerve tracts having nodes of Ranvier, which are known to be active centres of metabolic activity. Phosphorylase activity in all regions examined suggests that a mechanism for glycogen metabolism can function in most parts of the brain and in the cord.

SUMMARY

1. Six glycolytic enzymes of the anaerobic and aerobic routes of carbohydrate metabolism have been examined in human cerebral and cerebellar white and grey matter, dorsal and lumbar cord.

2. A comparison of activities in these regions shows that enzymes of the Embden-Meyerhof pathway are more active in grey matter than in white matter.

3. The very much lower absolute activities of the dehydrogenases of the direct oxidative pathways are relatively more active than other enzymes in white matter.

4. Phosphorylase activity in all regions suggests that glycogen metabolism probably functions in the brain and the cord.

5. Comparisons have been made with other mammalian brains where possible.

The authors thank Professor J. N. Cumings for his interest and encouragement throughout the work. Thanks are also due to Mr Wylie McKissock, F.R.C.S., for the frontal and temporal lobes, and the Whittington Hospital, Archway, for the post-mortem brains. N. R. is also indebted to the Knopp and Evelyn Staines Fellowships for financial assistance.

REFERENCES

- Adams, C. W. M., Davison, A. N. & Gregson, N. A. (1963). *J. Neurochem.* **7**, 234.
- Aronson, S. M., Saifer, A., Perle, G. & Volk, B. W. (1961). *Amer. J. Nutr. Sci.* **9**, 103.
- Breckenridge, B. M. & Crawford, E. J. (1961). *J. Neurochem.* **7**, 234.
- Breckenridge, B. M. & Norman, J. H. (1962). *J. Neurochem.* **9**, 383.
- Buell, M. V., Lowry, O. H., Roberts, N. R., Chang, M. W. & Kapphahn, J. I. (1958). *J. biol. Chem.* **232**, 979.
- De Moss, R. D. (1955). In *Methods in Enzymology*, vol. 1, p. 328. Ed. by Colowick, S. P. & Kaplan, N. O. New York: Academic Press Inc.
- Friede, R. L. (1961). *J. Neurochem.* **8**, 17.
- Friede, R. L. & Fleming, L. M. (1962). *J. Neurochem.* **9**, 179.
- Friedmann, M. M. & Lapan, B. (1958). *J. Lab. clin. Med.* **51**, 745.
- Glock, G. E. & McLean, P. (1953). *Biochem. J.* **55**, 400.
- Greville, G. D. (1962). In *Neurochemistry*, p. 238. Ed. by Elliott, K. A. C., Page, I. H. & Quastel, J. H. Springfield, Ill.: Charles C. Thomas.
- Hess, H. H. & Pope, A. (1961). *J. Neurochem.* **8**, 299.
- Kornberg, A. & Horecker, B. L. (1955). In *Methods in Enzymology*, vol. 1, p. 323. Ed. by Colowick, S. P. & Kaplan, N. O. New York: Academic Press Inc.
- Lazarus, S. S., Wallace, B. J. & Volk, B. W. (1962). *Amer. J. Path.* **41**, 579.
- Lohmann, K. (1933). *Biochem. Z.* **262**, 137.
- Mellwain, H. (1959). *Biochemistry and the Central Nervous System*, 2nd ed., p. 68. London: J. and A. Churchill Ltd.
- Marks, P. A. (1961). *Meth. med. Res.* **9**, 24.
- Naidoo, D. (1962). *J. Histochem. Cytochem.* **10**, 421.
- Naidoo, D. & Pratt, O. E. (1954). *Enzymologia*, **17**, 1.
- Pope, A. (1960). *J. Histochem. Cytochem.* **8**, 425.
- Robins, E. (1960). *J. Histochem. Cytochem.* **8**, 431.
- Robins, E. & Lowe, I. P. (1961). *J. Neurochem.* **8**, 81.
- Robins, E., Smith, D. E., Daesch, G. E. & Payne, K. E. (1958). *J. Neurochem.* **3**, 19.
- Robins, E., Smith, D. & Jen, M. K. (1957). *Progr. Neurobiol.* **2**, 205.
- Robinson, N. & Phillips, B. M. (1963). *Clin. chim. Acta*, **3**, 385.
- Schwartz, M. K. & Bodensky, O. (1961). *Meth. med. Res.* **9**, 5.
- Sidman, R. L. (1960). *J. Histochem. Cytochem.* **8**, 412.
- Wróblewski, F. & La Due, J. S. (1955). *Proc. Soc. exp. Biol., N.Y.*, **90**, 210.