SUMMARY

1. A water-soluble greenish fluorescent pigment from the culture broth of a strain of *Pseudomonas fluorescens-putida* intermediate has been crystallized after its extraction with water-saturated phenol, successive adsorption on Norit A and Dowex 50, and finally separation from other associated components by electrophoresis over a cellulose bed.

2. The pigment has been shown to be a brownish microcrystalline low-molecular-weight high-melting solid with a distinct powder-X-ray-diffraction pattern. Its different physicochemical properties have been described.

3. Chemical degradative studies and the assay of the pigment with *Crithidia fasciculata* suggest that the pigment is a pteridine derivative with a polyhydroxy moiety in the side chain.

We are extremely grateful to the Council of Scientific and Industrial Research, New Delhi, for sponsoring the work, to Dr S. Ganguly for the strain and to Dr A. K. Bose of Stevens Institute of Technology, New Jersey, for the infrared-absorption and the nuclear-magnetic-resonance spectra. The gift of a number of substituted pteridines from Dr Hugh S. Forrest is also gratefully acknowledged.

REFERENCES

- Albert, A. (1954). *Biochem. J.* **57**, x.
 Andes, J. E. & Myers, V. C. (1937). *J. biol. Chem.* **118**, 137.
 Broquist, H. P. & Albrecht, A. M. (1955). *Proc. Soc. exp. Biol., N.Y.*, **89**, 178.
 Buchanan, J. G., Dekker, C. A. & Long, A. G. (1950). *J. chem. Soc.* p. 3162.
 Chakrabarty, A. M. & Roy, S. C. (1964). *Biochem. J.* **92**, 105.
 Clemo, G. R. & Daglish, A. F. (1950). *J. chem. Soc.* p. 1481.
 Crammer, J. L. (1948). *Nature, Lond.*, **161**, 349.
 Dixon, J. S. & Lipkin, D. (1954). *Analyt. Chem.* **26**, 1092.
 Elliott, R. P. (1958). *Appl. Microbiol.* **6**, 241.
 Esposito, R. G. & Fletcher, A. M. (1961). *Arch. Biochem. Biophys.* **93**, 369.
 Forrest, H. S., Hanly, E. W. & Lagowski, J. M. (1961). *Biochim. biophys. Acta*, **50**, 596.
 Forrest, H. S. & McNutt, W. S. (1958). *J. Amer. chem. Soc.* **80**, 739.
 Forrest, H. S. & Mitchell, H. K. (1955). *J. Amer. chem. Soc.* **77**, 4865.
 Ganguly, S. (1955). *Ann. Biochem.* **15**, 215.
 Giral, F. (1936). *An. Soc. esp. Fis. Quim.* **34**, 667.
 Guttman, H. N. (1962). *Proc. 3rd int. Pteridine Symposium*. Ed. by Pfliederer, W. & Taylor, E. C. Stuttgart: Pergamon Press. (in the Press).
 Haynes, W. C., Stodola, F. H., Locke, J. M., Pridham, T. G., Conway, H. F., Sohns, V. E. & Jackson, R. W. (1956). *J. Bact.* **72**, 412.
 Hillemann, H. (1938). *Ber. dtsh. chem. Ges.* **71** B, 46.
 Kluyver, A. J. (1956). *J. Bact.* **72**, 406.
 Kogl, F. & Postowsky, J. J. (1930). *Liebigs Ann.* **480**, 280.
 Kuhn, R., Rudy, H. & Wagner-Jauregg, T. (1933). *Ber. dtsh. chem. Ges.* **66**, 1950.
 McNutt, W. S. (1956). *J. biol. Chem.* **219**, 365.
 Maley, G. F. & Plaut, G. W. E. (1959). *J. biol. Chem.* **234**, 641.
 Masuda, T. (1955). *Pharm. Bull., Tokyo*, **4**, 71.
 Masuda, T., Kishi, T. & Asai, M. (1957). *Pharm. Bull., Tokyo*, **5**, 598.
 Nathan, H. A. & Cowperthwaite, J. (1954). *Proc. Soc. exp. Biol., N.Y.*, **85**, 117.
 Stokstad, E. L. R., Hutchings, B. L., Mowat, J. H., Boothe, J. H., Waller, C. W., Angier, R. B., Semb, J. & Subba Row, Y. (1948). *J. Amer. chem. Soc.* **70**, 5.
 Sullivan, M. (1905). *J. med. Res.* **14**, 109.
 Turfitt, G. E. (1937). *Biochem. J.* **31**, 212.
 Wolf, F. T., Jones, E. A. & Nathan, H. A. (1958). *Nature, Lond.*, **182**, 475.

Biochem. J. (1964), **93**, 148

The Metabolism of Fructose Polymers in Plants

4. β -FRUCTOFURANOSIDASES OF TUBERS OF *HELIANTHUS TUBEROSUS* L.*

BY J. EDELMAN AND T. G. JEFFORD

Department of Plant Physiology, Imperial College of Science and Technology, London, S.W. 7

(Received 14 January 1964)

The Jerusalem artichoke (*Helianthus tuberosus* L.) contains in its stems and tubers a series of β -D-fructofuranosides, each member of which has the general formula $\text{Glc} \sim \text{Fru-Fru}_n$, where $\text{Glc} \sim \text{Fru}$ represents a sucrosyl group and n may be any number from zero (sucrose) to about 35 (inulin);

the fructose residues appear all to be (2 $\rightarrow$ 1')-linked, giving straight chains (Haworth & Learner, 1928); a similar situation occurs in other members of the Compositae. A chromatogram of this series of compounds from an extract of artichoke tubers is shown in Fig. 1. Each spot on paper chromatograms seems to represent a single substance (Bacon, 1960), and glucose-free fructose chains

* Part 3: Edelman, Recaladin & Dickerson (1963).

could not be detected by Quillet (1957). Apart from the cell walls, the tubers contain no other carbohydrates in significant amounts.

It is well known that there is a gradual decrease in the average degree of polymerization of the fructosides in the tuber during winter storage (Bacon & Loxley, 1952). Furthermore, most of the polymerized fructose is lost during sprouting of the tubers, much of it providing substance for the early growth of the daughter plants (Jefford & Edelman, 1961). Examination of the changes in the carbohydrate series in the course of this loss

showed that they were compatible with the hypothesis that the hydrolytic β -D-fructofuranosidase described by Edelman & Bacon (1951*a*) is involved in the breakdown of polymerized material.

This paper describes some characteristics of this hydrolytic activity from artichoke tubers, with a view to relating it to the physiological situation.

A preliminary report of some of this work has been presented to the Biochemical Society (Edelman & Jefford, 1963).

METHODS

Jerusalem artichoke tubers were obtained from the Botanical Supply Unit, University of London.

Estimations. Total fructose (free and combined) was measured colorimetrically by the method of S. W. Cole (see Bacon & Bell, 1948), and reducing sugar by the method of Somogyi (1952) and Nelson (1944). Soluble protein was measured colorimetrically by the method of Nielsen (1958).

Preparation of enzyme free of substrate. In a typical preparation tubers were disbudded, peeled, washed and dried, and the weighed tuber material (590 g.) was macerated in small batches in a Waring Blender in the cold room in a medium containing 0.1 M-cysteine, 0.05 M-sodium diethyldithiocarbamate and 0.02 M-phosphate buffer, each reagent adjusted to pH 7.2. The homogenate was filtered through four thicknesses of muslin and the filtrate returned, together with a little cold water and more tissue, to the Blender. The macerate finally yielded 610 ml. of extract and 95 g. of pulp. The extract was dialysed overnight against tap water containing sodium diethyldithiocarbamate and phosphate buffer, pH 7.2, at 2°, and then centrifuged at 2800g for 10 min. at 0°. Protein was precipitated from the supernatant liquid by addition of solid ammonium sulphate to 50% saturation, the solution being maintained at about pH 7.5. The precipitated protein was separated by centrifugation at 10000g for 10 min. at 0°, washed with 50% saturated ammonium sulphate solution, and then dissolved in 5 mM-phosphate buffer, pH 7.5. The solution was dialysed overnight against cold (London) tap water to inactivate transfructosylase activity [see Recaldin, (1961); preparations described by Edelman & Bacon (1951*b*) retained transfructosylase activity after dialysis against Sheffield tap water: we ascribe this difference to solutes in the water rather than to differences in the enzymes]. The suspension was centrifuged at 3000g for 10 min. at 0° and then poured into 12.5 vol. of acetone at -15°. A quantity of washed talc was added to the suspension as a diluent of the precipitated protein. After standing for 20 min. the mixture of precipitated protein and talc was separated by decantation and centrifugation at 8000g for 10 min. at -10°. The precipitate was washed twice with dry acetone at -15° and dried under vacuum over calcium chloride to yield 2.4 g. of pale-brown friable powder containing 35% (w/w) of soluble protein as calibrated against casein and estimated by the Nielsen (1958) method, and only 0.1% (w/w) of anhydrofructose.

Chromatography of protein. A column of DEAE-cellulose (Whatman DE 50, powder form: 1 g.; 11 mm. $\times$ 85 mm.) was washed with about 5 bed-vol. of N-sodium hydroxide and then with 5 mM-phosphate buffer, pH 7.5, until the pH of the effluent buffer was unchanged.

Fig. 1. Chromatogram of artichoke-tuber extract. Developed for 2 days with propan-1-ol-ethyl acetate-water (6:1:3); sprayed with urea-metaphosphoric acid.

Protein solutions were prepared for chromatography by suspending about 0.5 g. of acetone-dried powder in 5–10 ml. of 5 mM-phosphate buffer, pH 7.5; the suspension was centrifuged at 3000g for 1 min. at room temperature and the supernatant liquid dialysed overnight against 5 mM-phosphate buffer, pH 7.5, at 2°. A sample was removed for assay of hydrolase activity and the remainder put on to the column (amounts of up to 40 mg. of protein/100 mg. of DEAE-cellulose). After washing with 5 mM-phosphate buffer, pH 7.5, the column was eluted with solutions (pH 7.5) of successively greater ionic strength.

Assay for hydrolase activity. Unless stated otherwise, digests contained 50% (v/v) of enzyme preparation (effluent fraction from DEAE-cellulose column) and 0.1 M-acetate buffer, pH 5.0, and were incubated at 25°, usually for 4 hr.

In some experiments the liberation of fructose was measured by increase in total reducing-sugar values (method 1), but a more specific method (2), which also overcame high blank values, was normally used: mercuric chloride (final concentration 2 mM) was added to the digests to stop enzyme action, and free fructose was separated from the oligosaccharides by passage through small charcoal-Celite columns (3:1, w/w; 0.35 g.), and subsequently estimated by the total fructose method. Table 1 shows that the simpler but less accurate reducing-sugar method (1) gave reasonable results, but method 2 was used for most experiments, especially those involving oligosaccharides with reducing power.

Enzyme units. Following the recommendation of the International Union of Biochemistry, units are expressed as μ moles of hexose liberated/min. at 25°. For disaccharides giving two reducing hexoses on hydrolysis, 1 unit is taken as the liberation of 2 μ moles of hexose under the same conditions.

Chromatography of carbohydrates. Descending partition chromatography of sugars was carried out either on sheets of paper or, for separation of large amounts of mixed sugars (up to 1.5 g.), on a hollow cylindrical column of paper (Grycksbo Chromatographic no. 3, quality A; 25 mm. int.diam. to 58 mm. ext.diam. $\times$ 360 mm.) fitted with a pressure mantle (LKB 3502 ChroMax, LKB Produktor, Fabriksaktiebolag, Stockholm, Sweden). Whatman no. 1 paper was used for qualitative and quantitative chromatography, 3MM and 1.6MM for preparative separations, and no. 54 for the resolution of mixtures containing hexoses and hexitols. Chromatograms were developed at 25°: development of the paper column was carried out at room temperature. The solvents used most often were propan-1-ol-ethyl acetate-

water in the proportions 7:1:2 (by vol.) or 6:1:3 (by vol.). The former was used principally for sugars with D.P.* 1–4, which were separated satisfactorily after about 48 hr. development. The second mixture was used for separation of oligosaccharides of greater D.P., sugars with D.P. up to 10 being separated from mixtures by extending the development time to 4–5 days. Occasionally multiple development was used. Ethyl methyl ketone-acetic acid-aq. saturated boric acid soln. (9:1:1, by vol.) was used to resolve mixtures containing hexoses and hexitols (Rees & Reynolds, 1958).

Charcoal columns for the separation of monosaccharides from oligosaccharides were made from a mixture (1:1, w/w) of Activated Charcoal (British Drug Houses Ltd.) and Celite 535 (Johns-Manville, New York, N.Y., U.S.A.). The mixture was washed with water until the initially acid washings were about pH 5.5. The amount of sugar applied to the columns was never in excess of 1–1.5 g. of mixed oligosaccharides/20 g. of charcoal-Celite (Bacon, 1954). Monosaccharides were eluted with water, and the oligosaccharides with 25% (v/v) ethanol.

Columns of fuller's earth (cf. Binkley, 1955) were used for the separation of certain sugar mixtures which could not be resolved satisfactorily on paper or charcoal-Celite columns. After removal of the fines by decantation, the fuller's earth was mixed with about one-quarter of its weight of Celite 535 to accelerate the rate of flow. The columns were set up with absolute ethanol, and sugars were applied in a small volume of about 50% (v/v) ethanol; the column was eluted stepwise with ethanol containing progressively greater concentrations of water.

The eluates from columns were examined by chromatography on paper sheets.

The following spray reagents were used to detect carbohydrates on paper chromatograms: benzidine-trichloroacetic acid (Bacon & Edelman, 1951) as a general reagent; urea-metaphosphoric acid (Wise, Dimler, Davis & Rist, 1955) for free and combined fructose in the presence of aldoses; alkaline 2,5-diphenyl-3-(4-styrylphenyl)tetrazolium chloride (Bacon, 1959) for free or combined reducing fructose groups; sodium metaperiodate-benzidine (Cifonelli & Smith, 1954) for sugars and sugar alcohols.

Substrates

Inulin. Inulin was obtained commercially (Thomas Kerfoot and Co. Ltd., Vale of Bardsley, Lancs.; batch no. ED45348, from chicory). It was chromatographed to determine the lower limit of D.P.; successive developments in propan-1-ol-ethyl acetate-water (6:1:3) followed by spraying with urea-metaphosphoric acid revealed no fructosan of D.P. < 15. A sugar corresponding in R_F to that of D.P. 15 in the artichoke extract was present in trace amount; sugars between this and the origin (D.P. > about 20) were present in similar amounts, but at least 95% of the inulin was composed of polymers of D.P. > 20.

For comparative purposes, the average molecular weight was assumed to be 5000 ($\equiv$ D.P. 30; see Haworth, Hirst & Percival, 1932), although no experimental estimate of this was made.

Artichoke-tuber oligosaccharides. Individual members of the oligosaccharide series were prepared by paper chromatography [on Whatman 3MM or 1.6MM sheets or on a

Table 1. Comparison of methods for estimation of hydrolase activity

Results are expressed as milliunits of enzyme activity/ml. of digest. For description of methods see text.

Substrate	Enzyme	Method 1	Method 2
		1	2
(1F-Fructosyl) ₂ sucrose (10 mM)	None	0.22	0.83
	Hydrolase A	14.6	14.2
	Hydrolase B	37.0	31.8
Inulin (50 mg./ml. $\approx$ 10 mM)	None	1.89	1.67
	Hydrolase A	6.45	7.95
	Hydrolase B	19.2	21.5

* Abbreviation: D.P., degree of polymerization.

LKB ChroMax column, developed with propan-1-ol-ethyl acetate-water (6:1:3 or 7:1:2, by vol.) and rechromatographed before use. When the chromatographic mobilities of these sugars were plotted in the form $\log[R_F/(1-R_F)]$ against the assumed D.P. up to 10 (French & Wild, 1953) a straight line was obtained (Fig. 2).

A standard mixture of oligosaccharides (substrate M) for enzyme assay was prepared from tubers which had been stored at 2° for 8 months. The tubers were washed, dis-budded and 217 g. was disintegrated in a Waring Blender with about 100 ml. of 5 mM-sodium diethyldithiocarbamate, which was added to suppress polyphenol-oxidase activity. The homogenate was squeezed through muslin, boiled for 4 min. to denature enzymes, and the juice filtered while still hot. Chromatography showed that the oligosaccharides of lower D.P. predominated; there was some material with R_F zero, and also free fructose. Fractional precipitation with ethanol was carried out to rid the extract of free fructose (to decrease the initial reducing-sugar value), of sucrose (because of its inhibitory action, see below), and of the sugar of D.P. > 15 which were presumed to be attacked more slowly. Ethanol (3 vol.) was added to the 210 ml. of extract; the mixture was left at 2° overnight and then the precipitate was separated by centrifugation at 9000g for 10 min.; a residual opalescence was removed by filtration; a further 5 vol. of ethanol was added to the filtrate and the mixture allowed to stand overnight at 2° before separation of the precipitate by centrifugation at 9000g for 10 min. The precipitate was dissolved in water, 8 vol. of ethanol was added, and the material which precipitated overnight at 2° separated as above, dissolved in water and made up to 50 ml.: the solution contained 44.5 mg. of combined fructose/ml. Chromatography of a

sample of this solution showed that it was virtually free of fructose and sucrose, and that the bulk of the material comprised the oligosaccharides of D.P. 5–12.

Inulobiose series of oligosaccharides. Analogues of the natural artichoke series of sugars differing by the absence of a terminal glucose residue were prepared by partial acid hydrolysis of inulin, e.g. 2.45 g. of inulin was incubated with 11 ml. of 0.1N-oxalic acid for 30 min. at 65–70°. The hydrolysate was cooled and put on to a column (33 mm. × 145 mm.) composed of 40 g. of charcoal-Celite mixture, and the column washed with about 10 bed-vol. of water. Paper chromatography of a sample of the effluent showed that there was some disaccharide present with the glucose and fructose; this was recovered by paper chromatography. The sugars remaining on the charcoal-Celite column were eluted with 25% (v/v) ethanol and the effluent was evaporated to dryness. The dried eluate (1.02 g.) was put on to the LKB ChroMax paper column, which was developed with propan-1-ol-ethyl acetate-water (7:1:2); this column, like paper sheets, did not resolve sucrose and the reducing disaccharide *O*-β-D-fructofuranosyl-(2→1)-*O*-β-D-fructose (1-fructosylfructose; inulobiose) satisfactorily, but a column of fuller's earth did. The disaccharide fractions from the aqueous effluent of the charcoal-Celite column and from the paper column were bulked and applied to a fuller's earth-Celite column (28 mm. × 125 mm.); this was eluted successively with absolute, 95 and 90% (v/v) ethanol, which removed all the sucrose. The column was washed with 80% (v/v) ethanol and the inulobiose then recovered by elution with water.

The structure of the products of hydrolysis was deduced from the known structure of inulin and the following evidence. (1) A chromatogram of the inulin hydrolysate showed a series of spots each giving a yellow colour with benzidine-trichloroacetic spray, indicating the absence of glucose, and a colour reaction with the 2,5-diphenyl-3-(4-styrylphenyl)tetrazolium chloride reagent, indicating the presence of a reducing fructose group. A plot of $\log[R_F/(1-R_F)]$ against the assumed D.P. gave a straight line (Fig. 2), indicating that they were members of an homologous series. (2) Estimation of the quantity of various spots by the total reducing-sugar and total ketose methods supported the assumed structures (see Table 2).

The reducing disaccharide was named inulobiose by Pazur & Gordon (1953) and the same terminology used and extended to other members of the series, e.g. inulotriose.

Fructosides of raffinose. Analogues in which the fructose chain was terminated by a melibiose group instead of a glucose group were prepared by transfer of fructose to raffinose, mediated by Taka-diastase (Pazur, 1952). Raffinose (final concentration 0.3M) was incubated in 25 ml. of 4 mM-phosphate-citrate buffer, pH 5.5, with 2% (w/v) of Taka-diastase preparation (Parke, Davis and Co. Ltd., Hounslow, Middlesex) at 25° for 21 hr. The digest was then heated in a boiling-water bath for 4 min., filtered while hot, and the volume of the filtrate reduced in a rotary evaporator to about 1 ml. of thick reddish-brown syrup. The syrup was put on to the ChroMax column and eluted with propan-1-ol-ethyl acetate-water (6:1:3); chromatography of samples from the fractions on paper sheets showed that there was considerable overlapping of the sugars present. All the fractions containing sugars with R_F less than that for raffinose were combined, evaporated to dryness and finally collected in a tared beaker; the weight of

Fig. 2. Mobilities of oligosaccharides on chromatograms developed in propan-1-ol-ethyl acetate-water (6:1:3). Results were obtained by measurement of distance moved by the leading edge of each spot. ●, Artichoke series; ○, inulin hydrolysate (glucose-free products).

the sugar was 1.40 g. This material was run again on the column, propan-1-ol-ethyl acetate-water (7:1:2) being used for development: this solvent effected a satisfactory separation of raffinose from the presumed 1^F -fructosyl-raffinose, most of which was separated from a spot of lower R_F [$(1^F\text{-fructosyl})_2$ raffinose]. These oligosaccharides were finally purified by chromatography on paper sheets.

Oligofructosides terminated by hexitols. Analogues terminated by a sugar alcohol instead of a glucose molecule were prepared from individual oligosaccharides (D.P. 2-5) with a terminal reducing fructose (see above) by reduction with sodium borohydride at room temperature for 30 hr. The solutions were poured through charcoal-Celite columns, which were washed with about 5 bed-vol. of water and then with 25% (v/v) ethanol; the latter eluate was evaporated to dryness and further purified by chromatography on paper sheets. The chromatograms showed in each case the appearance of a single new spot with lower R_F than the starting material, and giving a colour reaction with the urea-metaphosphoric acid, but not with the 2,5-diphenyl-3-(4-styrylphenyl)tetrazolium chloride spray.

Total hydrolysis of the trifructosylhexitol by Taka-diastase gave fructose, sorbitol and mannitol only, the hexitols in about equal amounts as estimated visually on paper chromatograms: it is evident that the reducing fructosyl residues of the oligosaccharide were reduced to a mixture of both hexitols, as would occur with free fructose.

Methyl β -D-fructofuranoside. This was prepared by the action of dialysed yeast invertase (British Drug Houses Ltd.; final dilution 1:160) on sucrose (15%, w/v) in 25% (v/v) methanol (Bacon, 1952). The digest was incubated at 20° for 160 min. and then boiled for 4 min. Chromatography of samples from the digest showed the formation of a non-reducing substance containing ketose with R_{Fructose} 1.52 in propan-1-ol-ethyl acetate-water (7:1:2): this compound reached a maximum after about 120 min. The mixture was chromatographed on the ChroMax column, and the methyl β -fructofuranoside run again on a sheet of Whatman 1.6MM paper.

Irisin. This polysaccharide was extracted from a rhizome of *Iris pseudacorus* L. collected near Thornfalcon, Somerset. The rhizome (26 g.) was washed, peeled and sliced, and the slices were extracted with boiling 80% (v/v) ethanol and then three changes of boiling water. The extracts were

combined and a sample was chromatographed; chromatograms sprayed with benzidine-trichloroacetic acid and with urea-metaphosphoric acid showed a large amount of fructose at the origin, and small quantities of sucrose and free fructose. After storage at 2° overnight only a small precipitate had formed; the addition of 4 vol. of ethanol produced a heavy white precipitate which was separated by centrifugation after standing overnight at 2°. This was dissolved in hot water, and the solution centrifuged while hot; the small precipitate was discarded and the solution treated with 4 vol. of ethanol; the precipitated irisin was separated as before and dried in a vacuum desiccator to yield 2.73 g. of white powder, with an anhydrofructose content of 88% (w/w).

Bacterial levan. This was separated from the medium in which a culture of *Bacillus subtilis* had been grown (cf. Dedonder, 1961). A nutrient broth (5 g. of Lab Lemco, 10 g. of peptone, 5 g. of sodium chloride/l., pH 5) was mixed with 0.5 vol. of 30% (w/v) sucrose; the mixture was autoclaved, inoculated with *B. subtilis*, and kept at 37° for 24 hr. Levan was precipitated and purified by the method of Hestrin, Feingold & Avigad (1956). Samples of the precipitate and of the mother liquor were chromatographed in propan-1-ol-ethyl acetate-water (7:1:2) for 2 days; the precipitated material did not move from the origin, and the solution contained principally glucose and fructose, a small amount of sucrose, oligosaccharides with R_F values corresponding to those of *O*- β -D-fructofuranosyl-(2 $\rightarrow$ 6)-*O*-D-glucose (6-fructosylglucose), 1^F -fructosylsucrose and $(1^F\text{-fructosyl})_2$ sucrose, and some material at the origin. The levan was reprecipitated and dried after treatment with several changes of absolute ethanol and then with acetone. The anhydrofructose content of the dried powder was 74% (w/w).

6-Fructosylglucose. This disaccharide was separated from the medium in which *B. subtilis* had been grown (see above). It was present in the fraction soluble in 75% (v/v) ethanol, and was recovered by chromatography on paper. On paper chromatograms it reacted with the benzidine-trichloroacetic acid and urea-metaphosphoric acid sprays, but not with the 2,5-diphenyl-3-(4-styrylphenyl)tetrazolium spray. Estimation of a quantity of the sugar gave values of 367 μ g. with the total fructose method and 378 μ g. with the reducing-sugar method. These results, together with the R_F , were taken as confirmation of its structure.

Table 2. Reducing power of oligosaccharides obtained after partial acid hydrolysis of inulin

		For details of procedure see text.			
Oligosaccharide (assumed constitution)		Reducing power (mg. as free fructose)	Total fructose (mg. as mono- saccharide)	Ratio (c): (b)	Theoretical ratio (from a)
(a)	Preparation*	(b)	(c)		
Inulobiose	A	42.4	87.1	2.06	2
	A†	25.3	45.1	1.78	
	B†	32.0	59.6	1.86	
Inulotriose	A	35.4	90.6	2.56	3
Inulotetraose	A	22.5	88.6	3.94	4
	B	4.48	17.5	3.91	
Inulopentaose	A	14.2	66.5	4.68	5
	B	4.79	23.0	4.81	

* A, 2.30 g. of inulin treated; B, 2.45 g. of inulin treated.

† After removal of sucrose by chromatography on fuller's earth.

RESULTS

Protein preparations were fractionated on DEAE-cellulose columns and the fractions assayed for activity against substrate M (see above) and against sucrose, both in 0.1M-acetate buffer, pH 5. Fig. 3 shows that there was a separation of hydrolase from invertase activity: Fig. 4 shows the elution pattern in an experiment under similar conditions, where the specific activity of the enzyme is calculated from the soluble protein content. The two main fractions of hydrolase activity, A and B, were run again separately on DEAE-cellulose columns under conditions similar to those used previously. The resulting elution diagrams (Fig. 5) show that both fractions remained homogeneous under these conditions. Hydrolases A and B were further characterized with combined fractions corresponding to those indicated in Fig. 3, but collected from different columns of similar dimensions.

Invertase. The fractions containing the greatest activity against sucrose were combined and concentrated in a current of air at room temperature, and the rates of hydrolysis of sucrose, raffinose and melezitose were compared; the results (Table 3) show that the enzyme is a β -fructofuranosidase; this conclusion was supported by the appearance of melibiose and free fructose on chromatograms of

Fig. 3. Chromatography of protein preparation on DEAE-cellulose. The column (1 g.) was eluted with 5 mM-phosphate buffer, pH 7.5, changed stepwise as indicated by the arrows, to 20 mM, 50 mM, 0.1M-NaCl in 50 mM, 0.2M-NaCl in 50 mM, and finally with 0.3M-NaCl in 50 mM-phosphate buffer, pH 8.0. ●, Invertase activity (0.1M-sucrose as substrate); O, hydrolase activity (assayed with substrate M, containing 11.1 mg. of total fructose/ml.). Activity was estimated by method 1.

Fig. 4. Specific activity of hydrolase in fractions eluted from a DEAE-cellulose column. This was a different column from that described in Fig. 3, but the elution pattern was virtually identical: conditions were as described for Fig. 3, except that the final elution with the pH 8.0 solution was omitted. The broken line represents the specific activity of the unfractionated protein.

Fig. 5. Rechromatography of hydrolases A and B on DEAE-cellulose. Conditions were similar to those described for Fig. 3, but 300 mg. columns were used. O, Hydrolase A; ●, hydrolase B.

Table 3. Action of invertase on sucrose, raffinose and melezitose

Each digest contained 0.1M-substrate and 25% (v/v) of invertase preparation, concentrated fourfold; the other conditions were as described for assay of hydrolase activity by method 1.

Substrate	Reducing power (μ g./ml. of digest)		
	1 min.	4 hr.	24 hr.
Sucrose	22	721	4020
Raffinose	74	239	1090
Melezitose	22	46	39

samples from the raffinose digest. Little trans-fructosylation occurred with these substrates, there being only traces of material with R_f lower than that of sucrose or raffinose in the respective digests.

Characteristics of hydrolases A and B

The proteins corresponding to the two peaks A and B were assumed to be different species because of their separation; this was supported by other characteristics.

Effect of pH. After storage overnight at -25° at pH 7.5 the activities of A and B were 104 and 51 % of their original values; subsequently at room temperature the activity of B fell to almost zero within a few hours, but that of A was reduced by only 20 %. Both enzymes were stable at pH 5.0 for long periods at room temperature or at -25° . The effect of pH on the activity of the two enzymes is shown in Fig. 6.

Effect of temperature. Fraction B, when acting upon either oligo- or poly-saccharide, was more labile at high temperature than A: Table 4 gives results of a typical experiment. Table 5 (column headed 10 mM-inulotetraose) shows the effect of temperature upon the rates of reaction of the two activities with oligosaccharide as substrate; aver-

age values of Q_{10} (over the range $3-25^\circ$) were calculated from these data from the relationship

$$\log Q_{10} = \frac{10}{t_2 - t_1} \log \frac{k_2}{k_1}$$

(where t_1 and t_2 are temperatures, and k_1 and k_2 are the rates of reaction at the corresponding temperatures): they were 1.86 for A and 1.73 for B.

Comparison of specificities. Fig. 7 shows on a comparative basis the action of the two enzymes, each upon two series of substrates.

Table 4. *Effect of heat-treatment on activity of hydrolases A and B*

A was heated for 15 min. (AH) and B for 5 min. (BH) at 55° in 5 mM-acetate buffer, pH 5.0, and then cooled rapidly to 0° . Each digest contained 10 mM-substrate (50 mg. of inulin ml.); controls contained untreated enzyme, for B 25 % (v/v) of enzyme preparation. Activity was estimated by method 2.

Substrate	Enzyme	Free fructose liberated (μ m-moles/ml./min.)	Loss of activity from treated enzyme (% of control)
Inulopentaose	AH	11.0	29.8
	A	15.7	—
Inulin	AH	4.88	31.4
	A	7.11	—
Inulopentaose	BH	2.68	87.6
	B	21.9*	—
Inulin	BH	2.09	89.2
	B	16.5*	—

* Corrected to value expected with 50 % (v/v) of enzyme preparation.

Fig. 6. Influence of pH on activity of hydrolases A and B. Buffers were phosphate-citrate; enzyme preparation, 25 % (v/v); activity was assayed with substrate M (11.1 mg. of total fructose/ml.) and estimated by method 1. Activity of both enzymes at pH 5.0 was checked with 0.1 M-acetate buffer. ●, Hydrolase A; ○, hydrolase B.

Fig. 7. Comparison of activity of hydrolases A and B against various substrates. Each digest contained 10 mM-substrate. Activity was estimated by method 2. Hydrolase A: □, natural artichoke oligosaccharides; ■, inulin hydrolysate. Hydrolase B: ○, natural artichoke oligosaccharides; ●, inulin hydrolysate.

Comparison of mode of action. It is known that the primary action of artichoke hydrolase is to release fructose from (2→1′)-linked chains of this sugar (Edelman & Bacon, 1951*a*). Table 1 shows that both fractions A and B were similar in this respect; this Table, together with Fig. 8, is consistent with the concept that single fructose residues are hydrolysed successively from the ends of the chains, releasing the next lower homologue.

Comparative inhibition by sucrose. Action on fructosyl chains by unfractionated preparations of hydrolase is inhibited by sucrose (Recaldin, 1961). Such inhibition occurred with both fractions A and

B, but to different extents, the latter being considerably more susceptible (Table 5). Change in temperature over the physiological range has little effect on the inhibitory action of sucrose.

Further investigation of hydrolase B

Specificity. The action of the enzyme upon the artichoke series of carbohydrates and also upon hydrolysis products of inulin is shown in Fig. 7. Comparative activities upon other compounds containing fructosyl residues are shown in Table 6. The enzyme was inactive against maltose, α′-trehalose, melibiose, cellobiose and lactose.

Substrate concentration. Data plotted by the Lineweaver-Burk method (see for example, Fig. 9) gave the Michaelis constant for inulin as

$$3.3 \times 10^{-2} \pm 0.3 \times 10^{-2} \text{ M}$$

(11 determinations), assuming a molecular weight of 5000. K_m for inulopentase was $2.2 \times 10^{-2} \text{ M}$ (one determination). Determination of K_m for lower members of the artichoke series is complicated by the production of sucrose during the hydrolysis, as sucrose is a powerful inhibitor of the enzyme. As the inhibition of inulin hydrolysis by sucrose is non-competitive (see below) it was assumed that sucrose is a non-competitive inhibitor with respect to the other members of the series. This enabled K_m for 1^F-fructosylsucrose to be estimated by carrying out the reaction at various substrate concentrations in the presence of sucrose at constant concentration: a Lineweaver-Burk plot enabled K_m to be calculated, from the intercept on the 1/*S* axis, to be $3.3 \times 10^{-2} \text{ M}$.

It is known from quantitative analysis of artichoke-tuber extracts that the amounts of the sugars of D.P. 3–7 bear such relationship to one another that the total fructose present in each is the same: judged from paper chromatograms, this relationship appears to extend through most of the series of sugars. The rate of hydrolysis of some of these oligosaccharides was investigated to see if the hydrolase is instrumental in such distribution of fructose residues: oligosaccharides of D.P. 3, 4, 5 and 7 were therefore incubated with the enzyme in

Fig. 8. Action of hydrolase B on inulin and lower oligosaccharides of the natural artichoke series. Each digest contained 10 mm-substrate, as follows: (c) sucrose (D.P. 2); (d) 1^F-fructosylsucrose (D.P. 3); (e) (1^F-fructosyl)₂sucrose (D.P. 4); (f) D.P. 5; (g) D.P. 6; (h) D.P. 7; (i) inulin. Samples were applied to Whatman no. 1 paper, with marker mixtures: (a) raffinose, maltose, sucrose and glucose; (b) artichoke-tuber extract. Developed for 2 days with propan-1-ol-ethyl acetate-water (6:1:3); sprayed with benzidine-trichloroacetic acid. Free fructose is not shown on this chromatogram.

Table 5. *Effect of sucrose on the hydrolysis of inulotetraose by hydrolases A and B at various temperatures*

Activity was estimated by method 2 after 3 hr. incubation, and is expressed as μm-moles/ml./min.

Enzyme	Temp.	10 mm-Inulotetraose	10 mm-Inulotetraose + 10 mm-sucrose	10 mm-Sucrose	Inhibition by sucrose (%)
A	3°	5.56	4.50	0.31	24.5
	16	11.3	9.62	0.28	17.4
	25	17.9	15.0	0.43	17.0
B	3	13.5	5.34	0.31	66.6
	16	26.8	11.7	0.31	59.3
	25	42.0	19.1	0.52	56.2
None	25	1.30	—	0.28	—

Table 6. *Rate of hydrolysis of various sugars by hydrolase B*

Each digest contained 10 mM-substrate; activity was measured by method 2, and the results are compared with the rate at which inulin (50 mg./ml. $\approx$ 10 mM) was attacked.

Sugar	Rate (%)
Inulin	100
Sucrose	2.9
Raffinose	1.2
1 ^F -Fructosylraffinose	43.4
(1 ^F -Fructosyl) ₂ raffinose	57.1
1-Fructosylhexitol	2.2
1-(1 ^F -Fructosyl) ₂ hexitol	26.7
1-(1 ^F -Fructosyl) ₃ hexitol	78.7
O-Methyl β -D-fructofuranoside	3.6
6-Fructosylglucose	2.4
Melezitose	0.6

Table 7. *Action of hydrolase B on various oligosaccharides*

Each digest contained substrates at a concentration of 10 mg. of combined fructose/ml.; * activity was estimated by method 1.

Substrate	Substrate (μ moles/ml.)*	Increase in reducing sugar (μ moles/ml./4 hr.)
1 ^F -Fructosylsucrose	27.8	2.80
(1 ^F -Fructosyl) ₂ sucrose	18.5	1.29
(1 ^F -Fructosyl) ₃ sucrose	13.4	4.16
(1 ^F -Fructosyl) ₅ sucrose	9.3	0.94

* Equivalent to 55.6 μ moles of total fructose.

Table 8. *Effect of various sugars on the hydrolysis of inulin by hydrolase B*

Each digest contained 5 mM-inulin and 10 mM-inhibitor; activity was measured by method 2, and results are corrected for fructose produced (if any) from the inhibitor in control digests without inulin.

Inhibitor	Inhibition (%)
None	0
Sucrose	66.5
1 ^F -Fructosylsucrose	83.5
Melezitose	28.0
Raffinose	25.0
Inulobiose*	57.0
Methyl fructoside	18.3
Sucrose†	42.9
1-Fructosylhexitol†	44.3

* Sucrose-free.

† 5 mM.

such amounts that the concentration of total fructose in each digest was the same, namely 10 mg./ml. Results are given in Table 7.

Inhibitors. Of the sugars tested, those given in Table 8 were found to be inhibitory: under similar conditions, 6-fructosylglucose, melibiose, maltose, $\alpha\alpha'$ -trehalose, cellobiose, lactose, glucose and

fructose were without significant effect (< 10 %). The effect of sucrose upon the enzymic hydrolysis of inulotetraose and inulopentaose was found to be similar to its effect on inulin hydrolysis (Table 9).

When a mixture of 1^F-fructosylsucrose and inulin was incubated with the enzyme, the amount of fructose released was less than the sum of that liberated from the two substrates incubated separately. From the known effect of substrate concentration on rate, this result cannot have been due to saturation of the enzyme in the latter cases. Quantitative estimation of the products from these digests showed that this suppression was almost entirely due to inhibition by the trisaccharide of enzyme action on inulin. Thus Table 10 shows that the amount of sucrose liberated from the trisaccharide alone was almost identical with that liberated from the mixture of both substrates. The calculation included in the Table shows that the trisaccharide was the most inhibitory of all substances tested (cf. Table 8), whereas inulin had practically no inhibitory effect (< 5 %) on hydrolysis of the trisaccharide.

Table 9. *Effect of sucrose on the enzymic hydrolysis of polysaccharide or oligosaccharide*

Each digest contained 10 mM-sucrose, and activity was estimated by method 2. Results are expressed as percentage inhibition compared with control digests without sucrose.

Substrate concn. (mM) ...	5	10
Substrate		
Inulin	58.0	51.4, 59.4
Inulopentaose	56.7	—
Inulotetraose	—	56.2*

* Result from Table 5.

Table 10. *Inhibition by 1^F-fructosylsucrose of enzymic hydrolysis of inulin*

Fructose and sucrose were recovered from paper chromatograms (Whatman no. 1; developed with propan-1-ol-ethyl acetate-water, 7:1:2, for 2 days) of samples of digests described in Table 9, and estimated by the total fructose method. Results are expressed as μ moles/ml. of digest. [Assuming equimolar enzymic production of sucrose and fructose from 1^F-fructosylsucrose, *d* - *c* gives the contaminating sucrose in the trisaccharide preparation = 0.76 μ mole (5.1 %, w/w, of the sugar present). Thus from *f*, and assuming that no sucrose was liberated from inulin (see *b*), the sucrose (and hence the fructose) liberated from 1^F-fructosylsucrose in the presence of inulin = 3.39 μ moles. From *e*, the fructose produced from inulin in the presence of 1^F-fructosylsucrose = 4.03 - 3.39 = 0.64 μ mole. Therefore, from *a*, the inhibition is 79 %.]

Incubation mixture	Fructose	Sucrose
Inulin	3.09 (<i>a</i>)	0 (<i>b</i>)
1 ^F -Fructosylsucrose	3.55 (<i>c</i>)	4.31 (<i>d</i>)
Inulin + 1 ^F -fructosylsucrose	4.03 (<i>e</i>)	4.15 (<i>f</i>)

Fig. 9. Inhibition by sucrose of inulin hydrolysis: Lineweaver-Burk plot. Sucrose, 10 mM when present. Activity was estimated by method 2. O, Inulin; ●, inulin plus sucrose.

Fig. 11. Data of Fig. 10 plotted by the method of Dixon. ●, 5% inulin; O, 1% inulin.

Fig. 10. Sucrose inhibition of hydrolysis of 5% (≈ 10 mM) and 1% (w/v) (≈ 2 mM) inulin: activity was estimated by method 2, and results were corrected by the use of control digests without inulin. O, 5% inulin; ●, 1% inulin.

Experiments were carried out to determine the nature of the inhibition by sucrose. A Lineweaver-Burk plot of results from an experiment in which rate of hydrolysis was determined at various inulin concentrations in the presence and absence of

sucrose is given in Fig. 9. Lines of best fit were calculated from the formula:

$$y = \bar{y} + \frac{xy - x\bar{y}}{x^2 - x\bar{x}}(x - \bar{x})$$

The result indicates that sucrose is a non-competitive inhibitor (K_i 2×10^{-3} M). This conclusion is supported by results from another experiment in which various concentrations of sucrose were used (Fig. 10): a fivefold increase in inulin concentration did not decrease the degree of inhibition at any sucrose concentration tested. When these data are plotted by the method of Dixon (see Dixon & Webb, 1958) the result (Fig. 11) again indicates the non-competitive nature of sucrose inhibition; from this graph K_i is calculated to be 5×10^{-3} M.

Preincubation of enzyme with 10 mM-sucrose for 90 min. at 25° did not increase the inhibition of hydrolysis of 5 mM-inulin (as compared with a control preincubated without sucrose).

DISCUSSION

The work described here shows that artichoke-tuber extracts contain three β -fructofuranosidases. That which hydrolyses sucrose (invertase) was present in very small quantity, requiring prolonged incubation with high sucrose concentration for its detection; it is a β -fructofuranosidase, as it attacked raffinose (liberating melibiose and fructose) but not melezitose.

The other two enzymes (hydrolases A and B) that account for the bulk of the fructosidase activity are remarkable in that they are virtually inactive against sucrose. Both A and B acted on straight-chain oligo- and poly-saccharides containing β -(2 $\rightarrow$ 1')-linked fructosyl residues: they were practically inactive against bacterial levan [β -

(2→6′)-linked] and irisin [mixed β -(2→1′)- and β -(2→6′)-linkages]. In each case heat-treatment lessened the activity towards polysaccharide (inulin) and towards oligosaccharide to the same extent, showing that in both A and B the same enzyme is responsible for hydrolysis of material of high as well as of low D.P. Hydrolases A and B, however, represent at least two distinct enzymes, because they show marked differences in the following properties: (1) mobility on DEAE-cellulose columns at pH 7.5; (2) stability at pH 7.5; (3) stability at 55° (pH 5.0); (4) susceptibility to inhibition by sucrose. They show similar, but not identical, changes in reaction rate with changing pH (Fig. 6).

More detailed comparison of the activities of A and B showed that they both remove single fructose units by multichain attack, as all the reducing power liberated from members of the non-reducing artichoke series was accounted for by increase in free fructose, and (with B) initial action on a single substrate resulted in the appearance of the next lower homologue (Fig. 8). Other fructosidases that are inactive against sucrose have been described, for instance extracellular enzymes produced by *Pseudomonas* sp. (Avigad & Zelikson, 1963) and *Arthrobacter tumescens* (Zelikson & Hestrin, 1961). In other respects, however, these enzymes have little similarity to those of the artichoke: thus they are specific for β -(2→6′)-linked chains of levan; moreover, they do not liberate free fructose, but levanbiose (*Pseudomonas*) or a mixture of oligosaccharides (*Arthrobacter*).

The effect of chain length and of the presence of the terminal glucosyl group on enzyme action was investigated by comparing the natural artichoke series with the series obtained by partial acid hydrolysis of inulin. Fig. 7 shows: (1) that there is a rapid increase in rate of hydrolysis with increasing D.P. of substrate with the lower members of both series; (2) that the presence of the terminal glucosyl residue in general reduces the rate of enzyme action. This Figure shows further differences between the two enzymes: in comparison with the rates at which inulin is attacked, A attacks substrates of low D.P. faster than does B, the relative rates approaching a maximum of 2.5 for A and 1.5 for B. The optimum D.P. is also different for the two enzymes, being clearly about 5 for A, and less sharply defined but apparently about 6–8 for B.

Transfructosylation is known to occur in tuber extracts (Edelman & Bacon, 1951*b*) and the ability of the hydrolase preparations to transfer fructosyl residues from inulin to sucrose was investigated. Hydrolase A was free of this activity, but with B a trace of 1^F-fructosylsucrose was formed; however, the chromatographic properties of the trans-

fructosylase (A. G. Dickerson, personal communication) would allow this to be explained as well by the presence of contaminating amounts of this enzyme in the preparation as by an intrinsic property of hydrolase B.

As the nature of the activity of these two enzymes appeared to be fundamentally similar, only one of them, hydrolase B, was used to characterize the activity more fully.

In common with most hydrolases, K_m of the enzyme is high: values obtained were of the order of $3 \times 10^{-2}M$ for both oligo- and poly-saccharide.

As noted above, the natural series of oligosaccharides terminated by a glucosyl residue is less readily hydrolysed than are those terminated by a reducing fructose group; this difference decreases with increase in chain length until at about D.P. 6–8 the rates are similar. It may be that, at this value, the glucose molecule is too remote to affect the active centre of the enzyme; as the trend with D.P. is similar for both the natural and the glucose-free series, however, the effect of D.P. cannot be ascribed to the presence of the glucosyl (or sucrosyl) residue in the natural series. Furthermore, the inactivity against sucrose could not be accounted for by possible inhibition by the glucosyl moiety of the substrate, nor as merely an expression of its D.P. Thus the enzyme has little activity against the disaccharide 1-fructosylfructose (inulobiose) or its reduction products, the corresponding 1-fructosyl-hexitols. However, although inulobiose is attacked very slowly, it is hydrolysed about four times as fast as is sucrose, showing that the sucrose linkage itself is resistant. The sucrosyl linkage of raffinose was not attacked by the enzyme, showing that increasing the chain length by addition of another hexose residue (galactosyl) is not in itself sufficient to make the sucrose linkage susceptible [the presence of a raffinose end group did not prevent the enzyme acting upon an attached β -(2→1′)-linked fructosyl chain; see below]. We conclude that the enzyme does not attack the sucrose link itself; a similar conclusion can be drawn with regard to the susceptibility of the sucrose linkage to the *Pseudomonas* enzyme, which is reported by Avigad & Zelikson (1963) to produce levanbiose from levantriose slowly but to be inactive against 6^F-fructosylsucrose. Hydrolase B is also inactive against *O*- β -D-fructofuranosyl-(2→6)-D-glucose and *O*-methyl β -D-fructofuranoside.

Inulobiose can be considered to represent the susceptible end of the readily attacked oligosaccharides of both series. The effects of attaching various residues (glucose, fructose, melibiose and hexitol) to inulobiose through its reducing group are shown in Table 11: it is apparent that a considerable increase in rate resulted from the increase in D.P., but that this increase was not the same in

all cases. This Table also shows that the rates of hydrolysis of 1^F-fructosylsucrose (D.P. 3) and 1^F-fructosylraffinose (D.P. 4) are similar, indicating that while the addition of the glucosyl residue increases the effective D.P. of the substrate, further addition of the galactosyl residue does not further augment the rate of hydrolysis. These results together show that although D.P. has a marked effect upon the enzyme action, at least three residues being necessary for significant rate, the enzyme does show specificity for the residues which are attached to a β -(2 $\rightarrow$ 1')-linked fructofuranosyl chain, but which are not necessarily susceptible to its attack. So far we have found that only a β -(2 $\rightarrow$ 1')-link between fructofuranosyl residues at the non-reducing end of a substrate molecule is susceptible to hydrolysis by the enzyme.

Sucrose markedly inhibits the enzymic hydrolysis of inulin and of the two oligosaccharides of D.P. 4 and 5 tested as typical of members of the artichoke series. This effect cannot be ascribed to competition between the fructofuranoside residues of this sugar and those of higher members of the series, as the inhibition is found to be non-competitive, suggesting that the site sensitive to sucrose inhibition on the enzyme is different from the active centre. Raffinose and melezitose were each less inhibitory (to inulin hydrolysis) than was sucrose, but were not very different from one another, suggesting that the inhibition is due to the sucrose molecule as a whole and not to either monosaccharide moiety alone. This was borne out by the effect of various other sugars; thus, although the lack of inhibitory activity of melibiose, maltose, lactose, cellobiose and $\alpha\alpha'$ -trehalose points to the necessity for a fructoside group, this being supported by the activity of fructosylhexitol and 1-fructosylfructose, lack of inhibitory action by methyl fructoside and 6-fructosylglucose, as well as the considerably reduced activity of raffinose (as compared with sucrose) shows that the mere presence of a fructosyl residue is not enough. On the other hand, inhibitory action is slightly enhanced by the addition of another fructose to the sucrose molecule to give 1^F-fructosylsucrose, the most potent inhibitor of inulin hydrolysis yet found among the sugars [Hatch, Sacher & Glasziou (1963) have reported that this trisaccharide in-

hibits an invertase from sugar-cane]; however, as this compound is also a substrate for the enzyme its inhibitory action may be at least partly competitive; this was not investigated further.

Physiological implications

It has been shown (Jefford & Edelman, 1961) that the series of fructosides is converted into sucrose and fructose when artichoke tubers sprout; it is likely that the hydrolases described here are responsible for this depolymerization.

In the tuber, fructose is distributed equally (on a weight basis) among the lower oligosaccharides and by inference among all of the members of the fructosan series (D.P. > 3), and this distribution is maintained when the total carbohydrate content is falling during sprouting (see Fig. 5 in Jefford & Edelman, 1961); this implies that the enzyme systems which act upon members of the series do so in relation to their fructose contents rather than their molarities. However, Fig. 7 and Table 7 show that the rate of action of hydrolase B upon substrates of different D.P. is not related in a simple way to either their molar concentration or to their total fructose contents. In fact, the distribution of fructose among the various oligosaccharides found in the living tissue can be reproduced in a cell-free system by a transfructosylase without hydrolytic activity which can be isolated from artichoke tubers (Edelman & Dickerson, 1963). It is now possible therefore to account for the changes which occur during the depolymerization of the fructosans in the tissue by the concerted action of hydrolase and transferase activities.

As the hydrolases are inhibited by sucrose, and sucrose occurs in the tissue, this sugar may be expected to have some influence over depolymerization in the tuber. The overall concentration of sucrose in the cells ranges from about 0.03 to 0.1 M under various conditions (Table 12), which is high enough to inhibit both hydrolases A and B considerably in free solution.

During storage of the tubers at low temperature ($\sim 2^\circ$) fructosides of high D.P. disappear rapidly (2–4 weeks), but, in contrast with sprouting tubers, there is a concomitant increase in the lower members of the series including sucrose and no substantial change in total sugar content (Jefford & Edelman, 1963); at 20° little depolymerization occurs over long periods (2–3 months) if sprouting is prevented (see also Table 12). This result suggests that, if the control of depolymerization is basically an effect of sucrose upon the hydrolase activity, sucrose is less inhibitory at low temperatures than at high; Table 5 demonstrates that this is not so, and therefore other, as yet unknown, factors must be involved in determining the state of equilibrium of the fructosides at different temper-

Table 11. *Comparative rates of hydrolysis of various oligosaccharides by hydrolase B*

Data from Fig. 7 and Table 6.

1-Fructosylfructose	1.00
(1 ^F -Fructosyl) ₂ fructose	6.7
1 ^F -Fructosylsucrose	6.3
(1 ^F -Fructosyl) ₂ hexitol	3.4
1 ^F -Fructosylraffinose	5.2

Table 12. *Sucrose content of stored artichoke tubers*

Sucrose concentration is expressed as mg. of fructose/g. of water; data are derived from experiments described by Jefford & Edelman (1963).

Period of storage (weeks)	Storage temperature	Concn. of sucrose
0	—	5.6 5.3
7	2°	17.0 14.1
7	20	6.7 9.8

atures. Most of the sucrose will be contained in the vacuole, however, and the effective concentration at the enzyme site is unknown.

Classification

It would seem that hydrolases A and B can both be accommodated either under 3.2.1.7 (inulin 1-fructanohydrolase) or 3.2.1.26 (β -D-fructofuranoside fructohydrolase) in the classification proposed by the International Union of Biochemistry (*Report of the Commission on Enzymes*, 1961); however, most invertases can also be classified in both categories [e.g. yeast invertase will attack β -(2 $\rightarrow$ 1')-linked fructosyl chains] although they appear to be a different group from the artichoke hydrolases. The classification of these enzymes must therefore be considered unsatisfactory.

SUMMARY

1. Three β -fructofuranosidases were separated from a protein preparation of Jerusalem artichoke tubers by chromatography on diethylaminoethyl-cellulose.

2. One of these was characterized as an invertase as it hydrolysed sucrose. The other two (hydrolases A and B) were similar to one another in their overall action: they were practically inactive against sucrose, but attacked a series of straight-chain oligosaccharides [(1^F-fructosyl)_n-sucrose, prepared from artichoke tubers] by hydrolysing terminal fructose residues from the chains; they were practically inactive against bacterial levan [β -(2 $\rightarrow$ 6')-linkages] and against irislin [mixed β -(2 $\rightarrow$ 1')-linkages and (2 $\rightarrow$ 6')-linkages].

3. Hydrolases A and B differed in their physical characteristics, and in their susceptibility to inhibition by sucrose, B being more susceptible than A. They showed small, but definite, differences in their action.

4. Comparison of the action of hydrolases A and B on two series of substrates containing β -(2 $\rightarrow$ 1')-

linkages showed that the rate was largely determined by D.P., being negligible for D.P. 2 (sucrose, inulobiose) and increasing over the range D.P. 3 to about D.P. 6; there was an optimum chain length (undetermined) since oligosaccharides of D.P. 5–8 (the highest tested) were hydrolysed about twice as fast as was inulin (average D.P. about 30); it was shown that each enzyme attacks substrates of high and low D.P.

5. Comparison of the rate of action of hydrolase B on various substrates containing chains of β -(2 $\rightarrow$ 1')-linked fructose residues with various groups attached to the 'reducing' end showed that not all glycosyl residues contribute equally to the effective D.P. of a substrate.

6. Several sugars, notably sucrose, and all containing fructosyl residues, inhibited the activity of hydrolase B against inulin; free fructose was without effect. The nature of sucrose inhibition was investigated in detail and shown to be non-competitive.

7. The physiological significance of these enzymes is discussed. Their probable involvement in the mobilization of fructosans when artichoke tubers sprout has been proposed in an earlier paper. It is shown here that the properties of the hydrolases cannot alone account for the distribution of fructose among various oligosaccharide fractions nor for the depolymerization which occurs when artichoke tubers are stored at low temperature, and so the method of control of these processes is still unexplained.

This programme is supported by a grant from the Department of Scientific and Industrial Research.

REFERENCES

- Avigad, G. & Zelikson, R. (1963). *Bull. Res. Council. Israel*, **11** A, 253.
 Bacon, J. S. D. (1952). *Biochem. J.* **50**, xviii.
 Bacon, J. S. D. (1954). *Biochem. J.* **57**, 320.
 Bacon, J. S. D. (1959). *Biochem. J.* **73**, 507.
 Bacon, J. S. D. (1960). *Bull. Soc. Chim. biol., Paris*, **42**, 1441.
 Bacon, J. S. D. & Bell, D. J. (1948). *Biochem. J.* **42**, 397.
 Bacon, J. S. D. & Edelman, J. (1951). *Biochem. J.* **48**, 114.
 Bacon, J. S. D. & Loxley, R. (1952). *Biochem. J.* **51**, 208.
 Binkley, W. W. (1955). *Advanc. Carbohydr. Chem.* **10**, 55.
 Cifonelli, J. A. & Smith, F. (1954). *Analyt. Chem.* **26**, 1132.
 Dedonder, R. (1961). *Bull. Soc. Chim. biol., Paris*, **42**, 1745.
 Dixon, M. & Webb, E. C. (1958). *Enzymes*, p. 24. London: Longmans, Green and Co. Ltd.
 Edelman, J. & Bacon, J. S. D. (1951a). *Biochem. J.* **49**, 446.
 Edelman, J. & Bacon, J. S. D. (1951b). *Biochem. J.* **49**, 529.
 Edelman, J. & Dickerson, A. G. (1963). *Biochem. J.* **88**, 35f.
 Edelman, J. & Jefford, T. G. (1963). *Biochem. J.* **88**, 36f.

- Edelman, J., Recaladin, D. A. C. L. & Dickerson, A. G. (1963). *Bull. Res. Council. Israel*, **11**A, 275.
- French, D. & Wild, G. M. (1953). *J. Amer. chem. Soc.* **75**, 2612.
- Hatch, M. D., Sacher, J. A. & Glasziou, K. T. (1963). *Plant Physiol.* **38**, 338.
- Haworth, W. N., Hirst, E. L. & Percival, E. G. V. (1932). *J. chem. Soc.* p. 2384.
- Haworth, W. N. & Learner, A. (1928). *J. chem. Soc.* p. 619.
- Hestrin, S., Feingold, D. S. & Avigad, G. (1956). *Biochem. J.* **64**, 340.
- Jefford, T. G. & Edelman, J. (1961). *J. exp. Bot.* **12**, 177.
- Jefford, T. G. & Edelman, J. (1963). *J. exp. Bot.* **14**, 56.
- Nelson, N. (1944). *J. biol. Chem.* **153**, 375.
- Nielsen, H. (1958). *Acta chem. scand.* **12**, 38.
- Pazur, J. H. (1952). *J. biol. Chem.* **199**, 217.
- Pazur, J. H. & Gordon, A. L. (1953). *J. Amer. chem. Soc.* **75**, 3458.
- Quillet, M. (1957). *C.R. Acad. Sci., Paris*, **244**, 2177.
- Recaladin, D. A. C. L. (1961). Ph.D. Thesis: University of London.
- Rees, W. R. & Reynolds, T. (1958). *Nature, Lond.*, **181**, 767.
- Report of the Commission on Enzymes of the International Union of Biochemistry* (1961). Oxford: Pergamon Press Ltd.
- Somogyi, M. (1952). *J. biol. Chem.* **195**, 19.
- Wise, C. S., Dimler, R. J., Davis, H. A. & Rist, C. E. (1955). *Analyt. Chem.* **27**, 33.
- Zelikson, R. & Hestrin, S. (1961). *Biochem. J.* **79**, 71.

Biochem. J. (1964), **93**, 161

Studies on Histones

8. A DEGRADATION PRODUCT OF LYSINE-RICH HISTONE*

By E. W. JOHNS

Chester Beatty Research Institute, Institute of Cancer Research: Royal Cancer Hospital, London, S.W. 3

(Received 11 March 1964)

Previous work from this Laboratory (Johns, 1964) has shown that the lysine-rich histones, f1, can be prepared by extracting calf-thymus nucleoprotein with 5% (0.75*N*) perchloric acid. An acetone precipitation method was also described for the further fractionation of this group which gave a lysine-rich histone (A), with an analysis comparable with that of the fraction A produced by column chromatography by Crampton, Stein & Moore (1957). *N*-Terminal-amino acid analysis showed that this fraction had virtually no free *N*-terminal amino acids. Phillips (1963*a*) has shown that the *N*-terminal amino acids of the lysine-rich histones are probably acetylated.

Another lysine-rich fraction of unusual composition (about 80% of lysine, alanine and proline) reported to be present in calf-thymus histones is the $\alpha 1$ fraction first described by Cruft, Hindley, Mauritzen & Stedman (1957) and later as f1(b) by Johns (1962). The yield of fraction f1(b), however, was small (0–5% of the total yield of f1) and variable, unlike the other histone fractions, and as it had threonine as the main *N*-terminal amino acid it was thought that it might be a product of acid degradation. The lysine-rich histone (A) mentioned above has therefore been degraded with hydrochloric acid and the products were examined. A preliminary account of this work has been given (Johns, 1963).

* Part 7: Johns (1964).

EXPERIMENTAL AND RESULTS

Preparation of histone. Lysine-rich histone (A) (70 mg.), prepared from group f1 by acetone precipitation as described by Johns (1964), was dissolved in 4 ml. of water. Constant-boiling HCl (3 ml.) was then added with stirring. The solution was heated rapidly to 100° and maintained at this temperature for 30 sec. Water (10 ml.) was then added and the solution cooled rapidly to room temperature. Trichloroacetic acid solution (3.0*M*) was then added and the protein precipitated in stages at final concentrations of 0.3*M*, 0.6*M*, 0.9*M*, 1.2*M* and 1.8*M*. The precipitates were washed in 40 ml. of acetone containing 0.1 ml. of conc. HCl, then three times in acetone and dried under vacuum. The amino acid analyses and yields are given in Table 1, together with the analyses of the starting material, fraction $\alpha 1$ prepared by Cruft *et al.* (1957) and f1(b) prepared by Johns (1962).

Amino acid analyses. Total amino acid analyses were carried out as described by Phillips & Johns (1959). *N*-Terminal-amino acid analyses were carried out as described by Phillips (1958) by a modification of the fluorodinitrobenzene method of Sanger (1945).

DISCUSSION

Although the fractions described above are the products of acid hydrolysis of f1 they are of interest because of their bearing on the structure of deoxyribonucleoprotein.

Phillips & Simson (1962) have shown, by examination of peptides obtained from the arginine-rich

Bioch. 1964, **93**