The Separation of Major Components of the Casein of Bovine Milk by Electrophoresis in a Density Gradient

By W. MANSON

The Hannah Dairy Research Institute, Ayr

(Received 17 June 1964)

1. Zonal electrophoresis in a column stabilized by a density gradient has been applied to the small-scale fractionation of the proteins of the casein complex of cow's milk. 2. The α_s - and β -fractions from the milk of individual Ayrshire cows have been shown to behave as single homogeneous proteins on electrophoresis at two pH values in starch gels. 3. The α_s -fraction has been found to be indistinguishable from α_s -casein prepared by Ca^{2+} fractionation of the same milk samples. 4. On the evidence of their electrophoretic behaviour in starch gels and their elementary analyses, α_1 - and α_s -casein are concluded to be substantially the same protein.

The heterogeneous nature of the casein complex of bovine milk was deduced by Mellander (1939) from a study of the behaviour of casein during moving-boundary electrophoresis. Under these conditions there were discernible three well-defined fractions, designated, in order of decreasing mobility, α -, β - and γ -casein. The subsequent separation of these on a preparative scale was, however, achieved not by electrophoresis but by chemical methods involving fractional precipitation (Hipp, Groves, Custer & McMeekin, 1952). Further investigation by chemical means resulted in the subdivision of the α -fraction into α_s - and κ -casein (Waugh & Hippel, 1956), thereby increasing the number of components isolated to four. Since then continuing investigation has in general led to the identification and at least partial characterization of an increasing number of components; until recently it has been suggested that the total number may be as high as 20 (Wake & Baldwin, 1961).

However, the multiplicity of procedures applied to the fractionation of casein and the lack of a unified system of nomenclature makes it probable that certain at least of the components thus prepared differ from each other in name only (Lindqvist, 1963). It is also probable, as Waugh *et al.* (1962) have suggested, that, though certain of the components can reasonably be regarded as primary in the sense that their structures when isolated are unchanged from those originally synthesized within the mammary gland, some of the remainder are more properly considered as secondary products arising by subsequent degradation from these primary caseins. In the view of Waugh *et al.* (1962) the most abundant of the caseins, those of the α_s -, β - and κ -groups, can be regarded as primary products.

Whether or not this proves to be a final estimate, it is apparent that in any procedure designed to fractionate casein into its major components it is desirable to minimize opportunities for the production of secondary products. Such opportunities may be afforded by the presence in milk samples of proteolytic enzymes of bacterial origin, and also by the extended length of time required by many of the methods of isolation involving repeated precipitation. The present work is concerned with attempts to minimize the risks of denaturation from each of these sources by using milk samples from cows selected for freedom from bacterial infection, in conjunction with the technique of zonal electrophoresis in a column stabilized by a density gradient. The use of individual cows had the added advantage that the isolation of single genetic variants of the α_s - and β -groups (Kiddy, Johnston & Thompson, 1964; Aschaffenburg, 1961) became possible. A preliminary separation of components of the casein complex by using this technique has already been reported (Manson, 1962).

MATERIALS AND METHODS

Milk. Milk was obtained from individual Ayrshire cows of the Institute herd, selected as being free from mastitis and in the middle period of lactation. As soon as the samples were taken they were cooled to 4° and stored at this temperature for up to 1 hr. before the fat was removed by centrifugation for 20 min. at 1500 *g* and 15°.

Mixed casein solution. Skimmed milk (300 ml.) was treated at 25° with HCl, initially 1*N* and finally 0.1 *N*, until the pH was 4.6. The precipitated casein was separated by centrifugation for 10 min. at 700 *g* and washed by suspension in distilled water (400 ml.), care being taken to ensure that the pH of the mixture was maintained at 4.6 by the

occasional addition of small amounts of 0.1 N-HCl. The supernatant liquid was decanted and the washing procedure repeated a further five times, after which the casein was suspended in distilled water (100 ml.) adjusted to pH 4.0 with N-acetic acid. This mixture was stirred at room temperature for 5 hr. by the procedure recommended by McMeekin, Hipp & Groves (1959) for the removal of proteolytic enzymes and phosphatase. The washed casein was separated by centrifugation at 700 g, suspended in distilled water (600 ml.) and dissolved by the addition of N-NaOH in such a way that the pH did not rise above 7.0. The resulting opalescent solution of mixed caseins was adjusted in concentration to approx. 1% of protein before storage at -15° .

Preparation of α_1 -casein. α_1 -Casein was prepared from the mixed casein solution by the procedure described by McMeekin *et al.* (1959).

Preparation of α_s -casein. α_s -Casein was prepared from skimmed milk by the adaptation of the method of Waugh & Hippel (1956) described by Manson (1961).

Sucrose solutions for zone electrophoresis. All sucrose solutions used during zonal electrophoresis were derived by suitable dilution from a stock 65% (w/v) solution of sucrose in water, from which much of the ultraviolet-absorbing material present in A.R. sucrose had first been removed. This was achieved by stirring the solution of A.R. sucrose at 55° for 25 min. with 10–15 g. of charcoal/l. and subsequent filtration.

Density-gradient electrophoresis. Zonal electrophoresis in a column stabilized by a density gradient was performed in the apparatus shown diagrammatically in Fig. 1, which differs significantly from that described by Partridge & Elsdén (1961) only in the design of the enlarged right limb of the U-tube. This limb is 7 cm. in diameter and has an effective length of 22 cm.

Fig. 1. Apparatus for zonal electrophoresis in a column stabilized by a density gradient.

The density gradient was confined to that part of the right limb surrounded by the water jacket and was prepared by an adaptation of a procedure described by Manson (1962).

For electrophoresis at pH 7.1, the apparatus was filled initially to within 15 cm. of the tops of the electrode vessels with phosphate buffer, *I* 0.06, and of the following composition (per l.): 1.20 g. of $\text{NaH}_2\text{PO}_4 \cdot 2\text{H}_2\text{O}$; 2.02 g. of Na_2HPO_4 ; 0.58 g. of NaCl. At this stage all taps were closed except those connecting the U-tube with the electrode vessels. A solution of sucrose (50%, w/v) in this buffer was introduced, under a suitable head, into the bend of the U-tube through the tap at the foot of the apparatus, displacing the lighter buffer upwards into the electrode vessels. This process was continued until the bend had been filled and the level of the sucrose solution was approx. 2 cm. below the ball-joint at the lower end of the left limb, when further upward movement was prevented by closing the tap connecting the top of this limb with the left electrode vessel. The sucrose solution was permitted to rise in the right limb until the level of the side arm, A, was reached. The inlet tap in the bend of the U-tube was then closed and the line supplying the sucrose solution disconnected.

The density gradient was formed in the right limb by attaching to the side arm A, a closed mixing chamber containing a magnetic stirrer and a solution of sucrose (3.5%, w/v) in the phosphate buffer (520 ml.). A further solution of sucrose (58%, w/v) in buffer was pumped through this chamber, with continual stirring, at 200 ml./hr., causing buffer containing a smoothly increasing concentration of sucrose to be introduced into the column over the dense sucrose solution occupying the bend of the U-tube. As a consequence, the lighter buffer was expressed from the column through tap C, the tap connecting the U-tube with the right electrode vessel being closed. When all of the light buffer in the right limb had been replaced by sucrose solution, as shown by the appearance in the column of a sharp boundary opposite the lower end of tube B, further effluent from tap C was collected in three fractions (5, 20 and 5 ml.) labelled 1, 2 and 3 respectively. Pumping was then discontinued and tap C closed.

The sample for electrophoresis was prepared by dialysing the mixed casein solution (1.0–1.2% of protein) against the phosphate buffer, pH 7.1, to be used in the electrophoresis. The density of this casein solution (20 ml.; approx. 220 mg. of protein) was adjusted, by the dropwise addition of a solution of sucrose (50%, w/v) in the same buffer, to that of fraction 2 removed from the top of the column. A drop of the protein solution should sink slowly when added to fraction 1 and rise when added to fraction 3.

A small beaker containing the casein solution was placed below tube B so that the tip of the tube extended beneath the surface of the protein solution. With tap C open, liquid was allowed to drip slowly through the tap at the foot of the U-tube, thus sucking the protein solution on to the top of the density gradient where it was deposited as a well-defined layer approx. 1 cm. thick. When the addition was complete the capillary tube B was rinsed with phosphate buffer (2 ml.) in the same fashion. Silver-silver chloride electrodes were placed in the electrode vessels (Partridge & Elsdén, 1961) with the anode in the left vessel. After adjustment of the liquid levels in the electrode vessels, to compensate for the different densities of the contents of the U-tube limbs, the taps connecting the U-tube with the electrode vessels

were cautiously opened. It was arranged that this would cause the protein layer to move about 1 cm. downwards.

After 30 min., during which sucrose diffuses into the initial protein zone, thus stabilizing it, a current of 65 mA (600 v) was applied and electrophoresis continued for 65 hr. During this time the column was cooled by circulating tap water at 11–13° through the jacket. The column was emptied through the side arm *A*, with the tap connecting the left electrode vessel and the U-tube closed, in 10 ml. portions at about 10 ml./min. The protein content of each fraction was determined by measurement of its extinction at 278 m μ .

Density-gradient electrophoresis of casein mixtures was also performed at pH 2.35. The procedure followed for the preparation of the density gradient and for the application of the sample was as described for electrophoresis at pH 7.1, except that HCl-KCl buffer (6.5 ml. of N-HCl and 3.2 g. of KCl/l.), 1.05, was used in place of phosphate buffer. Current (80 mA; 300 v) was passed through the system in which the cathode now occupied the left electrode vessel. After 24 hr. the current was temporarily disconnected and the U-tube isolated from the electrode vessels by closing the connecting taps. The electrode vessels were drained, and refilled to the same liquid levels with saturated NaCl and HCl-KCl buffer. The U-tube was reconnected to the electrode vessels and electrophoresis continued for 24 hr., when the buffer solution was again replaced. After a further 18 hr., electrophoresis was concluded and the column emptied. Protein was again determined by measurement of extinction at 278 m μ .

Starch-gel electrophoresis. The vertical technique described by Smithies (1959) was used with a gel measuring 25 cm. $\times$ 12 cm. $\times$ 0.5 cm. For the preparation of acetic acid gels containing urea (6.5 M) the procedure of Waugh *et al.* (1962) was followed, and tris-citrate gels containing urea (7 M) were prepared according to the method of Wake & Baldwin (1961). With each system electrophoresis was performed at 4° for 16–24 hr. and at a current density of 4–5 mA/cm.² (170–200 v).

Preparation of analytical samples. Approx. 1% solutions of α_1 - and α_2 -casein in 0.1 M-NaCl were adjusted to pH 7 with NaOH and dialysed exhaustively against 0.1 M-NaCl. The concentration of protein was calculated after equal volumes of dialysis residue and diffusate had been dried to constant weight at 105°. For nitrogen and phosphorus analyses portions of the standardized protein solutions thus obtained were used. Specific extinction coefficient determinations were performed in the Unicam SP.500 spectrophotometer by using suitable dilutions of further portions of the same casein solutions.

Chemical analyses. Nitrogen was determined by the micro-Kjeldahl method, phosphorus by the method of Allen (1940) and sialic acid according to the procedure of Svennerholm (1957).

RESULTS

Samples of mixed caseins prepared from the milk of individual Ayrshire cows were subjected to zonal electrophoresis at pH 7.1 in a column stabilized by a density gradient. Invariably, three fractions were obtained resembling the α -, β - and γ -fractions normally resulting from moving-boundary electrophoresis. A typical separation involving approx.

Fig. 2. Density-gradient electrophoresis in phosphate buffer, pH 7.1, of (a) approx. 200 mg. of mixed caseins and of (b) the protein present in fractions 49–62 of the separation shown in (a). The electrophoreses were performed at 65 mA (600 v) for 65 hr.

200 mg. of total protein is shown in Fig. 2(a). The α - and β -casein were isolated from the combined fractions 29–45 and 49–62 respectively by precipitation with N-hydrochloric acid at pH 4.5. Each was redissolved, the α -casein in dilute hydrochloric acid, to give an approx. 1% solution at pH 2.35, and the β -casein in dilute sodium hydroxide yielding a solution of similar concentration at pH 7.1. Electrophoresis of this β -casein, again at pH 7.1, yielded substantially a single fraction (Fig. 2b), although traces of both a slower- and a faster-moving contaminant are apparent. When the α -fraction, adjusted to pH 2.35, was subjected to density-gradient electrophoresis at this pH it exhibited heterogeneity in a manner similar to that reported by McMeekin *et al.* (1959) to occur when their preparation of α -casein was subjected to moving-boundary electrophoresis. The separation obtained is shown in Fig. 3(a).

Repetition of the electrophoresis procedure at pH 2.35 with protein isolated from fractions 31–43 of the separation shown in Fig. 3(a) yielded a single band of material (Fig. 3b). This was tentatively designated α_1 -casein. The protein from fractions 44–56 of the separation shown in Fig. 3(a) on similar treatment also yielded substantially a single band (Fig. 3c), and was identified tentatively as κ -casein. No attempt was, however, made to isolate or further purify the third and slowest-migrating component of the mixture shown in Fig. 3(a). The two-stage

Fig. 3. Density-gradient electrophoresis in HCl-KCl buffer, pH 2.35, of (a) the α -casein- κ -casein complex isolated from fractions 29–45 of the separation shown in Fig. 2(a), of (b) α_1 -casein from fractions 31–43 of the separation illustrated in (a), and of (c) κ -casein from fractions 44–56 of the separation shown in (a). The electrophoreses were performed at 80 mA (300 v) for 64 hr.

separation procedure can therefore be regarded as producing from acid-precipitated casein three major constituents, α_1 -, β - and κ -casein, together with two minor components.

The homogeneity of the α_1 - and β -casein preparations was investigated by starch-gel electrophoresis. In tris-citrate gels, pH 8.7, containing urea (7 M) (Wake & Baldwin, 1961), α_1 -casein prepared by electrophoresis yielded a single sharp band at loadings of between 0.02 and 0.07 mg. per slot. At loadings above 0.1 mg. per slot this single band gradually became diffuse, but on no occasion was more than one band visible. The maximum loading employed was 0.8 mg., when the band had broadened into a zone about 15 mm. long. With 0.02 mg. the single band migrated 100 mm. from the slot, and was about 25 mm. behind the brown front produced in this system. At pH 3.1, in acetic acid gels containing urea (6.5 M) (Waugh *et al.* 1962), this α_1 -casein preparation again appeared homogeneous, only a single sharp band being produced with loadings of 0.05–0.10 mg. of protein per slot. Under the conditions used this single band migrated about 50 mm. in 22 hr.

β -Casein prepared by electrophoresis, when subjected to starch-gel electrophoresis at pH 8.7 in tris-citrate gels containing urea (7 M), yielded a single band at loadings of up to 0.10 mg. per slot. This band was located at approx. 73 mm. from the origin when the brown line marking the front had migrated 130 mm. At pH 3.1, in acetic acid gels containing urea (6.5 M), a single sharp band migrating 40 mm. in 22 hr. was obtained. In neither of these systems was more than a single band visible.

It is apparent that α_1 - and β -casein prepared by density-gradient electrophoresis are each homogeneous entities, at least as judged by their behaviour when subjected to both density-gradient and starch-gel electrophoresis.

The α_1 -casein preparation was further characterized by elementary analysis and by comparison with α_1 -casein prepared by the method of McMeekin *et al.* (1959) and with α_1 -casein prepared by the procedure of Waugh *et al.* (1962). For the purpose of comparison all three caseins were prepared from single samples of milk from individual cows, thus ensuring that any differences in properties of the products would be attributable to the method of preparation.

α_1 -Casein and α_2 -casein prepared by Ca^{2+} precipitation were subjected to electrophoresis in starch gels containing urea at pH 8.7 and 3.1. At each pH value each sample yielded a single band indistinguishable from that produced by the α_1 -casein prepared by density-gradient electrophoresis. In addition, the α_1 - and the α_2 -casein prepared by the method of Waugh *et al.* (1962) each yielded a single band when put through the density-gradient electro-

Table 1. *Comparison of analyses of caseins prepared by electrophoresis and by other methods*

Experimental details are given in the text. Nitrogen and phosphorus are expressed as g./100 g. of substance.

	N	P	$E_{1\text{cm}}^{1\%}$ at 278 m μ
α_s -Casein*	14.8	1.02	10.4
α_s -Casein†	14.7	1.04	10.7
α_1 -Casein	14.7	1.03	10.6
β -Casein*	14.6	0.56	4.8

* Prepared by electrophoresis.

† Prepared by Ca^{2+} fractionation.

phoresis procedure described above, and in this respect also behaved as did the α_s -casein prepared by this method.

All three preparations were analysed for nitrogen, phosphorus and sialic acid content, and their specific extinction coefficients determined at 278 m μ . In no preparation was it possible to detect the presence of sialic acid by the method of Svennerholm (1957). The remaining analytical results are set out in Table 1.

It is apparent from these results that the three preparative procedures yield products virtually indistinguishable from each other on the basis of their nitrogen and phosphorus contents and extinctions at 278 m μ .

The analytical values quoted for α_s -casein prepared by the Ca^{2+} precipitation method are in reasonable agreement with those reported by Waugh *et al.* (1962). However, the nitrogen and phosphorus values for α_1 -casein are not in agreement with those obtained by McMeekin *et al.* (1959). These authors reported a nitrogen content of 14.1% and a phosphorus content of 0.85%.

DISCUSSION

The isolation in a pure state of major components of the casein complex of cow's milk is complicated by the occurrence of genetically controlled variants of two at least of these components, α_s - and β -casein (Kiddy *et al.* 1964; Aschaffenburg, 1961), each group of which is sufficiently closely interrelated to have prevented, to date, their successful fractionation on other than an analytical scale. A fractionation procedure having as its aim the production of single components of the casein complex was considered more likely to be successful if such complications could be excluded at the outset. Such an opportunity was apparently afforded by using milk samples drawn from individual cows of the Ayrshire breed in which the probability is that one variant

only of each group would occur. With α_s -casein this was likely to be the *B*-variant (Kiddy *et al.* 1964).

The existence of further possible complicating conditions can be deduced from results obtained in a study of the effect of mastitis on the composition of milk (Waite & Blackburn, 1963). In the milk of cows thus affected, the portion of the total protein content measured as casein was considerably smaller than normal. The casein can therefore be considered as occurring in company with an increased amount of contaminating proteins. In these circumstances it was considered prudent to select for fractionation experiments milk from cows free from udder infections.

Mixtures of caseins obtained by acid precipitation from the milks of five cows selected in this way, when subjected to zonal electrophoresis in a column stabilized by a density gradient, yielded consistently two minor and three major fractions. One of the two minor components, γ -casein, was probably present in decreased amounts as a result of the repeated washing with water at pH 4.6 accorded the crude acid-precipitated casein, a process reported to dissolve an appreciable proportion of the γ -casein present (El-Negowmy, 1963). The small amounts encountered of the other minor fraction, electrophoretically the slowest-migrating constituent of the α -group, was considered to render its presence of doubtful significance. It was not investigated further.

Of the major fractions, the α_s -casein behaved on electrophoresis in starch gels containing urea (both at pH 8.7 and 3.1) as a single homogeneous protein. It displayed none of the heterogeneity reported by Waugh *et al.* (1962) to occur under the same circumstances with α_s -casein prepared from the milk of Guernsey cows by fractionation with Ca^{2+} .

In an attempt to reproduce the findings of Waugh *et al.* (1962), the Ca^{2+} fractionation procedure described by them was applied to milk samples from portions of which this α_s -casein had been derived by density-gradient electrophoresis. Again, however, the product appeared homogeneous as judged by starch-gel electrophoresis. It is possible that the heterogeneity reported by Waugh *et al.* (1962) had its origins in the breed of cow from which the casein was prepared, and that it was due, in part at least, to the presence of more than one genetic variant of α_s -casein.

Considerable agreement is apparent in the analytical values for α_s -caseins prepared by the two procedures from the same original milk sample, as is shown in Table 1, and it is concluded that under the described conditions a single α_s -casein, free from significant amounts of contaminants, is produced from a single sample of milk, irrespective of which of these two fractionation procedures is followed.

A comparison was also made of the properties and

behaviour of α_s -casein prepared by electrophoresis and of α_1 -casein prepared from the same original milk sample by the procedure of McMeekin *et al.* (1959). From the analytical results in Table 1 and from their behaviour on electrophoresis in starch gels it is apparent that α_s - and α_1 -casein prepared from a single milk sample are substantially the same protein. The lack of agreement between the analytical values reported above for α_1 -casein and those reported by McMeekin *et al.* (1959) for their preparation does, however, probably reflect real differences between the two products, which may be satisfactorily explained by reference to the different sources from which they were derived.

The density-gradient fractionation yielded a second major product, β -casein, which also behaved as a homogeneous protein. The analytical results obtained for this preparation are in agreement with those of Garnier, Ribadeau-Dumas & Mocquot (1964).

Though both the α_s - and β -casein can be obtained in a pure state and in about 100 mg. quantities from one milk sample by this fractionation procedure, for the third major component, κ -casein, to be produced in similar quantities the yields from more than one fractionation would require to be combined, thus necessitating the use of more than one sample for each preparation. This was considered undesirable owing to possible variation between samples, and for this reason characterization of the κ -fraction was not proceeded with.

I am greatly indebted to Dr S. M. Partridge for his assistance with the design of the apparatus for density-gradient electrophoresis.

REFERENCES

- Allen, R. J. (1940). *Biochem. J.* **34**, 860.
Aschaffenburg, R. (1961). *Nature, Lond.*, **192**, 431.
El-Negowmy, A. M. (1963). *J. Dairy Sci.* **46**, 768.
Garnier, J., Ribadeau-Dumas, B. & Mocquot, G. (1964). *J. Dairy Res.* **31**, 131.
Hipp, N. J., Groves, M. L., Custer, J. H. & McMeekin, T. L. (1952). *J. Dairy Sci.* **35**, 272.
Kiddy, C. A., Johnston, J. O. & Thompson, M. P. (1964). *J. Dairy Sci.* **47**, 147.
Lindqvist, B. (1963). *Dairy Sci. Abstr.* **25**, 257.
McMeekin, T. L., Hipp, N. J. & Groves, M. L. (1959). *Arch. Biochem. Biophys.* **83**, 35.
Manson, W. (1961). *Arch. Biochem. Biophys.* **95**, 336.
Manson, W. (1962). *J. electroanalyt. Chem.* **3**, 203.
Mellander, O. (1939). *Biochem. Z.* **300**, 240.
Partridge, S. M. & Elsdon, D. F. (1961). *Biochem. J.* **79**, 26.
Smithies, O. (1959). *Biochem. J.* **71**, 585.
Svennerholm, L. (1957). *Biochim. biophys. Acta*, **24**, 604.
Waite, R. & Blackburn, P. S. (1963). *J. Dairy Res.* **30**, 23.
Wake, R. G. & Baldwin, R. L. (1961). *Biochim. biophys. Acta*, **47**, 225.
Waugh, D. F. & Hippel, P. H. von (1956). *J. Amer. chem. Soc.* **78**, 4576.
Waugh, D. F., Ludwig, M. L., Gillespie, J. M., Melton, B., Foley, M. & Kleiner, E. S. (1962). *J. Amer. chem. Soc.* **84**, 4929.