The Alcohol Dehydrogenase Gene Is Nested in the *outspread* Locus of *Drosophila melanogaster*

S. McNabb, S. Greig and T. Davis¹

Department of Genetics, University of Cambridge, Cambridge CB2 3EH, England

Manuscript received December 30, 1995

Accepted for publication March 11, 1996

ABSTRACT

This report describes the structure and expression of the *outspread* (*osp*) gene of *Drosophila melanogaster*. Previous work showed that chromosomal breakpoints associated with mutations of the *osp* locus map to both sides of the alcohol dehydrogenase gene (*Adh*), suggesting that *Adh* and the adjacent gene *Adh'* are nested in *osp*. We extended a chromosomal walk and mapped additional *osp* mutations to define the maximum molecular limit of *osp* as 119 kb. We identified a 6-kb transcript that hybridizes to *osp* region DNA and is altered or absent in *osp* mutants. Accumulation of this RNA peaks during embryonic and pupal periods. The *osp* cDNAs comprise two distinct classes based on alternative splicing patterns. The 5' end of the longest cDNA was extended by PCR amplification. When hybridized to the *osp* walk, the 5' extension verifies that *Adh* and *Adh'* are nested in *osp* and shows that *osp* has a transcription unit of ≥ 74 kb. *In situ* hybridization shows that *osp* is expressed both maternally and zygotically. In the ovary, *osp* is transcribed in nurse cells and localized in the oocyte. In embryos, expression is most abundant in the developing visceral and somatic musculature.

THE *outspread* (*osp*) locus of *Drosophila melanogaster* was identified initially by an EMS-induced recessive mutation, *osp*¹, which resulted in the wings of adult flies being spread out laterally from the long axis of the body (E. H. GRELL, unpublished data). It was first referred to in a publication by AARON (1979) as a mutation associated with some mutations of the alcohol dehydrogenase (*Adh*) gene. Additional *osp* mutations have been induced by a variety of mutagens, 14 by EMS, 31 by X- or gamma ray, one by diepoxyoctane and two by formaldehyde (O'DONNELL *et al.* 1977; WOODRUFF and ASHBURNER 1979; ASHBURNER *et al.* 1982; CHIA *et al.* 1985). All fall into a simple genetic complementation group and result in essentially the same wing phenotype as *osp*¹. There is some allele-specific variation in the angle at which the wings are held out from the body and whether or not the wings curve back toward the body, or tent. Complete loss of *osp* function (*e.g.*, heteroallelic deficiency) results in a strong *osp* phenotype; however, the flies still are viable and fertile in both sexes.

Although deletions of *osp* are often mutant for *Adh*, the two loci can be mutated separately. *osp* was mapped just distal to *Adh* on the basis of recombination between *osp*^{76e} and *Adh*ⁿ², both presumed point mutations. However, CHIA *et al.* (1985) molecularly mapped five *osp* breakpoints distal to and two proximal to *Adh*. By this

criterion, *Adh* maps within *osp*. In addition *Adh'*, a gene of unknown function but related to *Adh* in sequence and structure, maps within *osp* by the same criterion (referred to as *Adh-dup* by SCHAEFER and AQUADRO 1987; KREITMAN and HUDSON 1991).

Working from the hypothesis that *Adh* is nested within the *osp* locus and that *osp* is a large gene, we undertook molecular studies to identify the limits and organization of the *osp* locus. We extended a chromosome walk in the region and mapped additional *osp* breakpoints to it. We localized a genomic fragment that encoded a portion of the *osp* transcript and used that fragment to characterize *osp* transcription and to isolate *osp* cDNAs. The longest cDNA, extended by PCR and hybridized back to the chromosomal walk, confirmed that *Adh* and *Adh'* are indeed nested in the *osp* locus. This also demonstrated that *osp* is a large gene with a minimum of four exons spanning at least 74 kb.

The finding that *Adh* and *Adh'* are nested within *osp* raised the issue of whether this organization has an effect on the regulation of transcription of the nested genes. To address this issue, the temporal expression and tissue distribution of the *osp* transcript was characterized. The *osp* gene produces a polyadenylated transcript of ~6 kb that is transcribed during all developmental stages with peaks of expression during embryonic and pupal periods. In the embryo, *osp* is present as a maternal product that is anteriorly localized. Its dynamic zygotic expression shows early elements of pair rule control and later is most abundant in the visceral and somatic musculature. The peaks of *osp* expression do not coincide with those of *Adh* nor are the two expressed in the same tissues in embryos.

Corresponding author: Susan McNabb, Department of Zoology, Box 351800, University of Washington, Seattle, WA 98195.

¹ Present address: Academic Unit of Medical and Community Genetics, Kennedy Galton Centre Level 8V, Northwick Park and St. Mark's NHS Trust, Watford Road, Harrow HA1 3UJ, United Kingdom.

TABLE 1
Chromosomes used in this study grouped by progenitor

Aberration	Previous name	Breakpoints	Molecular ^a
A. On <i>b Adh^f cn bw</i>			
<i>Df(2L)A47</i>		34E1;35B2	Proximal to +108 ^b
<i>Df(2L)A48</i>		35B2-3;35D5-7	0 to +0.7 ^c
<i>Df(2L)A63</i>		Not visible	-13.1 to -18.2 ^b ; proximal to +108 ^b
<i>Df(2L)A72</i>		35B2-3;35B7-8	+0.7 to +1.3 ^c
<i>Df(2L)A178</i>		35B1-2;35B1-2	-110.8 to -113 ^c ; +76.5 to +78.5 ^b
<i>Df(2L)A217</i>		34F5;35B3	proximal to +108 ^b
<i>Df(2L)A220</i>		35B1-2;35B9	-24.6 to -28.4 ^c
<i>Df(2L)A245</i>		34A4;35B2	-137.8 to -134.6 ^d ; proximal to +90 ^b
<i>Df(2L)A260</i>		35B1-2;35B1-2	~-144.4 ^d ; proximal to +90 ^b
<i>Df(2L)A266</i>		35B2-3;35B2-3	~-144.4 ^d ; proximal to +90 ^b
<i>Df(2L)A379</i>		35B1;35B3;40-41;57A8-10 ^e	-61.8 to -66.2 ^c ; +84.5 ^b
<i>Df(2L)A445</i>		Not visible	-73 ^c ; proximal to +90 ^b
<i>Df(2L)fn52</i>		Not visible	-58 ^c ; proximal to +108 ^b
<i>Df(2L)noc10</i>		34F1-2;35B1-2	Proximal to +90 ^b
<i>Df(2L)noc13</i>		35A1-2;35B2	Proximal to +90 ^b
B. On <i>Adh^f pr pk cn sp</i>			
<i>Df(2L)b84a1</i>		34D3;35B1-2	Proximal to +108 ^b
<i>Df(2L)b84a4</i>		34D3;35B1-2	-37.5 to -42.5 ^c
<i>Df(2L)b84a7</i>		34C1-2;35B1-2	Not determined
<i>Df(2L)b84a8</i>		34D3;35B1-2	-37 ^c
C. On <i>In(2LR)O, Cy dp³ pr cn</i>			
<i>Df(2L)noc20</i>		34F1-2;35B2	+11.8 to +13.7 ^c
D. On <i>al dp b noc^{TE35B} pr 1(2)pwn cn^f</i>			
<i>Df(2L)TE35B-8</i>	<i>Df(2L)TE146(Z)GW8</i>	34E4-5;35A2-4	-11.2 to -15.9 ^c
<i>Df(2L)TE35B-9</i>	<i>Df(2L)TE146(Z)GW9</i>	34F1;35B2	Proximal to +108 ^b
<i>Df(2L)TE35B-11</i>	<i>Df(2L)TE146(Z)GW11</i>	Not visible	-107.5 to proximal to +50 ^c
<i>Df(2L)TE35B-15</i>	<i>Df(2L)TE146(Z)GW15</i>	34F3;35B2	Proximal to +108 ^b
<i>Df(2L)TE35B-48</i>	<i>Df(2L)TE146(Z)SR48</i>	35B1-2;35B1-2	+51.8 to +57.3 ^b
<i>Df(2L)TE35B-54</i>	<i>Df(2L)TE146(Z)SR54</i>	35B1-2;35C1	-107.5; +11.7 to +30.0 ^c
<i>Df(2L)GT4</i>		34F3;35B2	-11.2 to -13.3 ^c
E. On <i>Adh^D pr cn</i>			
<i>Df(2L)fn2</i>		35A3;35B2	Proximal to +108 ^b
<i>Df(2L)fn7</i>		34E1-2;35B3-5	Proximal to +108 ^b
<i>Df(2L)fn36</i>		35A3;35B4	Proximal to +108 ^b
F. On <i>Adh^{W3} pr cn</i>			
<i>Tp(2;3)osp³</i>		35B3-4;36C11-D1;98E2-3 ^c	+39.6 to +44.2 ^b
<i>Df(2L)osp18</i>		35B1-2;35C4-5	-37.5 ^c
<i>In(2L)osp²²</i>		35B3;38D3-5	-1.7 to -4.1 ^c
<i>Df(2L)osp29</i>		35B3;35E6	+36.9 ^c
<i>In(2L)osp⁵⁹</i>		35B2-3;38B3-6	-1.7 to -4.1 ^c
<i>T(2;3)osp⁹⁰</i>		35B3-4;89B9-11	+14 to +18 ^c
<i>Df(2L)osp144</i>		Not visible	-67.1 to -68.3; -8.3 ^c
<i>Df(2L)osp201</i>		Not visible	-8.3 to -18.2 (1-kb deletion) ^c
G. On <i>ck^{TE35BC} pr pk cn sp^h</i>			
<i>Df(2L)TE35BC-4</i>	<i>Df(2L)TE36(R)GW4</i>	35B3-4;35C3	Proximal to +108 ^b
<i>Df(2L)TE35BC-28</i>	<i>Df(2L)TE36(R)GW28</i>	35B2;35B7	Proximal to +108 ^b
<i>Df(2L)TE35BC-35</i>	<i>Df(2L)TE36(R)GW35</i>	35B2;35D4	Proximal to +108 ^b

TABLE 1

Continued

Aberration	Previous name	Breakpoints	Molecular ^a
G. On <i>ck^{TE35BC}pr pk cn sp^h</i> (continued)			
<i>Df(2L)TE35BC-3^D</i>	<i>Df(2L)TE36(R)GV3^{2D}</i>	<i>Df(2L)35B5;35D1-2</i>	Proximal to +108 ^b
<i>Sco^{rv13P}</i>	<i>ScoR + 13^{2P}</i>	+T(2;3)35B;81	
<i>Df(2L)TE35BC-3</i>	<i>T(2;3)TE36(R)GV3</i>	35B5-10;81	Proximal to +108 ^b
H. No progenitor chromosomes available			
<i>Df(2L)el77</i>		35A1-3;35B3	+44.2 to +51.4 ^b
<i>Df(2L)W</i>		35A2-3;35B3-5	Proximal to +108 ^b
<i>In(2L)C158</i>		26D1-2;35B3 ^c	+100.4 to +105.4 ^b
<i>T(2;3)H47</i>		35AB;65F	+36 to +46 ^b
<i>T(2;3)Mpe</i>		35B2.3;86C1-2	-0.3 to -2.8 ^c
<i>T(2;3)pb³</i>		T(2;3)35B3;83E2-8 + In(3R)83E2-8;89A9-10 + T(2;3)50C1-4;80 ⁱ	+36.7 to +40.7 ^b

^a Molecular limits of breakpoints in the *Adh* region according to the coordinates of CHIA *et al.* (1985).^b This study.^c CHIA *et al.* (1985).^d DAVIS *et al.* (1990).^e New order 21-35B3/57A8.10-40.41/35B1.3-40/57A8.10-60F.^f Previously *ab dp b noc^{TE146(2)}pr l(2)pwn cn*.^g GUBB *et al.* (1990).^h Previously *ck^{TE36(R)}pr pk cn sp*.ⁱ New order 21A-35B3/83E2.8-89A9.10/83E2.8-80/50C14-60F;61A-80/50C14-35B3/89A9.10-100F.

However, both *osp* and *Adh* transcripts are expressed in the adult female ovary and are maternally transmitted to oocytes.

The nested relationship between *osp* and *Adh* may be significant with respect to the evolution of the two genes. A comparative analysis of known nested genes suggests that the introns of large genes may provide attractive sites in which small intronic genes can become established, be expressed, and evolve. The extensive research that has been done on *Adh* evolution can now be extended to a new framework, the evolution of nested genes.

MATERIALS AND METHODS

Drosophila stocks: Stocks used in this study are listed in Table 1. The A series of deficiency stocks [e.g., *Df(2L)A47*, *Df(2L)A48*] are X-ray-induced *Adh* null mutations (AARON 1979; ASHBURNER *et al.* 1982). The *fn* series are formaldehyde-induced *Adh* null mutations: *Df(2L)fn52* was originally obtained from W. SOFER; the rest were described by O'DONNELL *et al.* (1977) and ASHBURNER *et al.* (1982). *Df(2L)noc10* and *Df(2L)noc13* were isolated by G. HARRINGTON and M. ASHBURNER (Flybase). The *b* series of deficiencies were described by CHIA *et al.* (1985), except for *Df(2L)b84a7* (GUBB *et al.* 1988). The *Df(2L)TE35B* series was described by GUBB *et al.* (1985, 1986). The *Df(2L)TE35BC* series was described by GUBB *et al.* (1984). The *osp* aberrations were as described in CHIA *et al.* (1985). *Df(2L)el77* and *Df(2L)W* were described by WOODRUFF and ASHBURNER (1979), *In(2L)C158* by ASHBURNER and LEMEUNIER (1976), *T(2;3)H47* by HILLIKER and TRUSIS-COULTER (1987), *T(2;3)Mpe* by HUGHES and SHELTON (1980) and *T(2;3)pb³* by KAUFMAN (1978). Genetic analysis of the mutations is shown in Figure 1.

Screening of genomic libraries, mapping breakpoints by Southern analysis and *in situ* hybridization to chromosomes:

These procedures are as described in DAVIS *et al.* (1990) with the following exceptions. In this study the genomic libraries used were: a Canton S library (MANIATIS *et al.* 1978), Canton S and Oregon R libraries in λ EMBL4 (provided by M. GOLDBERG), a library constructed from a stock carrying markers *y; cn bw sp* and isogenic for all four chromosomes in λ EMBL3 (made by J. TAMKUN and J. KENNISON; provided by J. TAMKUN), and a library made from the stock *Df(2L)DC17-12/In(2-LR)O*, *Cy dp^{h1}pr cn² (CyO)* (a gift from D. GUBB).

To simplify the mapping of breakpoints by restriction enzymes to a single homologue, many chromosomes were used in heterozygous combinations with *Df(2L)A48*, a deletion with its distal breakpoint at 0 to +0.3 on the chromosomal walk (CHIA *et al.* 1985) and its proximal breakpoint ≥ 19 loci proximal to the *Adh* locus (ASHBURNER *et al.* 1982). Lethal derivatives of *TE35BC* (GUBB *et al.* 1984), *Tp(2;3)osp³*, *T(2;3)pb³* and *In(2L)C158* were balanced over *In(2LR)CyO*, *Cy dp^{h1}pr cn²*. Other aberrations were induced on known progenitor chromosomes (Table 1), thus changes in hybridization pattern between aberrations and their precursors were judged by direct comparison.

The position of all breakpoints that are not deletions have been confirmed by *in situ* hybridization to polytene chromosomes.

RNA isolation and Northern analysis: Total RNA was isolated from staged collections from a Canton S wild-type strain. RNA was extracted using guanidinium thiocyanate (CHOM-CYNSKI and SACCHI 1987), denatured and run in formaldehyde agarose gels (MANIATIS *et al.* 1982), then transferred to nylon filters (Amersham) following the manufacturer's instructions. For the analysis of *osp* mutants, RNA was treated in the same fashion except that it was blotted onto nylon-reinforced nitrocellulose (Schleicher and Schuell) using the Amersham protocol for transfer to nitrocellulose.

Filters were probed with randomly primed DNA fragments

FIGURE 1.—Correlating the genetic and molecular maps of *osp* region aberrations. The relative location of genetic loci *outspread* (*osp*), Alcohol dehydrogenase (*Adh*) and neighboring loci *no-ocelli* (*noc*) and *l(2)35Bb* are indicated by the heavy horizontal bars at bottom. The genetic breakpoints of aberrations associated with *osp* are indicated along the vertical lines. Aberrations are deficiencies unless otherwise indicated as inversions (*In*), translocations (*T*), or transpositions (*Tp*). Deficiencies that lie entirely within the region are represented by horizontal lines that terminate at vertical lines; those that extend beyond the boundaries of the genes extend beyond the vertical lines. Breakpoints crossed by DNA in this and previous studies (see Table 1) are indicated by double vertical lines.

(FEINBERG and VOGELSTEIN 1983) from portions of the genomic walk that correspond to *osp* lesions (see above) or were identified by reverse Northern analysis as possible transcription units (G. AL-ATIA; B. MENG, personal communications). These were: λ ob2-3.8, λ ob1.12-3.8, λ ob1.12-4.7, λ gAC3-2.4, λ W2-10 and λ W5.02-2.9 and -4.6-kb fragments (Figure 2). The RNA size markers were from Bethesda Research Laboratories.

Polyadenylated RNA was selected on oligo-dT cellulose columns (Collaborative Research) as described in MANIATIS *et al.* (1982).

osp mutant chromosomes were normally maintained over balancer chromosomes wild type for *osp*. To analyze the *osp* RNA encoded by the mutant chromosomes, mutant alleles were recovered as heterozygotes with *Df(2L)A178*, which is deleted for most of the *osp* region and is null for *osp* expression (Figure 6). RNA was extracted from heterozygous flies scored as adults for the *osp* wing phenotype. Blots of total RNA, 20

μ g per lane, were probed with the 4.7-kb *EcoRI* fragment from λ ob1.12 (referred to subsequently as ob1.4.7).

Isolation and characterization of cDNAs: Aliquots of a 0–3-hr embryonic cDNA library in λ gt10 (POOLE *et al.* 1985) was screened using standard techniques (MANIATIS *et al.* 1982). An aliquot of the 8–12-hr embryonic cDNA library in the plasmid vector pNB40 (BROWN and KAFATOS 1988) was screened as described in that paper using ob1.4.7 as a probe.

cDNAs were verified as corresponding to the *osp* transcript by hybridization to the expected 6-kb RNA on Northern blots (as above) and by hybridization to Southern blots of the walk phage DNA. cDNAs were first mapped by hybridization to bidirectional Southern transfers (SMITH and SUMMERS 1980) of *EcoRI*-digested DNA from phage which spanned the entire walk region. The phage that hybridized were mapped in greater detail using Southern blots of doubly digested phage DNA.

Partial sequences of the cDNAs were obtained by double strand sequencing using a US Biochemicals sequencing kit,

FIGURE 2.—Molecular map of the *osp* walk. This figure summarizes the restriction map of phage from a chromosomal walk which begins with the map of CHIA *et al.* (1985) and extends 70 kb proximal to λ W4.04. Map coordinates are given on the dark upper line. Restriction maps for *EcoRI*, *XhoI* and the *HindIII* sites used in this study are shown below the coordinates. Genomic and phage fragments referred to in the text are indicated by size of fragment. All fragments are generated by *EcoRI* except for X7.7 and X7.2, which are *XhoI* fragments. The triangle indicates a 5.2-kb insertion (see legend for Figure 3). The position of a restriction site polymorphism is indicated by a P. The phage inserts are represented by the descending horizontal lines; asterisks indicate those from CHIA *et al.* (1985). The origin of the map coordinates is the *EcoRI* site 1321 bp distal to the start of transcription of the proximal *Adh* transcript (BENYAJATI *et al.* 1983; CHIA *et al.* 1985).

essentially as described by the manufacturer. To prepare the plasmids, 2-ml overnight cultures were prepared by the lysozyme boiling technique (HOLMES and QUIGLEY 1981) with these alterations. After boiling and pelleting, supernatants were extracted twice with equal volumes of phenol/chloroform, then ethanol precipitated in 0.3 M sodium acetate, pH 5.2. Samples were treated with RNase A and denatured before use in sequencing reactions.

PCR amplification beyond the 5' end of the longest *osp* cDNA: A 12–24-hr embryonic cDNA plasmid library, which contains directed inserts (BROWN and KAFATOS 1988), was used as a template for PCR amplification to obtain the 5' portion of the *osp* cDNA. The 3' primer was the reverse complement of sequences at the 5' end of 9114, the *osp* cDNA that extended furthest 5'. To select an appropriate 9114-internal primer, several hundred bases of the 5' end of the insert were sequenced from the SP6 promoter. A 20-mer, 5'-ATCGCGA-GAGCTGATTGCGT-3', located ~200 bp from the cloning site was chosen as a 3' primer. A second internal sequence was selected for verification of the PCR product. The 5' primer was the SP6 promoter site in the cloning vector. Samples were initially denatured at 95° for 5 min, annealed at 55° for 2 min, extended at 72° for 2 min, then denatured at 94° for 40 s for 25 cycles. Reactions contained 100 ng library, primers at 0.1 μ M each, 2.5 mM $MgCl_2$ and 200 μ M dNTPs in a 50- μ l reaction volume. Cocktails were layered over with mineral oil, denatured, then 0.3 μ l Taq polymerase (Cetus) was added. Primers were synthesized in the Department of

Biochemistry, University of Cambridge, Taq polymerase was obtained from Cetus and dNTPs were obtained from BRL.

The major PCR product of 1.5 kb was verified as corresponding to an *osp* cDNA by hybridizing to the 6-kb RNA and by hybridizing to Southern blots of walk phage DNA as described above for the analysis of cDNAs. The 1.5-kb band was purified and one fraction was labeled for use as the probe by random priming (as above). The other was end-filled and cloned into pBluescript KS⁺ (Stratagene) for further analysis. Several clones, including *posp5'*-319, were partially sequenced and verified that the contents of the PCR band were homogeneous and overlapped 9114. Clone *posp5'*-319 was also used to probe chromosomes *in situ* to demonstrate that it mapped to its site of origin at 35B.

***In situ* hybridization to whole mounts:** Whole embryos aged 0–24 hr and adult ovaries were collected and prepared as described by TAUTZ and PFEIFLE (1989), except that ovaries were fixed in the absence of heptane and, after staining, preparations were dehydrated and mounted in Permount (BDH Chemicals). cDNAs 9114, 7131 and 7111 were each tested as hybridization probes. The three produced identical hybridization patterns but 9114 gave the strongest signal, and thus was used for all subsequent experiments.

RESULTS

The molecular map of the *outspread* region: This study extends the work of CHIA *et al.* (1985) on the

FIGURE 3.—Summary: molecular map of *osp* mutations. The molecular extent of the *osp* locus, as determined by Southern mapping of *osp* mutations, is indicated by the thick bar at the top. The outer limits of *osp* are defined on the distal side by the *noc⁻osp⁺* mutation *Df(2L)GT4* and *noc⁻osp⁻* *Df(2L)TE35B-8*. The aberrations that define the proximal side of *osp* are the *osp⁻* mutations, *Tp(2;3)osp³*, *T(2;3)pb³* and *T(2;3)H47*, and *osp⁺* *In(2L)C158*. Mutations were mapped to *EcoRI* fragments; error bars indicate the uncertainty of the location of the breakpoint within the *EcoRI* fragment. There is a large error bar for the proximal side of *osp* due to the lack of diagnostic mutations between *Tp(2;3)osp³*, *T(2;3)pb³* and *T(2;3)H47*, and *In(2L)C158*. Interstitial deficiencies and fragments within which inversions break are represented by dashed bars. Deficiencies that extend beyond the region shown here are indicated by lines ending in arrowheads. *Df(2L)A379* is a deficiency associated with an inversion, and is represented by a horizontal deficiency line and a vertical arrow for the inversion breakpoint. Aberrations mapped by others are (a) CHIA *et al.* (1985) and (b) GUBB *et al.* (1990). *T(2;3)H47* was diagnosed by *in situ* hybridization of λ W5.02 to chromosomes. *Adh* and *Adh^r* are indicated by arrows, which show their transcriptional orientation.

molecular mapping of the *Adh* region of the *D. melanogaster* genome. CHIA *et al.* (1985) cloned the distal portion of the *osp* locus as part of a 164-kb genomic walk in overlapping phage, ending 40 kb proximal to the *Adh* locus with λ W4.04. By Southern mapping genomic DNA from *osp* mutants, they found five *osp* lesions distal to and two proximal to *Adh*, in addition to 21 aberrations that spanned the *Adh* region. In this study, we have extended the walk to 110 kb proximal to *Adh* (Figure 2) and have mapped additional *osp* mutations and chromosome variants to define the outer limits of the *osp* locus.

CHIA *et al.* (1985) defined the distal limit of *osp* as lying between -11.2 and -37 kb on the molecular map. This limit was based on their mapping of the proximal breakpoints of the *noc⁻osp⁺* deficiency *Df(2L)b84a8* at -37 kb and the *noc⁻osp⁻* deficiency *TE35B-8*, which ends in the 4.7-kb *EcoRI* fragment from λ ob1.12, at -11.2 to -15.9. GUBB *et al.* (1990) redefined the distal end of *osp* by mapping a new *noc⁻osp⁺* deficiency, *Df(2L)GT4*, which breaks between -11.2 and -13.3 and extends distally (Figure 3). The distal limit is now defined as lying between -11.2 and -13.3 kb.

Four *osp* deficiencies were mapped to the interval +40.0 to +105.4 kb: *Df(2L)el⁷⁷*, *Df(2L)TE35B-48*, *Df(2L)-A178* and *Df(2L)A379*, *In(2L)A379*. Genomic DNA from each line was digested with *EcoRI* and hybridized with the following probes. *Df(2L)el⁷⁷/Df(2L)A48* DNA, probed with the 7.2-kb *XhoI* fragment of λ W7.012 (Figure 2), yielded the 6.4-kb parental *EcoRI* fragment and a fusion fragment of 9.7 kb (Figure 4, lane 1). Thus *Df(2L)el⁷⁷* is broken in the 7.2-kb *EcoRI* fragment between +44.2 and +51.4. When DNA from *Df(2L)TE35B-48/Df(2L)A48* was probed with the 7.7-kb *XhoI* fragment of λ W7.11, parental *EcoRI* fragments of 0.8, 1.4 and 4.9 kb were seen (Figure 4, lanes 2, 3). The 6.4-kb fragment was replaced by a fusion fragment of 5.5 kb (Figure 4, lane 3). From this we conclude that *Df(2L)TE35B-48* is broken between +51.8 and +57.3, close to the *EcoRI* site at +57.3. When probed with λ W9.06, DNA from *Df(2L)A178/Df(2L)A48* lacked the parental fragment of 17.7 kb (Figure 4, lane 4) and instead had a fusion fragment of 15.0 kb (Figure 4, lane 5). Therefore the proximal limit of *Df(2L)A178* is in the 17.0-kb *EcoRI* genomic fragment at +76.5 to +78.5. Together with data from CHIA *et al.* (1985), this defines *Df(2L)A178* as

FIGURE 4.—Southern analysis of genomic DNA from *osp* mutants. Samples were digested with *Eco*RI unless noted. Fusion fragments are marked “F”. (a) *Df(2L)el⁷⁷/Df(2L)A48, b cn bw* probed with the 7.2-kb *Xho*I fragment of λ W7.012 (lane 1). (b) Parental strain *al dp b noc^{TE35B(Z)} pr l(2)pwn cn/Df(2L)A48, b cn bw* (lane 2) and derivative *Df(2L)TE35B-48, al dp b pr l(2)pwn cn/Df(2L)A48, b cn bw* (lane 3) probed with the 7.7-kb *Xho*I fragment of λ W7.11. (c) Parental strain *b Adh^f cn bw* (lane 4) and derivatives *Df(2L)A178, b cn bw/Df(2L)A48, b cn bw* (lane 5) and *Df(2L)A379, In(2LR)A379, b cn bw/Df(2L)A48, b cn bw* (lane 6) probed with λ W9.06. (d) Parental strain *Adhuf3cn/CyO* (lane 7) and derivative *Tp(2;3)osp³, Adh^{uf3} cn/CyO* (lane 8) probed with the 4.6-kb *Eco*RI fragment of λ W5.02. (e) *T(2;3)pb³/CyO* digested with *Hind*III and probed with 2.9- and 4.6-kb *Eco*RI fragments of λ W5.02 (lane 9). (f) *Xho*I-digested DNA from Canton S (lane 10) and *In(2L)CI58/CyO* (lane 11) probed with the proximal 5.0 kb of phage λ W11.01. (g) Parental strain *al dp b noc^{TE35B(Z)} pr l(2)pwn cn/CyO* probed with the 7.2-kb *Eco*RI fragment of λ W6.05 (lane 12). (h) *al dp b noc^{TE35B(Z)} pr l(2)pwn cn/CyO* probed with the 5.2-kb *Eco*RI fragment of λ W6.05 (lane 13). The 7.2-kb fragment is due to incomplete removal of the previous 7.2-kb *Eco*RI probe (see lane 12). Several strains appear to have an insertion at map position +51.4 kb. When the 5.2-kb *Eco*RI fragment of λ W6.05 was hybridized to *Eco*RI-digested DNA from either *al dp b noc^{TE35B(Z)} pr l(2)pwn cn/CyO* (lane 13) or a Canton S stock, it yielded a fragment of 11.5 kb in addition to the expected 6.4-kb fragment. This fragment was not observed (lane 12) when probed with the adjacent 7.2-kb fragment of λ W6.05. The 11.5-kb fragment is probably due to an insertion of 5.1 kb at about +51.4 kb, rather than a polymorphism, as the *Eco*RI fragments adjacent to the 6.4-kb genomic fragment would not, in combination with the 6.4-kb fragment, add up to an 11.5-kb fragment. This insertion appears to be associated with a small repeated element (faint bands lane 1) and is present in about half of the chromosomes used in this study. λ W6.05, λ W7.011 and λ W7.012 also appear to contain small repetitive elements.

a deletion of ~ 190 kb. DNA from *Df(2L)A379, In(2LR)A379/Df(2L)A48*, probed with λ W9.06, yielded only a 5.0-kb fragment (Figure 4, lane 6). The absence of a fusion fragment suggests that this aberration is broken close to the *Eco*RI site at +84.5.

Three additional aberrations, two *osp⁻* and one *osp⁺*, have been mapped to this region. DNA from *Tp(2;3)osp³/CyO*, digested with *Eco*RI and probed with the 4.6-kb *Eco*RI fragment of λ W5.02 (Figure 2), yielded four fragments (Figure 4, lane 8). Fragments of 3.3 and 1.3 kb are from the *CyO* balancer chromosome (Figure 4, lane 8) and are due to a polymorphism at +42.6 (Figure 2). The 4.6-kb band from the parental chromosome (Figure 4, lane 8) has been replaced by 4.8- and 1.9-kb fusion fragments, thus *Tp(2;3)osp³* appears to be broken in the λ W5.02–4.6-kb *Eco*RI fragment, between +39.6 and +44.2. This result was confirmed by *in situ* hybridization to *Tp(2;3)osp³* chromosomes. When λ W5.02 was used as probe, the mutant second and third chromosomes hybridized, in addition to the *CyO* balancer chromosome.

DNA from *T(2;3)pb³/CyO*, digested with *Hind*III and

probed with the 2.9- and 4.6-kb *Eco*RI fragments of λ W5.02, yielded parental fragments of 4.0 and 3.5 kb (Figure 4, lane 9). In addition, fusion fragments of 10.5 and 1.9 kb were found. The intensity of signal from the parental 4.0-kb fragment is weak, consistent with the translocation being broken within this fragment, between +36.7 and +40.7. This was confirmed by *in situ* hybridization to *T(2;3)pb³* chromosomes.

The breakpoint of *T(2;3)H47* was shown to lie within the DNA contained in λ W5.02, between +34.4 and +50.6, by *in situ* hybridization. Genomic DNA from λ W5.02 hybridized to 35B and also to a site on the third chromosome.

The *osp⁺* inversion *In(2L)CI58* defines the proximal outer limit of *osp*. *In(2L)CI58*, found in a natural population (ASHBURNER and LEMEUNIER 1976), breaks between *osp* and the locus *l(2)35Bb*. DNA from *In(2L)CI58/CyO* was digested with *Xho*I and probed with the proximal 5.0 kb of phage λ W11.01. In addition to the 15.0-kb parental fragment present in Canton S (Figure 4, lane 10), two novel fragments of 16.0 and 7.1 kb were obtained (Figure 4, lane 11). We tentatively concluded

that this inversion is broken between +100.4 and +105.4 and confirmed it by hybridizing λ W11.01 *in situ* to *In(2L)C158* chromosomes (data not shown). These data demonstrate that the proximal end of *osp* lies to the left of the *In(2L)C158* proximal breakpoint, *i.e.*, left of +105.4 on the molecular map.

The remaining aberrations have proximal limits beyond λ W11.01. Locations of the *osp* mutations are summarized in Figure 3.

The mapping of mutants defines the proximal end of *osp* as lying between +39.6, as defined by *Tp(2;3)osp³*, *T(2;3)pb³* and *T(2;3)H47* and +105.4, as defined by *In(2L)C158*. Taken together with previous work (GUBB *et al.* 1990), we conclude that the *osp* locus lies between -11.2 to -13.3 and +39.6 to +105.4 on the walk map, and is 52.8 to 118.7 kb in length.

Identification of the *osp* transcript and its developmental expression: Fragments of cloned genomic DNA to which *osp* lesions mapped were used to probe developmental Northern blots. Although six different fragments were used as probes (see MATERIALS AND METHODS), only the *EcoRI* 4.7-kb fragment from the proximal end of λ ob1.12 (Figure 2) was successful in detecting RNA. It hybridized a major band of ~6 kb. The transcript is most abundantly expressed in embryonic and pupal stages of development, although it is detected at all stages (Figure 5). The control reprobing of the blot with an actin 5A clone showed that similar amounts of RNA were present in each lane (data not shown). This 6-kb RNA is detected in polyadenylated RNA fractions from pupae (data not shown).

The 6-kb transcript is absent in *osp* mutants: The evidence that the 6-kb band represents the *osp* transcript is that the 6-kb transcript is absent in 14 different *osp* mutants and defective in one, but present in stocks that are not mutant for *osp*. Adult flies that were heterozygous for one of 15 *osp* alleles over *Df(2L)A178* were tested for accumulation of the 6-kb transcript (Figure 6). Flies with deficiencies distal to *osp*, *e.g.*, *GT4*, *b84a4* and *b84a8* (CHIA *et al.* 1985), produce the 6-kb transcript. In contrast, *osp* mutants lack the 6-kb transcript. These include deficiencies such as *A220*, *osp18*, *osp29*, *A63*, *A72*, *A48* and *A379*; inversions *osp⁵⁹* and *osp²²*; translocations *pb³* and *osp⁹⁰*; and transpositions *osp³* and *H47*. The inversion proximal to the *osp* region, *C158*, which is *osp⁺* but inverted between *osp* and the adjacent gene *l(2)35Bb*, does not disrupt production of the 6-kb transcript.

For *Df(2L)osp144*, novel transcripts of 4.6, 3.1, 2.4, 2.0 and 1.3 kb were detected. This deficiency extends from approximately -10 to -70 (CHIA *et al.* 1985) and likely removes part of the *osp* 3' exon (described in further detail later). The origin of the *osp144* transcripts is not known. However, *osp144* has a strong *osp* phenotype, which indicates that these short RNAs are not sufficient for *osp* function.

Characterization of the *osp* cDNAs: One cDNA homologous to *obl-4.7*, 211, was obtained from the 0–3-

FIGURE 5.—Developmental profile of the *osp* transcript. This Northern blot contains total RNA from a Canton S wild-type strain probed with the *EcoRI* 4.7-kb fragment from λ ob1.12. The stages indicated at the top of each lane are: EE, early embryos (0–3 hr of development); E, embryos (3–24 hr); L₁, first instar larvae (2–3 days); L₂, second instar larvae (3–4 days); EL₃, early third instar larvae (4–5 days); LL₃, larvae from the second half of third instar (5–6 days); PE, day 6–8 pupae; PL, day 8–9 pupae; and A, adults (9–12 days). The *osp* transcript is 6 kb in length (large arrow). Numbers to the left indicate the migration of RNA size markers in kilobases. The highest molecular weight band (small arrow) is genomic DNA. Smudges and faint bands at larger or smaller than 6 kb may represent RNA processing byproducts. This background is absent in the sample from early embryos (lane EE).

hr embryonic cDNA library (POOLE *et al.* 1985). Two cDNAs, 7111 and 9114, were isolated from the 12–24-hr embryonic cDNA library (BROWN and KAFATOS 1988). cDNA 9114 contained the longest cDNA insert of 4.2 kb, short of the 6 kb required for a complete *osp* cDNA. When used to probe Southern blots of the walk region, the cDNAs all mapped to the left of *Adh* (summarized in Figure 7).

The structures of the *osp* cDNAs provide evidence of alternative splicing of *osp* transcripts. When mapped to Southern blots of *osp* region phage DNA digested with *EcoRI* (summarized in Figure 7), cDNA 211 hybridized only to the 4.7-kb fragment of λ ob1.12, at -15.8 to -11.2; 7111 hybridized to the 4.7-kb λ ob1.12 fragment, the 2.4-kb λ gAC3 fragment, and the 2.0-kb fragment of λ gAC2 (which overlaps completely the 2.4-kb fragment of λ gAC3). cDNA 9114 hybridized the regions covered by 7111 plus an additional portion of λ gAC3. These results suggest that there are at least two *osp* RNA splice variants that share a common 3' segment.

FIGURE 6.—Northern analysis of *osp* mutants. Total RNA from adults was probed with the 4.7-kb *Eco*RI fragment from λ ob1.12 (top) or *rp49* (bottom). All *osp* mutants were tested in heteroallelic combination with *Df(2L)A178*. Lanes (1) Canton S, (2) *Df(2L)A220*, (3) parental strain *Adh^{u3} pr cn*, (4) *Df(2L)-osp29*, (5) *Df(2L)osp³*, (6) *T(2;3)osp⁹⁰*, (7) *In(2L)osp⁵⁹*, (8) *Df(2L)osp²⁰¹*, (9) *Df(2L)osp144*, (10) *Df(2L)osp¹⁸*, (11) *Df(2L)A63*, (12) *Df(2L)A72*, (13) *Df(2L)-A379*, (14) *In(2L)C158*, (15) Canton S, (16) *T(2;3)H47*, (17) *Df(2L)A48*, (18) *T(2;3)pb³*, (19) *In(2L)osp²²*, (20) *Df(2L)-GT4*, (21) *Df(2L)b84a4*, (22) *Df(2L)-b84a8*. Sizes of the bands are indicated to the left.

Some of the variation observed between the cDNAs may be due to incomplete splicing, however, the cDNAs all detect a 6-kb band on Northern blots. The 6-kb band must represent a set of transcripts of approximately the same length. The possibility of some of the variants arising from an *osp*-related gene is unlikely. Reduced stringency genomic Southern blots probed with 9114 yielded no bands aside from those representing the *osp* walk region (data not shown).

The PCR 5' extension of the longest *osp* cDNA hybrid-

izes to restriction fragments both proximal and distal to *Adh*: The cDNAs failed to provide direct molecular evidence that *Adh* was nested in *osp*. Since the cDNAs were not full length, it was possible that by extending them further, such evidence would be obtained. Thus, the region 5' to the longest *osp* cDNA, 9114, was extended by PCR. Since *osp* is abundant in embryos and an excellent embryonic cDNA library was available, we amplified the 5' extension from the pNB40 12–24-hr embryonic cDNA library (BROWN and KAFATOS 1988). The primers con-

FIGURE 7.—Summary of the structure of the *osp* cDNAs. cDNAs were hybridized to *osp* walk phage DNA digested with either *Eco*RI or *Eco*RI in combination with a second enzyme. Walk fragments that were hybridized are represented by lines corresponding to the different cDNAs above the genomic map. The names of the cDNAs are to the right of the lines and their actual lengths to the left. *Adh* and *Adh'* are indicated by arrows just above the genomic map. The walk phage and *Eco*RI fragments hybridized are shown below.

FIGURE 8.—Hybridization of the 5' PCR extension of 9114 to the *osp* walk. *posp5'-319* was hybridized to a Southern blot of *EcoRI*-digested DNA from *osp* walk phage. Lanes (1) λ ob1.12, (2) λ gAC3, (3) λ gAC2, (4) λ W2, (5) λ W3.13, (6) λ W4.04, (7) λ W5.02, (8) λ W6.05, (9) λ W7.11, (10) λ W8.061, (11) λ W9.06, (12) λ W10.03, and (13) λ W11.01. Migration of the λ HindIII markers is indicated on the left; the sizes of the walk phage bands on the right. All other bands are due to partial digestions. Additional mapping (not shown) indicates that the apparent 2.1-kb band in lane 4 corresponds to the distalmost 2.0-kb fragment of λ gAC2.

sisted of a 3' primer internal to the 5' end of 9114 and a 5' primer internal to the pNB40 vector.

The amplification yielded a major band of 1.5 kb, bringing the total *osp* cDNA length up to 5.5 kb, close to full length. The identity of this band was verified by hybridization to an internal probe. It was then cloned to make *posp5'-319* and used to probe DNA from the *osp* walk phage (Figure 8). As expected, it hybridized to the *EcoRI* 2.4-kb fragment of λ gAC3 and to λ gAC2-*EcoRI* 2.0 kb, from which the primer originated, distal to *Adh*. In addition, it hybridized to fragments proximal to *Adh*, to λ W3-*EcoRI* 15 kb and to the *EcoRI* 2.7-kb fragment shared by λ W7.11 and λ W8.061 at +63.0 to +65.7 kb (summarized in Figure 9). The 5' extension identifies two new exons, indicating that *osp* is composed of at least four exons.

To characterize the 5' extension further, *posp5'-319* was used to probe Northern blots and polytene chromosomes. As a Northern probe, it detected only the 6-kb transcript. The only site of chromosomal hybridization was 35B2-3, where *osp* and *Adh* are located. In summary, hybridization to a predicted internal probe, to its site of origin on the walk, to the *osp* RNA, and to the chromosomal location of *osp* confirms that the 1.5-kb PCR amplification product corresponds to a portion of the

osp gene product. The 5' extension represents at least two new exons proximal to *Adh* and *Adh'*. The cDNAs extend from -15.8 to -11.2 to +63.0 to +65.7 kb, demonstrating that the *osp* transcription unit is ≥ 74 kb in length.

Characterization of maternal and zygotic expression of *osp*: The presence of *osp* transcripts in 0–3-hour embryos (Figure 5) indicates that *osp* is expressed in ovaries. Expression in ovaries of wild-type flies was examined further by *in situ* hybridization (Figure 10). The *osp* transcript is present in multiple nurse cells in the germarium. By stage 3, it is localized principally to the presumptive oocyte where it continues to be detected through stage 10. The *osp* transcript is present in a broad band in the anterior end of late stage oocytes. The anterior localization is retained until after cellular blastoderm (Figure 10A). In addition, *osp* appears to be expressed by follicle cells (Figure 10B). Follicle cells may contribute to an accumulation of *osp* transcript around the periphery of the oocyte (Figure 10A, stage 10), however, most of the transcript detected at the periphery is internal to the oocyte (Figure 10B). In ovaries from heteroallelic *osp*⁻ females *Df(2L)A63/Df(2L)A178*, no transcription was detected.

In wild-type embryos, *osp* displays a highly dynamic pattern early in embryogenesis. From the earliest stages of embryogenesis through early syncytial blastoderm, maternal *osp* transcript is localized at the anterior end of the embryo (Figure 11A). As the ventral furrow forms, *osp* is expressed at low levels lateral to the furrow. Later it is expressed in several prominent stripes (Figure 11B) that extend across the entire germ band, visible only briefly as the germ band elongates. At stage 7, *osp* is expressed strongly in the maxillary and labial segments and appears to be expressed in faint segmental stripes (Figure 11C). By stage 8, *osp* expression in the mandibular and maxillary segments is intensified (Figure 11D). In addition, its segmental expression resembles pairs of spheres within the mesodermal layer (Figure 11E). Segmental staining in areas of mesodermally derived tissues is prominent throughout the period of germ band extension. As the germ band retracts, the pattern along the ventral surface is refined and a few cells along the ventral midline are prominent (Figures 11F) in a pattern similar to aspects of *achaete-scute* (CABRERA *et al.* 1987) and *nautilus* (MICHELSON *et al.* 1990) expression. Subsequently, the staining parallels the shape of the somatic and visceral musculature (M. BATE, personal communication; Figure 11, G–L). Some tissues that are not of mesodermal origin, including the salivary glands (Figure 11, F–J) and later the malpighian tubules (Figure 11, K and L) also stain intensely. *osp* transcripts are detected through stage 17. In control embryos produced by heterozygous *osp* parents *Df(2L)A63/Df(2L)A178*, no staining was detected at any stage.

DISCUSSION

***Adh* and *Adh'* are nested in the *osp* locus:** *osp* is a large gene of up to 119 kb, with a minimum transcription unit

FIGURE 9.—Summary of the molecular characterization of the *osp* locus. The *osp* exons are represented by thick horizontal bars, the most 3' of which ends in an arrow to indicate the direction of transcription. Asterisks indicate portions of *osp* hybridized exclusively by the 1.5-kb 5' cDNA extension. *Adh* and *Adh'* are symbolized by smaller arrows below. The distance in kilobases is given along the thick central line. Restriction fragments to which *osp* mutations are localized are indicated below it. The walk phase are shown at the lowest level.

of 74 kb that encodes a set of 6-kb transcripts. It appears, on the basis of the cDNAs recovered, to possess at least four introns and to encode at least two classes of RNAs distinguishable as splice variants. We conclude that *Adh* and *Adh'* are nested within *osp*, since they are encompassed by *osp* cDNAs (summarized in Figure 9).

In addition to *Adh'*, the open reading frame (ORF) 3' of *Adh*, a 5' ORF was previously identified 4.2 kb upstream from the *Adh* adult (distal) promoter (KREITMAN and AGUADE 1986). This 5' ORF lies within a restriction fragment which we have shown contains an *osp* exon. The 5' ends of cDNAs 9114 and 7111 overlap this 5' ORF. The known sequence of 9114 is virtually identical to the 5' ORF throughout its length of 247 bp; thus this ORF is almost certainly part of the *osp* gene product. The conceptual translation of the 5' ORF does not yield a protein with significant homology to any known protein. The region between -660 and -5000 bp 5' of the *Adh* adult promoter, in which this ORF is located, contains an enhancer for larval *Adh* (POSAKONY *et al.* 1985; CORBIN and MANIATIS 1989). It remains to be determined if alterations to this portion of the *osp* locus have an effect on expression of larval *Adh*.

Characteristics of nested genes: In addition to the *osp-Adh* nested locus, six nested gene sets have now been described. In *Drosophila*, these are the gene for guanine-adenosine ribosyl transferase (*GART*) and intronic pupal cuticle gene (*Pcp*) (HENIKOFF *et al.* 1986a), the *dunce* gene and intronic genes *Sgs-4*, *PIG-1*, *ng-1*, *ng-2*,

ng-3 and *ng-4* (CHEN *et al.* 1987; FURIA *et al.* 1990, 1991, 1993), the R1 locus of cAMP dependent kinase (*Pka-R1*) and two intronic genes of unknown function (KALDERON and RUBIN 1988), and the *doublesex* (*dsx*) locus and the intronic adult male specific transcription unit (CHEN 1995). Two sets of nested genes have been identified in humans: the neurofibromatosis type 1 gene (*NFI*) and intronic oligodendrocyte myelin glycoprotein gene (*OMGP*), *EVI2A* and *EVI2B* (XU *et al.* 1990), and *factor VIII* and an intronic gene of unknown function (LEVINSON *et al.* 1990). *NFI* also has a nested structure in mice (CAWTHON *et al.* 1990). A few generalizations can be drawn from these cases.

1. Nested genes are broadly distributed but they may not be stable over extended periods of evolutionary time. The presence of nested genes in the human, mouse and *Drosophila* genomes suggests that they may be distributed throughout the Metazoa. For *GART*, the nested arrangement is present in *D. pseudoobscura* (HENIKOFF and EGHTEGARZADEH 1987), separated from *D. melanogaster* by 5 million years, but absent in *Chironomus tentans*, a dipteran separated from *D. melanogaster* by 50 million years (CLARK and HENIKOFF 1992). For *factor VIII*, the mouse gene possesses no intron and the homologue of the intronic gene is located distantly, unlike the human locus (LEVINSON *et al.* 1990).
2. Intronic genes are small genes located in large genes. Intronic genes range in size from the 0.6-kb

FIGURE 10.—Expression of *osp* in ovarioles. Dissected, whole mount ovarioles were hybridized *in situ* with 9114. (A) Two ovarioles are shown. The ovariole that extends toward the lower right has stretched lengthwise to show the developing oocyte cysts arrayed in order of increasing maturity: the germarium (G) and stages 3, 4/5, 6, 8, 8/9 and 10. *osp* is expressed in the germarium and in more mature cysts (arrows). In stages 8 and 9, *osp* RNA is most abundant in the posterior of the oocyte, but at stage 10, it is concentrated in the anterior (arrow). (B) A stage 10 cyst from a different ovariole, in which the *osp* transcript is concentrated at the anterior end of the oocyte (arrow). (C) The follicle cells that surround the stage 12 cyst in (A) appear to express some *osp* (arrow).

ng-1 (FURIA *et al.* 1993) to the 2.7-kb *OMGP* (XU *et al.* 1990). However, intronic genes may possess introns, as in the cases of *Adh* (BENYAJATI *et al.* 1983), *Pcp* (HENIKOFF *et al.* 1986a) and *EVI2A* and *EVI2B* (CAWTHON *et al.* 1990).

Nest genes range from the 10-kb *GART* (HENIKOFF *et al.* 1986a) to the 240 kb *NFI* (XU *et al.* 1990; GOLDBERG and COLLINS 1991). Large genes with sizeable introns appear to provide big targets or residences for intronic genes. With the exception of *dsx*, all nest genes encode enzymes. Known nested genes are associated with genetic disorders or mutations, and since large genes are more easily mutated than are short ones, they are probably overrepresented in this survey.

3. The number of genes per intron and per locus, and their transcriptional orientations, are not constrained. *GART* (HENIKOFF *et al.* 1986a) and *factor VIII* (LEVINSON *et al.* 1990) have only one intronic gene each, *osp* and *Pka-R1* (KALDERON and RUBIN 1988) each have two, *NFI* contains three (XU *et al.* 1990). *dunce*, with the largest number of intronic genes at a single locus, has at least six: four in one intron and two in another (FURIA *et al.* 1993). In-

tronic genes may be in the same or opposite orientation to the nest gene and to each other. The *dunce* locus includes examples of both (FURIA *et al.* 1993).

4. A number of intronic genes share sequence similarity or tissue specificity with other intronic genes at the same locus. Nested in *osp*, *Adh* and *Adh'* are similar in sequence and appear to be related by an ancient gene duplication (SCHAEFFER and AQUADRO 1987; KREITMAN and HUDSON 1991). The *EVI2A* and *EVI2B* genes that nest in *NFI* are unrelated in sequence but are both expressed in peripheral blood mononuclear cells (CAWTHON *et al.* 1990, 1991). Of the six intronic genes of *dunce*, four are expressed in third instar larval salivary glands, although they differ somewhat in temporal specificity. Two reside in each of the two introns (FURIA *et al.* 1993). The nest genes do not share sequence similarity with their intronic genes, and in only two cases, those of *Adh* and *osp* and *NFI* and *OGMP* (XU *et al.* 1990), is there a suggestion of overlapping tissue specificity.

The presence of neighboring intronic genes with shared origins or transcriptional regulation suggests that the introns of nest genes can be accommodating environments in which other genes can

FIGURE 11.—Distribution of *osp* RNA in embryos. Whole mount embryos were hybridized *in situ* with cDNA 9114. (A) Early cellular blastoderm; (B) stage 7 as the germ band extends; (C) early stage 8; (D and E) stage 8, lateral section (D) and medial longitudinal section (E); (F) stage 12, early germ band retraction, ventral surface; (G–I) stage 14, rotated slightly so that ventral is toward the plane of the viewer: lateral section (G), more medial section (H), and ventral surface (I); (J–L) stage 17: medial section (J), dorsal section (K), and medial transverse section (L). Arrows and labels indicate regions that show high levels of expression. Frames are arranged with the anterior at left, dorsal at top except where noted. amg, anterior midgut; es, esophagus; hg, hindgut; la, labial segment; md, mandibular segment; mg, midgut; mt, Malpighian tubules; mx, maxillary segment; ph, pharynx; pmg, posterior midgut; sg, salivary gland; sm, somatic musculature; sms, somatic mesoderm; vm, visceral musculature.

survive, can be expressed and can evolve. It is important to know what features distinguish nest genes and their introns as permissive residences. One feature may be an open chromatin conformation. Three of the five nested genes found in *Drosophila*, *dunce*, *GART* and *dsx*, are located at chromosomal regions that undergo puffing in the salivary gland chromosomes (ASHBURNER 1967, 1969). The other two, *osp* and *Pka-R1*, are located in regions composed of interbands and faint bands. These cytological features are consistent with local decondensation of chromatin.

Transcriptional interference between *osp* and *Adh*:

For nested genes, the question of transcriptional interference between the genes arises. Transcriptional interference can be inferred through competing patterns of transcription. At the cellular level, *osp* and *Adh* have

been examined most closely in embryos, where expression of the two genes does not overlap. *osp* is expressed primarily in mesodermal tissues, but also in the salivary glands and Malpighian tubules (but not other gut-derived tissues). *Adh*, on the other hand, is expressed in fat body and in gut derivatives not including the Malpighian tubules in embryos (LOCKETT and ASHBURNER 1989). However, in adult females *Adh* and *osp* are both expressed in the ovary, thus defining a tissue and a time when transcriptional interference may occur.

The nature of the *osp* gene product: The maternal *osp* transcript is expressed in nurse cells of developing egg chambers in the ovary and is packaged into the oocyte, where it is localized at the anterior end. It is also expressed in the follicle cells.

The zygotic *osp* transcript is detected very early in embryogenesis, by ~3.7 hr into development. In general, zygotic transcription in the embryo begins at ~2

hr of development. Since the rate of transcription in *Drosophila* is 1.1–1.4 kb/min (THUMMEL *et al.* 1990; SHERMOEN and O'FARRELL 1991; IRVINE *et al.* 1991) and the probe used here detects portions of the *osp* RNA ≥ 60 kb from its transcriptional start site, *osp* transcription must begin at ~ 3 hr of development, just after the embryo completes cellularization. *osp* transcripts are detected throughout the embryonic period. *osp* is strongly expressed in a segmental pattern as the germ band elongates, extends and retracts, predominantly in tissues of mesodermal origin.

The expression pattern of *osp* suggests that, while it may have a complex role in development, it may be particularly important in development of muscle. Since it is most abundantly expressed in embryos and pupae and is present very early in the embryo, it may play a role in formation or structuring of the musculature. The *outspread* wing phenotype could result from a structural defect or a deficit in muscle function. Remarkably, null mutations of *osp* are not lethal, the mutants are fertile and no defects have been observed among the progeny aside from the wing phenotype. Future studies will focus on defining the role of *osp* through analysis of its sequence, its protein localization, and its interactions with other genes that play roles in mesoderm and muscle development.

We thank M. ASHBURNER, in whose lab this work was done, for financial support and comments on this manuscript. We thank BARBARA WAKIMOTO, JOHN ROOTE, NICK BROWN, E. MICHAEL BATE, DAVID GUBB, MADELEINE CROSBY, A. RUS HOELZEL, the late MIKE WILCOX, ROBERT WEINZEL, ANNIE ROWE, GUY TEAR, PETE JEFFS, BUO MENG, GHADA AL-ATIA and DANIEL ST. JOHNSTON for practical help and fruitful discussions. Thanks to PAM ANGEL, BELINDA DURRANT, GLYNNIS JOHNSON and JILL REGAN for fly stocks and to TAMSIN MAJERUS for *in situ* hybridizations. Thanks to BARBARA WAKIMOTO, CELESTE BERG and MARGRIT SCHUBIGER for comments on this manuscript. This work was supported by a programme grant from the Medical Research Council to M. ASHBURNER.

LITERATURE CITED

- AARON, C. S., 1979 X-ray induced mutations affecting the level of the enzyme alcohol dehydrogenase in *Drosophila melanogaster*: frequency and genetic analysis of the null-enzyme mutants. *Mutat. Res.* **63**: 127–137.
- ASHBURNER, M., 1967 Patterns of puffing activity in the salivary gland chromosomes of *Drosophila*. I. Autosomal puffing patterns in a laboratory stock of *Drosophila melanogaster*. *Chromosoma* **21**: 398–428.
- ASHBURNER, M., 1969 Patterns of puffing activity in the salivary gland chromosomes of *Drosophila*. II. X chromosome puffing patterns in *Drosophila melanogaster* and *D. simulans*. *Chromosoma* **21**: 47–63.
- ASHBURNER, M., and F. LEMEUNIER, 1976 Relationships within the melanogaster species subgroup of the genus *Drosophila* (Sophophora). I. Inversion polymorphisms in *Drosophila melanogaster* and *Drosophila simulans*. *Proc. R. Soc. Lond. B Biol. Sci.* **193**: 137–157.
- ASHBURNER, M., C. S. AARON and S. TSUBOTA, 1982 The genetics of a small autosomal region of *Drosophila melanogaster*, including the structural gene for alcohol dehydrogenase. V. Characterization of X-ray-induced *Adh* null mutations. *Genetics* **102**: 421–435.
- BENYAJATI, C., A. R. PLACE and W. SOFER, 1983 Formaldehyde mutagenesis in *Drosophila*. Molecular analysis of ADH-negative mutants. *Mutat. Res.* **111**: 1–7.
- BROWN, N., and F. KAFATOS, 1988 Functional cDNA libraries from *Drosophila* embryos. *J. Mol. Biol.* **203**: 425–437.
- CABRERA, C. V., A. MARTINEZ-ARIAS and M. BATE, 1987 Expression of three members of the *achaete-scute* gene complex correlates with neuroblast segregation in *Drosophila*. *Cell* **50**: 425–433.
- CAWTHON, R. M., P. O'CONNELL, A. M. BUCHBERG, D. VISKOCHIL, R. B. WEISS *et al.*, 1990 Identification and characterization of transcripts from the neurofibromatosis 1 region: the sequence and genomic structure of *EV12* and mapping of other transcripts. *Genomics* **7**: 555–565.
- CAWTHON, R. M., L. B. ANDERSEN, A. M. BUCHBERG, G. XU, P. O'CONNELL *et al.*, 1991 cDNA sequence and genomic structure of *EV12B*, a gene lying within an intron of the neurofibromatosis type 1 gene. *Genomics* **9**: 446–460.
- CHEN, C., T. MALONE, S. K. BECKENDORF and R. L. DAVIS, 1987 At least two genes reside within a large intron of the *dunce* gene of *Drosophila*. *Nature* **329**: 721–724.
- CHEN, H.-J., 1995 Functional domains of the doublesex proteins in *Drosophila melanogaster*. Ph.D. Thesis, University of California, Davis.
- CHIA, W., R. KARP, S. MCGILL and M. ASHBURNER, 1985 Molecular analysis of the *Adh* region of the genome of *Drosophila melanogaster*. *J. Mol. Biol.* **186**: 689–706.
- CHOMCZYNSKI, P., and N. SACCHI, 1987 Single-step method of RNA isolation by acid guanidinium thiocyanate-phenol-chloroform extraction. *Anal. Biochem.* **162**: 156–159.
- CLARK, D. A., and S. HENIKOFF, 1992 Unusual organizational features of the *Drosophila GART* locus are not conserved within Diptera. *J. Mol. Evol.* **35**: 51–59.
- CORBIN, V., and T. MANIATIS, 1989 Role of transcriptional interference in the *Drosophila melanogaster Adh* promoter switch. *Nature* **337**: 279–282.
- DAVIS, T., J. TRENEAR and M. ASHBURNER, 1990 The molecular analysis of the *el-noc* complex of *Drosophila melanogaster*. *Genetics* **126**: 105–119.
- FEINBERG, A. P., and B. VOGELSTEIN, 1983 A technique for radiolabeling DNA restriction endonuclease fragments to high specific activity. *Anal. Biochem.* **132**: 6–13.
- FURIA, M., F. A. DIGLIO, D. ARTIACO, E. GIORDANO and L. C. POLITO, 1990 A new gene nested within the *dunce* genetic unit of *Drosophila melanogaster*. *Nucleic Acids Res.* **18**: 5837–5841.
- FURIA, M., F. A. DIGLIO, D. ARTIACO, P. P. D'AVINO, D. CAVALIERE *et al.*, 1991 Molecular organization of the *Drosophila melanogaster Pig-1* gene. *Chromosoma* **101**: 49–54.
- FURIA, M., P. P. D'AVINO, S. CRISPI, D. ARTIACO and L. C. POLITO, 1993 Dense cluster of genes is located at the ecdysone-regulated 3C puff of *Drosophila melanogaster*. *J. Mol. Biol.* **231**: 531–538.
- GITSCHIER, J., W. I. WOOD, T. M. GORALKA, K. L. WION, E. Y. CHEN *et al.*, 1984 Characterization of the human factor VIII gene. *Nature* **312**: 326–330.
- GOLDBERG, N. S., and F. S. COLLINS, 1991 The hunt for the neurofibromatosis gene. *Arch. Dermatol.* **127**: 1705–1707.
- GUBB, D., M. SHELTON, J. ROOTE, S. MCGILL and M. ASHBURNER, 1984 The genetic analysis of a large transposing element of *Drosophila melanogaster*. The insertion of a *w⁺ rst⁺* TE into the *ck* locus. *Chromosoma* **91**: 54–64.
- GUBB, D., J. ROOTE, G. HARRINGTON, S. MCGILL, B. DURRANT *et al.*, 1985 A preliminary analysis of TE146, a very large transposing element of *Drosophila melanogaster*. *Chromosoma* **92**: 116–123.
- GUBB, D., J. ROOTE, S. MCGILL, M. SHELTON and M. ASHBURNER, 1986 Interactions between *white* genes carried by a large transposing element of *Drosophila melanogaster*. *Genetics* **112**: 551–575.
- GUBB, D., S. MCGILL and M. ASHBURNER, 1988 A selective screen to recover chromosomal deletions and duplications in *Drosophila melanogaster*. *Genetics* **119**: 377–390.
- GUBB, D., M. ASHBURNER, J. ROOTE and T. DAVIS, 1990 A novel transvection phenomenon affecting the *white* gene of *Drosophila melanogaster*. *Genetics* **126**: 167–176.
- HENIKOFF, S., and M. EGHTEGARZADEH, 1987 Conserved arrangement of nested genes at the *Drosophila Gart* locus. *Genetics* **117**: 711–725.
- HENIKOFF, S., M. A. KEENE, K. FECHTEL and J. W. FRISTOM, 1986a Gene within a gene: nested *Drosophila* genes encode unrelated proteins on opposite DNA strands. *Cell* **44**: 33–42.
- HENIKOFF, S., M. A. KEENE, J. S. SLOAN, J. BLESKAN, R. HARDS 1986b Multiple purine pathway enzyme activities are encoded at a single genetic locus. *Proc. Natl. Acad. Sci. USA* **83**: 720–724.
- HILLIKER, A. J., and S. N. TRUSIS-COULTER, 1987 Analysis of the

- functional significance of linkage group conservation in *Drosophila*. *Genetics* **117**: 233–244.
- HOLMES, D. S., and M. QUIGLEY, 1981 A rapid boiling method for the preparation of bacterial plasmids. *Anal. Biochem.* **114**: 193.
- HUGHES, M. A., and M. SHELTON, 1980 Report of new mutants—*Drosophila melanogaster*. *Dros. Inf. Serv.* **55**: 204–205.
- IRVINE, K. D., S. L. HELFAND and D. S. HOGNESS, 1991 The large upstream control region of the *Drosophila* homeotic gene *Ultrabithorax*. *Development* **111**: 407–424.
- KALDERON, D., and G. M. RUBIN, 1988 Isolation and characterization of *Drosophila* cAMP dependent protein kinase genes. *Genes Dev.* **2**: 1539–1556.
- KAUFMAN, T. C., 1978 Cytogenetic analysis of chromosome 3 in *Drosophila melanogaster*: isolation and characterization of four new alleles of the *proboscipedia* (*pb*) locus. *Genetics* **90**: 579–596.
- KREITMAN, M., and R. H. HUDSON, 1991 Inferring the evolutionary histories of the *Adh* and *Adh-dup* loci in *Drosophila melanogaster* from patterns of polymorphism and divergence. *Genetics* **127**: 565–582.
- KREITMAN, M., and M. AGUADE, 1986 Excess polymorphism at the *Adh* locus in *Drosophila melanogaster*. *Genetics* **114**: 93–110.
- LEVINSON, B., S. KENWICK, D. LAKICH, G. HAMMONDS, JR. and J. GITSCHER, 1990 A transcribed gene in a intron of the human factor VIII gene. *Genomics* **7**: 1–11.
- LOCKETT, T. J., and M. ASHBURNER, 1989 Temporal and spatial utilization of the *alcohol dehydrogenase* gene promoters during the development of *Drosophila melanogaster*. *Dev. Biol.* **134**: 430–437.
- MANIATIS, T., R. C. HARDISON, E. LACY, J. LAUER, C. O'CONNELL *et al.*, 1978 The isolation of structural genes from libraries of eucaryotic DNA. *Cell* **15**: 687–701.
- MANIATIS, T., E. F. FRITSCH and J. SAMBROOK, 1982 *Molecular Cloning*. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
- MICHELSON, A. M., S. M. ABMAYR, M. BATE, A. MARTINEZ-ARIAS and T. MANIATIS, 1990 Expression of a MyoD family member prefigures muscle pattern in *Drosophila* embryos. *Genes Dev.* **4**: 2086–2097.
- O'DONNELL, J., H. C. MANDEL, M. KAUSS and W. SOFER, 1977 Genetic and cytogenetic analysis of the *Adh* region of *Drosophila melanogaster*. *Genetics* **86**: 553–566.
- POOLE, S. J., L. KAUVAR, B. DREES and T. KORNBERG, 1985 The *engrailed* locus of *Drosophila*: structural analysis of an embryonic transcript. *Cell* **40**: 37–43.
- POSAKONY, J. W., J. A. FISCHER and T. MANIATIS, 1985 Identification of DNA sequences required for the regulation of *Drosophila* alcohol dehydrogenase gene expression. Cold Spring Harbor Symp. Quant. Biol. **50**: 515–520.
- SAVAKIS, C., M. ASHBURNER and J. H. WILLIS, 1986 The expression of the gene coding for alcohol dehydrogenase during the development of *Drosophila melanogaster*. *Dev. Biol.* **114**: 194–207.
- SCHAEFFER, S. W., and C. F. AQUADRO, 1987 Nucleotide sequence of the *Adh* gene region of *Drosophila pseudoobscura*: evolutionary change and evidence for an ancient gene duplication. *Genetics* **117**: 61–73.
- SHERMOEN, A. W., and P. H. O'FARRELL, 1991 Progression of the cell cycle through mitosis leads to abortion of nascent transcripts. *Cell* **67**: 303–310.
- SMITH, G. E. and M. D. SUMMERS, 1980 The bidirectional transfer of DNA and RNA to nitrocellulose or diazobenzyloxymethyl paper. *Anal. Biochem.* **109**: 123–129.
- TAUTZ, D., and C. PFEIFLE, 1989 A non-radioactive *in situ* hybridization method for the localization of specific RNAs in *Drosophila* embryos reveals translation control of the segmentation gene *hunchback*. *Chromosoma* **98**: 81–85.
- THUMMEL, C. S., K. C. BURTIS and D. S. HOGNESS, 1990 Spatial and temporal patterns of *E74* transcription during *Drosophila* development. *Cell* **61**: 101–111.
- WOODRUFF, R. C., and M. ASHBURNER, 1979 The genetics of a small autosomal region of *Drosophila melanogaster*, including the structural gene for alcohol dehydrogenase. I. Characterization of deficiencies and mapping of *Adh* and visible mutations. *Genetics* **92**: 117–132.
- XU, G. F., P. O'CONNELL, D. VISKOCIL, R. CAWTHON, M. ROBERTSON *et al.*, 1990 The neurofibromatosis type 1 gene encodes a protein related to GAP. *Cell* **62**: 599–608.

Communicating editor: J. A. BIRCHLER