THE MUTATIONAL BEHAVIOR OF THE R^r :JANA ALLELE IN ZEA MAYS¹

M. H. EMMERLING

Cornell University, Ithaca, New York

Received December 14, 1959

PREVIOUS studies (STADLER and NUFFER 1953; STADLER and EMMERLING 1954) have shown that the action of the standard R^r allele upon anthocyanin is due to two closely linked genes which are separable by crossing over. The identification of a sufficient number of crossovers within the R complex, together with evidence on the sequence of mutational steps, proved the existence of independent components.

However, most of the experimental evidence on which this conclusion is founded is derived from studies with the allele R^r :Cornell. STADLER (1946), working with various R^r alleles of diverse origin, found that spontaneous mutation usually affects seed color and plant color independently. But in none of the former studies had there been a check on crossing over. Consequently, it remains to be determined whether all R^r alleles possess separable components.

The most common type of mutational sequence observed by STADLER (1948) is illustrated in Figure 1.

The R^r alleles which mutate regularly to r^r (seed color mutation) and R^g (plant color mutation) are Cornell, Boone, Quapaw, Ponca, and Black. Both common types of mutants, r^r and R^g , have given mutations to r^g (loss of seed color and plant color effect). In this group of R^r alleles, mutations of R^r to r^g were rarely found, indicating that the action of the R segment is due to two separate genes rather than to the action of a single gene.

However, this same investigation revealed that the stepwise course of spontaneous mutation shown by R^r :Cornell, Ponca, etc., is not characteristic of all R^r alleles. STADLER (1948) reported that two R^r alleles with dilute pigmentation (R^r :Catspaw and Winnebago) mutate directly to r^g and not to r^r . In the case of these two alleles, he concluded that both plant and seed color are dependent upon a single genetic component.

The present report is concerned with a third type of R^r allele which mutates regularly to r^g and less frequently to r^r . In contrast to R^r :Catspaw and Winne-

FIGURE 1.—Mutational sequence of R^r :Cornell, Boone, Quapaw, Ponca, and Black.

¹ Paper No. 379. Department of Plant Breeding, Cornell University, Ithaca, N.Y.

bago, this allele which is known as R^r :Jana is identified by strong plant color both in the seedling and the flowering stages.

The first indication of a striking difference between R^r :Jana and R^r :Cornell was the finding of a r^g mutant in the heterozygote R^r :Jana/ R^r :Cornell. Since this type of mutant was not expected from mutation of R^r :Cornell, it was suspected that it originated from R^r :Jana. Consequently, a more extensive analysis was made of the types of seed color mutants occurring in stocks homozygous for R^r :Jana. The present study gives no indication of the frequency of mutations affecting plant color.

MATERIAL AND RESULTS

The seed mutation data from the cross of ♀♀ $g R^r$:Jana $K/G R^r$:Jana $k \times \delta \delta g r^g k/g r^g k$ are summarized in Table 1. The stocks of homozygous R^r :Jana were marked on either side of the R complex with g (golden plant; 14 units from R) and K (abnormal knob 10; one unit from R).

Out of a total population of 245,515 female gametes tested, 24 colorless seed mutants were analyzed, and of these all but two were r^g ; none of these cases exhibited defective pollen. Of the 22 r^g mutants produced, 13 were $g r^g K$, seven were $G r^g k$, and two were $G r^g K$ in constitution. The simplest explanation of the origin of these cases is that the plant and seed color determiners of R^r :Jana mutated simultaneously to the double recessive or to r^g . On this assumption the $g r^g K$ mutants would be attributed to mutations in the $g R^r K$ chromosome, and the $G r^g k$ mutants to mutations in the $G R^r k$ chromosome. The origin of the two $G r^g K$ cases in which a crossover occurred may be ascribed to mutations in the $g R^r K$ chromosome with a coincidental crossover between g and R , since the number of crossovers expected by coincidence is about three (or 14 percent). The possibility that these two cases arose from mutations in the $G R^r k$ chromosome with a simultaneous crossover between R and K is unlikely because of the low frequency of crossing over in this region (one percent).

It will be seen in Table 1 that homozygous R^r :Jana also produced two seed

TABLE 1

The frequencies and types of seed color mutants observed in progenies from homozygous R^r :Jana. ♀♀ $gR^rK/GR^rk \times \delta \delta gr^gk/gr^gk$

Culture	Number of seeds examined	Number of colorless seeds	Mutant fraction determined	Phenotype of mutant individuals			
				gr^gk	gr^gk	Gr^gK	Gr^gk
56:825	75,525	6	4/6	1	3	0	0
826	66,970	9	7/9	0	4	0	3
827	23,460	5	3/5	0	2	1	0
828	26,480	1	1/1	0	0	0	1
829	31,135	6	5/6	1	2	1	1
830	21,200	3	3/3	0	1	0	2
58F 3245	745	1	1/1	0	1	0	0
Total	245,515*	31	24/31	2	13	2	7

* Adjusted mutation frequency $24/31 \times 245,515 = 24/190,076$.

color mutants of type $g r^r K$. These are mutants that would be expected if the plant and seed color effects of the R^r segment were due to independent components. On this basis the origin of the two $g r^r K$ mutants would be ascribed to mutations of the seed color determiner R to r in the $g R^r K$ chromosome since the mutant chromosomes are of the same g and K constitution as the $g R^r K$ chromosome.

The most striking demonstration of the difference in mutability comes from a comparison of the types of seed color mutants occurring in stocks homozygous for R^r :Jana and R^r :Cornell. The mutation data of R^r :Cornell along with the data of Table 1 are shown in Table 2. It is evident from this comparison that the r^g class of mutation does not occur in the progeny of R^r :Cornell in spite of the large numbers tested. Of the 110 mutants analyzed in the progeny of R^r :Cornell, approximately 44 percent were the result of oblique crossing over, and 56 percent were the result of intragenic alterations, designated as noncrossover mutations. In the case of homozygous R^r :Jana, all of the mutants tested were mutations of the noncrossover type with the possible exception of the two $G r^g K$ mutants. As mentioned previously, these two cases could be due to noncrossover mutation with a coincidental crossover in the $G-R$ segment.

It is also apparent from the data of Table 2 that the total mutation rate is higher in R^r :Cornell than in R^r :Jana. The calculated frequency of mutation for R^r :Cornell is 3.17×10^{-4} , while the rate determined for R^r :Jana is only 1.26×10^{-4} . This difference could be explained as being due to modifying genes since the data were obtained from different stocks and also in separate growing seasons.

In general it is evident that R^r :Jana mutated more frequently to r^g (rate = 1.16×10^{-4}) than to r^r (rate = 0.11×10^{-4}), and that the mutations were chiefly the result of noncrossover alterations.

A further study was made of R^r :Jana in compounds with R^g noncrossover-1, a plant color mutant derived from R^r :Cornell. In a previous publication (EMMERLING 1958), it was reported that the R segment of R^g noncrossover-1 possesses a recessive plant color component (designated p) and a dominant seed color component (designated S), and that p and S are synaptically homologous. This resulted in plants heterozygous for $p S K$ and PS :Cornell k , in the production of two classes of seed color crossovers, as illustrated in Figure 2. The occurrence of oblique crossing over produced the apparent mutants $g p^{(s)}k$ (or $g r^g k$) and $G P^{(s)}K$ (or $G r^r K$). In addition to the crossover mutants, the compound $g p S K/G P S$:Cornell k yielded two classes of noncrossover mutants namely, $g p s K$

TABLE 2

Comparison of the types and frequencies of aleurone color mutants from homozygous R^r :Cornell and R^r :Jana

Parent	Population	Mutants	Rate (10^{-4})	Noncrossovers		Crossovers		Exceptional cases		
				$gr^r K$	$Gr^r k$	$Gr^r K$	$gr^r k$	$gr^g K$	$Gr^g K$	$Gr^g k$
$g R^r$:Cornell $K/G R^r$:Cornell k	346,794	110	3.17	28	34	25	23	0	0	0
$g R^r$:Jana $K/G R^r$:Jana k	190,076	24	1.26	2	0	0	0	13	2	7

FIGURE 2.—Oblique crossover mutants resulting in loss of seed color.

(or $g r^g K$) resulting from mutation in the $g P S K$ chromosome, and $G P s k$ (or $G r^r k$) resulting from mutation in the $G P S k$ chromosome.

Thus if R^r :Jana were like R^r :Cornell in its mutational behavior, the heterozygote R^r :Jana k/R^g noncrossover-1 K , should yield noncrossover mutants of types $r^r k$ and $r^g K$, and also oblique crossover mutants of types $r^r K$ and $r^g k$. But in view of the results reported herein, the data from the heterozygote are complicated by the fact the R^r :Jana mutates to r^g in one step. In other words, the origin of the $r^g k$ mutant could be ascribed either to mutation in the R^r :Jana k chromosome, or to the occurrence of oblique crossing over (Figure 2, type A). However, the information obtained from tests of the heterozygote does provide a basis for estimating the frequency of r^r mutations from R^r :Jana, and for determining whether the plant and seed color effects of R^r :Jana are separable by crossing over.

Two stocks of R^r :Jana/ R^g noncrossover-1 were used as female parents in backcrosses to r^g/r^g (Table 3). Section A includes the results from the cross $G R^r$:Jana $k/g R^g$ noncrossover-1 $K \times g r^g k/g r^g k$, in which the stock carried both g and K , and section B similarly includes tests of R^r :Jana k/R^g noncrossover-1 K but without the g marker.

In culture A, 20 mutants were analyzed, and of these 18 were r^g and two were r^r . Of the 18 r^g mutants produced, eight were of type $g r^g K$ resulting from mutations in the $g R^g K$ chromosome, and one was of type $G r^g K$ originating from mutation in the $g R^g K$ chromosome with a coincidental crossover in the $g-R$ region. The other nine r^g cases are of the type which could be produced by mutation of R^r :Jana to r^g as well as by oblique crossing over. However, the most probable mechanism involved in the production of six of the nine mutants, the $G r^g k$ cases, is that they arose from gene mutations in the $G R^r$:Jana k chromosome to $G r^g k$. If these six cases were ascribed to the occurrence of oblique crossing over (Figure 2, type A), it would be necessary to assume that a crossover occurred between P and S as well as a crossover between G and P . The expected frequency of this double crossover is approximately 0.005 percent. In the case of the three remaining mutants of type $g r^g k$, oblique crossing over could account for the phenotype observed without postulating the occurrence of a second crossover in the $g-R$ segment. Another possibility to consider in regard to the origin of the three $g r^g k$ cases is that they arose from gene mutations in the $G R^r$:Jana k chromosome with a coincidental crossover in the $g-R$ region.

TABLE 3

The frequencies and types of seed color mutants observed in progenies from heterozygous R^r :Jana. ♀♀ R^r Jana/ R^g Cornell × ♂♂ r^g/r^g

Culture	Number of seeds examined	Number of colorless seeds	Mutant fraction determined	Phenotype of mutant individuals								Df†
				<i>r^rk</i>		<i>r^gK</i>		<i>r^rK</i>		<i>r^gk</i>		
A. <i>GR^r</i> :Jana <i>k/gR^g</i> :Cornell <i>K</i>												
56: 822	47,665	12	11/12	<i>g</i>	<i>G</i>	<i>g</i>	<i>G</i>	<i>g</i>	<i>G</i>	<i>g</i>	<i>G</i>	
75:1308	8,137	4	4/4	0	0	5	0	0	1	2	3	0
1309	7,262	2	0/2	1	0	1	1	0	0	0	1	0
1317B	2,157	1	1/1	0	0	0	0	0	0	0	0	0
1415	3,354	1	0/1*	0	0	0	0	0	0	0	0	0
77:1563	10,897	3	3/3	0	0	1	0	0	0	0	2	1
1564A	3,249	1	1/1	0	0	0	0	0	0	1	0	0
1564B	2,405	0	0/0	0	0	0	0	0	0	0	0	0
Total	85,126	24	20/24	1	0	8	1	0	1	3	6	1
B. <i>R^r</i> :Jana <i>k/R^g</i> :Cornell <i>K</i>												
56: 820	48,628	4	4/4	0		3		0		1		1
821	80,243	20	17/20	0		12		0		5		1
823	62,590	13	13/13	1		6		1		5		2
824	25,539	8	6/8	0		5		0		1		0
75:1318	2,855	1	1/1	0		0		0		1		0
Total	219,855	46	41/46			26		1		13		4
Sum total	304,981†	70	61/70	2		35		2		22		5

* One mutant not analyzed.

† Adjusted frequency 61/265,769.

‡ Number of visible cytological deficiencies.

The origin of the one $G r^r K$ mutant may be attributed to two alternative mechanisms. The most probable one is that it originated as the result of oblique crossing over (Figure 2, type B). Another less likely possibility is that it arose by mutation in the $G R^r$:Jana k chromosome to $G r^r k$ followed by a crossover in the $R-K$ region. However, this event is expected only in one percent of the mutants analyzed. In the case of the one mutant of type $g r^r k$, the most probable mechanism that could account for its origin is by gene mutation of $G R^r$:Jana k to $G r^r k$ with a simultaneous crossover in the $g-R$ segment.

In culture B, 41 mutants were analyzed, and of these one was of the $r^r K$ class resulting from either oblique crossing over or mutation with a coincidental crossover in the $R-K$ segment (one percent), and one was of type $r^r k$ originating from mutation of R^r :Jana k to r^r . Of the 39 r^g mutants produced, 26 were $r^g K$ in constitution and 13 were $r^g k$ in type. The $r^g K$ cases may be ascribed to mutations of the $R^g K$ chromosome, while the $r^g k$ mutants may be attributed to either mutations of R^r :Jana to r^g or to the occurrence of oblique crossing over (Figure 2, type A).

The totalled data for cultures A and B in Table 3 show that 57 of the 61 mutants analyzed were r^g in constitution, and of these, 22 could be ascribed to mutations

of R^r :Jana if oblique crossing over of type A were excluded. The total mutation rate of R^r :Jana to r^r was 0.08×10^{-4} ($2/265,769$). The frequency of oblique crossing over was $2/265,769$ (0.08×10^{-4}). In addition to mutations, five plants were found to have cytological deficiencies and abortive pollen.

CONCLUSIONS

It seems clear from the comparison of the homozygous R^r :Jana and homozygous R^r :Cornell that a large proportion of the seed color mutants from R^r :Jana were of the type that would be expected if (P) and (S) were a single entity with two kinds of action. Whether, however, the R segment of R^r :Jana actually consists of a single element cannot be definitely concluded on the basis of the present results since a small proportion of the mutants were of the type expected from the action of two independent genes. For the moment, the significance of this finding still remains unclear. Although it is possible that the seed color factor of R^r :Jana mutated independently of the plant color element, it is also conceivable that the origin of the r^r mutants may be attributed to some other mechanism which is not apparent at the present time. It is also puzzling, in view of the high frequency of oblique crossing over in stocks of R^r :Cornell, to find so few oblique crossovers in the progeny of R^r :Jana. It may be that the structure of R^r :Jana is a complex of two spatially segregated parts so oriented as to be nearly incapable of separation by crossing over. Furthermore, the available data on Jana give no evidence of the frequency of mutations involving loss of the plant color effect (R^g) without loss of the seed color effect. To test this possibility it would be essential to examine a large number of seedlings from homozygous R^r :Jana in order to identify the plant color mutants. Consequently, it would be decidedly premature to conclude from the evidence on hand that the action of R^r :Jana is due to a single element.

One may suggest from these results that the origin of the r^g mutants is due to a suppressive type of mechanism, inhibiting both plant and seed color. However, there is no evidence that indicates such a mutator system is present in the stocks of R^r :Jana. Approximately 200 seeds of each of the r^g mutants were closely examined for dominant mutations, but no spots or sectors of colored aleurone were found. A similar examination was made of the plant tissues, both in the seedling and flowering stages and also in the seedling stage of the F_1 progeny. Again no evidence of reverse mutations was found. It might be relevant to mention here that another R^r allele was found whose mutational behavior is similar to R^r :Jana, but its mutant progeny were highly unstable. This allele, which is known as R^r :Nanson, mutates directly to r^g in one mutational step and also to R^g . However, in succeeding generations reverse mutations were identified in the aleurone and plant tissues.

Another way of explaining the observed r^g mutants is on the assumption that in R^r :Jana, null elements are present either to the left or right of the R^r segment. This would result in the production of gametes with allelic types equivalent to r^g if crossing over occurred following oblique synapsis of the null element with either (P) or (S) (see STADLER and EMMERLING 1956 for discussion of null elements). However, if oblique crossing over were the sole cause of the r^g mutations,

all of the mutants should show the occurrence of crossing over in this interval but not in the $G-R$ segment except for the possible occurrence of double crossing over, which is improbable within this short segment. To account for the origin of the r^o mutants observed in terms of null elements, the occurrence of double crossing over would have to be assumed in the parental stocks of 20 of the 22 r^o mutants identified. Thus, it does not seem likely that oblique crossing over was the major source of the exceptional r^o mutants.

SUMMARY

The allele R^r :Jana mutates regularly to r^o (colorless seed, colorless plant) and less frequently to r^r (colorless seed, colored plant). These results were compared with those of a previous experiment involving the allele R^r :Cornell, in which the R^r segment mutates with appreciable frequency to r^r and rarely to r^o . The pronounced difference between the two R^r alleles may be ascribed to the apparent absence of two separable seed and plant color components in the case of R^r :Jana, and the existence of two independent elements within the R segment of R^r :Cornell. Alternative explanations were discussed.

LITERATURE CITED

- EMMERLING, M. H., 1958 An analysis of intragenic and extragenic mutations of the plant color component of the R^r complex in *Zea mays*. Cold Spring Harbor Symposia Quant. Biol. **23**: 393-407.
- STADLER, L. J., 1946 Spontaneous mutation at the R locus in maize. I. The aleurone-color and plant-color effects. Genetics **31**: 377-394.
- 1948 Spontaneous mutation at the R locus in maize. II. Race differences in mutation rate. Am. Naturalist **82**: 289-314.
- STADLER, L. J., and M. H. EMMERLING, 1954 Problems of gene structure. III. Relationship of unequal crossing over to the interdependence of R^r elements (S) and (P). Science **119**: 585.
- 1956 Relation of unequal crossing over to the interdependence of R^r elements (P) and (S). Genetics **41**: 124-137.
- STADLER, L. J., and M. G. NUFFER, 1953 Problems of gene structure. II. Separation of R^r elements (S) and (P) by unequal crossing over. Science **117**: 471-472.