THE CYTOGENETICS OF "FLAKED," A VARIATION IN TOMATO AFFECTING TWO CELL LAYERS¹

J. W. LESLEY AND MARGARET M. LESLEY

*Department of Horticulture, University of California Citrus Experiment Station,
Riverside, California*

Received February 6, 1961

THERE are at least six current explanations of variegation: plastid differences; position effect (LEWIS 1950) in some manner related to heterochromatin; controlling units including *Ac-Ds* (McCLINTOCK 1950), *Mp* (BRINK and NILAN 1952), and *En* (PETERSON 1960); somatic crossing over (STERN 1936; JONES 1937) revealing hidden heterozygosity; a gene for stickiness of the chromosomes (BEADLE 1932); and irregular distribution in the soma of a ring or rod chromosome. The flaked variegation appears to be due to irregular distribution of a small chromosome fragment.

Somatic lagging and loss occur when a ring or rod becomes dicentric; this may also result when a fragment is telocentric. Like ring chromosomes, telocentrics may break or double their size. The functioning of a centromere may be either weak or strong. In general, a chromosome with a very short arm has a weakly functioning centromere; therefore bridge loss tends to occur (PTASHNE 1960). An extra telocentric fragment has been described as inducing variegation in maize (RHOADES 1940) and in wheat (SEARS 1952b).

A ring chromosome causing variegation may be part of a normal chromosome complement (McCLINTOCK 1932, 1938) or it may be present in addition to a normal complement (FROST, LESLEY and LOCKE 1959).

A chromosome fragment with a centromere at or near one end is the cause of variegation in maize, wheat, tobacco (CLAUSEN 1930) and stock. The fragment producing variegation may be either a ring or a rod and it may be as large as half a chromosome or so small as to be barely perceptible. It seems possible that weakly staining minute fragments may account for other unsolved cases of variegation.

MATERIALS AND METHODS

Seeds of the tomato variety Canary Export were soaked in a solution containing the radioactive isotope P^{32} as phosphate. The initial activity per seed was $10.9\mu\text{C}$ (LESLEY and LESLEY 1956). An R_1 plant which was slow-growing and nearly sterile but otherwise like the parent type produced 11 plants from self-pollination. Eight of these R_2 plants were moderately fruitful. The other three

¹ Paper No. 1292, University of California Citrus Experiment Station, Riverside, California.

plants were less fruitful and included a new phenotype 54.27G1, which was termed "flaked."

The chromosomes were studied in aceto-orcein smears or, with the best results, in acetocarmine by the Walters method (BROWN 1949). For genetic study in-breeding for four generations and outcrossing with linkage testers were employed. The common tomato is normally self-pollinating. For better control, unless otherwise noted, all plants were hand-pollinated and, if emasculated in the field, protected with paper bags.

Description of flaked: The name flaked was suggested by the external appearance of the leaves. They are composed of small, nearly white flake-like areas surrounded by normal green tissue (Figure 1). The "white" areas usually have a green tinge due to underlying green tissue and the term "white" is only relative. White flakes are scattered irregularly and are more conspicuous on the upper surface of the lamina. The calyx may be variegated. A similar greenish white variegation occurs on the stem but consists of longitudinal stripes varying in width and length but often clearly defined. The unripe fruit has green and white sectors extending from basal to distal end, varying in width, and resembling the stem in arrangement. Ripening is abnormally slow, and as it proceeds, the white sectors, which are usually narrower than the green, fail to form lycopene in the outer cortex so that the ripe fruit is striped with red and nonred sectors. The skin color of the fruit is not affected. The cotyledons usually are slightly variegated but less so than the first leaves.

Flaked plants are smaller and have smaller leaves than normal plants of the variety in which the mutation originated. As a rule the greater part of the leaf and stem surface is normal green. The more extensive the white tissue the smaller the leaf. Leaflets which are predominately white may be reduced sometimes to mere vestiges. The lamina is sometimes distorted due to the slower growth of the white tissue. The white stem stripes are slightly sunken. The fruit is malformed if wide white sectors are present.

Variegation of another kind occurs in which the epidermal hairs are absent. Normal plants are hairy, having the trichomes and glandular hairs typical of the parent type. The epidermis of flaked is usually composed of hairy and of glabrous areas which are without trichomes and glandular hairs; the hairy areas tend to be more extensive. The pattern is similar to that of the green and white variegation. The two kinds of variegation seem to occur independently so that the four combinations, green hairy, green glabrous, white hairy, white glabrous, are found on the stems and leaves (Figure 2). Occasionally a whole shoot occurs of which the epidermis of stem, leaves, and flower is glabrous (Figure 3). The underlying tissue has the usual green and white variegation. The leaves and flowers tend to be small and the anthers are dialytic due to the absence of interlacing hairs.

The extent of white and of glabrous tissue varies in different parts of the same plant. A wholly glabrous shoot conspicuous by its glossy appearance occurs on about one fifth of the unpruned flaked plants. Some shoots appear to be wholly hairy but close examination discloses small glabrous stripes. The amount of varie-

FIGURE 1.—Leaf of flaked tomato showing green and white variegation. Lamina more distorted on right side.

gation sometimes differs considerably in sibs from selfing a variegated plant. The white areas especially may be large and the lamina distorted or small and inconspicuous. Figure 1 shows a moderately variegated leaf.

FIGURE 2.—Section of fresh stem of flaked showing variegation in chlorophyll and hairiness in outer cortex and epidermis, and the four combinations green (dark tissue) hairy, green glabrous, nongreen hairy, and nongreen glabrous.

Flaked is male fertile but subnormal in the amount of stainable pollen and is female fertile although it produces less fruit and seed from self-pollination than a normal plant. Even glabrous shoots produce some stainable pollen but no fruit even after the application of normal pollen. The dialytic staminal column partially accounts for this unfruitfulness. The dialytic sparsely hairy *dl* mutant described by RICK (1956) produced a normal crop only when artificially self-pollinated.

In sections of fresh leaves of flaked (Figure 4) the normal green palisade cells

FIGURE 3.—Shoot of flaked tomato with green and white variegation, and glabrous lateral shoot (right).

of the upper surface are absent in white areas and are more or less completely replaced by nongreen spongy tissue. The spongy tissue of the lower surface may also be deficient in chlorophyll. Under the white stem stripes the cell layers of the outer cortex contain no chlorophyll.

Flaked shoots were grafted on two normal plants and a normal shoot on flaked. The scions and stocks were unchanged, indicating no graft transmission and that flaked is not due to a virus disease.

FIGURE 4.—Section of fresh leaf of flaked with palisade layer to one side of midrib interrupted by nongreen tissue.

Cytology: The original flaked R_2 plant, 54.27G1, was small and had very short internodes. Its R_2 sibs were rather pale green and of reduced fertility but were not variegated or dwarfed. That 54.27G1 was unfruitful was not surprising since it was cytologically very abnormal. The p.m.c. contained 12 pairs of chromosomes and one small fragment or nine pairs, two trivalents and one small fragment at diakinesis and M1. Rarely a ring of six was found. It is evident that the two members of one pair had undergone reciprocal translocation with two different chromosomes. Since a fragment was formed in a diploid (12 pair) R_1 plant, the chromosome of which the fragment was a part must have been deficient. The fragment was present in many p.m.c. It therefore presumably had an attachment constriction. At A1 it often lagged and was lost like other unpaired chromosomes in meiosis. A bridge was sometimes found at A1 indicating that an inversion was present as well. Since the bridge was often complex the inversion was probably in one of the chromosomes that forms a trivalent.

The variegated R_2 plant produced seeds from pollination with a normal plant. Plants from this backcross were nonvariegated. The only plant from selfing was sterile and variegated in the same manner as its parent. It was cytologically similar also, but was much more vigorous and less dwarfed than 54.27G1. It produced no progeny.

An F_2 from 54.27G1 by a normal plant of the same variety contained 14 plants of various types. One was albino, another resembled the gene mutant wiry (w_s) but none was flaked. A nonflaked F_2 plant was pollinated by 54.27G1. The resulting family 56.6 varied considerably in appearance, fruitfulness, and cytological condition. Two plants resembled the wiry mutant. It is evident from Table

TABLE 1
Family 56.6 from F_2 55.105.8 (normal plant) $\times$ 54.27G1 (flaked R_2 plant)

Plant number	Plant type	Pollen* type	Chromosome type	Fruitfulness†
56.6.1	flaked	+ or $\pm$	12 pairs + 1 fragment	—
56.6.2	flaked	+ or $\pm$	12 pairs + 1 whole + 1 fragment or trivalent	$\pm$
56.6.3	normal	+ or $\pm$	12 pairs (1 tetravalent)	+
56.6.4	flaked	$\pm$	12 pairs + 1 whole + 1 fragment	—
56.6.5	flaked	+	12 pairs + 1 fragment	1 fruit
56.6.6	mutant	+	12 pairs (1 weakly joined pair)	$\pm$
56.6.7	mutant	0	12 pairs + 1 fragment (1 tetravalent)	—
56.6.8	normal	0	12 pairs + 1 whole	—
56.6.9	flaked	0	12 pairs + 1 fragment	1 fruit
56.6.10	flaked	$\pm$	12 pairs + 1 trivalent + 1 fragment	1 fruit
56.6.11	flaked	0	12 pairs + 1 fragment (1 tetravalent)	—

* Pollen scale: 50 percent stainable (+), 25 percent stainable ($\pm$), 0 percent (0).

† Fruitfulness: good (+), fair ($\pm$), poor (—).

1 that all flaked plants in this family had 12 pairs of chromosomes and a small fragment and that tetravalence may occur in either flaked or nonflaked plants.

Another family (56.98) similar in origin to 56.6, from a nonflaked parent selfed, gave two nonflaked and three flaked plants. The flaked plants had 12 pairs and a small fragment.

The discussion of flaked which follows refers to plants obtained by selfing or by crossing with nonflaked lines. These plants were free from multivalence and no longer produced either albino or wiry mutants.

Flaked plants always have 12 pairs and one or two small fragments in some of their p.m.c. The glabrous sterile shoots also had this chromosome complement so far as tested. Since the fragment is so small (Figures 5 and 6) and tends to be weakly stained, it is impossible to be sure that it is not present in cells in which it cannot be seen. It is most easily seen at lateral M1 since it was often above or below the metaphase alignment. It has also been seen lagging or dividing belatedly at A1 and M2. It is sometimes attached to a pair of chromosomes at diakinesis and M1. When one fragment is present it is often lost in meiosis; when two fragments are present they are often weakly paired but one of them may be attached to the pair referred to above, the other free. As far as could be ascer-

FIGURE 5.—First metaphase in p.m.c. of flaked. Fragment is exceptionally deeply stained.

FIGURE 6.—Second metaphase in p.m.c. of flaked; undivided fragment going belatedly to one plate.

tained the size of the fragment has remained very much the same for eight generations of transmission. One pair of chromosomes is often weakly associated; this may result from competition for pairing. It might also indicate an injured centromere region or a deficiency in one or both homologues.

In F_1 , 12 pairs of chromosomes and a fragment are usually present but in one plant no fragment was found. Since F_1 plants always appear normal, it is assumed that flaked plants contain a pair of deficient chromosomes and a fragment. A deficient chromosome and a fragment are complementary, making production of normal-looking tissue possible. In F_1 the presence of one unaltered chromosome results in a normal-looking plant. Plants containing two deficient chromosomes but lacking a fragment are nonviable.

At pachytene the fragment has been seen attached by one end near the centromere region of chromosome 6 (BARTON 1950). This chromosome has three large heterochromatic regions (Figure 7) and can therefore be identified in many

FIGURE 7.—Pachytene in p.m.c. of flaked. The fragment is attached near the centromere of chromosome 6.

FIGURE 8.—Mitosis in tapetal cells of flaked. The undivided fragment lags. In one plate the central chromosomes were not countable.

cells. The fragment is rod-shaped with two small darker-staining regions. In F_1 pairing seems normal in chromosome 6 except for a possible nonconformity within the heterochromatic portion of the short arm.

In tapetal cells the fragment has been seen to lag (Figure 8). The lagging of the fragment in mitosis accounts for the somatic instability of flaked and for the probable absence of the fragment in some p.m.c. of F_1 plants.

Genetics of flaked: Flaked is self-fertile. Eleven families containing 231 plants which originated from controlled self-pollination of flaked parents were all flaked. F_1 flaked $\times$ nonflaked was nonflaked in ten families and the reciprocal cross gave eight families of nonflaked. Flaked was completely recessive. The F_2 data (Table 2) from crosses of flaked with unrelated lines containing various recessive genes seem to be conflicting. Flaked plants were always in a minority and were entirely absent in nine families. In 56.163.1 the observed ratio 186:0 differs very significantly from 15:1, with $\chi^2 = 12$, $P < 0.01$. In all F_2 families combined, the nonflaked:flaked ratio is 1673:161 and fits neither a 3:1 nor a 15:1 ratio. With a 3:1 expectation the heterogeneity is $\chi^2 = 58$, D.F. = 19 and $P < 0.01$, indicating that the F_2 deviations are not due to random sampling.

An arbitrary separation of the F_2 families containing flaked from those not containing it was made. In the 11 families containing flaked the ratio is 998:161 or 6.2:1 so that flaked was transmitted to about one seventh of the F_2 generation. By the method of BRAND and SNEDECOR quoted by MATHER (1938), with the expected ratio 6.2:1, the heterogeneity $\chi^2 = 18.5$, with D.F. = 10, and $P = 0.05$, is not significant. With a 3:1 expectation the deviation χ^2 is 76, D.F. = 10 and $P < 0.01$ and with 15:1, the deviation is even greater.

TABLE 2
F₂ from flaked $\times$ nonflaked and the reciprocal cross

F ₁ parents*	P ₁	F ₂		Seed germination (percent)
		Nonflaked	Flaked	
55.24.10	Flaked $\times$ nonflaked	5	3	45
55.24.11	Flaked $\times$ nonflaked	22	1	85
57.103.1	Flaked $\times$ nonflaked	360	55	85
58.181.4	Flaked $\times$ nonflaked	261	56	99
59.092.1	Flaked $\times$ nonflaked	53	7	94
59.091.4	Flaked $\times$ nonflaked	55	8	97
60.030.1	Flaked $\times$ nonflaked	51	0	89
56.137.1	Flaked $\times$ nonflaked	30	0	91
56.163.1	Flaked $\times$ nonflaked	186	0	97
59.131.1	Flaked $\times$ nonflaked	64	0	100
59.131.2	Flaked $\times$ nonflaked	56	0	88
60.069.5	Flaked $\times$ nonflaked	51	6	86
60.025.1	Flaked $\times$ nonflaked	50	3	83
57.102.1	Nonflaked $\times$ flaked	77	16	98
59.129.1	Nonflaked $\times$ flaked	56	5	95
59.136.1	Nonflaked $\times$ flaked	10	1	35
56.139.1	Nonflaked $\times$ flaked	55	0	86
59.060.2	Nonflaked $\times$ flaked	53	0	83
59.134.1	Nonflaked $\times$ flaked	72	0	75
59.130.1	Nonflaked $\times$ flaked	62	0	97

* All F₁ parents were nonflaked.

No significant difference was found in F_2 from reciprocal crosses. Flaked plants occurred in three families from F_1 nonflaked $\times$ flaked. This evidence indicates that inheritance is not cytoplasmic. In F_2 families 58.181.4 and 59.091.4 seed germination was high but only 17 percent were flaked; in F_2 families 56.163.1 and 59.131.1 (Table 2) originating from flaked $\times$ nonflaked and the reciprocal no flaked occurred. The deficiency of flaked was not due to low seed viability.

The genetic evidence is consistent with the theory that flaked contains a pair of deficient chromosomes and one or more fragments. Controlled self-pollination of flaked gave only flaked progeny. F_1 plants were all nonflaked. All flaked plants contain one or more fragments and are somatically unstable. The sporocytes of F_1 plants contain a normal chromosome, a deficient chromosome, and may or may not contain a fragment. Only those containing a fragment will transmit flaked and probably only gametes containing a normal chromosome or a deficient chromosome and a fragment are viable. The proportion of flaked in F_2 will depend on the presence or absence of the fragment in the sporocytes. Figure 9 shows that from sporocytes containing one fragment the expected F_2 ratio, nonflaked:flaked, is 8:1. The observed ratios from parents 59.092.1 and 59.091.4 (Table 2) approach this but other ratios deviate from it in both directions. The proportion of flaked is expected to be higher if one or both sporocytes contained two fragments and it

FIGURE 9.—The possible combinations of gametes of F_1 normal $\times$ flaked having one normal, one deficient, and one fragment chromosome, if only gametes containing a whole chromosome or a deficient chromosome and a chromosome fragment are viable.

Fragment is shaded, chromosome lacking shaded portion is deficient.

would be lowered by loss of the fragment, which is known to occur in meiosis or if the sporocytes lacked the fragment. The karyotype (Figure 9) containing two normal chromosomes and one or two additional fragments has not been identified. It would not be flaked and would be trisomic for a portion of chromosome 6.

DISCUSSION

Pachytene study suggests that the deficient chromosome and fragment of flaked are components of chromosome 6, which contains linkage group IV. There is, in this chromosome, a recessive gene *md* (*mo*) which originated from thermal neutron radiation. The gene *md* causes small white spots on the leaves, and, in this respect, seems to be similar to flaked. Unlike flaked *md* is temperature sensitive (BURDICK 1959) and has normal gametic and zygotic viability. Mendelian inheritance of flaked is certainly not indicated on any conventional basis. Moreover, in flaked plants variegation in chlorophyll and in hairiness are always associated. Lack of hairs can occur without white flaking in unrelated material but this is not true in the flaked line. It therefore seems probable that in flaked the fragment contains two loci which are seldom, if ever, separated. The original line in which flaked occurred did not contain the recessive alleles *yg2*, *yg3* and *md*. It is possible that a mutation similar to *md* occurred when the fragment was broken off.

The variegation in chlorophyll and in hairiness may be due simply to the different effect of the presence or absence of the fragment in subepidermal and epidermal layers. If the fragment is present, the epidermis is hairy, if absent, it is glabrous. In the subepidermal layer, chlorophyll and palisade cells are normal if the fragment is present, but, if absent, no chlorophyll or palisade cells are found. Many geneticists believe and cytological observations confirm that the chromosome content of these layers is normally the same. If so, the deficient chromosome has a very different effect in different cell layers. Presumably different loci are active in the epidermal and subepidermal layers.

The observed variation in the degree of flaking in different plants is explained by variation in frequency of somatic fragment loss. It might also be due to breakage of the fragment and consequent gene loss.

That telocentric fragments tend toward loss in the soma and misdivision, as well as toward union to form isochromosomes, was demonstrated by DARLINGTON (1939). RHOADES (1940) and SEARS (1952 a, b) showed that similar behavior of telocentric fragments results in variegation in rye and in wheat. The centromere of the fragment in flaked is at or near one end. It is possible that neither the altered chromosome 6 nor the fragment has a complete centromere. It was noted above that one pair of chromosomes in flaked is often more or less separate at M1.

In chromosome 6, a large heterochromatic region lies on each side of the centromere, and the fragment must have been derived from one of these. In F_1 , the only observed difference in pachytene pairing appears to be within the heterochromatic mass in the short arm, and the fragment has been found attached to chromosome 6 near this region. Changes in position and amount of

heterochromatin have been adduced as inducing variegation (LEWIS 1950; McCLINTOCK 1950).

The fragment in flaked is very hard to find in some smears. It tends to be much paler than the other chromosomes and, except in very well stained material, could easily be missed. It seems conceivable that fragments too small to be demonstrated with certainty may cause variegation.

SUMMARY

"Flaked," a variegated tomato plant, occurred in R_2 from a seed soaked in a solution of P^{32} as phosphate. Normal green areas are irregularly variegated with small whitish flakes in which the palisade cells are replaced by nongreen parenchyma. Variegation due to loss of hairs also occurs but it is distributed independently and may give rise to glabrous shoots.

Flaked has rather poor pollen and produces few fruits, which vary greatly in seed production. Selfed, it produced only flaked. All flaked plants have either one or two small fragments in addition to 12 pairs of chromosomes. The fragment tends to be lost both in meiosis and mitosis.

All F_1 plants look normal; usually they have 12 pairs of chromosomes and one fragment but occasional F_1 plants lack the fragment.

At pachytene the fragment is a single strand and is sometimes found attached to chromosome 6. Recessive genes affecting chlorophyll are known to be present in this chromosome but the line in which flaked occurred was homozygous for the normal alleles.

Since all flaked plants have the fragment and all F_1 plants from crossing non-flaked and flaked are normal, it follows that flaked plants lack a normal chromosome 6. The normal parts of flaked plants contain a fragment in addition to the deficient chromosome. The white flakes are believed to lack the fragment and are therefore deficient. The partial male sterility of flaked suggests that only gametes containing the fragment in addition to a deficient chromosome are viable. In the epidermis the loss of the fragment results in loss of hairs. Buds from glabrous shoots may have 12 pairs of chromosomes and one fragment. They are always sterile.

Loss of the fragment in the soma of an F_1 plant would give rise to germ cells that lack it, but would not affect the appearance of the plant since all cells contain a normal chromosome 6. F_2 families contained normal and flaked plants in the mean ratio of six normal to one flaked. Several F_2 families, however, contained no flaked plants. Neither a 3:1 nor a 15:1 ratio is indicated. It is believed that loss of the fragment in mitosis may explain the irregular transmission of flaked.

LITERATURE CITED

- BARTON, D. W., 1950 Pachytene morphology of the tomato chromosome complement. *Am. J. Botany* **37**: 639-643.
- BEADLE, G. W., 1932 A gene for sticky chromosomes in *Zea Mays*. *Z. Ind. Abst. Vererb.* **63**: 195-217.
- BRINK, R. A., and R. A. NILAN, 1952 The relation between light variegated and medium variegated pericarp in maize. *Genetics* **37**: 519-544.
- BROWN, S. W., 1949 The structure and meiotic behavior of the differentiated chromosomes of tomato. *Genetics* **34**: 437-461.
- BURDICK, A. B., 1959 Tomato Genetics Cooperative **9**: 21.
- CLAUSEN, R. E., 1930 Inheritance in *Nicotiana Tabacum*. X. Carmine-coral variegation. *Cytologia* **1**: 358-368.
- DARLINGTON, C. D., 1939 Misdivision and the genetics of the centromere. *J. Genet.* **37**: 341-364.
- FROST, H. B., MARGARET M. LESLEY, and W. F. LOCKE, 1959 Cytogenetics of a trisomic of *Matthiola incana*, involving a ring chromosome and somatic instability of singleness (*versus* doubleness) of flowers and shape of leaves. *Genetics* **44**: 1083-1099.
- JONES, D. F., 1937 Somatic segregation and its relation to atypical growth. *Genetics* **22**: 484-522.
- LESLEY, J. W., and MARGARET M. LESLEY, 1956 Effect of seed treatments with X-rays and phosphorus 32 on tomato plants of first, second, and third generations. *Genetics* **41**: 575-588.
- LEWIS, E. B., 1950 The phenomenon of position effect. *Advances in Genet.* **3**: 73-116.
- MCCLEINTOCK, BARBARA, 1932 A correlation of ring-shaped chromosomes with variegation in *Zea Mays*. *Proc. Natl. Acad. Sci. U.S.* **18**: 677-681.
- 1938 The production of homozygous deficient tissues with mutant characteristics by means of the aberrant mitotic behavior of ring-shaped chromosomes. *Genetics* **23**: 315-376.
- 1950 The origin and behavior of mutable loci in maize. *Proc. Natl. Acad. Sci. U.S.* **36**: 344-355.
- MATHER, K., 1938 *The Measurement of Linkage in Heredity*. Chemical Publ. Co. New York.
- PETERSON, P. A., 1960 A mutable pale green mutable system in maize. *Genetics* **45**: 115-133.
- PTASHNE, M., 1960 The behavior of strong and weak centromeres at second anaphase of *Drosophila melanogaster*. *Genetics* **45**: 499-506.
- RHOADES, M. M., 1940 Studies of a telocentric chromosome in maize with reference to the stability of the centromere. *Genetics* **25**: 483-520.
- RICK, C., 1956 Cytogenetics of the tomato. *Advances in Genet.* **8**: 267-382.
- SEARS, E. R., 1952a Misdivision of univalents in common wheat. *Chromosoma* **4**: 535-550.
- 1952b The behavior of isochromosomes and telocentrics in wheat. *Chromosoma* **4**: 551-562.
- STERN, C., 1936 Somatic crossing over and segregation in *Drosophila melanogaster*. *Genetics* **21**: 625-730.