THE SALIVARY CHROMOSOMES OF *DROSOPHILA MICROMELANICA* AND *DROSOPHILA MELANURA*

HARRISON D. STALKER

Department of Zoology, Washington University, St. Louis, Missouri

Received November 16, 1964

THE utilization of salivary gland chromosomes in determining phylogenetic relationships of species belonging to the genus *Drosophila* has been carried out with considerable success in the *D. obscura* group (DOBZHANSKY and EPLING 1944), the *D. virilis* group (STONE, GUEST and WILSON 1960; STONE 1962), and the very large *D. repleta* group (WASSERMAN 1960, 1962, 1963). During the past few years the author has been working on the salivary gland chromosomal polymorphism of the members of the *D. melanica* group, with a view to determining their phylogenetic relationships.

The *D. melanica* group includes seven Nearctic species, of which all but *D. melanissima* have been reared in the laboratory and studied cytologically. Of the six species which have been cultured, *D. melanica* was studied cytologically by WARD (1952), who found the species highly polymorphic, with most of the chromosomal variability confined to chromosome 2, a situation that is characteristic of the species group as a whole. The chromosomal polymorphism of *D. paramelanica*, as well as the associated sex-ratio condition, was described by STALKER (1960, 1961) who also studied the chromosomal polymorphism of *D. euronotus* (STALKER 1964a) and *D. nigromelanica* (STALKER 1964b). The present paper deals with the chromosomes of *D. micromelanica* and *D. melanura*, as well as the evolutionary relationships of these two species with each other and with *D. nigromelanica*.

MATERIALS AND METHODS

The known distributions of *D. micromelanica* and *D. melanura* are indicated in Figure 1. It will be noted that while both species show a markedly disjunct distribution, *D. micromelanica* is clearly southern, while *D. melanura* may be classified as a northern species. Neither species is abundant in collections, and since in both of them the material available to the author was limited, it is probable that some of the chromosomal polymorphism within each of these species is still unknown. The populations sampled are indicated in the tables.

In the case of *D. micromelanica* all laboratory stocks or wild-caught individuals were compared with a structurally homozygous, standard stock derived from the Smithville, Texas strain. In *D. melanura*, a number of different structurally homozygous strains were obtained from pair matings, but all proved to be so weak that only one of them (from Webster Groves, Missouri) survived long enough to be used as a standard in test crosses.

Both species are attracted to fruit, and are characteristically found in deciduous woods. Although neither is a domestic species in the usual sense, *D. melanura* may be taken in resi-

¹ This work has been carried out with the financial assistance of National Science Foundation Grant G20107.

FIGURE 1.—Geographical distributions of *Drosophila melanura* and *D. micromelanica*.

dential areas which support large trees. The larval food is unknown for both species, but some other members of the species group are known to breed on infected tree sap.

D. micromelanica is a small species, (somewhat smaller than *D. melanogaster*). It is relatively easy to rear in the laboratory, and breeds well in pair matings. In many strains mating occurs in adults that are less than 12 hours old. *D. melanura* is a large species, comparable in this respect to *D. euronotus*, but readily distinguished from that species by the coloration, male genitalia and egg-filament shape. MILLER (1944), who described *D. melanura*, stated that it was not easy to culture in the laboratory, and the present author agrees. In this species mating rarely occurs before adults are three days old.

THE CHROMOSOMES

The metaphase chromosomes of *Drosophila melanura* were first described by MILLER (1944). They are essentially like those of most other species of the group, and consist of a pair of long V-shaped chromosomes (XX or XY pair), two pairs of long rods (chromosomes 2 and 3), a pair of short V's (chromosome 4), and a pair of microchromosomes (chromosome 5). A strain derived from the St. Louis area and studied by the author showed oogonial chromosomes of the type described by MILLER (see Figure 2). WARD (1949) has described a somewhat different karyotype for the species, based on a strain from Guarete, Maine. However, he has recently pointed out (personal communication), that there is now some doubt that the Maine strain was in fact *D. melanura*.

FIGURE 2.—The metaphase chromosome configuration of *Drosophila melanura* (to the left), and the four configurations reported for *D. micromelanica* (Micro I, II, III and IV on the right).

The metaphase chromosomes of *D. micromelanica* differ from those of all other species in the group in one important way. In *D. micromelanica* the X chromosome is a rod rather than a large V, and it can be shown by comparison of the salivary chromosome banding patterns that the X of *D. micromelanica* corresponds to the X-right element of the other group members, while the X-left element of the other species is represented by a rod-shaped autosome in *D. micromelanica*. In order to simplify the nomenclature used in chromosome arm identification in the various members of the group, the autosomal arm in *D. micromelanica* which corresponds to X-left in other members of the group is designated in this paper as "XL" or "X-left" rather than being given a special autosomal designation.

The evolutionary separation of a metacentric chromosome into two rods would require the acquisition of an extra centromere, while the reverse change, involving the formation of a V from two rods, would not. For this reason, and because in fact, there are numerous instances in *Drosophila* evolution in which fusions have occurred, while the reverse is known to have occurred only once, (PATTERSON and STONE 1952), it is assumed that the rod-shaped X in *D. micromelanica* indicates an ancestral rather than a derived condition. It is for this reason that in the present paper the gene sequences in *D. micromelanica* are taken as a starting point for the series of inversions which have occurred in the evolution of this species group as a whole.

While all studied strains of *D. micromelanica* have rod-shaped X chromosomes, the various strains have been reported to differ markedly in their autosomal karyotype. These karyotypes, modified from WARD (1949), are shown in Figure 2. It will be noted that fusion of the long autosomal rods has been reported (type II), a centromere shift has been reported in chromosome 4 (compare type I with the other three types), and the microchromosomes may appear either as rods or as dots, presumably due to heterochromatin losses or gains.

Different workers, studying *D. micromelanica* strains from the same locality, have reported quite different karyotypes, suggesting considerable karyotypic variability, not only within the species, but within specific localities, or possibly within populations. The situation, in so far as it has been possible to work it out from the literature and through personal correspondence, is summarized in Table 1.

TABLE 1
Karyotypes reported in D. micromelanica

Locality	Strain No.	Karyotype	Authority
Smithville, Texas	unknown	I	STURTEVANT and NOVITSKI (1941)
Smithville, Texas	unknown	IV	WARD (1949)
Smithville, Texas	unknown	III or IV	This paper
Austin, Texas	unknown	I	STURTEVANT and NOVITSKI (1941)
Perry, Florida	PEF-11	IV	This paper
Inverness, Florida	TAF-10	IV	This paper
Buckingham Co., Va.	VWM-36	IV	This paper
Huachuca Mts., Ariz. (2 strains)	unknown	I	STURTEVANT and NOVITSKI (1941)
Cave Creek, Ariz.	U.T. 1853.9	III	WARD (1949); This paper
Kingman, Ariz.	U.T. 1867.4	III	WARD (1949)
Coronado National Forest, Ariz.	U.T. 2160.12	IV	This paper

See figure 2 for karyotype illustrations. U. T. refers to University of Texas strain numbers. See text in regard to Karyotype II.

In addition (and not included in the table) is the report of WHARTON (1943), who analyzed two strains, one from Austin, Texas, and the other from the Huachuca Mountains, Arizona. One strain was described as type I, the other type II. Unfortunately the author does not indicate which was which, nor are the strain numbers given, thus it is impossible to tell whether one or both of WHARTON's strains were those used by STURTEVANT and NOVITSKI, and thus whether a real disagreement exists.

The Smithville, Texas strain studied by the author could not be classified as belonging to either type III or type IV, since the microchromosomes generally appeared slightly elongated, but less strikingly so than those in type IV of figure 2.

Some ganglion cells of all strains studied by the author showed a rather interesting stickiness of the centromere ends of nonhomologous rod-shaped chromosomes, resulting in what appeared to be V-shaped chromosomes, resembling those of WHARTON's type II. A few cells were found which had double fusions of this sort, but in most cases only two rods were involved. However in all strains studied, some individuals were found within which both "fusion" and "non-fusion" type cells appeared.

On the basis of genuine chromosome arm fusion being likelier than separation, it would appear that WHARTON's type II, with the fused autosomal arms, is derived from one of the other three types, in which such fusion does not occur. Whether or not the rod-shaped chromosome 4 (type I), or the V-shaped 4 (types II, III and IV) is the derived one cannot be decided. Since all other species in the *D. melanica* group have V-shaped fourth chromosomes and no fusion of autosomal arms, it would appear that the other species were derived from type III or type IV of *D. micromelanica*, rather than from type I or type II.

Hybridization tests: Attempts to form hybrids involving the species *D. melanura*, *D. micromelanica*, or both, have been reviewed by STALKER (1960, 1964a, b). The situation is as follows: *D. melanura* has been found to form hybrids (female

fertile, male sterile) with *D. euronotus*, *D. paramelanica* and *D. melanica*. *D. micromelanica* has not been found to form hybrids with any other species. However, STURTEVANT and NOVITSKI (1941), in making crosses between Texas and Arizona strains of *D. micromelanica*, showed that when Texas females were crossed to Arizona males, the F_1 hybrid males were sterile, although the females were fertile. The reciprocal crosses yielded fertile hybrids of both sexes. These authors found that the sterility of the male hybrids was associated with the presence of a Texas X and an Arizona Y chromosome. Crosses made by the present author between the various strains of *D. micromelanica* available to him are summarized in Table 2.

It will be seen from this table that no hybrid male sterility was found between the Texas and Arizona strains used. Male sterility was found, however, when crosses were made between females from Florida or Virginia and males from either Texas or Arizona. No sterility was produced from the reciprocal crosses, nor was there any sterility in the progeny of crosses between any of the three strains from Florida and Virginia.

Thus, on the basis of the tests with these strains, the species could be divided up into an Eastern race from Florida and Virginia, and a Western race from Texas and Arizona, with crosses of Eastern females by Western males always yielding sterile male hybrids.

These results may be reconciled with those of STURTEVANT and NOVITSKI if it is assumed that Texas is an overlap zone for the Eastern and Western races, and that the earlier workers used Eastern strains from Texas, while the present author used Western Texas strains. It is, of course, understood that this may be an oversimplification of the situation, and that in fact the species may be divided up into more than two genetically semi-isolated races.

The inversions of D. micromelanica: The limits of the inversions found to be heterozygous within *D. micromelanica* are indicated by the brackets in Figures 3, 6, 7, 8 and 9; the geographical distributions are shown in Table 3. The data from Texas, Arizona and Virginia come from laboratory stocks; the four Virginia stocks each originated from a single inseminated wild female. The Florida data

TABLE 2

Tests for hybrid male-fertility in interstrain crosses of D. micromelanica.
See Table 3 for more complete locality descriptions

Females	Males					
	Smithville, Texas	2160.12, Arizona	1853.9, Arizona	No. 36, Virginia	Perry, Florida	Inverness, Florida
Smithville, Texas		Fertile	Fertile	Fertile*	Fertile	Fertile
2160.12, Arizona	Fertile		Fertile	Fertile	Fertile	Fertile
1853.9, Arizona	Fertile	Fertile		Fertile	Fertile	Fertile
No. 36, Virginia	Sterile	Sterile	Sterile		Fertile	Fertile
Perry, Florida	Sterile	Sterile	Sterile	Fertile		Fertile
Inverness, Florida	Sterile	Sterile	Sterile	Fertile	Fertile	

* In this cross five different Virginia strains were used. The F_1 males were fertile in all cases.

TABLE 3
Gene sequences in the chromosome arms of *D. micromelanica*

Locality—Strain	Chromosome						
	"X-left"	X	3	4		2	
Smithville, Texas	+ A	+	+ A	+	+	K	GJ
Coronado National Forest, Ariz., U.T. 2160.12	+	+ A	+ A	+	+	D K	GJ
Ramsay Canyon, Ariz., U.T. 2157.3	+ A	+	+	+	+	K	
Cave Creek, Ariz., U.T. 1853.9	+ A	+	+ A	+	+	D K	GJ
Buckingham Co., Va. Strain 36	+ A	+ A AB	+	A	+	A B D	BGJK
Strain 20	+ A	+	+	+	+	B D	BGJK EC
Strain 11	+	+	+	A	+	A B AB D	F FK EC
Strain 9	+	+ A AB	+	A	+	K	GJ
Perry, Florida	+	+ A	+ A	+	+	D DK	DGJK
Inverness, Florida Strain 10	+	A	+	+	+	K DK	DGJK
Wild male	+	A	+	+	+		DGJK

"X-left" refers to the autosomal arm of *D. micromelanica* which is homologous to the left arm of the X in all other members of the species group.

are based on the F_1 progeny of inseminated wild females, and (in the case of Inverness) a single wild male. Each laboratory stock was sampled by analyzing seven to fifteen larvae from each of ten stock females crossed to Standard males. In the case of the wild Florida females, five F_1 daughters from each were crossed to Standard males, and seven to ten larvae analyzed from each daughter. The Florida male was crossed to Standard females, and 21 larvae examined.

In Table 3 various gene sequences are recorded as present or absent. The symbol + refers to the Standard sequence in each case. The data in the table indicate that while in this species all chromosomes show inversion heterozygosity, chromosome 2 is the most variable, as in other members of the species group. *D. micromelanica* is the only group member in which inversion heterozygosity has been found in chromosome 3, although as will be shown below and in a later paper, this chromosome has not remained structurally unchanged during the evolution of the group as a whole.

The data permit no generalization except that apparently the Virginia population is distinct from all others in having seven inversions that all other populations lack (B in the X; A in chromosome 4; A, B, F, E, and C in chromosome 2).

The inversions of D. melanura: *D. melanura* is rarely abundant in collections, and stocks are difficult to maintain in the laboratory. As a result, the inversion heterozygosity uncovered within the species is based on material from only four localities, two of them in the St. Louis, Missouri area (see Table 4). All data presented in the table are based on crosses of wild-caught flies to the Webster Groves, Missouri standard strain. Most of the early Webster Groves material was used to determine the limits of the inversions, but owing to the initial lack of a viable homozygous standard strain, could not be used to determine inversion frequencies. The limits of the inversions are indicated in Figures 3, 6 and 9. Obviously more extensive sampling would be expected to uncover more variability, and the inversions presently known may constitute only a fraction of those actually existing in the species. For reasons that will become clear below, the inversion symbols used in all chromosome arms except X-right are chosen so as not to duplicate those used in *D. micromelanica*, and as a result, the standard sequences in X-left and chromosome 2 of *D. melanura* are given the symbols

TABLE 4

Gene sequences in chromosomes X and 2 in D. melanura, showing inversion differences from the species standard for each arm. See text and footnotes for explanations of symbols

Locality	X-left*		X-right		Chromosome 2†							
	Stand.	F	Stand.	A	Stand.	C'	C'E'	D'	D'E'	C'D'	F'	C'D'E'
Brentwood, Mo.	4	1	4	1	4	1	1	.	.	.	.	.
Webster Groves, Mo.	11	5	14	2	8	1	6	1	1	1	1	.
Terre Haute, Ind.	3	1	4	.	4	.	1	.	1	.	.	.
Bear Camp Gorge, N.C.	1	.	1	.	.	.	1	.	.	.	.	1

* In terms of the species group as a whole, X-left standard sequence carries the inversions BCDE. Thus F in X-left would be BCDEF.

† In terms of the species group as a whole, chromosome 2 standard sequence carries the inversions WXYZA'B'. Thus C' in chromosome 2 would be WXYZA'B'C', etc.

BCDE and WXYZA'B' respectively, rather than the usual +. The inversion symbols listed in Table 4 thus represent single inversion differences from these standard sequences.

Consideration of interspecific changes in banding pattern: The intraspecific variability in the chromosomes of *D. micromelanica* and *D. melanura* has been described above. The chromosomes of the closely related *D. nigromelanica* have been described by STALKER (1964a). We will now consider the structural changes which have occurred during the evolutionary divergence of these species, based on interspecific differences presently found in their banding patterns.

In the earlier cytological work on the members of the *D. melanica* group, it was necessary to assign symbols for the inversions and gene sequences in any given chromosome arm independently for each species, with the result that at present a given symbol (such as B in chromosome 2) has a different meaning in each of the five species in which it has been used. In this and the following paper on the group, a new symbolism will be used in which for any given chromosome arm, a given symbol will be used only once for the group as a whole. In each arm (except X-right) the symbol (+) will refer only to the standard gene sequence in *D. micromelanica*. In the present paper, this means that the symbols of STALKER (1964a) for *D. nigromelanica* would have the following equivalents, with the old symbols listed first. X-left, + = GH; Chromosome 2, + = HLMN-OPQRS, B = HLMNOPQR, A = HLMNOPQRST, C = HLMNOPQRSU; Chromosome 3, + = CB; Chromosome 4, + = AEFghi, A = AEFghij. With the new symbol system, when the alphabet is used up for a given chromosome arm, A', B', C', etc. are employed. It should be emphasized that symbols such as A and A' or B and B' are in no way related.

In the case of the X-right chromosome arm, no attempt has been made to introduce a single system of symbols for the species group, since it has not been possible to work out extensive interspecific homologies for that arm.

To compare the species banding patterns maps must be used, since hybrids cannot be obtained between any of the three forms. Because of the single-armed X in *D. micromelanica* and the compound X in the other *D. melanica* group species, it may be assumed that *D. micromelanica* is the more nearly ancestral member of the group, and the question to be considered is: by what series of structural changes could the banding patterns of *D. micromelanica* have been converted into those found in the other two species? If such changes occurred through a series of inversions, then those gene sequences which still survive as part of the chromosomal polymorphism of any of the three species are of course known, but the presumably extinct intermediate gene sequences must be inferred. If the banding patterns of the living members of any two species are markedly different, then a variety of different series of inversions could lead from one species to the other, and while with sufficient patience and energy it is possible to determine the *minimum* number of inversions by means of which the transformation might have occurred, the actual number will not be known, and may be higher than the estimated minimum. Also the order in which the independent (nonoverlapping) inversions appeared in the series cannot be established.

Reconstructions of interspecific inversion series may be used in two ways. First, they give an easily grasped estimate of the minimum amount of chromosomal rearrangement that has taken place in the evolutionary divergence of any two species. Secondly, they may indicate branched evolutionary pathways. Thus, if the changes in a given chromosome follow the pathway: Species A $\longleftrightarrow$ Species B $\longleftrightarrow$ Species C, and if the chromosomally intermediate species B is ancestral, then a branching phylogeny leading from B to both A and C is indicated.

When photomaps are used for interspecific comparisons of banding pattern they must be supplemented by a very large number of additional photographs of each chromosome region, since the maps themselves show the chromosome in only one condition of bending, puffing, staining and stretching, and short regions that appear to have the same sequence when two species maps are compared, may be shown to differ when a more complete set of photographs is consulted. Conversely, regions that appear to be very different in species maps may be shown to be alike when enough supplementary photographs are studied.

Recent work of LEVINE and VAN VALEN (1962) has indicated that a given section of *Drosophila* salivary gland chromosome may be so changed by a gene mutation that even though no structural rearrangement has occurred, the apparent pattern of the region is entirely different. In the light of this finding, it may be questioned whether any reliability can be placed on comparisons of photomaps of different species, since interspecific genic differences could conceivably result in striking alterations in the appearance of the chromosomes. Such gene-controlled differences in the chromosomes of the *D. melanica* group would certainly introduce a complication in making map comparisons, since even regions which were structurally identical might be impossible to homologize because of apparent differences. On the other hand, the probability that genic changes would cause nonhomologous segments which contained many bands to appear to be homologous, is so remote as to be of little concern, since in the recognition of homologous sections in species maps, very short regions cannot be homologized with assurance under any circumstances, and longer regions are homologized by consideration of their total pattern. Thus while genic modification of the appearance of the banding pattern may cause difficulty, it will not lead to assumptions of homology where in fact none exists.

Comparison of species photomaps is difficult at best, and the method presently used by the author is presented in the hope that it may be of help to others who are interested in using the technique. In starting the comparison of two species, large map prints are prepared on glossy paper which is dried in blotter rolls, care being taken to leave plenty of blank space on each print for making notations. The two photomaps are mounted parallel to each other on heavy cardboard with Magic Tape. After study of the maps themselves it is usually possible to indicate probable homologous regions with grease pencil. Using these preliminary clues as a guide, the collection of reference prints is consulted to test the suspected homology in a given region. The reference numbers of prints used to establish an homology are then noted in ink on Magic Tape next to the pertinent regions,

and the grease pencil marks replaced by ink lines. At the start all "difficult" regions are ignored, since their homologies are frequently suggested after major regions of the maps have been worked out.

Even in those cases in which all chromosome regions can apparently be homologized, two types of error may occur. First, very short inversion differences can be missed entirely. Secondly, the exact limits of homologous regions are difficult to determine in certain cases, and as a result, interspecific inversions which are in fact slightly overlapping may be interpreted as adjacent but independent.

In many of the species comparisons of highly polymorphic chromosomes such as chromosome 2 in the *D. melanica* group, it has been impossible to establish homologies throughout the whole length of the chromosome. The nonhomologized regions are short, and altogether rarely involve more than about five percent of the total length of the chromosome. They may be the result of genic transformation of the apparent pattern of the chromosome, or more likely are sections which have been shifted repeatedly by inversions and are too short to recognize. In the interpretations which follow, such short regions are ignored.

The errors of interpretation introduced by ignoring such doubtful regions have not been investigated mathematically, but a fairly large series of empirical tests was carried out on chromosome models made up to differ by four inversions, and the general conclusion was that inability to homologize short regions may (but need not) result in an underestimation of the number of inversions by which the chromosomes differ, but apparently will not result in an overestimate. In summary then, it appears that the divers errors involved in the use of photomaps for interspecific comparisons will tend to result in an underestimate of the structural differences involved.

The interspecific homologies in chromosome 2: In Figure 3 Standard maps of chromosome 2 are shown for all three species. Homologous regions are indicated by numbered brackets. It will be noted that the map of *D. micromelanica* is numbered and bracketed twice, once above, once below the map. The upper numbering system is to be compared to that of *D. nigromelanica*, the lower to that of *D. melanura*; the two systems are quite independent. Regions which could not be homologized are neither numbered nor bracketed. In those cases in which homologous regions are inverted with respect to each other, this fact is indicated by reversed arrows.

Inspection of the homologous regions in any pair of photomaps sometimes suggests the homology indicated, often the homology is apparent only after inspection of additional photographs. As an illustration of the way in which such additional photographs may be useful, Figure 4 has been prepared, dealing with the chromosome 2 homology of *D. micromelanica* and *D. melanura*. In this figure the first four distal regions (1-6, 7-12, 13-19, and 21-25) are considered. For each region a sample of three photographs from *D. micromelanica* is compared with three from *D. melanura*. Careful study of this plate will indicate that while any pair of photographs gives very dubious evidence, consideration of all three photographs for each species leaves little doubt that the regions are in fact homologous. Study of the many additional photographs available makes the homologies even more convincing.

FIGURE 3.—The Standard gene sequences in chromosome 2 of *Drosophila nigromelanica*, *D. micromelanica* and *D. melanura*. The proximal ends of the three arms are indicated by heavy arrows. The lettered brackets indicate the limits of the intraspecific inversions known in each of the three species. The numbers indicate regions of the chromosomes showing interspecific homology in banding pattern. The system of numbers *below* the map of *D. micromelanica* should be compared with that of *D. melanura*. The numbers *above* the *D. micromelanica* map should be compared with those of *D. nigromelanica*. The two numbering systems in *D. micromelanica* are independent. Regions for which no homologies could be established are unnumbered. In those cases in which the banding sequence runs in opposite directions in the two species, this is indicated by reversed arrows.

FIGURE 4.—A sample of voucher photographs for the first four sections of chromosome 2 in *D. micromelanica* compared with the corresponding sections in *D. melanura*. In each set of six photographs the upper three are *D. micromelanica*, the lower three *D. melanura*. The upper right-hand photograph for region 7—12 in *D. micromelanica* is incomplete on the right end. See text.

Several different series of inversions might transform the Standard chromosome 2 of *D. micromelanica* into that of *D. melanura*. At least six inversions would be required in all, and six such inversions are indicated in Figure 5A.

The structural differences between Standard chromosome 2 of *D. micromelanica* and that of *D. nigromelanica* would require at least nine inversions, indicated in Figure 5B. Of these, inversion S is presently heterozygous within *D. nigromelanica* populations.

It will be noted from examination of Figure 3 that the homologous regions for *D. micromelanica* and *D. melanura* have entirely different limits from those homologous for *D. micromelanica* and *D. nigromelanica*, and that the *D. micromelanica* chromosome 2 is a structural intermediate between those of the other two species.

Homologies in X-left: The homologies between the autosome "X-left" of *D. micromelanica* and X-left in the other two species are indicated in Figure 6.

The transition from Standard in *D. micromelanica* to that in *D. nigromelanica* may be accomplished by the occurrence of two inversions, starting with *D. micro-*

FIGURE 5.—(A). The series of six interspecific inversions which will convert the Standard + gene sequence of chromosome 2 in *D. micromelanica* into the Standard WXYZA'B' in *D. melanura*.

(B). The series of nine inversions which will convert the standard + gene sequence in *D. micromelanica* chromosome 2 into the Standard HLMNOPQRS in *D. nigromelanica*. The first eight inversions in this latter series are interspecific, the last (S) is intraspecific in *D. nigromelanica*.

FIGURE 6.—The Standard gene sequences in chromosome X-left in *D. nigromelanica* and *D. melanura* and “X-left” in *D. micromelanica*. The meaning of the symbols is the same as in Figure 3.

melanica Standard; first, inversion G (34–40 41–55), then the overlapping inversion H (31–32 55–41). Thus Standard X-left of *D. nigromelanica* is GH.

The transition from “X-left” Standard in *D. micromelanica* to that in *D. melanura* may be accomplished by the occurrence of four inversions; starting with *D. micromelanica* Standard, inversions: B (10–22), C (33–38 --- 50–54), D

(54-50 --- 46-39) and E (47-49). Thus Standard X-left of *D. melanura* is BCDE.

In this chromosome, as in chromosome 2, it is clear that "X-left" of *D. micromelanica* forms a structural intermediate between X-left of the other two species.

Homologies in chromosome 3: The homologies involving chromosome 3 in the three species are indicated in Figure 7. The transition from Standard in *D. micromelanica* to that in *D. nigromelanica* may be accomplished by the occurrence of two inversions; starting with *D. micromelanica* Standard, these are: B (6-12) and C (26-40). Thus *D. nigromelanica* Standard is BC.

The transition from Standard in *D. micromelanica* to that in *D. melanura* may be accomplished by the occurrence of two inversions; starting with *D. micromelanica* Standard, these are: C (26-30 31-40) and E (26-30). *D. melanura* Standard is accordingly CE.

It will be noted that in the case of chromosome 3, the Standard of *D. micromelanica* does not form a structural intermediate between the Standards of the other two species, instead, the structural intermediate for all three species is C, a gene sequence which is not presently known in any of them.

Homologies in 4-left: The homologies for this arm are shown in Figure 8. The gene sequence for *D. nigromelanica* Standard differs from that of *D. micromelanica* by the inversion A (10-19). *D. melanura* Standard differs from that of *D. micromelanica* by the inversions B (16-26) and C (27-30). The three Standard sequences are: *D. micromelanica*, +; *D. nigromelanica*, A; *D. melanura*, BC. Again, Standard of *D. micromelanica* forms a structural intermediate between those of the other two species.

Homologies in 4-right: The homologies for this arm are shown in Figure 8. The Standard sequence of *D. nigromelanica* may be derived from that of *D. micromelanica* by the five inversions: E (4-8), F (9-11), G (12-20 23-28), H (28-23) and I (20-12). Accordingly the *D. nigromelanica* Standard is EFGHI.

The Standard of *D. melanura* may be derived from that of *D. micromelanica* by the three inversions: K (13-15 --- 22-30), L (4-12 30-22 21-16) and M (22-30 12-24), and *D. melanura* Standard is KLM. Again, Standard of *D. micromelanica* stands in an intermediate position to those of the other two species.

As indicated above, no complete analysis of homologies was possible in X-right. It is not clear whether this indicates very extensive rearrangement during the course of evolution, or whether gene changes resulting in apparent alteration in banding pattern are responsible.

The very short salivary arm associated with chromosome 5 rarely appears in a satisfactorily stained and fixed condition, and for this reason no serious attempt has been made to study its intraspecific and interspecific variability. Examples of this chromosome in the three species are shown in Figure 9. Apparently the banding pattern is either the same, or nearly the same in all three species. In *D. melanura* two different forms of this chromosome exist, one with, and the other without the extreme terminal puffing. Although both types have been found in a single locality (Webster Groves, Missouri), no intermediates between them have been observed.

FIGURE 7.—The Standard gene sequences in chromosome 3 of *D. nigromelanica*, *D. micromelanica* and *D. melanura*. The meaning of the symbols is the same as in Figure 3.

DISCUSSION

The total analyzed inversion differences within and between the three species are summarized in Figure 10. If it is granted, on the basis of the one-armed X chromosome, that *D. micromelanica* is the more ancestral of the three species,

FIGURE 8.—The Standard gene sequences in chromosome arms 4-left and 4-right in *D. nigromelanica*, *D. micromelanica* and *D. melanura*. The meaning of the symbols in the same as in Figure 3.

then the banding patterns for arms X-left, 2, 4-left and 4-right all indicate a separate origin of *D. melanura* and *D. nigromelanica* from the former species. That this origin is not completely independent is indicated by two facts: first both of the derived species share the inversion C in chromosome 3, an inversion not presently known in *D. micromelanica*; second, both derived species have the same X-autosome fusion, a fusion unknown in *D. micromelanica*.

The total homozygous interspecific inversion differences between *D. micro-*

FIGURE 9.—The Standard gene sequences in chromosome arm X-right in *D. melanura*, X in *D. micromelanica* and chromosome 5 in all three species. The lettered brackets indicate the limits of intraspecific inversions.

melanica and the two derived species is about the same in each case; 17 inversions to *D. melanura* and 18 to *D. nigromelanica*. Although it is unlikely that further study of *D. nigromelanica* would change these figures much, since that species is already well known cytologically, further study of *D. melanura* and *D. micromelanica* might bring about a slight reduction, since some inversions now considered to be interspecific might be found to be heterozygous within one or the other of the two species. It should be remembered however, that the X-right

FIGURE 10.—The intraspecific and interspecific inversions in *D. micromelanica*, *D. nigromelanica* and *D. melanura*. The letters in parentheses in each species box refer to the intraspecific (heterozygous) inversions differing from the Standard of that species. The letters which are on the arrows, or in the species boxes but not in parentheses, represent interspecific inversions. The symbol + represents Standard sequences for the various arms in *D. micromelanica*. Thus, for example, in chromosome 4, *D. micromelanica* has only one intraspecific inversion (D) which differs from the Standard +. The Standard chromosome 4 of *D. nigromelanica* differs from that of *D. micromelanica* by the six interspecific inversions AEFGHI. Thus *D. nigromelanica* Standard is AEFGHI; it has in addition the intraspecific inversion (J). It will be noted that with the exception of chromosome 3, the *D. micromelanica* Standard sequences always form structural intermediates between the Standards of the other two species.

interspecific inversions are not included in the above total, and judging from the difficulty in determining homologies in this arm, the number of such inversions may not be negligible.

It is of course recognized that when we speak of *D. nigromelanica* and *D. melanura* as being derived from *D. micromelanica* we are speaking loosely, and mean that the ancestors of the present day species are involved in all three cases, ancestors which may have been genetically quite different from the named species of today.

The complex that is called *D. micromelanica* today is not a single monotypic species, but a group of incompletely isolated subspecies. The present disjunct distribution and small population sizes within this species would be expected to lead to just such subspeciation. If at the time of origin of the branches leading to *D. nigromelanica* and *D. melanura* a similar situation existed, it could very well explain the rather involved two-step origin of the two derived species. The first step would involve the development of a race or population characterized by the presence of the X-autosome fusion, and inversion C in chromosome 3. The second step would involve the splitting of this race into two additional races, one leading to *D. nigromelanica*, the other to *D. melanura*. Ultimately each of these latter races acquired its own constellation of homozygous (interspecific) and heterozygous (intraspecific) inversions that we observe today.

SUMMARY

The salivary gland chromosomes of *Drosophila micromelanica* and *D. melanura* are illustrated by Standard photomaps. The limits of the known intra-specific inversions are indicated in each species, and the geographical distribution of the inversion polymorphism is described.

Both of the above species belong to the *D. melanica* species group. *D. micromelanica* is found in the United States south of 40° N. latitude and in northern Mexico, is rare in collections, has a markedly disjunct distribution, exhibits considerable variability in its metaphase chromosomes, and shows structural polymorphism in all major salivary chromosomes. A total of 14 inversions differing from Standard is known in this species, of which nine are on chromosome 2. The species is divisible into an Eastern (Virginia, Florida and possibly Texas) and Western (Texas, Arizona) subspecies on the basis of the production of sterile male hybrids in the cross of Eastern females $\times$ Western males. All strains of this species have a single-element rod-shaped X chromosome, unlike the V-shaped two-element X found in all other members of the *D. melanica* group. It is accordingly indicated that *D. micromelanica* is the most nearly ancestral member of the group.

D. melanura is found primarily in the eastern United States north of 34° N. latitude. It has an apparently disjunct distribution, and is generally not abundant in collections. A total of six inversions differing from Standard is known, four of these are in chromosome 2. Thus both *D. micromelanica* and *D. melanura* agree with the other four members of the species group in showing a concentration of inversion polymorphism in the second autosome.

Comparisons of banding patterns in the three species *D. micromelanica*, *D. melanura* and *D. nigromelanica* were carried out with the aid of Standard photomaps and a large collection of supplementary photographs. Homology of banding pattern is presented in detail for all major chromosome arms except that which is the X in *D. micromelanica* and X-right in the other two species. In chromosome arms X-left, 2, 4-left and 4-right it could be demonstrated that the chromosomes of *D. micromelanica* are structural intermediates between those of the other two species. However chromosome 3 of *D. melanura* and of *D. nigromelanica* carry an inversion in common which is not found in *D. micromelanica*. It is suggested that *D. nigromelanica* and *D. melanura* arose from ancestral forms of *D. micromelanica*, and that their origin occurred in at least two steps: the first step involved the development of a race or population with the inversion C in chromosome 3 and the X-autosomal fusion; the second step involved the separation of the above race into two additional races, one leading to each of the two derived species.

LITERATURE CITED

- DOBZHANSKY, TH., and C. EPLING, 1944 Contributions to the genetics, taxonomy and ecology of *Drosophila pseudoobscura* and its relatives. Carnegie Inst. Wash. Publ. 554.

- LEVINE, L., and L. VAN VALEN, 1962 A gene in *Drosophila* that produces a new chromosomal pattern. *Science* **137**: 993-994.
- MILLER, D. D., 1944 *Drosophila melanura*, a new species of the melanica group. *Jour. N. Y. Ent. Soc.* **52**: 85-97.
- PATTERSON, J. T., and W. S. STONE, 1952 *Evolution in the Genus Drosophila*. Macmillan, New York.
- STALKER, H. D., 1960 Chromosomal polymorphism in *Drosophila paramelanica* Patterson. *Genetics* **45**: 95-114. — 1961 The genetic systems modifying meiotic drive in *Drosophila paramelanica*. *Genetics* **46**: 177-202. — 1964a Chromosomal polymorphism in *Drosophila euronotus*. *Genetics* **49**: 669-687. — 1964b The salivary gland chromosomes of *Drosophila nigromelanica*. *Genetics* **49**: 883-893.
- STONE, W. S., 1962 The dominance of natural selection and the reality of superspecies (species groups) in the evolution of *Drosophila*. *Univ. Texas Publ.* **6205**: 507-537.
- STONE, W. S., W. C. GUEST, and F. D. WILSON, 1960 The evolutionary implications of the cytological polymorphism and phylogeny of the virilis group of *Drosophila*. *Proc. Natl. Acad. Sci. U.S.* **46**: 350-361.
- STURTEVANT, A. H., and E. NOVITSKI, 1941 Sterility in crosses of geographical races of *Drosophila micromelanica*. *Proc. Natl. Acad. Sci. U.S.* **27**: 392-394.
- WARD, C. L., 1949 Karyotype variation in *Drosophila*. *Univ. Texas. Publ.* **4920**: 70-79. — 1952 Chromosome variation in *Drosophila melanica*. *Univ. Texas Publ.* **5204**: 137-157.
- WASSERMAN, M., 1960 Cytological and phylogenetic relationships in the repleta group of the genus *Drosophila*. *Proc. Natl. Acad. Sci. U.S.* **46**: 842-859. — 1962 Cytological studies of the repleta group of the genus *Drosophila*: Parts III, IV, V and VI. *Univ. Texas Publ.* **6205**: 63-134. — 1963 Cytology and phylogeny of *Drosophila*. *Am. Naturalist* **97**: 333-353.
- WHARTON, L. T., 1943 An analysis of the metaphase and salivary chromosome morphology within the genus *Drosophila*. *Univ. Texas Publ.* **4313**: 282-319.