dihydroxyacetone phosphate ratio in the muscles was increased by butyrate and decreased by diabetes and starvation.

6. It is concluded: that the oxidation of pyruvate in rat heart and rat diaphragm muscles may be inhibited in diabetes because of the greater availability of fatty acids and ketone bodies for respiration in this condition; that glycolysis in these muscles is not reversible to any significant extent; that the NADH_2/NAD ratio in cell sap is increased by glucose, anoxia, salicylate, fatty acids and ketone bodies, and diminished in alloxan-diabetes and starvation.

7. The possibility that the inhibitory effects of diabetes, fatty acids and ketone bodies on pyruvate oxidation are due to inhibition of pyruvate dehydrogenase by a rise in the acetyl-CoA/CoA ratio is discussed.

We thank the British Diabetic Association and the British Insulin Manufacturers for grants that contributed to the cost of these investigations, Evans Medical Ltd. for a generous gift of heparin, and Mrs W. Thorne for skilled technical assistance.

REFERENCES

- Bing, R. J. (1954-55). *Harvey Lect.* **50**, 27.
 Bloom, B., Stetten, R. & Stetten, D. (1953). *J. biol. Chem.* **204**, 681.
 Campbell, P. N. & Stone, N. E. (1956). *Biochem. J.* **62**, 9P.
 Fajans, S. S. (1961). *Metabolism*, **10**, 951.
 Fry, I. K. & Butterfield, W. J. H. (1962). *Lancet*, ii, 66.
 Garland, P. B., Newsholme, E. A. & Randle, P. J. (1962). *Nature, Lond.*, **195**, 381.
 Garland, P. B. & Randle, P. J. (1962). *Biochem. J.* **84**, 78P.
 Garland, P. B. & Randle, P. J. (1964a). *Biochem. J.* **91**, 6C.
 Garland, P. B. & Randle, P. J. (1964b). *Biochem. J.* **93**, 678.
 Glock, G. E. & McLean, P. (1955). *Biochem. J.* **61**, 397.
 Goodman, D. S. (1958). *J. Amer. chem. Soc.* **80**, 3887.
 Green, D. E., Herbert, D. & Subrahmanyam, V. (1941). *J. biol. Chem.* **138**, 327.
 Hales, C. N. & Randle, P. J. (1963). *Biochem. J.* **88**, 137.
 Henneman, D. H. & Bunker, J. P. (1957). *Amer. J. Med.* **23**, 34.
 Hiatt, H. H., Goldstein, M., Lareau, J. & Horecker, B. L. (1958). *J. biol. Chem.* **238**, 303.
 Hohorst, H. J., Kreutz, F. H. & Bücher, Th. (1959). *Biochem. Z.* **332**, 18.
 Holzer, E., Söling, H. D., Goedde, H. W. & Holzer, H. (1963). In *Methods of Enzymatic Analysis*, p. 602. Ed. by Bergmeyer, H. U. New York: Academic Press Inc.
 Krebs, H. A. & Henseleit, K. (1932). *Hoppe-Seyl. Z.* **210**, 33.
 Morgan, H. E., Henderson, M. J., Regen, D. M. & Park, C. R. (1961). *J. biol. Chem.* **236**, 253.
 Newsholme, E. A. & Randle, P. J. (1962). *Biochem. J.* **83**, 387.
 Newsholme, E. A. & Randle, P. J. (1964). *Biochem. J.* **93**, 641.
 Pearson, O. H., Hsieh, C. K., Du Toit, C. H. & Hastings, A. B. (1949). *Amer. J. Physiol.* **158**, 261.
 Randle, P. J. (1954). *Brit. med. J.* **i**, 1237.
 Randle, P. J., Newsholme, E. A. & Garland, P. B. (1964). *Biochem. J.* **93**, 652.
 Shepp, M. C., Nickerson, R. J., Dagenais, M. M., Steinke, J., Martin, D. B. & Renold, A. E. (1960). *J. clin. Invest.* **39**, 1499.
 Short, D. J. & Parkes, A. S. (1949). *J. Hyg., Camb.*, **47**, 209.
 Synder, J. (1961). *J. Lipid Res.* **2**, 195.
 Trout, D. L., Estes, E. H., jun. & Friedberg, S. J. (1960). *J. Lipid Res.* **1**, 199.
 Vallance-Owen, J., Dennes, E. & Campbell, P. N. (1958). *Lancet*, ii, 336.
 Villee, C. A. & Hastings, A. B. (1949). *J. biol. Chem.* **181**, 131.
 Weil, R., Altszuler, N. & Kessler, J. (1961). *Amer. J. Physiol.* **201**, 251.
 Wollenberger, A., Ristau, O. & Schoffa, G. (1960). *Pflüg. Arch. ges. Physiol.* **270**, 399.

Biochem. J. (1964), **93**, 678

Regulation of Glucose Uptake by Muscle

10. EFFECTS OF ALLOXAN-DIABETES, STARVATION, HYPOPHYSECTOMY AND ADRENALECTOMY, AND OF FATTY ACIDS, KETONE BODIES AND PYRUVATE, ON THE GLYCEROL OUTPUT AND CONCENTRATIONS OF FREE FATTY ACIDS, LONG-CHAIN FATTY ACYL-COENZYME A, GLYCEROL PHOSPHATE AND CITRATE-CYCLE INTERMEDIATES IN RAT HEART AND DIAPHRAGM MUSCLES

By P. B. GARLAND* AND P. J. RANDLE*
Department of Biochemistry, University of Cambridge

(Received 3 February 1964)

The experimental observations summarized in the preceding papers have shown that certain abnormalities of carbohydrate metabolism in heart

* Present address: Department of Biochemistry, University of Bristol.

and diaphragm muscles of alloxan-diabetic or starved rats could be explained if the provision of fatty acids and ketone bodies for respiration is increased (Newsholme & Randle, 1964; Randle, Newsholme & Garland, 1964; Garland, Newsholme

& Randle, 1964). These abnormalities of carbohydrate metabolism are seen not only *in vivo* (where the plasma concentrations of fatty acids and ketone bodies are elevated) but also *in vitro*, where the muscle is removed from the influence of circulating fatty acids and ketone bodies. It was pointed out that the provision of ketone bodies for respiration in muscles of diabetic or starved rats *in vitro* is unlikely to be increased significantly, but that provision of fatty acids for respiration could be increased if the net breakdown of muscle glycerides is accelerated by diabetes or starvation (Randle *et al.* 1964). Direct evidence on this point has now been obtained by measurements of the intracellular concentrations of fatty acyl-CoA, free fatty acids and glycerol phosphate and of the output of glycerol in hearts and diaphragms from fed, starved and alloxan-diabetic rats.

Two abnormalities of carbohydrate metabolism in muscles of diabetic or starved rats that may be due to increased provision of fatty acids and ketone bodies for respiration are diminished rates of the phosphofructokinase and pyruvate-dehydrogenase reactions. It has been shown that citrate is an inhibitor of phosphofructokinase and that acetyl-CoA is a competitive inhibitor with respect to CoA of pyruvate dehydrogenase. Because of the possibility that inhibitory effects of fatty acids, ketone bodies, diabetes and starvation on these enzymes might be mediated by an increase in the intracellular concentration of citrate and of the acetyl-CoA/CoA ratio, we have investigated the effects of these agents on the concentrations of these intermediates in the perfused rat heart. Preliminary accounts of some of these observations have been published (Garland & Randle, 1962*a*, *b*, 1963, 1964; Garland, Randle & Newsholme, 1963). Hitherto in comparing the effects of fatty acids and ketone bodies on carbohydrate metabolism in muscle with those of diabetes and starvation we have referred only to the possibility that increased provision of fatty acids for respiration might be responsible for these changes in carbohydrate metabolism in diabetes and starvation.

With the demonstration that accumulations of acetyl-CoA and of citrate may contribute to two of these changes it has become important to know specifically whether it is an increase in the rate of oxidation of fatty acids that leads to an accumulation of acetyl-CoA and citrate in diabetes and of citrate in starvation. The experimental evidence relevant to this point is also considered below.

MATERIALS

Rats. Hearts were obtained from male albino Wistar rats (200–300 g.), hemidiaphragms from male or female albino Wistar rats (100–150 g.) and epididymal fat pads

from male albino Wistar rats (200–240 g.). Details of diet, induction of alloxan-diabetes, hypophysectomy, adrenalectomy and treatment of diabetic rats with insulin, growth hormone or corticosteroids were as described by Newsholme & Randle (1964).

Chemicals and enzymes. 2',7'-Dichlorofluorescein was obtained from Eastman Kodak Co. Silica gel G for thin-layer chromatography (Merck) and Desaga apparatus were obtained from Camlab (Glass) Ltd., Cambridge. Citrate lyase (EC 4.1.3.6) was prepared for us by Dr D. Kerridge from a culture of *Aerobacter aerogenes* by the method of Dagley (1963). Malate dehydrogenase, isocitrate dehydrogenase and L-glutamate dehydrogenase were obtained from C. F. Boehringer und Soehne G.m.b.H. The sources of other chemicals and enzymes were as stated by Newsholme & Randle (1964), Garland *et al.* (1964) and Tubbs & Garland (1964).

Media. The perfusion and incubation media were as described by Newsholme & Randle (1964) and Garland *et al.* (1964). The concentrations of various additions to the media are given in the text or Tables.

PROCEDURE

Muscles. For the assay of metabolic intermediates in muscle, hearts were perfused by continuous flow, except in experiments in which the medium contained albumin, when recirculation was used. For measurement of the glycerol output hearts were perfused by recirculation. The details of perfusion of hearts and incubation of hemidiaphragms were as given by Newsholme & Randle (1964) and Randle *et al.* (1964). Fat pads were incubated under the same conditions as were hemidiaphragms. For the assay of substances in muscle, hearts were frozen with the tissue clamp of Wollenberger, Ristau & Schoffa (1960) cooled in liquid N₂; hemidiaphragms were frozen in acetone–solid CO₂. The muscles were powdered in a percussion mortar cooled in solid CO₂ and extracted with 5% (w/v) perchloric acid (2 ml./g. of powder). The extract separated by centrifugation was adjusted to pH 7 and perchlorate was precipitated with 6*N*-KOH and 5*M*-K₃PO₄ by using wide-range indicator paper (British Drug Houses Ltd.). The final extract is referred to as perchloric acid extract. The details of extraction of muscle powder for assay of long-chain fatty acyl-CoA were as described by Tubbs & Garland (1964).

METHODS

Glycerol. Glycerol was assayed in perfusion and incubation media by the method of Garland & Randle (1962*b*).

Free fatty acids. The concentrations of free fatty acids in rat heart and diaphragm muscles were estimated spectrophotometrically by the method of Duncombe (1962) after extraction followed by thin-layer chromatography, as described by Garland & Randle (1963).

Long-chain fatty acyl-CoA. The concentration of long-chain fatty acyl-CoA in rat heart was estimated as described by Tubbs & Garland (1964).

Glycerol phosphate. Glycerol phosphate was assayed in perchloric acid extracts of rat heart and diaphragm by the method of Hohorst, Kreutz & Bücher (1959). The production of NADH₂ was followed either by the change in extinction at 340 mμ in a Beckman spectrophotometer or

by the change in fluorescence in an Eppendorf fluorimeter fitted with a primary filter transmitting at 366 m μ and a secondary filter at 420–3000 m μ with a mercury-vapour lamp.

Citrate. Citrate was assayed in perchloric acid extracts with citrate lyase by the method of Dagley (1963); in this method citrate is cleaved to acetate and oxaloacetate and the latter converted into pyruvate by oxaloacetate decarboxylase present in the preparation of citrate lyase. The pyruvate formed was then assayed by NADH₂ oxidation with lactate dehydrogenase. Some assays were made also by adding NADH₂ and malate dehydrogenase, when oxaloacetate formed is converted quantitatively into malate and NADH₂ into NAD. In the latter instance it was necessary to run a blank (omitting the perchloric acid extract) to correct for the activity of NADH₂ oxidase in the citrate lyase. This blank was not necessary in the assay run through pyruvate, because the pyruvate is formed before the addition of NADH₂ and the lactate-dehydrogenase reaction proceeded much more rapidly than the oxidation of NADH₂ by NADH₂ oxidase.

Isocitrate. Isocitrate was assayed spectrophotometrically with NADP and isocitrate dehydrogenase [assay medium: tris-HCl buffer, pH 7.4 (70 mM), EDTA (7 mM) and MnSO₄ (2 mM)].

α -Oxoglutarate. α -Oxoglutarate was assayed spectrophotometrically or fluorimetrically by the method of Bergmeyer & Bernt (1963).

Oxaloacetate. Oxaloacetate was assayed on 1 ml. of muscle extract in a Beckman DK2 recording spectrophotometer with NADH₂ and malate dehydrogenase [assay medium: tris-HCl buffer, pH 7.0 (50 mM), EDTA (1 mM) and NADH₂ (0.2 mM)]. Oxaloacetate is unstable in solution, and the loss in perchloric acid extracts of muscle amounted to approx. 15%/hr. at 0°. Assays of oxaloacetate were therefore begun and completed as soon as possible after perfusion (always within 2 hr.). No attempt has been made to correct the values obtained for possible losses within this time.

Glutamate and malate. Glutamate and malate were assayed spectrophotometrically with glutamate dehydrogenase and malate dehydrogenase and a hydrazine trap as described by Bernt & Bergmeyer (1963) and Hohorst (1963).

CoA. CoA (thiol) was assayed by the change in extinction in a 4 cm. light-path at 366 m μ (in an Eppendorf photometer) on adding α -oxoglutarate dehydrogenase (0.5 unit) to a mixture containing (in 2.2 ml.): potassium phosphate buffer, pH 6.8 (100 μ moles), MgCl₂ (2 μ moles), EDTA (1 μ mole), glutathione (0.5 μ mole), 3-acetylpyridine-adenine dinucleotide (Sigma Chemical Co.) (0.4 μ mole), α -oxoglutarate (2 μ moles) and muscle extract (0.5–1 ml. containing 0–0.02 μ mole of CoA). With this amount of CoA and with 3-acetylpyridine-adenine dinucleotide, CoA is assayed quantitatively.

Acetyl-CoA. Acetyl-CoA was assayed as described by Garland & Randle (1964).

Oxygen uptake. The oxygen uptake of the heart was measured in the apparatus described by Randle *et al.* (1964). The medium issuing from the heart in the 5–10 min. period of perfusion was collected under heptane. The oxygen concentration in the medium before and after passage through the heart was then measured with an oxygen electrode calibrated with perfusion medium saturated at 36° with O₂ + CO₂ (95:5).

Calculations. Glycerol outputs were calculated and expressed as μ moles of glycerol/g. wet wt. of tissue/hr., and oxygen uptakes as ml. of oxygen/g. wet wt. of tissue/hr. The concentrations of substances in muscle were calculated as μ moles/100 g. wet wt. of tissue as described by Newsholme & Randle (1961, 1964). They may be converted into intracellular concentrations (μ M) by multiplying by 23.2 for heart and by 17 for diaphragm.

RESULTS AND DISCUSSION

Glycerol output and intracellular concentrations of long-chain fatty acyl-CoA, free fatty acids and L-glycerol 3-phosphate. The substrate for β -oxidation of fatty acids in muscle is presumably the CoA derivative. The results given in Table 1 show that the concentration of long-chain fatty acyl-CoA is increased in hearts from alloxan-diabetic rats (by about 70 %) or from starved rats (by about 50 %) perfused with medium containing glucose and insulin. Treatment of alloxan-diabetic rats with insulin decreased the concentration to normal levels. In hearts from normal rats perfusion with 0.75 mM-palmitate (carried by 2 % bovine plasma albumin) increased the concentration to levels similar to those seen in starvation but lower than those seen in diabetes. The concentration of long-chain fatty acyl-CoA was not increased by β -hydroxybutyrate or pyruvate (which decreased the concentration), showing that the effects of these substrates on glucose metabolism are not due to an increase in the concentration of long-chain fatty acyl-CoA. The concentration of free fatty acids in hearts perfused with medium containing glucose and insulin was likewise increased in diabetes (by about 130 %) and by starvation (though the effects of the latter were not significant). In rat hemidiaphragms incubated in medium containing glucose the concentration of free fatty acids was also increased by diabetes and starvation. The addition of insulin to the incubation medium (containing glucose) lowered the concentration of free fatty acids in hemidiaphragms, but the increased concentrations induced by diabetes and starvation were maintained. On the other hand, no increase in the concentration of free fatty acids in hemidiaphragms was seen when alloxan-diabetes was induced in hypophysectomized rats. These measurements show that the provision of fatty acids for respiration is increased in heart and diaphragm muscles by diabetes and starvation, and that this increase may be prevented in diabetes by prior hypophysectomy or by treatment of the rat with insulin. These findings are consistent with the view advanced in the preceding papers that it is the increased provision of fatty acids for respiration that leads to a number of abnormalities of carbohydrate metabolism in these muscles in diabetes and starvation.

Table 1. *Effects of alloxan-diabetes and starvation, and of fatty acids and ketone bodies, on the glycerol output and concentrations of free fatty acids, long-chain fatty acyl-CoA and glycerol phosphate in rat heart and rat diaphragm*

For schedules of treatment of rats with growth hormone, cortisol or insulin see Newsholme & Randle (1964). Other details are given in footnotes and in the text. D, Diaphragm; F, fat pad; H, heart. The results are given as means $\pm$ S.E.M., the minimum number of observations being five.

Source of tissue	Addition(s) to perfusion or incubation medium			Glycerol output (μ moles/g. wet wt. of tissue/hr.)			Concn. (μ moles/100 g. wet wt. of tissue)		
	Glucose	Insulin	Others	Tissue...	Heart	Diaphragm	Epididymal fat pad	Free fatty acids	
								Heart	Diaphragm
Normal fed rat	{ 1.5 (D, F) or 1.0 (H) mg./ml. }	None	None	1.77 $\pm$ 0.15	0.41 $\pm$ 0.05	0.66 $\pm$ 0.06	0.66 $\pm$ 0.06	35 $\pm$ 5.9 $\dagger$	5.7 $\pm$ 0.40 $\dagger$
Starved (40 hr.) rat		None	None	2.68 $\pm$ 0.17*	0.45 $\pm$ 0.04	1.30 $\pm$ 0.19*	1.30 $\pm$ 0.19*	14 $\pm$ 2.4 $\dagger$	3.9 $\pm$ 0.66 $\dagger$
Alloxan-diabetic rat	{ 1.5 (D, F) or 1.0 (H) mg./ml. }	None	None	4.10 $\pm$ 0.53*	0.66 $\pm$ 0.04*	1.15 $\pm$ 0.10*	1.15 $\pm$ 0.10*	8.5 $\pm$ 1.9 $\dagger$	4.3 $\pm$ 0.26 $\dagger$
Normal fed rat		0.1	None	2.66 $\pm$ 0.21	0.66 $\pm$ 0.05	0.87 $\pm$ 0.05	0.87 $\pm$ 0.05	4.5 $\pm$ 0.5	15.6 $\pm$ 1.5
Starved (48 hr.) rat	{ 1.5 (D, F) or 1.0 (H) mg./ml. }	0.1	None	3.81 $\pm$ 0.27*	0.72 $\pm$ 0.09	1.27 $\pm$ 0.15 $\dagger$	1.27 $\pm$ 0.15 $\dagger$	5.3 $\pm$ 0.9	11.4 $\pm$ 0.48
Alloxan-diabetic rat		0.1	None	4.62 $\pm$ 0.22*	1.15 $\pm$ 0.13*	1.26 $\pm$ 0.03*	1.26 $\pm$ 0.03*	8.3 $\pm$ 0.66*	8.0 $\pm$ 0.42*
Hypophysectomized rat	{ 1.5 (D, F) or 1.0 (H) mg./ml. }	0.1	None	1.46 $\pm$ 0.07	0.45 $\pm$ 0.04	0.45 $\pm$ 0.04	0.45 $\pm$ 0.04	10.3 $\pm$ 1.8*	9.6 $\pm$ 1.3*
Hypophysectomized alloxan-diabetic rat		0.1	None	1.17 $\pm$ 0.08 $\dagger$	0.53 $\pm$ 0.03	0.58 $\pm$ 0.03	0.58 $\pm$ 0.03	52 $\pm$ 4.3 $\dagger$	—
Hypophysectomized alloxan-diabetic rat treated with growth hormone + cortisol	{ 1.5 (D, F) or 1.0 (H) mg./ml. }	0.1	None	—	0.86 $\pm$ 0.12*	1.67 $\pm$ 0.37 $\dagger$	1.67 $\pm$ 0.37 $\dagger$	46 $\pm$ 6.6 $\dagger$	—
Adrenalectomized rat		0.1	None	0.96 $\pm$ 0.07	0.40 $\pm$ 0.05	0.62 $\pm$ 0.06	0.62 $\pm$ 0.06	—	—
Alloxan-diabetic rat	{ 1.5 (D, F) or 1.0 (H) mg./ml. }	0.1	None	1.21 $\pm$ 0.14	0.36 $\pm$ 0.03	0.70 $\pm$ 0.09	0.70 $\pm$ 0.09	—	—
Alloxan-diabetic rat treated with insulin		0.1	None	4.62	—	—	—	10.7 $\pm$ 1.3	9.6 $\pm$ 1.2
Normal fed rat	{ 1.5 (D) or 1.0 (H) mg./ml. }	0.1	Albumin [3% (H) or 0.85% (D)]	2.80	—	—	—	5.7 $\pm$ 0.6*	17.8 $\pm$ 2.3
Normal fed rat		0.1	Albumin [3% (H) or 0.85% (D)] + palmitate [0.7 mM (H) or 0.6 mM (D)]	—	—	—	—	4.8 $\pm$ 0.17	—
Normal fed rat	{ 1.5 mg./ml. }	None	None	—	0.47 $\pm$ 0.03	0.81 $\pm$ 0.06	0.81 $\pm$ 0.06	—	—
Normal fed rat		0.1	None	—	0.47 $\pm$ 0.03	0.58 $\pm$ 0.06 $\dagger$	0.58 $\pm$ 0.06 $\dagger$	—	—
Normal fed rat	{ 1.5 (D) or 1.0 (H) mg./ml. }	0.1	DL- β -Hydroxybutyrate (70 mg./100 ml.)	—	—	—	—	3.6 $\pm$ 0.26	28.6 $\pm$ 1.9
Normal fed rat		0.1	Butyrate (40 mg./100 ml.)	—	—	—	—	3.6 $\pm$ 0.29	28.6 $\pm$ 1.7
Normal fed rat	{ 1.5 (D) or 1.0 (H) mg./ml. }	0.1	Pyruvate (40 mg./100 ml.)	—	—	—	—	—	—
Normal fed rat		0.1	—	—	—	—	—	2.5 $\pm$ 0.27 $\dagger$	21.7 $\pm$ 0.17*

* $P < 0.01$; † $P < 0.05$; for other differences from control, $P > 0.05$.

§ These tissues were frozen with acetone-solid CO₂; other tissues were frozen with the tissue clamp of Wollenberger *et al.* (1960) (see the Methods section).

|| In this experiment the whole diaphragm was used; in other experiments hemidiaphragms were used.

¶ Insulin omitted from incubation medium.

The increased concentrations of long-chain fatty acyl-CoA and of free fatty acids that have been detected in these muscles in diabetes and starvation could arise either as a result of accelerated hydrolysis of muscle glycerides (lipolysis) or from a diminished rate of re-esterification of long-chain fatty acyl-CoA (e.g. with glycerol phosphate). Since glycerol kinase has not been detected in rat heart or skeletal muscles (Wieland & Suyter, 1957), we have used the output of glycerol by these tissues as an index of the rate of lipolysis, as suggested by Vaughan (1961). The rate of re-esterification of long-chain fatty acyl-CoA has not been determined in these studies, but some measurements of the concentration of glycerol phosphate (which may be one of the controlling factors) have been made.

In rats hearts perfused with medium containing glucose in the presence and in the absence of insulin the output of glycerol was increased by starvation and by alloxan-diabetes. The effects of diabetes were greater than those of starvation. On the other hand, in hypophysectomized rats or adrenalectomized rats the induction of alloxan-diabetes did not increase the output of glycerol. In alloxan-diabetic rats treated with insulin the output of glycerol was markedly decreased (compared with the untreated diabetic rats). In isolated rat hemidiaphragms incubated in medium containing glucose in the presence and in the absence of insulin, glycerol output was increased by diabetes but not by starvation. In hypophysectomized rats the induction of alloxan-diabetes did not accelerate glycerol output unless the animals were injected with growth hormone and cortisol. Moreover, in adrenalectomized rats the induction of alloxan-diabetes also failed to increase the output of glycerol by hemidiaphragms. These results are given in Table 1, where the outputs of glycerol by rat epididymal adipose tissue are also included for comparison. The changes in glycerol output in the adipose tissue in diabetes and starvation, and with hypophysectomy, adrenalectomy and treatment with growth hormone and cortisol, paralleled the changes seen in muscle. The output of glycerol by the adipose tissue on a wet-weight basis was similar to that of diaphragm but lower than that of heart. In view of this it seems very unlikely that the glycerol released by heart and diaphragm was derived solely from adipose-tissue cells in these muscle preparations (i.e. that the output of glycerol by these tissues does provide an estimate of the rate of hydrolysis of muscle glycerides). These studies show that lipolysis in muscle, like that of adipose tissue, is increased by diabetes and by starvation, and that this increase is dependent on pituitary growth hormone and adrenal corticosteroids. Moreover, adrenaline, which accelerates

lipolysis in adipose tissue, also accelerates lipolysis in muscle (Garland & Randle, 1962*b*, 1963). The only difference thus far detected between the two tissues concerns the action of insulin *in vitro*. In adipose tissue insulin *in vitro* diminished the output of glycerol, in confirmation of findings by Jungas & Ball (1962); insulin *in vitro* did not, however, diminish the output of glycerol by hemidiaphragms.

The concentration of glycerol phosphate in rat heart and diaphragm muscles perfused or incubated with medium containing glucose in the presence and in the absence of insulin was diminished by alloxan-diabetes and by starvation. Treatment of diabetic rats with insulin raised the concentration of glycerol phosphate in the perfused heart towards normal. In hypophysectomized rats, on the other hand, the induction of alloxan-diabetes did not lower the concentration of glycerol phosphate. In the heart *in vivo* the concentration of glycerol phosphate was lowered by diabetes and starvation (from a control value of $12.0 \pm 1.2 \mu\text{moles}/100 \text{ g. wet wt.}$ to 1.5 ± 0.41 and $4.0 \pm 1.6 \mu\text{moles}/100 \text{ g. wet wt.}$ respectively). Factors that may lead to lower concentrations of glycerol phosphate in these tissues in diabetes and starvation are inhibition of phosphofructokinase, which may diminish the formation of dihydroxyacetone phosphate (the precursor of glycerol phosphate), and the diminished cytoplasmic NADH_2/NAD ratio, which may decrease the proportion of triose phosphates converted into glycerol phosphate. In rat hearts perfused with fatty acids or ketone bodies the concentration of glycerol phosphate is maintained in spite of the fact that these substrates lead to inhibition of phosphofructokinase. This may be accounted for by the increase in the cytoplasmic NADH_2/NAD ratio that they induce, and this change may account for the increase in concentration of glycerol phosphate induced by butyrate in rat hemidiaphragms.

These findings suggest that the increased concentrations of long-chain fatty acyl-CoA and free fatty acids in heart and diaphragm in diabetes are caused in part by an increased rate of lipolysis and in part by a diminished rate of re-esterification (assuming that the diminished concentration of glycerol phosphate will lead to lower rates of re-esterification). The raised concentrations in starvation may be similarly explained, though it is possible that alterations in the rate of re-esterification may be of particular importance in starvation (the glycerol output by diaphragm was not increased by starvation). Two other mechanisms that may alter the concentrations of fatty acyl-CoA and of free fatty acids need to be considered for purposes of exclusion. The concentration of fatty acyl-CoA could increase if in diabetes and starvation the rate of oxidation of fatty acid were decreased. This is

extremely unlikely in view of evidence, summarized in the next sections, showing that the rate of fatty acid oxidation is increased in these conditions. It was also conceivable that loss of fatty acid to perfusion or incubation media could influence the intracellular concentration of free fatty acid; such a loss has not been detected in hearts perfused with 2 % bovine plasma albumin (to trap any fatty acids that might be released).

Effects of fatty acids and ketone bodies, and of alloxan-diabetes and starvation, on the concentrations of acetyl-CoA and of CoA in perfused rat heart. The concentrations of acetyl-CoA and of CoA (thiol) in the perfused heart were measured for two reasons. In the preceding paper (Garland *et al.* 1964) evidence was presented that the oxidation of pyruvate (derived from glucose) by pyruvate dehydrogenase in rat heart is inhibited by alloxan-diabetes and by perfusion with medium containing fatty acids and ketone bodies but not by starvation. In experiments with partially purified pig-heart pyruvate dehydrogenase it has been shown that the oxidation of pyruvate to acetyl-CoA by the enzyme is inhibited by an increase in the concentration of acetyl-CoA relative to that of CoA (Garland & Randle, 1964). Further, evidence has been given that the phosphorylation of fructose 6-phosphate by phosphofructokinase is inhibited in rat heart by diabetes and starvation and by perfusion with fatty acids or ketone bodies (News-holme & Randle, 1964). In extracts of rat heart the activity of phosphofructokinase is inhibited by citrate (Garland *et al.* 1963; Randle *et al.* 1964), and these agents increase the concentration of citrate in rat heart (see the next section). A rise in the acetyl-CoA/CoA ratio might be expected to increase the concentration of citrate through the citrate-synthase reaction (see the next section).

The results of these measurements are given in Table 2. In hearts from normal fed rats perfused for 10 min. with medium containing glucose (1 mg./ml.) and insulin (0.1 unit/ml.) in the presence and in the absence of 2 % bovine plasma albumin the concentration of CoA was 6.7 μ moles/100 g. wet wt. and that of acetyl-CoA less than 0.3 μ mole/100 g. wet wt. (the amount was too small for reliable assay). The concentration of acetyl-CoA was increased after perfusion with 0.75 mM-palmitate carried by 2 % albumin (more than 1.8-fold), by 5.5 mM-DL- β -hydroxybutyrate (more than 6.7-fold), by 4 mM-butyrate (more than 6.9-fold), by 5 mM-acetate (more than 7-fold) and by alloxan-diabetes (more than 2.9-fold), but not by starvation for 40 hr. The concentration of CoA was diminished by palmitate to 46 % of that of the control, by β -hydroxybutyrate to 45 % and by butyrate to 42 %, but not by diabetes or starvation. The acetyl-CoA/CoA concentration ratio in

hearts from normal fed rats was less than 0.05. This ratio was increased by palmitate (4-fold), by β -hydroxybutyrate (15-fold), by butyrate (16-fold) and by diabetes (2.9-fold), but not by starvation. These observations are consistent with the view that a rise in the acetyl-CoA/CoA ratio may contribute to the inhibition of pyruvate dehydrogenase induced by these agents. The observation that starvation, which leads to little if any inhibition of pyruvate oxidation, did not increase the acetyl-CoA/CoA ratio is also consistent with this view.

The concentrations of acetyl-CoA measured in the present study in hearts frozen with the tissue clamp of Wollenberger *et al.* (1960) are lower than those obtained by freezing with acetone-solid carbon dioxide (Garland & Randle, 1964). Since the former method leads to more rapid freezing of muscle the values obtained with it are presumably closer to concentrations in the cell. The concentrations of acetyl-CoA and of CoA that have been measured are the total muscle concentrations, and their relation to the concentrations that obtain in the vicinity of pyruvate dehydrogenase in the cell has yet to be ascertained. It is presumably the latter that is of importance in controlling the activity of the enzyme.

Effects of fatty acids and ketone bodies, and of alloxan-diabetes and starvation, on the concentrations of citrate and oxaloacetate in perfused rat heart. The results of these measurements are given in Table 2. The concentration of citrate in hearts from normal fed rats perfused with medium containing glucose (1 mg./ml.) and insulin (0.1 unit/ml.) was 22 μ moles/100 g. wet wt. The concentration of citrate was increased by diabetes (2.4-fold), by starvation (1.5-fold) and by perfusion with 5.5 mM- β -hydroxybutyrate (3-fold), 4 mM-butyrate (2.3-fold), 0.75 mM-palmitate (carried by 2 % albumin) (3-fold) or acetate (2.5-fold). Treatment of the diabetic rat with insulin lowered the intracellular concentration of citrate to normal. The concentration of citrate measured in the heart *in vivo* was likewise increased in alloxan-diabetes (from 20 to 42 μ moles/100 g. wet wt.). These findings are consistent with the view that the inhibitory effects of these agents on the phosphofructokinase reaction in rat heart may be mediated by an increase in the intracellular concentration of citrate. They are also consistent with the observations that the inhibitory effects of diabetes on the phosphofructokinase reaction are corrected by treatment with insulin, and that the inhibitory effects of starvation are less than those of the other agents (starvation induces a smaller rise in citrate concentration). The citrate concentration measured in the present studies is the total heart concentration, and it is not known how this citrate is distributed between mitochondrial and cytoplasmic fractions. If, in company with other

Table 2. *Effects of alloxan-diabetes and starvation and of DL-β-hydroxybutyrate, palmitate, butyrate and acetate, on the concentrations of acetyl-CoA, CoA, citrate, oxaloacetate and isocitrate in perfused rat heart*

Source of heart	Addition(s) to perfusion medium	Concn. (μmoles/100 g. wet wt. of heart)					Acetyl-CoA/CoA ratio in tissue	Citrate/isocitrate ratio in tissue
		Acetyl-CoA	CoA (thiol)	Citrate	Oxaloacetate	Isocitrate		
<i>Experiments in vitro</i>								
Normal fed rat	None	< 0.3	6.66 ± 0.15	22 ± 2	1.1 ± 0.24	—	< 0.05	—
Starved (40 hr.) rat	None	< 0.3	6.72 ± 0.20	34 ± 4†	3.7 ± 0.90†	—	< 0.05	—
Alloxan-diabetic rat	None	0.86 ± 0.13*	6.81 ± 0.75	53 ± 9*	2.5 ± 0.13*	3.8 ± 0.40*	0.13	14
Alloxan-diabetic rat treated with insulin	None	—	—	13 ± 4*	—	0.9 ± 0.24*	—	14
Normal fed rat	Albumin (2%)	< 0.3	6.30 ± 0.51	24 ± 4	—	1.7 ± 0.13	< 0.05	14
Normal fed rat	Albumin (2%) + palmitate (0.75 mm)	0.54 ± 0.13	2.88 ± 0.33*	70 ± 5*	—	4.7 ± 0.51*	0.19	15
Normal fed rat	—	—	—	22 ± 2	—	2.0 ± 0.13	—	11
Normal fed rat	DL-β-Hydroxybutyrate (5.5 mm)	2.01 ± 0.30*	3.0 ± 0.43*	62 ± 4*	1.1 ± 0.22	4.0 ± 0.50*	0.67	15
Normal fed rat	Butyrate (4 mm)	2.07 ± 0.24*	2.8 ± 0.37*	49 ± 3*	—	—	0.74	—
Normal fed rat	Pyruvate (4 mm)	—	—	90 ± 4*	—	5.7 ± 0.67*	—	16
Normal fed rat	Acetate (4 mm)	2.10 ± 0.21*	—	59 ± 1*	—	—	—	—
<i>Experiments in vivo</i>								
Normal fed rat	—	—	—	37 ± 6	—	—	—	—
Alloxan-diabetic rat	—	—	—	78 ± 4.3*	—	—	—	—

* $P < 0.01$; † $P < 0.02$; for other differences from appropriate control $P > 0.05$.

glycolytic enzymes, phosphofructokinase is located in cytoplasm, then inhibition of phosphofructokinase is presumably dependent on the transfer of citrate from mitochondria to cytoplasm, evidence for which has been presented by Lowenstein (1963).

Two factors that may contribute to the raised concentration of citrate induced by diabetes, starvation, fatty acids and ketone bodies are a rise in the acetyl-CoA/CoA concentration ratio and a rise in the concentration of oxaloacetate, since the equilibrium constant for the citrate-synthase reaction is given by:

$$K = \frac{[\text{citrate}][\text{CoA}]}{[\text{acetyl-CoA}][\text{oxaloacetate}]}$$

Evidence summarized in the preceding section has shown that the acetyl-CoA/CoA concentration ratio is increased by each of these agents that increase the concentration of citrate (except for starvation). The results of measurements of oxaloacetate concentration are given in Table 2. The concentration of oxaloacetate in hearts from normal fed rats perfused with medium containing glucose (1 mg./ml.) and insulin (0.1 unit/ml.) was 1.1 μ -moles/100 g. wet wt. The concentration of oxaloacetate was increased by starvation (3.4-fold) and by diabetes (2.3-fold) but not by perfusion with β -hydroxybutyrate. The increased citrate concentration in starvation may thus be due to the increased concentration of oxaloacetate.

The greater rise in citrate concentration in muscle from diabetic animals (in which the concentration of oxaloacetate was similar to that in starvation) may be due to the additional increase in the acetyl-CoA/CoA ratio in the muscle from the diabetic animal. The concentration of oxaloacetate in hearts perfused with glucose and insulin was markedly increased by perfusion with 4 mM-pyruvate (3-fold), and this could contribute to the increase in citrate concentration induced by pyruvate (Table 2).

The mechanism of the increased oxaloacetate concentration in diabetes and starvation is not known. Reactions that might contribute to the elevated concentration in muscle perfused with pyruvate are synthesis from pyruvate through actions of the malic enzyme and malate dehydrogenase and a coupled transamination reaction between pyruvate and aspartate catalysed by glutamate-pyruvate transaminase and glutamate-oxaloacetate transaminase. Pyruvate (4 mM) increased the concentration of malate in rat heart from 23 ± 3 to 128 ± 7 μ moles/100 g. wet wt. (means $\pm$ S.E.M. for six hearts in each group), increased that of α -oxoglutarate from 6.7 ± 0.8 to 25 ± 2.4 μ moles/100 g. wet wt. (means $\pm$ S.E.M. for six hearts in each group), and decreased that of glutamate from 323 ± 10 to 248 ± 9 μ moles/100 g.

wet wt. (means $\pm$ S.E.M. for five hearts in each group).

Mechanism of the effects of fatty acids and ketone bodies, and of alloxan-diabetes, on the concentrations of citrate and acetyl-CoA in perfused rat heart. The increased concentrations of acetyl-CoA and of citrate that we have observed in rat heart with diabetes and in perfusions with fatty acids and ketone bodies are explicable in terms of an increased rate of oxidation of one or other of these substrates to acetyl-CoA. This conclusion is suggested by the following points of evidence.

In hearts from diabetic rats or hearts from normal fed rats perfused with acetate, butyrate, octanoate, palmitate, β -hydroxybutyrate or acetoacetate the oxidation of glucose is depressed (Williamson & Krebs, 1961; Williamson, 1961; Garland *et al.* 1963, 1964). The oxygen consumption of the heart was not decreased by ketone bodies in the experiments of Fisher & Williamson (1961) nor in our experiments by alloxan-diabetes or by perfusion with 4 mM-acetate [results (ml. of oxygen/g. wet wt./hr.) were: for control heart, 6.0 ± 0.6 ; for heart from diabetic rat, 6.7 ± 0.5 ; for heart perfused with acetate, 5.4 ± 0.6 (means $\pm$ S.E.M. for five hearts in each group)]. In view of these findings it seems reasonable to suggest that the oxidation of fatty acids and ketone bodies are increased in hearts perfused with these substrates. Moreover, in view of these findings, and the demonstration of increased intracellular concentrations of free fatty acids and long-chain fatty acyl-CoA in the heart in diabetes, it seems reasonable to suggest that the respiration of endogenous fatty acids is accelerated in muscle from diabetic animals. Since in hearts perfused with fatty acids and ketone bodies or in hearts from diabetic rats the concentrations of acetyl-CoA and citrate are increased, it seems reasonable to conclude that the rate of formation of acetyl-CoA from these substrates (or from endogenous fatty acids in diabetes) is greater than the rate of oxidation of acetyl-CoA and citrate through the citrate cycle, until higher steady-state concentrations of acetyl-CoA and citrate are attained. The alternative explanation, that the accumulation of citrate and acetyl-CoA is due to impaired utilization of citrate, is unlikely for the following reasons.

The reactions that might be involved in the utilization of citrate are reaction with ATP to yield oxaloacetate and acetyl-CoA (catalysed by citrate-cleavage enzyme) and oxidation (after conversion into isocitrate) through the reactions catalysed by aconitase and isocitrate dehydrogenase. A diminution in the rate of citrate breakdown to acetyl-CoA and oxaloacetate by citrate-cleavage enzyme could not account for increases in the concentrations both of citrate and acetyl-CoA

observed with fatty acids and ketone bodies, nor for increases in the concentrations of citrate, acetyl-CoA and oxaloacetate observed in diabetes. Moreover, we have been unable to detect any ATP-dependent cleavage of citrate by extracts of rat heart measured under the conditions described by Srere (1959), from which we conclude that there is little, if any, citrate-cleavage enzyme in rat heart. Measurements of the intracellular concentrations of citrate and of isocitrate have shown that their concentrations increase in proportion in hearts from alloxan-diabetic rats or in hearts perfused with fatty acids or ketone bodies (see Table 2). The citrate/isocitrate ratios observed (11:1 to 16:1) are consistent with the 15:1 ratio observed with sheep-heart aconitase (Krebs, 1953). It seems unlikely therefore that the accumulation of citrate is due to any delay in the formation of isocitrate. It seems equally unlikely that the rise in citrate and isocitrate concentrations is due to any impairment in the oxidation of isocitrate. The oxygen consumption of the heart is not diminished by ketone bodies or diabetes or acetate, and, though some oxygen may be consumed in the formation of acetyl-CoA, citrate and isocitrate from β -hydroxybutyrate or fatty acids of C_4 or above, this is not so with acetate, which also increases the concentration of citrate. In view of these considerations it seems reasonable to conclude that the increased concentrations of acetyl-CoA and of citrate in rat heart in diabetes are due to the increased rate of oxidation of fatty acids to acetyl-CoA, and that this is a primary event in the development of abnormalities of glucose metabolism in muscle from diabetic animals described in the preceding papers.

In starvation of the rat for 40 hr. the concentration of acetyl-CoA was not increased, although the concentration of long-chain fatty acyl-CoA was increased. The cause of this difference in starvation (from that seen in diabetes or in hearts perfused with palmitate) has yet to be ascertained.

SUMMARY

1. The concentration of long-chain fatty acyl-CoA in perfused rat heart is increased by alloxan-diabetes, by starvation and by the addition of palmitate *in vitro* to the perfusion medium. In diabetes the concentration of fatty acyl-CoA can be returned to normal levels by treatment of the diabetic rat with insulin.

2. The concentration of free fatty acids in the perfused rat heart is increased by diabetes, and in rat hemidiaphragms by diabetes and starvation. In hypophysectomized rats the concentration of free fatty acids in hemidiaphragms was not increased by the induction of diabetes.

3. The output of glycerol by perfused rat heart was increased by diabetes and starvation. Prior hypophysectomy or adrenalectomy or treatment with insulin prevented diabetes from increasing the glycerol output. In rat hemidiaphragms the glycerol output was similarly increased by diabetes in normal but not in hypophysectomized rats. Treatment of hypophysectomized alloxan-diabetic rats with growth hormone and cortisol led to an increase in the glycerol output by hemidiaphragms.

4. The concentration of glycerol phosphate in rat heart and rat hemidiaphragms was decreased by diabetes and by starvation. Prior adrenalectomy or treatment of the rat with insulin prevented diabetes from lowering the concentration of glycerol phosphate.

5. It is concluded that there is increased provision of fatty acids for respiration in heart and diaphragm muscle of alloxan-diabetic and starved rats *in vitro*, due in part to an increased rate of lipolysis and possibly in part to a diminished rate of re-esterification of fatty acid.

6. In perfused hearts from diabetic rats or in hearts from normal fed rats perfused with medium containing fatty acids or ketone bodies the concentrations of acetyl-CoA, citrate and isocitrate are increased. These changes in concentrations are attributed to an increased rate of oxidation of fatty acids and ketone bodies to acetyl-CoA.

7. These findings are consistent with the view that an increased rate of oxidation of fatty acids and ketone bodies is responsible for abnormalities of glucose phosphorylation, fructose 6-phosphate phosphorylation and pyruvate oxidation in muscles of diabetic rats.

We thank the British Diabetic Association and the British Insulin Manufacturers for grants that helped to defray the expense of these investigations, the Royal Society for an equipment grant, and Mrs W. Thorne for skilled technical assistance.

REFERENCES

- Bergmeyer, H. U. & Bernt, E. (1963). In *Methods of Enzymatic Analysis*, p. 324. Ed. by Bergmeyer, H. U. Berlin: Verlag Chemie.
- Bernt, E. & Bergmeyer, H. U. (1963). In *Methods of Enzymatic Analysis*, p. 384. Ed. by Bergmeyer, H. U. Berlin: Verlag Chemie.
- Dagley, S. (1963). In *Methods of Enzymatic Analysis*, p. 313. Ed. by Bergmeyer, H. U. Berlin: Verlag Chemie.
- Duncombe, W. G. (1962). *Biochem. J.* **83**, 6 p.
- Fisher, R. B. & Williamson, J. B. (1961). *J. Physiol.* **158**, 86.
- Garland, P. B., Newsholme, E. A. & Randle, P. J. (1964). *Biochem. J.* **93**, 665.
- Garland, P. B. & Randle, P. J. (1962a). *Biochem. J.* **84**, 78 p.

- Garland, P. B. & Randle, P. J. (1962*b*). *Nature, Lond.*, **196**, 987.
- Garland, P. B. & Randle, P. J. (1963). *Nature, Lond.*, **199**, 381.
- Garland, P. B. & Randle, P. J. (1964). *Biochem. J.* **91**, 60.
- Garland, P. B., Randle, P. J. & Newsholme, E. A. (1963). *Nature, Lond.*, **200**, 169.
- Hohorst, H. J. (1963). In *Methods of Enzymatic Analysis*, p. 328. Ed. by Bergmeyer, H. U. Berlin: Verlag Chemie.
- Hohorst, H. J., Kreutz, F. H. & Bücher, Th. (1959). *Biochem. Z.* **332**, 18.
- Jungas, R. L. & Ball, E. G. (1962). *Fed. Proc.* **21**, 202.
- Krebs, H. A. (1953). *Biochem. J.* **54**, 78.
- Lowenstein, J. M. (1963). In *The Control of Lipid Metabolism*, p. 57. Ed. by Grant, J. K. London: Academic Press (Inc.) Ltd.
- Newsholme, E. A. & Randle, P. J. (1961). **80**, 655.
- Newsholme, E. A. & Randle, P. J. (1964). *Biochem. J.* **93**, 641.
- Randle, P. J., Newsholme, E. A. & Garland, P. B. (1964). *Biochem. J.* **93**, 652.
- Srere, P. A. (1959). *J. biol. Chem.* **234**, 2544.
- Tubbs, P. K. & Garland, P. B. (1964). *Biochem. J.* **93**, 550.
- Vaughan, M. (1961). *J. Lipid Res.* **2**, 293.
- Wieland, O. & Suyter, M. (1957). *Biochem. Z.* **329**, 320.
- Williamson, J. R. (1961). *Biochem. J.* **81**, 16r.
- Williamson, J. R. & Krebs, H. A. (1961). *Biochem. J.* **80**, 540.
- Wollenberger, A., Ristau, O. & Schoffa, G. (1960). *Pflüg. Arch. ges. Physiol.* **270**, 399.