APPENDIX

X-Ray-Diffraction and Electron-Microscope Observations on Soluble Derivatives of Feather Keratin

By B. K. FILSHIE, R. D. B. FRASER, T. P. MACRAE AND G. E. ROGERS
*Division of Protein Chemistry, C.S.I.R.O., Wool Research Laboratories, Parkville N. 2,
 Melbourne, Victoria, Australia*

(Received 10 September 1963)

It has been shown (Fraser & MacRae, 1963) that solutions of unfractionated feather-keratin derivatives can be dried down into films that give well-oriented X-ray-diffraction patterns containing many of the features associated with the microfibrillar structure of the native material. It was found that similar films could be obtained from the fractions described by Harrap & Woods (1964). In most cases spontaneous birefringence developed around the edge of the film during drying. The X-ray-diffraction patterns yielded by these films when the X-ray beam passed parallel to the surface were similar to those of the unfractionated derivatives at low angles of diffraction, with a prominent meridional reflexion at 23 Å and equatorial reflexion at 33 Å. At wider angles the fractionated materials show strong layer lines at 4.8 and 2.4 Å and a series of equatorials which index as orders of a 33 Å spacing.

The electron micrographs shown in Plate 1 were obtained by spraying a 1:1 (v/v) mixture of 0.05% protein solution and 1% phosphotungstate, pH 5.6, on to grids covered with a carbon-collodion film and using the negative-staining method of Brenner & Horne (1959). It is apparent from Plate 1 that, on drying, the fractionated derivatives spontaneously polymerize to form fibrils. The apparent

thickness of the fibrils varies from 40 to 60 Å at their narrowest part (arrow in Plate 1b) to approx. 130 Å at their widest part, suggesting that they are helical with a pitch of approx. 1500 Å. At first sight the fibrils appear to consist of two filaments coiled around each other to give a two-strand rope, but a ribbon containing three strands cannot be excluded at present. A feature is the tendency of the fibrils to aggregate laterally in precise register.

At the present time it is not clear whether the individual filaments about 40 Å in diameter in the fractionated proteins are structurally similar to the microfibrils in the native feather material (Rogers & Filshie, 1962; Filshie & Rogers, 1962; Fraser & MacRae, 1963), although the similarity between the low-angle X-ray patterns supports this view.

REFERENCES

- Brenner, S. & Horne, R. W. (1959). *Biochim. biophys. Acta*, **34**, 103.
 Filshie, B. K. & Rogers, G. E. (1962). *J. Cell Biol.* **13**, 1.
 Fraser, R. D. B. & MacRae, T. P. (1963). *J. molec. Biol.* **7**, 272.
 Harrap, B. S. & Woods, E. F. (1964). *Biochem. J.* **92**, 8.
 Rogers, G. E. & Filshie, B. K. (1962). *Proc. 5th int. Congr. Electron Microscopy*, 0-2. New York: Academic Press Inc.

Biochem. J. (1964), **92**, 19

Soluble Derivatives of Feather Keratin

2. MOLECULAR WEIGHT AND CONFORMATION

By B. S. HARRAP AND E. F. WOODS
*Division of Protein Chemistry, C.S.I.R.O., Wool Research Laboratories, Parkville N. 2,
 Melbourne, Victoria, Australia*

(Received 10 September 1963)

The preceding paper (Harrap & Woods, 1964) details the preparation, amino acid analysis and electrophoretic heterogeneity of feather proteins. Previous estimates of homogeneity with regard to size (Rougvie, 1954; Woodin, 1954) have suggested

that feather keratin is composed of fairly homogeneous units of mol.wt. 10000. Since, in contrast with Woodin (1954), we have found soluble feather proteins to be electrophoretically heterogeneous, it was important also to re-examine the molecular-

weight homogeneity of these proteins. The preparation of soluble derivatives by Rougvie's (1954) procedure involved the use of peracetic acid; this reagent has been shown for wool to give incomplete oxidation and hydrolysis of peptide bonds (O'Donnell & Thompson, 1959). The present paper reports the physical properties of derivatives obtained by milder and more specific means. We have critically examined the homogeneity by means of the sedimentation-equilibrium method, which has not previously been applied to these proteins.

To give further insight into the conformation of these proteins in solution, measurements of optical rotatory dispersion have been made both in aqueous solution and in several non-aqueous solvent mixtures.

EXPERIMENTAL

Materials

Soluble feather derivatives were prepared as described by Harrap & Woods (1964). Most of the work reported in the present paper has been done with *S*-carboxymethyl-rachis; a few studies were also done on rachis keratose and *S*-sulpho-rachis. The partially fractionated *S*-carboxymethyl-rachis samples were also prepared, as described by Harrap & Woods (1964), by the ethanol-zinc acetate method.

2-Chloroethanol was purified as described by Goldstein & Katchalski (1960); all other chemicals were of A.R. quality.

Methods

Sedimentation studies. The Spinco model E ultracentrifuge equipped with RTIC control unit and phase plate was used for all sedimentation experiments. Both schlieren and Rayleigh interference optics were employed. For interference optics a symmetrical upper mask assembly of slit width 0.75 mm. was used, and the optics were aligned by the procedures outlined in the Spinco service manual and by LaBar & Baldwin (1962). Interference patterns were photographed on type IIG emulsion spectroscopic plates (Eastman Kodak). These were aligned on the stage of a two-dimensional comparator (Nikon Shadowgraph, model 6B) by means of the fringe system from the air reference holes. Schlieren patterns from sedimentation-velocity experiments were recorded on metallographic plates.

Velocity sedimentation was carried out at 59780 rev./min. in single-sector capillary-type synthetic-boundary cells (Kegeles, 1952). Velocity runs were of 70–80 min. duration and photographs were taken at 4 min. intervals for the calculation of sedimentation coefficients. Double-sector cells with epoxy resin centre pieces and quartz windows were used for equilibrium experiments at speeds from 13000 to 25000 rev./min. Perfluorotributylamine (F.C. 43; Minnesota Mining and Manufacturing Co., Minn., U.S.A.) or fluorocarbon oil (Kel F1; M. W. Kellogg Co., N.J., U.S.A.) was used as a base for the liquid columns in the equilibrium studies. To find the reference base line corrected for any cell distortion, the cell was flushed with water at the end of an experiment, the two sides were

refilled with water to the previous levels, and the reference base line was recorded at the same speed. The correction to the fringe numbers was made as described by LaBar & Baldwin (1962). The values of initial protein concentration, c_0 , required for the calculations of molecular weight for both schlieren and interference optics were determined by using a double-sector synthetic-boundary cell at speeds of less than 8000 rev./min.; generally two runs were made on one solution and the value obtained for Δn (the refractive index difference between the solution and its diffusate) was checked with a Brice-Phoenix differential refractometer. Subsequent dilutions were made by weight and c_0 was calculated. Fractional fringe numbers were determined from two or three frames photographed immediately after reaching speed and when solvent and solution menisci were coincident. Column heights in equilibrium studies were in the range 0.8–2.5 mm. The time for equilibrium to be established was calculated from the equation of Van Holde & Baldwin (1958). In all cases equilibrium was attained in less than 16 hr. at 25°.

Molecular weights were calculated by several methods:

(1) Schlieren optics. For column heights greater than 1.2 mm. the apparent weight-average molecular weight, M_w (app.), over the entire column was calculated from the equation (Svedberg & Pedersen, 1940):

$$M_w(\text{app.}) = \frac{RT}{(1 - \bar{v}\rho)\omega^2} \cdot \frac{2\Delta c}{c_0(r_b^2 - r_m^2)} \quad (1)$$

where ω is the angular velocity of the rotor, $\bar{v}$ the partial specific volume of the protein, ρ the density of the solvent, r_m and r_b the distances from the centre of rotation to the top and bottom menisci respectively, c_0 the initial concentration of the protein, and Δc (i.e. $c_b - c_m$) the difference in the concentrations of protein at the top (c_m) and bottom (c_b) menisci. The quantity Δc was calculated from the integral:

$$\int_{r_m}^{r_b} \Delta r \frac{dc}{dr}$$

Both Δc and c_0 were expressed in area units and calculated by trapezoidal integration.

The M_w (app.) value at any point in the sedimenting column was calculated from the equation:

$$M_w(\text{app.}) = \frac{RT}{(1 - \bar{v}\rho)\omega^2} \cdot \frac{dc}{rc} \quad (2)$$

where dc/dr is the concentration gradient and c the concentration at any point, r , in the cell, calculated as described by O'Donnell & Woods (1962). For columns sufficiently short such that c_r equals c_0 at the centre of the column (less than 1 mm.), M_w (app.) was calculated from measurements of the concentration gradient at the mid-point by using eqn. (2) (Yphantis, 1960).

The *Z*-average molecular weight, M_z , was calculated by using eqn. (3) (Van Holde & Baldwin, 1958):

$$M_z(\text{app.}) = \frac{RT}{(1 - \bar{v}\rho)\omega^2} \cdot \frac{\left(\frac{1}{r} \cdot \frac{dc}{dr}\right)_b - \left(\frac{1}{r} \cdot \frac{dc}{dr}\right)_m}{\Delta c} \quad (3)$$

(2) Interference optics. M_w (app.) was calculated by using eqn. (1), where the concentrations Δc and c_0 are expressed in terms of fringe numbers. To evaluate Δc ,

which represents the total fringe count across the column, it is necessary to evaluate the fractional fringe accurately. To do this another form of eqn. (2) was employed:

$$M_w(\text{app.}) = \frac{2RT}{(1 - \bar{v}\rho)\omega^2} \cdot \frac{d(\ln c)}{d(r^2)} \quad (4)$$

A plot of $\log c$ against r^2 , where r is the distance of each fringe from the centre of rotation, extrapolated to r_m^2 and r_b^2 gives Δc directly. However, to do this it is necessary to be able to assign a concentration (in terms of fringe number) to one of the fringes in the cell. By counting the fringes across the plates an approximate value of Δc could usually be determined to within one fringe number. From this value of Δc , a first approximation, c'_m , to the concentration at the meniscus, was calculated from the approximate equation (Van Holde & Baldwin, 1958):

$$c_m = \frac{\Delta c}{e\Delta c/c_0 - 1} \quad (5)$$

Thus, from the value c'_m , fringe numbers, c' , were assigned to all the fringes across the column. A plot of $\log c'$ against r^2 was then made and Δc calculated from the extrapolated values of c'_m and c'_b . The quantity Δc is not significantly affected by a small departure of the c' values from the true values. From the Δc values so obtained the true c_m was calculated from the equation (Richards & Schachman, 1959):

$$c_m = c_0 - \frac{r_b^2 \Delta c - \int_{r_m}^{r_b} r^2 dc}{r_b^2 - r_m^2} \quad (6)$$

and the true concentrations throughout the cell followed directly. To calculate $M_z(\text{app.})$ the equation derived by Lansing & Kraemer (1935) was used:

$$M_z(\text{app.}) = \frac{M_{w,b}c_b - M_{w,m}c_m}{c_b - c_m} \quad (7)$$

where $M_{w,b}$ and $M_{w,m}$ were evaluated from the limiting slopes of the plot of $\log c$ against r^2 at the bottom and top menisci respectively by using eqn. (4).

Diffusion coefficient. This was calculated by the reduced height-area method from the rate of spreading of a boundary in a synthetic-boundary cell in the ultracentrifuge at low speed (7928 rev./min.). It was also determined from the rate of approach-to-sedimentation equilibrium as described by Sophianopoulos, Rhodes, Holcomb & Van Holde (1962). According to this method the graph of $\log[(dc/dr)_{eq} - (dc/dr)_t]$ against time, t , should be linear with slope given by:

$$\frac{-\pi^2 D}{r_b^2 - r_m^2} \left\{ 1 + \left(\frac{r_b^2 - r_m^2}{2\pi} \right)^2 \left[\frac{\omega^2 M(1 - \bar{v}\rho)}{2RT} \right]^2 \right\} \quad (8)$$

where $(dc/dr)_{eq}$ and $(dc/dr)_t$ are the values of the concentration gradients at the mid-point at equilibrium and t respectively.

Partial specific volume. The apparent specific volumes, $\bar{v}'$, were calculated from the densities of the dialysed protein solutions and their diffusates. The densities were measured in 25 ml. pycnometers at 25°. The relation between the thermodynamic partial specific volume $\bar{v}$ and $\bar{v}'$ in three-component systems has been discussed by Casassa & Eisenberg (1961). The insertion of $\bar{v}'$ into eqns. (1)–(4) gives

the anhydrous molecular weight on extrapolation to infinite dilution.

Intrinsic viscosity. Simple Ostwald viscometers with a water time of 200 sec. were used to measure the viscosities at $25 \pm 0.02^\circ$.

Refractive index increment. Measurements were made at 25° with a Brice-Phoenix differential refractometer at 546 m μ . For *S*-carboxymethyl-rachis dn/dc is 0.00186 (c in g./100 ml.) in 0.01 M-borate–0.2 M-NaCl buffer, and 0.00158 in 8 M-urea–0.2 M-NaCl.

Optical rotatory dispersion. A spectropolarimeter, which was built up from a photoelectric polarimeter (W. F. Stanley and Son), a quartz monochromator (Carl Leiss), and a mercury-arc light source, was used for these determinations. Measurements were made at the following wavelengths: 578, 546, 486, 436, 405, 365 and, occasionally, 334 m μ . Solutions were clarified by filtration through sintered-glass filters (Jena; porosity 4). Water-jacketed (25°) polarimeter tubes (1 dm.) with fused quartz end plates were used.

Protein concentrations. These were determined by measuring the Kjeldahl nitrogen and taking the nitrogen content as 16.1% (Harrap & Woods, 1964), and where possible these were checked refractometrically. When protein concentrations in 8 M-urea were required the method used by Herriott (1935) to measure the tyrosine content of pepsin with Folin reagent was applied. A calibration curve was first determined with an aqueous solution of the feather protein and then compared with a similar curve obtained with the same protein dissolved in 8 M-urea. Urea did not interfere with the colour development. Extinctions were measured at 750 m μ .

RESULTS

Sedimentation velocity. *S*-Carboxymethyl-rachis in 0.01 M-borate–0.2 M-sodium chloride buffer, pH 9.1, gave a single sedimenting peak at all concentrations in the ultracentrifuge. Fig. 1 presents a plot of $1/S_{20,w}$ (as s⁻¹) against c (as g./100 ml.), which gives on least-squares analysis the following equation:

$$1/S_{20,w} = 0.883 + 0.199c; \quad S_{20,w}^0 = 1.13 \pm 0.02 \text{ s.}$$

Intrinsic viscosity. Fig. 2 presents a plot of the specific viscosity, i.e. $(\eta_{rel.} - 1)/c$, against c for *S*-carboxymethyl-rachis in 0.01 M-borate–0.2 M-

Fig. 1. Sedimentation velocity as function of concentration for *S*-carboxymethyl-rachis in 0.01 M-sodium borate–0.2 M-NaCl buffer, pH 9.1.

sodium chloride buffer, pH 9.1, and also in 8M-urea-0.02M-tris-hydrochloric acid-0.2M-sodium chloride buffer, pH 7.5. Least-squares analysis of the results leads to values for the intrinsic viscosities, i.e. $[\eta]$, of $0.14_8 \pm 0.004$ dl./g. in aqueous buffer at pH 9 and $0.19_2 \pm 0.001$ dl./g. in 8M-urea.

Partial specific volume. The mean of three determinations at different protein concentrations gave a value of $\bar{v}'$ of 0.703 ml./g. in 0.01M-borate-0.2M-sodium chloride buffer, pH 9.1. Two determinations in 8M-urea-0.02M-tris-hydrochloric acid-0.2M-sodium chloride buffer, pH 7.5, also gave a mean value of 0.703 ml./g.

Diffusion coefficient. Two measurements were made by the reduced height-area method in the ultracentrifuge and gave values for $D_{20,w}$ of 8.2×10^{-7} and 8.0×10^{-7} cm.²sec.⁻¹ at protein concentrations of 0.97 and 0.58 % respectively. Fig. 3 presents the diffusion coefficients calculated from the rate of approach to equilibrium at various protein concentrations. In each experiment the plot of $\log[(dc/dr)_{eq} - (dc/dr)_t]$ against t was linear. Least-squares analysis of the results in Fig. 3 gives a value for $D_{20,w}^0$ of 7.7×10^{-7} cm.²sec.⁻¹.

Molecular weight from Svedberg equation. By combining the value of $S_{20,w}^0$ with $D_{20,w}^0$ determined by the approach-to-equilibrium method, a value of 12000 is obtained for the molecular weight. If we take the mean of the two reduced height-area measurements for $D_{20,w}$ and combine this with $S_{20,w}^0$, a value of 11400 is obtained for the molecular weight.

Fig. 2. Viscosity of *S*-carboxymethyl-rachis in: ○, 0.01M-sodium borate-0.2M-NaCl buffer, pH 9.1; △, 8M-urea-0.02M-tris-HCl-0.2M-NaCl buffer, pH 7.5.

Fig. 3. Diffusion coefficient as a function of concentration for *S*-carboxymethyl-rachis, calculated from approach to equilibrium.

Sedimentation equilibrium. In the initial sedimentation-equilibrium experiments the schlieren patterns showed a sharp upward curvature towards the bottom meniscus. Calculations of the molecular weight by means of eqn. (2) showed that this was constant over three-quarters of the column and sharply increased over the remainder of the column. In these experiments the amount of aggregated material was estimated to be approx. 10 % of the total; this quantity was even greater if silicone oil or carbon tetrachloride was used as base liquid. The aggregated material was largely removed by centrifugation in the Spinco model L preparative ultracentrifuge at 40000 rev./min. for 15 hr. Fig. 4 shows the change in the schlieren patterns after the treatment, and the following results refer to experiments carried out on protein solutions that had been subjected to a preliminary centrifugation. It was essential to redialyse after the preliminary centrifugation as there is appreciable redistribution of salts during the preparative centrifugation. Failure to do this gave a low c_0 value, and the values for the molecular weight given by Woods (1961) were high for this reason.

Table 1 presents the molecular-weight data for *S*-carboxymethyl-rachis in 0.01M-borate-0.2M-sodium chloride buffer, pH 9.1, calculated for both schlieren and interference optics. In each series the most concentrated solution was given a preliminary centrifugation as described above. For series 1 the

Fig. 4. Sedimentation equilibrium of *S*-carboxymethyl-rachis (a) before (c 0.91 g./100 ml.) and (b) after (c 0.71 g./100 ml.) preliminary centrifuging in the Spinco model L centrifuge at 13410 rev./min.; the phase-plate angle was 70°; the solvent was 0.01M-borate-0.2M-NaCl buffer, pH 9.1.

upper three-quarters of the supernatant was studied and for series 2 only the top one-third was used. It is evident from the results that the apparent molecular weight is dependent on concentration. This complicates any analysis for heterogeneity.

When employing interference optics homogeneity may be assessed from plots of $\log c$ against r^2 across the column. For an ideal monodisperse two-component system these plots should be linear. For series 1 of Table 1 the plots of $\log c$ against r^2 were slightly concave upwards, with the exception of the solution of highest concentration, which gave a linear plot. For series 2 the plots for the two highest concentrations were slightly concave downwards, and for the two lowest concentrations they were linear over most of the column but curved upwards at r values close to the bottom meniscus. Upward curvature of the plots of $\log c$ against r^2 is an indication of a continuing increase in molecular weight throughout the column, whereas downward curvature arises from non-ideality of the solution and has been reported for myosin (Woods, Himmelfarb & Harrington, 1963). In the system studied in the present work it seems that both effects are present and may compensate each other. Which of the two effects is predominant depends on the concentration difference between the top and bottom menisci. At high initial concentrations the difference in concentration, i.e. $c_b - c_m$, between the two menisci is greater than at lower initial concentrations, and the non-ideality effect predominates; at lower initial concentrations heterogeneity is the dominating factor. Similar conclusions were drawn from the molecular weight as a function of r calculated from eqn. (2) by using schlieren optics, and from the form of the plots of $(1/r)(dc/dr)$ against c_r . It is difficult in this situation to make an exact estimate of the heterogeneity, but from the form of

these functions series 2 is less heterogeneous than series 1. The molecular weights from schlieren optics (Table 1) are in agreement with this conclusion; however, from interference optics there is little difference between the two series.

The molecular-weight data presented in Table 1 indicate that the values of $M_z(\text{app.})$ determined from schlieren optics are in most cases greater than those determined from interference optics. The determination of M_z involves the slopes of the plots of $\log c$ against r^2 at the ends of the column in interference optics and of the values of dc/dr at the menisci in schlieren optics, both of which are subject to large errors. The extrapolated M_z and M_w values (averaging the results from the two optical systems) and the constancy of the molecular weights calculated at different points in the column are inconsistent with a continuous distribution of molecular weights. Further, the functions $\log c$ against r^2 and $(1/r)(dc/dr)$ against c_r were linear over more than 80% of the column, showing a steep rise only in the vicinity of the bottom meniscus, indicating that the system consists of a monomer with a small percentage of material of much higher molecular weight than the main component. We estimate from this upward curvature at the base that series 1 contains about 2.5% of aggregate and series 2 about 1.5%. To determine the molecular weight of the monomer we chose a value for the monomer molecular weight which best satisfied the values for M_w and M_z for the two series of measurements and at the same time agreed with the estimates of the amount of aggregate present. By this means we arrived at a value of 10600 ± 300 for the monomer molecular weight and about 40000 for the molecular weight of the aggregate.

Since the protein carries a net charge under the conditions of our measurements we must apply the

Table 1. *Molecular weight of S-carboxymethyl-rachis*

Experimental details are given in the text.

	Initial c (g./100 ml.)	Schlieren optics			Interference optics		
		$M_w(\text{app.})$	$M_z(\text{app.})$	M_z/M_w	$M_w(\text{app.})$	$M_z(\text{app.})$	M_z/M_w
Series 1	1.214	9990	11630	1.16	10090	10950	1.11
	0.964	10120	10740	1.06	9660	9050	0.94
	0.730	10280	11280	1.10	10120	10340	1.02
	0.477	10510	13140	1.25	10450	11340	1.08
	0.239	10920	14740	1.35	10730	11170	1.04
	0*	11050	15160	1.37	10950	11420	1.04
Series 2	0.796	9830	10240	1.04	9780	9570	0.98
	0.554	10200	10210	1.00	10310	10390	1.01
	0.399	10530	10980	1.04	10240	10180	0.99
	0.226	10680	11040	1.03	10400	11240	1.08
	0*	11140	11480	1.03	10720	11790	1.10

* Extrapolated values from least-squares analysis of $1/M(\text{app.})$ as a function of c , where c for M_w refers to $[(c_m + c_b)/2]$ and for M_z to $(c_m + c_b)$ (Van Holde & Baldwin, 1958).

Table 2. *Optical rotatory dispersion of S-carboxymethyl-rachis*

Experimental details are given in the text. The P₂ and S₂ fractions were prepared as described by Harrap & Woods (1964).

Solvent	Unfractionated		P ₂ fraction		S ₂ fraction	
	$[\alpha]_D$	b_0	$[\alpha]_D$	b_0	$[\alpha]_D$	b_0
Water	-136°	0	-132°	0	-149°	0
8M-Urea	-139	0	—	—	—	—
2-Chloroethanol-formic acid (4:1, v/v)	—	—	-47	-220	-58	-240
Acetic acid-formic acid (4:1, v/v)	—	—	-75	-150	-83	-180
Dioxan-formic acid (1:1, v/v)*	—	—	-59	-50	-70	-70
Methanol-formic acid (4:1, v/v)	—	—	-86	0	-89	0

* Higher concentrations of dioxan could not be investigated because of precipitation of the protein.

equations of Williams, Van Holde, Baldwin & Fujita (1958) for the sedimentation of strong electrolytes to correct the molecular weight according to the equation:

$$\lim_{c \rightarrow 0} M(\text{app.}) = M_P \left[1 - \frac{Z \cdot M_B \cdot (1 - \bar{v}_B \rho)}{2 M_P (1 - \bar{v}_P \rho)} \right] \left(1 - \frac{Z \cdot M_B \cdot \theta_B}{2 M_P \theta_P} \right)$$

where Z is the net charge on the protein, the subscripts B and P refer to the supporting electrolyte and protein respectively, and the quantities θ_B and θ_P are the specific refractive increments. From the amino acid analysis (Harrap & Woods, 1964) the net charge at pH 9 is -6; assuming the supporting electrolyte to be entirely sodium chloride the product in brackets is 0.965 and M_P is therefore 10980. This molecular weight will also include any bound ions since the components of the system have been defined on the basis of dialysis equilibrium (Casassa & Eisenberg, 1961).

Less detailed measurements were also carried out on rachis keratose and *S*-sulpho-rachis. The results indicated that the molecular weights of these two derivatives are essentially the same as that of the *S*-carboxymethyl derivative.

Further measurements on the *S*-carboxymethyl-rachis were also made in 8M-urea to determine if any further decrease in molecular weight occurred in this solvent. To obtain anhydrous molecular weights in this three-component system we have made use of the treatment of Casassa & Eisenberg (1961), which has been discussed in detail in its application to solutions of myosin in concentrated guanidine hydrochloride solutions by Woods *et al.* (1963). To decrease the time taken to reach equilibrium the short-column method of Yphantis (1960) was used, as well as some longer-column experiments. Molecular weights in 8M-urea-0.02M-tris-hydrochloric acid-0.2M-sodium chloride buffer, pH 7.5, were about four times as dependent on concentration as those in 0.01M-borate-0.2M-sodium chloride buffer, pH 9.1, and they were

extrapolated to give a value for $M_w(\text{app.})$ of 11400. Control experiments with urea alone employing interference optics showed that the results were not affected by the distribution of urea in the centrifugal field. The schlieren-equilibrium patterns were similar in appearance to that in Fig. 4(a), showing that aggregates were still present and were not dispersed in urea. In view of this the agreement between the infinite dilution values of $M_w(\text{app.})$ in the two solvent systems is satisfactory and it is clear that 8M-urea does not further disaggregate the protein.

Optical rotatory dispersion. The optical-rotatory-dispersion data were treated according to the Moffitt & Yang (1956) equation:

$$m' = \frac{3}{n^2 + 2} \cdot \frac{M}{100} \cdot [\alpha] = a_0 \left(\frac{\lambda_0^2}{\lambda^2 - \lambda_0^2} \right) + b_0 \left(\frac{\lambda_0^2}{\lambda^2 - \lambda_0^2} \right)^2$$

where m' is the reduced residue rotation (obtained by correcting $[\alpha]$ for the refractive index, n , of the solvent), M is the mean residue weight (which was taken as 115), λ_0 is a dispersion constant (which was taken as 212 mμ; Urnes & Doty, 1961), and a_0 and b_0 are parameters related to the conformation of the protein. Values of b_0 and $[\alpha]_D$ are shown in Table 2 for both aqueous and non-aqueous solutions of *S*-carboxymethyl-rachis and of two fractions derived from it. The $[\alpha]_D$ values were obtained by extrapolation of results at lower wavelengths. The two fractions, P₂ and S₂, were obtained by ethanol precipitation as described by Harrap & Woods (1964), and represent 43 and 50 % respectively of the soluble protein. Only one measurement was made on the P₁ fraction (7 % of the total) because of the small amount available, and it yielded a value of zero for b_0 .

DISCUSSION

Molecular weights from sedimentation and diffusion measurements are somewhat higher than those from sedimentation-equilibrium experiments. The agreement, however, can be considered satis-

factory, since the sedimentation-velocity runs were made on solutions that had received no preliminary preparative ultracentrifugation and must have contained some aggregate; equilibrium patterns of such solutions were similar to Fig. 4(a). Also, the use of eqn. (8) to calculate diffusion coefficients from the approach to equilibrium assumes homogeneity of the preparation and no variation of D with concentration. This may account for the higher values of $D_{20,w}$ obtained by the reduced height-area method. We therefore take the best molecular weight to be that obtained from sedimentation equilibrium; this gives 10980 for the monomer after correcting for charge effects. This molecular weight refers to the *S*-carboxymethyl-protein, and as we are interested in the subunit as it exists in the unreduced state in the native keratin we subtract the increase in weight due to the conversion of each $-S-S-$ bond (four per molecule) into two $-S\cdot CH_2\cdot CO_2^-$ groups, and the molecular weight of the cations (Na^+) associated with the six negatively charged groups present at the pH of measurement. We thus arrive at a molecular weight of 10400 for the fundamental subunit of feather keratin. The dissolved protein accounts for 90 % of the feather rachis, and 90 % of the protein extracted is in the monomeric form. Since aggregates form on aging at the expense of the monomer we believe that the aggregates initially present consist of polymers of the subunit.

Other comparable studies on soluble feather derivatives are those of Woodin (1954) and Rougvie (1954), in both of which whole feather was used. In general our results are in agreement with theirs in that the molecular weight of the feather-keratin monomer is about 10000. Woodin's (1954) light-scattering measurements were made in 5*M*-urea solutions and he used a value of 0.00208 for dn/dc in the light-scattering equation to calculate the molecular weight. We have found a value of 0.00169 for dn/dc of *S*-carboxymethyl-rachis and rachis keratose in 5*M*-urea. Since $(dn/dc)^2$ appears in the light-scattering equation, Woodin's (1954) weight-average molecular weights should be considerably increased. On this basis his weight-average molecular weights would no longer agree with the number-average molecular weight obtained from osmotic-pressure measurements, and thus his solutions were not homogeneous and probably contained aggregates. The sedimentation-equilibrium measurements reported above allow a much more critical estimate of the homogeneity of feather keratin solutions than the previous studies. The high degree of homogeneity with regard to molecular weight contrasts with the electrophoretic heterogeneity reported by Harrap & Woods (1964).

The values of $[\alpha]_D$ in aqueous urea solution are somewhat more negative than those encountered

for most proteins even when denatured. This probably reflects the large negative residue rotation contributed by the high proline content of the proteins. Dweltz & Mahadevan (1961) have reported a value of $[\alpha]_D$ of -430° for a soluble feather protein in concentrated urea solution and have used this value to support a triple-helical model of feather keratin based on a similar model for collagen (Ramachandran & Dweltz, 1962). Their value of $[\alpha]_D$ is considerably different from that reported in Table 2. Since our values for $[\alpha]_D$ are of the same order as those quoted by Westover, Tiffany & Krimm (1962), this discrepancy may be connected with a gross error in Dweltz & Mahadevan's (1961) procedure of determining protein concentrations in concentrated urea solutions which involved removal of urea by dialysis against water for several days and applying a correction for the observed diffusion of protein through the bag. In our opinion this method must necessarily be subject to large errors. In any case it does not seem reasonable to interpret the value of $[\alpha]_D$ of -430° in terms of a triple-helix structure by analogy with collagen since the large laevo-rotation in collagen is connected with its polyproline II-type structure (Harrington & Hippel, 1961) rather than to its triple-helix structure.

The zero values for b_0 for the soluble feather proteins in aqueous buffer solutions (Table 2) indicate that under these conditions the proteins are in the random-coil conformation (Urnes & Doty, 1961). Consequently we have calculated the β -function of Scheraga & Mandelkern (1953) from the hydrodynamic results. The value obtained, 2.52×10^6 , is also consistent with the protein being in a random-coil conformation. There is no change in b_0 or $[\alpha]_D$ in 8*M*-urea; this supports the view of a random coil in dilute buffer solutions and indicates that the increase in $[\eta]$ in 8*M*-urea (Fig. 2) does not reflect a transition from a helical to more random conformation. Westover *et al.* (1962) have postulated from optical-rotatory-dispersion studies that solubilized feather calamus also exists in the random-coil form in solution.

Values for b_0 in several solvent mixtures for the two fractions S_2 and P_2 (Table 2) show that under suitable conditions these proteins can develop apparent helical contents ($-100b_0/640$) of as much as 40 %. Any further development in helical content of these proteins is probably limited by their high percentage of proline, since this residue cannot be accommodated in the α -helix structure. Szent-Györgyi & Cohen (1957) have indicated for aqueous solutions that if the proline content is greater than about 8 % and if the proline is randomly distributed then helix formation in proteins is probably prevented. They have also pointed out that this limitation is not necessarily true for non-aqueous

solutions, and this is supported by our results, which show that helical structures can still be formed even with a proline content as high as 11 %.

Schor & Krimm (1961*a, b*) have postulated that a proline residue occurs regularly at every eighth or ninth residue along the polypeptide chains of feather keratin. In view of the optical-rotation data for the soluble feather keratin in organic solvents this seems unlikely, since Goodman, Schmidt & Yphantis (1960) have shown that even for oligopeptides of high helix-forming ability stable α -helices are not formed for chains shorter than the nonapeptide.

SUMMARY

1. A detailed study has been made of the molecular weight of *S*-carboxymethyl-rachis of fowl feathers by sedimentation-equilibrium measurements. Both the schlieren and interference optical systems were used.

2. Most (85–90 %) of the proteins of feather-rachis keratin consist of uniformly sized species of mol.wt. 10400. As studied in solution higher-molecular-weight species may occur but these are probably aggregates of the subunit. No further decrease in molecular weight of the *S*-carboxymethyl derivatives occurs in 8*M*-urea.

3. From sedimentation-velocity and viscosity measurements the dissolved proteins appear to be in the random-coil conformation in aqueous solution, and this is supported by optical-rotatory-dispersion measurements.

4. A considerable degree of α -helix formation in the soluble proteins can be induced by organic solvent mixtures, and the significance of this is discussed in relation to physical models proposed for feather keratin.

The authors acknowledge the capable technical assistance of Mr D. J. A. Evans.

REFERENCES

- Casassa, E. F. & Eisenberg, H. (1961). *J. phys. Chem.* **65**, 427.
- Dweltz, N. E. & Mahadevan, V. (1961). *Biochem. J.* **81**, 134.
- Goldstein, L. & Katchalski, E. (1960). *Bull. Res. Council. Israel*, **9A**, 138.
- Goodman, M., Schmidt, E. E. & Yphantis, D. (1960). *J. Amer. chem. Soc.* **82**, 3483.
- Harrap, B. S. & Woods, E. F. (1964). *Biochem. J.* **92**, 8.
- Harrington, W. F. & Hippel, P. H. von (1961). *Advanc. Protein Chem.* **16**, 1.
- Herriott, R. M. (1935). *J. gen. Physiol.* **19**, 283.
- Kegeles, G. (1952). *J. Amer. chem. Soc.* **74**, 5532.
- LaBar, F. E. & Baldwin, R. L. (1962). *J. phys. Chem.* **66**, 1952.
- Lansing, W. D. & Kraemer, E. O. (1935). *J. Amer. chem. Soc.* **57**, 1369.
- Moffitt, W. & Yang, J. T. (1956). *Proc. nat. Acad. Sci., Wash.*, **42**, 596.
- O'Donnell, I. J. & Thompson, E. O. P. (1959). *Aust. J. biol. Sci.* **12**, 294.
- O'Donnell, I. J. & Woods, E. F. (1962). In *Modern Methods of Plant Analysis*, p. 291. Ed. by Paech, K. & Tracy, M. V. Berlin: Springer-Verlag.
- Ramachandran, G. N. & Dweltz, N. E. (1962). In *Collagen*, p. 147. Ed. by Ramanathan, N. New York: Interscience Publishers Inc.
- Richards, E. G. & Schachman, H. K. (1959). *J. phys. Chem.* **63**, 1578.
- Rougvie, M. A. (1954). Ph.D. Thesis: Massachusetts Institute of Technology.
- Scheraga, H. A. & Mandelkern, L. (1953). *J. Amer. chem. Soc.* **75**, 179.
- Schor, R. & Krimm, S. (1961*a*). *Biophys. J.* **1**, 467.
- Schor, R. & Krimm, S. (1961*b*). *Biophys. J.* **1**, 489.
- Sophianopoulos, A. J., Rhodes, C. K., Holcomb, D. N. & Van Holde, K. E. (1962). *J. biol. Chem.* **237**, 1107.
- Svedberg, T. & Pedersen, K. D. (1940). *The Ultracentrifuge*, p. 346. Oxford: Oxford University Press.
- Szent-Györgyi, A. G. & Cohen, C. (1957). *Science*, **126**, 697.
- Urnas, P. J. & Doty, P. (1961). *Advanc. Protein Chem.* **16**, 402.
- Van Holde, K. E. & Baldwin, R. L. (1958). *J. phys. Chem.* **62**, 734.
- Westover, C. J., Tiffany, M. L. & Krimm, S. (1962). *J. molec. Biol.* **4**, 316.
- Williams, J. W., Van Holde, K. E., Baldwin, R. L. & Fujita, H. (1958). *Chem. Rev.* **58**, 715.
- Woodin, A. M. (1954). *Biochem. J.* **57**, 99.
- Woods, E. F. (1961). *Abstr. 5th int. Congr. Biochem., Moscow*, no. 2–37–757.
- Woods, E. F., Himmelfarb, S. & Harrington, W. F. (1963). *J. biol. Chem.* **238**, 2374.
- Yphantis, D. (1960). *Ann. N.Y. Acad. Sci.* **88**, 586.