bmj.com news roundup

Full versions of these stories are available at: bmj.com/content/vol331/issue7517/#NEWS_ROUNDUP

UK agency recommends better monitoring of multidrug resistant *E coli*

The Health Protection Agency, the body that monitors and advises on infectious diseases in the United Kingdom, has recommended that monitoring of multidrug resistant *Escherichia coli* be improved. Its report, published this week, shows a big increase over the past four years in the number of infections associated with the bacterium.

The agency found that the number of infections caused by *E coli* bacteria that produce extended spectrum β lactamase (ESBL) was growing. Extended spectrum β lactamase is an enzyme that hydrolyses and confers resistance to modern cephalosporin antibiotics. The bacteria normally live harmlessly in the gut but are also one of the commonest causes of urinary tract infections.

The agency said that more than 1000 isolates of one particular type of ESBL producing *E coli* had been reported to its Antibiotic Resistance Monitoring and Reference Laboratory since 2003 and that these strains had not been seen in the UK before 2000.

Susan Mayor *London*

Investigations into Multi-drug Resistant ESBL-producing Escherichia coli Strains Causing Infections in England is accessible at www.hpa.org.uk.

Poor management, not drug costs, is hampering implementation of NICE guidance

Poor financial management and not the cost of drugs and treatment is preventing clinicians in England from doing their jobs properly, says the Audit Commission.

The watchdog of public spending claims that much of the guidance issued by the National Institute for Health and

MARC CARTER/STOCKGETTY IMAGES

Eight in 10 dancers have an injury each year, survey shows

Eight in 10 dancers in the United Kingdom have at least one injury a year that affects their ability to perform. Almost one in 10 are underweight, and a quarter report having had an eating problem at some time. These are among the findings of the second UK national survey of dancers' health, injuries, and lifestyle.

It shows that 80% of the 1056 professional dancers surveyed experience at least one injury each year. Professional dancers, such as the one pictured, had an average of 11.5 days off classes, 7.9 days off rehearsal, and 6.6 days off performance because of injury in the previous year. Fourteen per cent of dancers had experienced a longer term injury that kept them off class, rehearsal, and performance for up to 18 months.

The highest rates of injury were among professional ballet dancers, and the most common site of injury was the lower back, followed by the ankles and knees. One in five female professional dancers (21%) had experienced their periods stopping for six months or more at some point in the past.

The report recommends that companies and schools put greater emphasis on body composition and health than on weight and thinness.

Roger Dobson *Abergavenny*

Fit to Dance 2 is available from Dance UK, price £8.50. Details are at www.danceuk.org.

Clinical Excellence (NICE) is not implemented because trusts do not organise their finances well enough.

Better communication between clinical and financial staff will help the situation, says the commission in a report, which it published last week in conjunction with the institute.

Hospitals and primary care trusts could boost clinical excellence if they followed all of NICE's guidance, which is aimed at increasing levels of fairness and cost effectiveness in the NHS.

The Audit Commission surveyed and got responses from 71 NHS bodies in 10 strategic health authority areas. It also carried out in-depth interviews at 16 sites.

It found that 33% of respondents said NICE had issued guidance in 2002-3 that they were unable to fund, including guidance on drugs used to treat arthritis that reduce the effect of tumour necrosis factor α .

Adrian O'Dowd *London*

Managing the Financial Implications of NICE Guidance is available at www.audit-commission.gov.uk.

US study shows that folic acid fortification decreases neural tube defects

The incidence of neural tube defects in infants fell by about a third in some ethnic groups when the United States gradually increased enrichment of grain products with folic acid in the 1990s, a study has found (*Pediatrics* 2005;116:580-6).

The study looked at 21 population based surveillance systems that recorded trends in the prevalence of spina bifida and anencephaly between 1995 and 2002. The studies covered about 11 million births. There were 4468 cases of spina bifida and 2625 cases of anencephaly.

The prevalence of neural tube defects declined from 1995 and 1996, before fortification with folic acid was introduced, to 1998, a period when fortification was optional. The prevalence declined further from 1998 to 2002, when fortification became mandatory.

The decline in the prevalence of neural tube defects was significant in Hispanic and non-Hispanic white ethnic groups (respective declines of 26% and 29% for anencephaly, and 36% and 34% for spina bifida).

Janice Hopkins Tanne *New York*

Pathologist faces GMC hearing over altered autopsy report

A Home Office pathologist has admitted to the General Medical Council that he altered the autopsy report of a patient who died during surgery before the surgeon stood trial for her manslaughter.

Kenneth Shorrock's initial report appeared to exonerate the surgeon, Hurais Ramis Syed, of blame, but at the trial he submitted a second report in which that conclusion was missing. He is contesting the GMC's charge of serious professional misconduct.

Mr Syed, a locum consultant urologist at Dewsbury and Dis-

trict Hospital, West Yorkshire, operated on 78 year old Gladys Allen to remove a cancerous kidney in January 2000. Mrs Allen died after the operation, from blood loss and cardiac arrest.

Dr Shorrock, then a consultant pathologist at the Royal Halifax Hospital, wrote an autopsy report that described the operation as "necessary surgery" and in which he concluded: "In my opinion there is no definite evidence of any avoidable deficiency in the medical or surgical treatment that she received."

But in December 2000 Dr Shorrock produced a second report in which the word "necessary" had disappeared, along with the final conclusion.

Then in October 2001, the GMC heard, Dr Shorrock signed a witness statement in which he said it was "likely that death was contributed to by inadvertence, both before and during the operation."

The GMC has postponed its judgment.

Owen Dyer *London*

Scotland doesn't have enough high dependency beds

A lack of high dependency beds in Scottish hospitals is compromising the care of patients after surgery, says the first review of anaesthesia services in Scotland.

The report, produced by NHS Quality Improvement Scotland, found that the provision of adequately staffed and equipped high dependency beds in the country was inadequate. This lack is leading to the premature discharge of patients from high dependency units to general wards, where they may not get the same level of postoperative care.

The report was based on a comparison of the performance of services with agreed standards and is said to be the first ever national review of anaesthesia services undertaken anywhere in the United Kingdom. Overall, it concluded that standards of care for patients receiving anaesthesia are high and have improved over recent years.

It expressed concern, how-

ever, that high standards are not being achieved in all areas. In particular, it identified small and remote sites, accident and emergency units, electroconvulsive treatment suites, and maternity units as areas where standards of practice of anaesthesia could be improved.

Bryan Christie *Edinburgh*

Anaesthesia: Care Before, During and After Anaesthesia can be found at www.nhshealthquality.org.

European MPs want companies to develop drugs for children

European Union legislation designed to encourage pharmaceutical companies to develop drugs specifically tailored to children's needs has moved closer to the statute book after members of the European parliament (MEPs) voted to endorse it.

The measures, which are expected to be finally approved by EU governments before the end of the year, have also been welcomed by Françoise Grossetête, the French Christian Democrat member who steered the bill through the parliament, and by pharmaceutical associations.

The legislation provides for the creation of a paediatrics committee within the London based European Medicines Agency. Its role would be to determine which drugs are of potential use for children and to require manufacturers to carry out the necessary research to develop a formula specifically for children.

The legislation offers drug companies an incentive to produce drugs specifically for children by proposing to extend by six months the life of patents or supplementary protection certificates for products they develop.

This, it is estimated, could give each manufacturer extra profits of between €0.8m (£0.5m; \$1m) and €9m, offsetting the costs of clinical trials, which can be as high as €4m.

Rory Watson *Brussels*

Senior doctors admit mistakes in campaign for more open culture

Rebecca Coombes *London*

More than a dozen senior British medical figures have publicly owned up to past clinical errors as part of a campaign to encourage other doctors to report incidents and improve the safety of patients.

The doctors have contributed their stories to a handbook for junior doctors, published last week by the National Patient Safety Agency.

The agency is calling on all doctors to report mistakes so that the root causes can be addressed. Research has shown that doctors are less likely than other health professionals to report when things go wrong because they do not have time or do not feel they will be treated fairly.

Graeme Catto, president of the General Medical Council, is one of 14 senior doctors who gave a personal account of a medical error. As a general consultant physician he missed the

The agency's medical director, John Lilleyman, said steps had been taken to make the patients discussed in the anecdotes unidentifiable. "These incidents happened in the distant past, and it is unlikely any patient will recognise themselves. There is a small risk, but it was one we were prepared to take. The verisimilitude gives impact. We wanted to emphasise that if you do something wrong you should put your hand up."

The handbook, *Medical Error*, tells doctors that as well as reporting an incident locally—in which case the report automatically goes on to the National Patient Safety Agency—they now also have the option of reporting directly but anonymously to the agency (<http://npsa.nhs.uk/staffreports>).

Professor Lilleyman added that some doctors did not report an incident because of a lack of confidence that they will be dealt with fairly. "We understand that until doctors feel they are working in a more open culture we will need to have an anonymous reporting system."

Stephen Green, head of risk management at the Medical Defence Union (MDU), said: "Our members often tell us of the terrible distress they feel when they make a mistake. They are often surprised to learn that a significant proportion of incidents reported to the MDU can be traced back to lack of procedure or systems failing, rather than to lack of individual clinicians' knowledge. It is vital that we learn from incidents and near misses."

To improve patient safety the National Patient Safety Agency has focused on a number of changes to the systems that doctors work in, including asking NHS acute trusts to standardise the "crash call" number to 2222. □

Medical Error, which is supported by the BMA's Junior Doctors Committee, has been sent to all junior doctor members. It is also available at www.saferhealthcare.org.uk.

fact that a patient had a blood infection. The man developed meningitis and died.

Professor Catto said the mistake "will always live with me" and that he had learned the need to pay endless attention to detail.

Other contributors include James Johnson, the BMA chairman, Carol Black, president of the Royal College of Physicians, and the editors of the *BMJ* and the *Lancet*.