- Fiske, C. H. (1921). *J. biol. Chem.* **47**, 59.
 Folch, J. & Lees, M. (1951). *J. biol. Chem.* **191**, 807.
 Friedberg, F. & Greenberg, D. M. (1947). *J. biol. Chem.* **168**, 411.
 Friedberg, F., Tarver, H. & Greenberg, D. M. (1948). *J. biol. Chem.* **173**, 355.
 Gaitonde, M. K. & Richter, D. (1955). (In the Press.)
 Greenberg, D. M. & Winnick, T. (1948). *J. biol. Chem.* **173**, 199.
 Henriques, F. C., Kistiakowsky, G. B., Margnetti, C. & Schneider, W. G. (1946). *Industr. Engng Chem. (Anal.)* **18**, 349.
 May, R. L. (1948). *Rev. canad. Biol.* **7**, 642.
 Pinajian, J. J. & Cross, J. M. (1951). *Analyt. Chem.* **23**, 1056.
 Pirie, N. W. (1932). *Biochem. J.* **26**, 2041.
 Rutman, R., Dempster, E. & Tarver, H. (1949). *J. biol. Chem.* **177**, 491.
 Simpson, M. V. & Tarver, H. (1950). *Arch. Biochem.* **25**, 384.
 Tarver, H. & Morse, L. M. (1948). *J. biol. Chem.* **173**, 53.
 Tarver, H. & Schmidt, C. L. A. (1939). *J. biol. Chem.* **130**, 67.
 Williams, R. B. & Dawson, R. M. C. (1952). *Biochem. J.* **52**, 314.
 Young, L., Edson, M. & McCarter, J. A. (1949). *Biochem. J.* **44**, 179.

The Specific Polysaccharides of some Gram-negative Bacteria

By D. A. L. DAVIES

Microbiological Research Department, Porton, Wilts.

(Received 5 November 1954)

In their work on the specific capsular polysaccharides of the Pneumococci, Goebel and Avery and their collaborators (see Goebel & Avery, 1931; Goebel, 1939) showed the kind of chemical relationship which existed between some of the materials and which accounted for their serological cross-reactivity. A number of specific polysaccharides elaborated by organisms in the Gram-negative group have been examined chemically in spite of the difficulty in obtaining adequate amounts of starting material and of identifying sugars by isolation from complex mixtures in polysaccharide hydrolysates. The development of chromatographic methods of analysis has much simplified the task of identification and much smaller quantities of material are required.

From the specific degraded polysaccharide of the 'Smooth' form of *Shigella dysenteriae*, Morgan (1936, 1938) isolated derivatives of *N*-acetylglucosamine, galactose and rhamnose; Jesiatz & Goebel (1952) found that the *Shigella sonnei* specific lipopolysaccharide contained acetylglucosamine, galactose, glucose and a heptose; Goebel, Binkley & Perlman (1945) isolated specific polysaccharides from some strains of *Shigella flexneri* and demonstrated the presence of acetylglucosamine, glucose and rhamnose. These three *Shigella* species thus appeared to contain acetylglucosamine and the three possible combinations of two from the other three hexose sugars, galactose, glucose and rhamnose. Aldoheptose sugars have recently been found, not only in *Sh. sonnei* products but also in a specific polysaccharide material obtained from *Sh. flexneri* (Slein & Schnell, 1953) and in the specific protein-polysaccharide complex of *Sh. dysenteriae* (Davies & Morgan, 1954). None of these heptoses has yet been identified by isolation of the sugar or of a character-

istic derivative. Although some family relationship is revealed by the qualitative composition of the *Shigella* polysaccharides, a group of materials which do not generally cross-react serologically (Weil, 1947), the situation has been somewhat complicated by the discovery of a 'phage-resistant mutant of *Sh. sonnei* whose specific lipopolysaccharide appears to contain one unidentified hexosamine and no other sugar residue (Goebel & Jesiatz, 1952).

In the *Salmonella* group, galactose, glucose and mannose were identified as components of the specific degraded polysaccharides of *Salm. typhosum*, (Freeman, 1942) and *Salm. typhimurium* (Freeman, 1943). These three sugars were also found in the specific polysaccharide of *Salm. abortus equi* by Lüderitz & Westphal (1952a) with the addition of hexosamine, rhamnose and abequose.

In two strains of *Escherichia coli* examined by Lüderitz & Westphal (1952b), and belonging to different 'O' groups, acetylhexosamine, galactose, glucose and rhamnose were present in the polysaccharides; one material also contained xylose whereas the other contained mannose.

Single strains of organisms from other taxonomic groups have yielded polysaccharides composed of yet other groups of sugars, e.g. galactose and mannose in *Haemophilus pertussis* (Akiya, Takahashi, Kuriyama & Ogawa, 1951), hexosamine, glucose and methylpentose in *Serratia marcescens* (Hartwell, Shear, Adams & Perrault, 1943), acetylhexosamine, galactose and mannose in *Proteus vulgaris* (Bendich & Chargaff, 1946), glucose and glucuronic acid in *Klebsiella pneumoniae* (Goebel & Avery, 1927), glucose, galactose, arabinose and glucuronic acid in *Vibrio* spp. (Linton, 1940) and glucosamine, glucose, xylose and rhamnose in *Trichomonas foetus* (Feinberg & Morgan, 1952).

These data suggest, as would be suspected on serological grounds, that different groups of sugars may be characteristic of different taxonomic groups of bacteria. To what extent the specific polysaccharides differ in structure in serologically closely related bacteria is, however, not known, except in the case of *Pneumococcus* Types III and VIII (Goebel, 1935).

In this communication some data on the composition of the specific polysaccharides of a number of Gram-negative bacteria are recorded. The materials were isolated, not primarily for this purpose, by extraction of acetone-dried cells with trichloroacetic acid (Boivin, Mesrobianu & Mesrobianu, 1933). This method was chosen for several reasons; material having the specificity of the 'O' somatic antigen can be extracted in this way from the 'Smooth' forms of all Gram-negative bacteria which have been tested (Boivin, Mesrobianu, Mesrobianu & Nestorescu, 1934; Boivin & Mesrobianu, 1937), except *Pasteurella pestis* (Girard, 1941). *P. pestis* does, however, contain a somatic antigen which will be described in a subsequent communication. Some milder methods of extraction which are more effective in certain cases are more restricted in their applicability, e.g. diethylene glycol is a very good solvent for the recovery of the somatic antigenic material of *Sh. dysenteriae* (Morgan, 1937; Morgan & Partridge, 1940, 1941) but it fails to extract the analogous material from some strains of *Sh. flexneri* (Goebel *et al.* 1945). Although the yield by the trichloroacetic acid method is usually less than that obtained by other methods when these are applicable, its use does avoid the necessity of searching for a suitable solvent for each organism investigated. The product of trichloroacetic acid extraction is highly antigenic whereas by some other methods a specific but non-antigenic material is obtained. The warm 45 % phenol method of Westphal, Lüderitz & Bister (1952), which appears to be of general applicability, yields a specific lipopolysaccharide which is pyrogenic and toxic (Westphal, Lüderitz, Eichenberger & Keiderling, 1952), such a material can be a Forssman heterophile antigen without inducing the formation of specific agglutinins or precipitins (Davies & Morgan, 1954). Trichloroacetic acid extracts are not grossly impure and after fractionation with ethanol are composed principally of two components; one is the protein-polysaccharide-phospholipid antigen, the other is free specific undegraded polysaccharide hapten. The antigenic material can be readily separated from the polysaccharide hapten and other relatively small molecular species by sedimentation in a preparative ultracentrifuge at 100 000 *g* (Davies & Morgan, 1953).

Antigens prepared in this way have been hydrolysed with dilute acetic acid to liberate the degraded

polysaccharide haptens which were further hydrolysed and examined chromatographically. Data are recorded for eighteen different *Salmonellas* and several strains of *Shigella* and *Chromobacterium violaceum*.

MATERIALS AND METHODS

Organisms investigated. The strains of bacteria used in this investigation are detailed in Table 1.

Materials for analysis. Samples of extraction products for analysis were thoroughly dialysed against distilled water at 0–2° and dried from the frozen state. They were further dried to constant weight *in vacuo* at 78° before making into solutions of known concentration in water.

Total nitrogen. The Kjeldahl method was used with the distillation apparatus of Markham (1942) and the mixed bromocresol green-methyl red indicator of Ma & Zuazaga (1942).

Phosphorus. Determinations were made on samples containing 5–20 µg. of P by a modification of the method of Fiske & Subbarow (1925).

Hexosamine. This was estimated by a modification of the colorimetric method of Elson & Morgan (1933) on samples which had been hydrolysed, except where otherwise stated, with 0.5 N-HCl for 8 hr. at 100°.

Ultracentrifuging. High-speed centrifuging was carried out in a 'Spinco' model 'L' preparative ultracentrifuge at 2–4°. Figures given for the force in *g* refer to the force exerted at the centres of the tubes and represent average values.

Chromatography. Samples of polysaccharides were hydrolysed in sealed ampoules at 100° with 0.5 N-H₂SO₄ for

Table 1. *Strains of bacteria used for the extraction of specific polysaccharides*

Organism	Strain
<i>Salmonella paratyphosum</i> A	H.A.6.
<i>Salm. typhimurium</i>	M.R.D.
<i>Salm. stanley</i>	5716
<i>Salm. salinatis</i>	6247
<i>Salm. budapest</i>	5724
<i>Salm. choleraesuis</i>	N.C.T.C. 5736
<i>Salm. montevideo</i>	5747
<i>Salm. newport</i>	5751
<i>Salm. kottbus</i>	5753
<i>Salm. typhosa</i>	'O' 901
<i>Salm. dar-es-salaam</i>	5773
<i>Salm. gallinarum</i>	5775
<i>Salm. london</i>	5777
<i>Salm. senftenberg</i>	N.C.T.C. 3158
<i>Salm. meleagridis</i>	6023
<i>Salm. nitrosea</i>	5789
<i>Salm. grumpensis</i>	6533
<i>Shigella dysenteriae</i>	K 624 'S'
<i>Sh. schmitzii</i>	N.C.T.C. 8212
<i>Sh. dysenteriae</i>	N.C.T.C. 7921
<i>Sh. alcalescens</i>	Brig. Archer, R.A.M.C.
	'Lewitus'
	N.C.T.C. 7917
	'Birch'
	'Brown'
	'Mentekab'
	'Frazer's Hill'
	'Amethystinum'
	'Metropolitan Water Board'
<i>Chromobacterium violaceum</i>	

8 hr., except where otherwise stated, and neutralized with $\text{Ba}(\text{OH})_2$. The neutral salt-free hydrolysates were evaporated to dryness *in vacuo* and redissolved at suitable concentrations for application to paper chromatograms. Whatman no. 1 paper was used with a variety of solvent systems (phenol, collidine-lutidine-water, ethyl acetate-pyridine-water and butanol-pyridine-water). Papers were sprayed, after drying, with 3% anisidine hydrochloride in water-saturated butanol (Hough, Jones & Wadman, 1950) which distinguishes by colour the different classes of sugars, and with ninhydrin or AgNO_3 to reveal hexosamines.

EXPERIMENTAL

Preparation of organisms. Cultures were checked for smoothness of colonial morphology and salt-stability. Strains of *Salmonella* and *Shigella* were grown for 22 hr. at 37° on CCY agar (Gladstone & Fildes, 1940) in enamelled steel trays each containing 500 ml. of medium, or in 'Thompson' bottles containing 250 ml. of CCY broth. Strains of *Chromobacterium violaceum* were grown on tryptic meat digest agar in Roux bottles. The growths were washed off agar surfaces or separated by centrifuging in the cold from broth cultures and washed in the cold with saline and with water; the sedimented cells were resuspended in cold distilled water and poured into 10 vol. of acetone at -10°. The acetone was changed several times and the cells finally dried *in vacuo* over H_2SO_4 and stored at 0° over P_2O_5 until required.

Extraction with trichloroacetic acid. Dry bacteria were suspended in distilled water at 0-2° at 2% (w/v) concentration and an equal volume of cold *n* trichloroacetic acid was added. The mixture was stirred in the cold for 2-3 hr. and centrifuged at 12 000 *g* to separate the cells. The supernatant solution was dialysed in the cold until largely free from acid and concentrated under reduced pressure (at 16°) to a concentration of about 1% (w/v) of total solid material. The solution was then fractionated with cold ethanol.

When an extract of *Sh. dysenteriae* was made in this way (Davies, 1953), the material which was precipitated from solution between 50 and 84% (v/v) ethanol concentration was found on examination in the analytical ultracentrifuge to be composed of two substances. One was a rapidly sedimenting polydisperse component and the other showed a relatively slow-moving sharp peak. These materials had the same serological specificity but the heavy component was an antigenic protein-polysaccharide-phospholipid complex, whereas the component of smaller molecular size was a non-antigenic undegraded specific polysaccharide hapten. These materials could be separated by fractional high-speed centrifuging, the polysaccharide remaining in solution when the antigen was sedimented at 100 000 *g*; repeated centrifuging was necessary to free each material from the other.

Trichloroacetic acid extracts of the other organisms examined behaved in much the same way as that of *Sh. dysenteriae*. The amount of ethanol required to precipitate the antigenic material varied in different cases and the more soluble preparations which required larger amounts of ethanol for precipitation were contaminated with a polyglucose which remained in admixture with the specific polysaccharide in the 100 000 *g* supernatant fluid after ultracentrifugation.

One batch of cells of *Salm. poona* was extracted with warm 45% phenol according to the method of Westphal, Lüderitz & Bister (1952). Lipopolysaccharide was obtained by

ethanol precipitation from the aqueous phase which separated on cooling the suspension to 2°. The yield by this method was substantially higher than that obtained by trichloroacetic acid extraction but the product was contaminated with nucleic acid, the last traces of which were only removed with difficulty.

Preparation of the specific degraded polysaccharide haptens. The purified antigens were dissolved at 1% (w/v) concentration in 1% (v/v) acetic acid and heated for 4 hr. at 100° in an atmosphere of nitrogen. The hydrolysis products were extracted with ether to remove phospholipid and the protein component was separated by centrifuging, washed with 1% acetic acid and freeze-dried. The acetic acid supernatant solution and washings of the sedimented protein were combined and reduced to about a third of their volume by evaporation under reduced pressure at 16°, and the polysaccharide was then precipitated by the addition of several volumes of ethanol; precipitation with ethanol was repeated at least once and the material finally redissolved in water, centrifuged at 20 000 *g* for 30 min. to remove traces of insoluble material, dialysed and freeze-dried.

Physical examination of the polysaccharide haptens. It has not been possible to examine all of the specific polysaccharide preparations for homogeneity. Three typical specimens were examined in the analytical ultracentrifuge (Spinco model 'E'); *Salm. kottbus* and *Salm. niloese* polysaccharides were run at pH 4.1 (1% (w/v) concentration in acetate buffer, *I* = 0.1, containing 0.1 M-NaCl) and *Salm. salinatis* material at pH 8 (0.44% (w/v) concentration in phosphate buffer, *I* = 0.2). The three preparations all behaved in the same way and only a single sedimenting peak was revealed in each case; a typical photograph is shown in Fig. 1a.

Fig. 1. Ultracentrifugal and electrophoretic examination of some *Salmonella* polysaccharide preparations. (a) *Salm. niloese* sedimentation pattern, concentration 1% (w/v) in acetate buffer pH 4.1, *I* = 0.2, force 260 000 *g*, exposure at 2.5 hr. (b), (c) and (d) electrophoresis diagrams showing the ascending limb of enantiographic pictures; (b) *Salm. montevideo* concentration 0.5% (w/v) in acetate buffer pH 4, *I* = 0.2, exposure at 8 hr., potential gradient 1.06 v/cm. (c) *Salm. budapest* concentration 0.45% (w/v) in acetate buffer pH 4, *I* = 0.1, exposure at 17.5 hr., potential gradient 2.83 v/cm. (d) *Salm. salinatis* concentration 0.44% (w/v) in phosphate buffer pH 8, *I* = 0.2, exposure at 13 hr., potential gradient 2.2 v/cm. Direction of movement is shown by arrows.

Examination in the Tiselius electrophoresis apparatus showed, however, that more than one component was present in some cases. *Salm. montevideo* polysaccharide showed one component after running for 8 hr. at pH 4 in acetate buffer ($I=0.2$), and the ascending and descending peaks were enantiographic; a photograph of the ascending limb is shown in Fig. 1b. *Salm. budapest* polysaccharide preparation showed two peaks both at pH 4 and at pH 8 (acetate and phosphate buffers, $I=0.1$ and 0.2 , respectively). The patterns were enantiographic and a photograph of the ascending limb is shown in Fig. 1c. *Salm. salinatis* polysaccharide showed at least three components after running for 13 hr. at pH 8 (phosphate buffer, $I=0.2$); a picture of the ascending limb is shown in Fig. 1d.

The significance of the electrophoretic results is not known, and further work will be necessary in order to find what the several components represent.

RESULTS

Different numbers of trays or bottles were used when growing the various organisms but some strains gave much better growth than others; *Salm. gallinarum* always gave very small growths.

Antigens. Details of the antigenic protein-polysaccharide-phospholipid complexes are shown in Table 2; the yields were very variable the lowest being from *Salm. stanley* (0.25%) and the highest from *Salm. newport* (6.7%). The N contents varied between 1.5 and 4.8% and the P values were between 1.0 and 5.6% among the *Salmonella* products; these variations were related to differences in the proportions of the components in the complexes, the high N figures being found in preparations having a high protein content except where substantial amounts of hexosamine were present in the polysaccharide components as, for example, in *Salm. poona* and *Sh. alkalescens*.

The analytical figures found for the *Shigella* antigens were not very dissimilar from those previously recorded for the *Sh. dysenteriae* antigenic complex (Davies, Morgan & Mosimann, 1954). The *Chr. violaceum* strains yielded antigens with low P contents.

Polysaccharides. The conditions of hydrolysis employed to separate the degraded polysaccharide hapten from the antigenic materials were those which had been found most suitable for the recovery of the degraded polysaccharide of *Sh. dysenteriae*. The conditions are probably not ideal in the case of some of the other materials examined. The N content of some of the polysaccharides is considerably higher than can be accounted for by their hexosamine contents; this was due to the presence of amino acids, which did not, however, interfere with the chromatographic examination of the materials. The yields and N contents of the degraded polysaccharides are shown in Table 3; the P contents were very low in most cases and all were below 1%.

The sugars composing the specific polysaccharides of the *Salmonellas* are shown in Table 4. Traces of hexosamine were present in most of the samples but significant amounts were found only in *Salm. paratyphosum A* (7%), *Salm. montevideo* (13%), *Salm. poona* (16%) and *Salm. grumpensis* (16%). All of the *Salmonella* materials contained galactose, glucose and mannose except for those of *Salm. poona*, *Salm. grumpensis* and *Salm. choleraesuis* which contained no mannose. Other sugars present in some of the preparations were arabinose, xylose, fucose, rhamnose, abequose, tyvelose and one unidentified sugar.

Abequose and tyvelose have recently been described by Westphal, Lüderitz, Fromme & Joseph (1953) and appear to be bisdeoxyhexoses.

Table 2. The 'O' antigenic complexes of some Gram-negative bacteria

Organism	Extract (mg.)	Yield (%)	N (%)	P (%)
<i>Salmonella paratyphosum A</i>	376	0.3	4.2	1.1
<i>Salm. typhimurium</i>	720	1.7	4.6	2.3
<i>Salm. stanley</i>	104	0.25	2.9	5.6
<i>Salm. salinatis</i>	1160	2.0	2.14	2.8
<i>Salm. budapest</i>	443	1.7	2.4	1.9
<i>Salm. choleraesuis</i>	33	2.5	—	—
<i>Salm. montevideo</i>	2800	2.8	1.5	1.3
<i>Salm. newport</i>	6680	6.68	2.4	1.25
<i>Salm. kottbus</i>	712	2.8	2.2	1.85
<i>Salm. typhosa</i>	450	1.3	2.6	1.7
<i>Salm. dar-es-salaam</i>	294	1.1	1.84	2.1
<i>Salm. gallinarum</i>	39	1.0	—	3.6
<i>Salm. meleagridis</i>	115	0.8	—	—
<i>Salm. senftenberg</i>	1070	0.85	3.7	1.0
<i>Salm. niloese</i>	864	1.8	1.64	1.8
<i>Salm. grumpensis</i>	157	0.43	2.3	4.2
<i>Salm. poona</i>	1200	1.5	4.8	1.8
<i>Shigella schmitzii</i>	5880	6.3	4.7	1.6
<i>Sh. alkalescens</i>	320	1.0	5.2	1.2
<i>Sh. dispar</i>	980	2.1	4.9	1.8
<i>Chromobacterium violaceum</i> 'Brown'	22	1.0	5.8	0.03
'Birch'	111	2.7	6.1	0.6

The unidentified sugar was found in *Salm. paratyphosum A* and *Salm. choleraesuis*; like the bis-deoxysugars this travelled very rapidly on paper chromatograms. In butanol-pyridine-water as solvent, R_{R} values (representing rate of movement compared with that of rhamnose) found were: abequose 1.18, tyvelose 1.30 and the unidentified component 1.44; this latter substance gave a rela-

tively small spot which was only clearly visible when a larger amount of polysaccharide hydrolysate was applied to the paper than was used in the runs shown in Figs. 2 and 3; when spots were eluted from paper chromatograms and examined by the sulphuric acid-cysteine reaction (Dische & Shettles, 1948; Dische, Shettles & Osnos, 1949), the same absorption spectrum was obtained as with tyvelose and abequose.

Table 3. *The degraded polysaccharide haptens of some Gram-negative bacteria*

Organism	Polysaccharide recovered (mg.)	Yield (% wt. of antigen)	N (%)
<i>Salmonella paratyphosum A</i>	193	22	3.16
<i>Salm. typhimurium</i>	122	31	3.4
<i>Salm. stanley</i>	6.3	16	—
<i>Salm. salinatis</i>	167	34	1.03
<i>Salm. budapest</i>	183	42	1.2
<i>Salm. choleraesuis</i>	4.0	20	—
<i>Salm. montevideo</i>	491	49	1.3
<i>Salm. newport</i>	1270	43	1.0
<i>Salm. kottbus</i>	159	32	1.03
<i>Salm. typhosa</i>	131	44	—
<i>Salm. dar-es-salaam</i>	62	27	0.7
<i>Salm. gallinarum</i>	8.1	32	—
<i>Salm. london</i>	2160	—	0.25
<i>Salm. meleagridis</i>	17	28	—
<i>Salm. senftenberg</i>	103	42	2.3
<i>Salm. niloese</i>	406	47	0.5
<i>Salm. grumpensis</i>	34	35	1.96
<i>Salm. poona</i>	213	42	3.5
<i>Chromobacterium violaceum</i> ('Brown')	1.37	55	3.8

'O' Group:

Fig. 2. Paper chromatogram showing the sugar components of some *Salmonella* specific polysaccharides ('O' groups A, B and C). Run with butanol-pyridine-water and sprayed with anisidine. 1, *Paratyphosum A*; 2, *typhimurium*; 3, *budapest*; 4, *salinatis*; 5, *stanley*; 6, *montevideo*; 7, *choleraesuis*; 8, *kottbus*; 9, *newport*.

'O' Group:

Fig. 3. Paper chromatogram showing the sugar components of some *Salmonella* specific polysaccharides ('O' groups D, E and G). Run with butanol-pyridine-water and sprayed with anisidine. 1, *Typhosa*; 2, *gallinarum*; 3, *dar-es-salaam*; 4, *meleagridis*; 5, *london*; 6, *niloese*; 7, *senftenberg*; 8, *poona*; 9, *grumpensis*.

Table 4. The sugar components of some specific polysaccharides of the *Salmonellas*

Organism	Somatic antigens	'O' group	Hexosamines	Hexoses	Pentoses	Methyl-pentoses	Others
<i>Salmonella paratyphosum A</i>	I, II, XII	A	Glucosamine	Galactose Glucose Mannose	Xylose	Rhamnose	Tyvelose, one unidentified
<i>Salm. typhimurium</i>	I, IV, V, XII	B	—	Galactose Glucose Mannose	—	Rhamnose	Abeyose
<i>Salm. budapest</i>	I, IV, XII	B	—	Galactose Glucose Mannose	—	Rhamnose	Abeyose
<i>Salm. salmatis</i>	IV, XII	B	—	Galactose Glucose Mannose	—	Rhamnose	Abeyose
<i>Salm. stanley</i>	IV, V, XII	B	—	Galactose Glucose Mannose	Xylose	Rhamnose	Abeyose
<i>Salm. montevideo</i>	VI, VII	C ₁	Glucosamine	Galactose Glucose Mannose	—	—	—
<i>Salm. choleraesuis</i>	VI, VII	C ₁	—	Galactose Glucose	Xylose Arabinose	—	One unidentified
<i>Salm. kotbus</i>	VI, VIII	C ₂	—	Galactose Glucose Mannose	—	Rhamnose	Abeyose
<i>Salm. newport</i>	VI, VIII	C ₂	—	Galactose Glucose Mannose	—	Rhamnose	Abeyose
<i>Salm. typhosa</i>	IX, XII	D	—	Galactose Glucose Mannose	—	Rhamnose	Tyvelose
<i>Salm. gallinarum</i>	I, IX, XII	D	—	Galactose Glucose Mannose	—	Rhamnose	Tyvelose
<i>Salm. dar-es-salaam</i>	I, IX, XII	D	—	Galactose Glucose Mannose	—	Rhamnose	Tyvelose
<i>Salm. melagris</i>	III, X, XXVI	E ₁	—	Galactose Glucose Mannose	—	Rhamnose	—
<i>Salm. london</i>	III, X, XXVI	E ₁	—	Galactose Glucose Mannose	—	Rhamnose	—
<i>Salm. nitrose</i>	I, III, XIX	E ₂	—	Galactose Glucose Mannose	—	Rhamnose	—
<i>Salm. senftenberg</i>	I, III, XIX	E ₂	—	Galactose Glucose Mannose	Xylose	Rhamnose	—
<i>Salm. poona</i>	XIII, XXII	G	Glucosamine	Galactose Glucose	—	Fucose	—
<i>Salm. grumpensis</i>	XIII, XXII	G	Chondrosamine	Galactose Glucose	—	Fucose	—

Table 5. The sugar components of some specific bacterial polysaccharides

Organisms	Hexosamines	Hexoses	Pentoses	Methylpentose	Others
<i>Shigella schmitzi</i>	Glucosamine	Galactose Glucose	—	Rhamnose	—
<i>Sh. alkalescens</i>	Glucosamine	Galactose Glucose	—	—	—
<i>Sh. dysenteriae</i>	Glucosamine	Galactose Glucose	—	Rhamnose	—
<i>Chromobacterium violaceum</i>	—	Galactose Glucose	Xylose Arabinose	—	Galacturonic acid
'Lewittus'	Hexosamine	Glucose	—	Fucose	Manno-gala-heptose
'Brown'	Hexosamine	Glucose	—	Rhamnose	Manno-gala-heptose
'Birch'	Hexosamine	Glucose	—	Rhamnose	Heptose
N.C.T.C. 7917	Hexosamine	Galactose Glucose	—	—	—
'Mentekab'	—	Galactose Glucose	—	Rhamnose	Fucose
'Frazer's Hill'	Hexosamine	—	—	Rhamnose	—
'Amethystinum'	—	Glucose	—	—	—
'Metropolitan Water Board'	—	Galactose Glucose	Xylose	Rhamnose	—

The relative amounts of the various sugars present can be roughly judged by examination of the photographs shown in Figs. 2 and 3, when it is borne in mind that deoxysugars react somewhat less strongly with the anisidine spray reagent than the hexoses and that the pentoses react more strongly than hexoses. Although mannose and arabinose cannot be distinguished in the photograph (Fig. 2), they show on the papers as brown and pink spots respectively; the *Salm. choleraesuis* polysaccharide spot in the mannose position is actually due to arabinose. Similarly, xylose and fucose cannot be readily distinguished in Fig. 3, although they appeared on the paper as pink and yellow-green spots, respectively; *Salm. senftenberg* contains xylose and '0' group *G* organisms fucose (overlapping a trace of ribose in *Salm. poona* due to nucleic acid contamination). Hexosamines are not revealed by anisidine under the conditions used in Figs. 2 and 3; they give a yellow colour on longer heating with the spray reagent but this treatment impairs the characteristic colours by which the other groups of sugars are distinguished.

Details of some *Shigella* polysaccharides are shown in Table 5; *Sh. schmitzii* and *Sh. dispar* specific polysaccharides contained all of the four hexoses previously found in *Shigella* species (glucosamine, galactose, glucose and rhamnose), while that of *Sh. alkalescens* contained, like *Sh. sonnei*, no rhamnose. *Sh. alkalescens* differed from the other *Shigella* species in that both glucosamine and chondrosamine were present and in approximately equal amounts; the total hexosamine content amounted to 35 % of the weight of the material. The usual conditions of hydrolysis which had been employed to separate the sugars from the various polysaccharides (0.5N acid for 8 hr.), failed to give complete hydrolysis of the *Sh. dispar* and *Sh. alkalescens* preparations, which required treatment with N acid for 12 hr. to produce discrete spots on paper chromatograms. These polysaccharides all differ from those of most of the *Salmonella* group in not containing mannose, but also differ from those *Salmonellas* containing no mannose in that *Salm. choleraesuis* contains two pentoses and the group *G* organisms two hexosamines and fucose.

The *Chr. violaceum* strains yielded polysaccharides which appeared to have little in common with one another except for 'Brown' and 'Birch' which were qualitatively identical and cross-reacted strongly serologically. Two aldoheptose sugars were found (Table 5), one in strains 'Brown' and 'Birch' present in substantial amounts, and another in strain N.C.T.C. 7917 in smaller, but still considerable amount. These sugars have not yet been isolated but were identified by their characteristic colour on paper chromatograms sprayed with anisidine and, after elution from paper, by their

absorption spectrum given after treatment with sulphuric acid and cysteine (Dische, 1953) and other reactions described by this author. The 'Brown' and 'Birch' heptose travelled on paper chromatograms in the position of D-manno-D-gala-heptose and differed from five other aldoheptoses examined. In the N.C.T.C. 7917 material the heptose travelled at the same rate as D-gala-D-manno-heptose. The only uronic acid found was galacturonic acid in *Chr. violaceum* strain 'Lewitus'.

DISCUSSION

The yield of '0' specific material obtained by trichloroacetic acid extraction differed considerably in different strains, due possibly to differences of 'Smoothness' and to the fact that since strains had varying growth rates organisms were unavoidably reaped at different stages of growth. Differences in the relative amount of antigenic complex and free specific undegraded polysaccharide were also found in the crude extracts and analysis of the antigenic fractions showed that the relative amounts of protein and polysaccharide components in the complexes varied.

The extent of the electrophoretic examination of the degraded polysaccharides is insufficient for the results to be interpreted at present. The several components detected by electrophoresis failed to separate on ultracentrifugation. Molecules of the same order of size as these degraded polysaccharide haptens had been removed from the parent antigenic preparations by high-speed centrifuging and the components must all have arisen by mild acetic acid hydrolysis of the antigens.

Mild acetic acid hydrolysis of the somatic antigenic complex of *Sh. dysenteriae* was considered to release polysaccharide by the splitting of a relatively labile link between acetylglucosamine and other sugar residues (Davies & Morgan, 1954). As such a labile link may not be present in the polysaccharides examined this might account for the incomplete separation of polysaccharide from antigenic complex found in some cases, the N content of some of the polysaccharides being 1-2 % higher than could be accounted for by their hexosamine content.

Among the *Salmonellas*, all strains contained galactose and glucose and, except in '0' group *G* (Kauffmann, 1951) and *Salm. choleraesuis*, mannose was present. These three sugars have not been found together in any of the other polysaccharides extracted but were present in one *Esch. coli* strain recorded by Lüderitz & Westphal (1952b). Those *Salmonellas* not containing mannose differ in some other respect from any other polysaccharide examined. One strain from '0' group 'A' contained eight different sugar residues. A group 'B' strain,

Salm. abortus equi, was examined by Westphal, Lüderitz, Eichenberger & Keiderling (1952) and its specific polysaccharide was found by these workers to contain five hexoses. The four group 'B' strains detailed in Fig. 2 contained the same sugar residues as the *Salm. abortus equi* preparation but *Salm. stanley* also contained a small amount of xylose. Two organisms in group 'C' did not match, *Salm. montevidео* having galactose, glucose, mannose and hexosamine while *Salm. choleraesuis* contained galactose, glucose, arabinose, xylose and only traces of hexosamine. Pairs of organisms in groups 'C₂' and 'E₁' and three in group 'D' corresponded well within their groups. In group 'E₃' *Salm. senftenberg* differed from its partner *Salm. niloese* in containing a small amount of xylose. Two group 'G' organisms corresponded well and both contained glucosamine and chondrosamine; the latter sugar has not been recorded from any other bacterial source (Davies, 1954), but is present in larger amount in the *Sh. alkalescens* polysaccharide.

Either tyvelose or abequose (Westphal *et al.* 1953) was present in ten of the eighteen strains examined. In *Salm. paratyphosum A* and *Salm. choleraesuis* a spot with a very high *R_F* value was found on chromatograms and appeared to be due to a similar deoxy sugar.

The distribution of xylose is scattered; significant amounts were present in the products of four *Salmonellas* all of which belonged to different '0' groups. A faint spot on chromatograms in the xylose position was found in several other polysaccharide preparations but these probably represented very small amounts as the anisidine spray reagent is very sensitive for this sugar.

In the *Shigella* group the polysaccharides examined and those recorded previously in the literature contain different hexoses from among a group of only four; glucosamine, galactose, glucose and rhamnose. The heptose sugar found by Slein & Schnell (1953) was phosphorylated and the heptose found by Jesiatz & Goebel (1952) was present in a material containing phosphorus. In *Shigella dysenteriae* heptose is present in the antigenic complex and the toxic phospholipopolysaccharide but not in the degraded polysaccharide which is free of phosphorus. The heptose in *Sh. dysenteriae* is present in relatively small amount, unlike those now found in several bacterial polysaccharides, including three strains of *Chr. violaceum*. In these cases the polysaccharides are free of phosphorus and the heptose sugar is a major component. *Chr. violaceum* strains appear to be a serologically heterogeneous group, and this is borne out by the finding that their specific polysaccharides had little in common with one another. One exception, however, was the case of strains 'Brown' and 'Birch' isolated from different sources, but serologically related and con-

taining the same sugar residues. Another strain, 'Frazer's Hill', contained a polysaccharide composed of hexosamine, glucose and rhamnose, the same hexoses as those found in the *Sh. flexneri* specific polysaccharide.

SUMMARY

1. Materials having the properties of the '0' somatic antigens have been extracted with trichloroacetic acid from acetone-dried cells of a number of strains of *Salmonella*, *Shigella* and *Chromobacterium violaceum*.

2. The antigenic preparations, after purification by ethanol precipitation and high-speed centrifuging, were hydrolysed with dilute acetic acid and the specific degraded polysaccharide haptens recovered from the solutions by ethanol precipitation after removal of the insoluble protein and lipid components.

3. The sugars present in the polysaccharides were identified by chromatographic methods.

4. In the *Salmonella* group the number of different sugars in the polysaccharides varied from four (*Salm. montevidео*) to eight (*Salm. paratyphosum A*). Galactose and glucose were present in all strains examined; mannose was found in fifteen of eighteen strains. Polysaccharides recovered from organisms in the same '0' group were, with a few exceptions, qualitatively the same but quantitative differences among the component sugars were apparent in many cases.

5. Organisms in the *Shigella* group all contained hexosamine and in addition, either two or three of the hexose sugars galactose, glucose and rhamnose.

6. *Chromobacterium violaceum* polysaccharides showed little in common with one another with the exception of one pair of strains. Two heptose sugars were present, one of which appeared to be D-manno-D-gala-heptose.

7. Chondrosamine was found in *Salmonellas* of '0' group G (*Salm. poona* and *Salm. grumpensis*) and *Sh. alkalescens*. An unidentified sugar was encountered in two *Salmonella* strains, this may be a bisdeoxy sugar similar to tyvelose and abequose.

I wish to thank Brig. F. E. Buckland, R.A.M.C., for providing strains of *Chromobacterium violaceum*. Ultracentrifugal and electrophoretic examinations were kindly carried out by Dr B. R. Record and Mr K. A. Cammack. This work is published by permission of the Chief Scientist, Ministry of Supply.

REFERENCES

- Akiya, S., Takahashi, H., Kuriyama, M. & Ogawa, N. (1951). *Jap. med. J.* **4**, 325.
- Bendich, A. & Chargaff, E. (1946). *J. biol. Chem.* **166**, 283.
- Boivin, A. & Mesrobian, L. (1937). *C.R. Soc. Biol., Paris*, **124**, 1176.

- Boivin, A., Mesrobianu, I. & Mesrobianu, L. (1933). *C.R. Soc. Biol., Paris*, **114**, 307.
- Boivin, A., Mesrobianu, I., Mesrobianu, L. & Nestorescu, B. (1934). *C.R. Soc. Biol., Paris*, **115**, 306.
- Davies, D. A. L. (1953). Thesis: University of London.
- Davies, D. A. L. (1954). *J. gen. Microbiol.* **10**, ii.
- Davies, D. A. L. & Morgan, W. T. J. (1953). *6th Int. Congr. Microbiol. Rome, Abstr.* vol. I, p. 457.
- Davies, D. A. L. & Morgan, W. T. J. (1954). *Biochem. J.* (In the Press.)
- Davies, D. A. L., Morgan, W. T. J. & Mosimann, W. (1954). *Biochem. J.* **56**, 572.
- Dische, Z. (1953). *J. biol. Chem.* **204**, 983.
- Dische, Z. & Shettles, L. B. (1948). *J. biol. Chem.* **175**, 595.
- Dische, Z., Shettles, L. B. & Osnos, M. (1949). *Arch. Biochem.* **22**, 169.
- Elson, L. A. & Morgan, W. T. J. (1933). *Biochem. J.* **27**, 1824.
- Feinberg, J. G. & Morgan, W. T. J. (1952). *2nd Int. Congr. Biochem. Paris, Abstr.* p. 388.
- Fiske, C. H. & Subbarow, Y. (1925). *J. biol. Chem.* **66**, 375.
- Freeman, G. G. (1942). *Biochem. J.* **36**, 340.
- Freeman, G. G. (1943). *Biochem. J.* **37**, 601.
- Girard, G. (1941). *C.R. Soc. Biol., Paris*, **135**, 1577.
- Gladstone, G. P. & Fildes, P. (1940). *Brit. J. exp. Path.* **21**, 161.
- Goebel, W. F. (1935). *J. biol. Chem.* **110**, 391.
- Goebel, W. F. (1939). *J. exp. Med.* **69**, 353.
- Goebel, W. F. & Avery, O. T. (1927). *J. exp. Med.* **46**, 601.
- Goebel, W. F. & Avery, O. T. (1931). *J. exp. Med.* **54**, 431.
- Goebel, W. F., Binkley, F. & Perlman, E. (1945). *J. exp. Med.* **81**, 315.
- Goebel, W. F. & Jesiatitis, M. A. (1952). *J. exp. Med.* **96**, 425.
- Hartwell, J. L., Shear, M. J., Adams, J. R. & Perrault, A. (1943). *J. nat. Cancer Inst.* **4**, 107.
- Hough, L., Jones, J. K. N. & Wadman, W. H. (1950). *J. chem. Soc.* p. 1702.
- Jesiatitis, M. A. & Goebel, W. F. (1952). *J. exp. Med.* **96**, 409.
- Kauffmann, F. (1951). *Enterobacteriaceae*. Copenhagen: Ejnar Munksgaard.
- Linton, R. W. (1940). *Bact. Rev.* **4**, 261.
- Lüderitz, O. & Westphal, O. (1952a). *Z. Naturf.* **7b**, 136.
- Lüderitz, O. & Westphal, O. (1952b). *Z. Naturf.* **7b**, 548.
- Ma, T. S. & Zuazaga, G. (1942). *Industr. Engng Chem. (Anal.)*, **14**, 280.
- Markham, R. (1942). *Biochem. J.* **36**, 790.
- Morgan, W. T. J. (1936). *Biochem. J.* **30**, 909.
- Morgan, W. T. J. (1937). *Biochem. J.* **31**, 3003.
- Morgan, W. T. J. (1938). *Helv. chem. Acta*, **21**, 469.
- Morgan, W. T. J. & Partridge, S. M. (1940). *Biochem. J.* **34**, 169.
- Morgan, W. T. J. & Partridge, S. M. (1941). *Biochem. J.* **35**, 1140.
- Slein, M. W. & Schnell, G. W. (1953). *Proc. Soc. exp. Biol., N.Y.*, **82**, 734.
- Weil, A. J. (1947). *J. Immunol.* **55**, 363.
- Westphal, O., Lüderitz, O. & Bister, F. (1952). *Z. Naturf.* **7b**, 148.
- Westphal, O., Lüderitz, O., Eichenberger, E. & Keiderling, W. (1952). *Z. Naturf.* **7b**, 534.
- Westphal, O., Lüderitz, O., Fromme, I. & Joseph, N. (1953). *Angew. Chem.* **65**, 555.