We wish to thank Professor Haddow for his encouragement, and Professor Desreux and M. de la Varr for facilities and help in the experiments on flow birefringence.

This investigation has been supported by grants to the Chester Beatty Research Institute from the British Empire Cancer Campaign, the Jane Coffin Childs Memorial Fund for Medical Research, the Anna Fuller Fund, and the National Cancer Institute of the National Institutes of Health, U.S. Public Health Service.

REFERENCES

- Alexander, P. & Fox, M. (1954). *J. Polymer Sci.* **12**, 533.
 Alexander, P. & Stacey, K. A. (1952). *Proc. Roy. Soc.* **212 A**, 274.
 Alexander, P. & Stacey, K. A. (1955). *Trans. Faraday Soc.* **51**, 299.
 Badger, R. M. & Blaker, R. H. (1950). *J. Amer. Chem. Soc.* **72**, 3129.
 Benoit, H. & Doty, P. (1953). *J. Phys. Chem.* **57**, 958.
 Cecil, R. & Ogston, A. G. (1948). *J. Chem. Soc.* p. 1382.
 Conway, B. E. & Butler, J. A. V. (1952). *J. Chem. Soc.* p. 3075.
 Crick, F. H. C. & Watson, J. D. (1953). *Nature, Lond.*, **171**, 737.
 Crick, F. H. C. & Watson, J. D. (1954). *Proc. Roy. Soc.* **223 A**, 80.
 Doty, P., Bunce, B. H. & Reichmann, M. E. (1953). *J. Polymer Sci.* **10**, 109.
 Doty, P., Reichmann, M. E., Rice, S. A. & Thomas, C. A. (1954). *J. Amer. Chem. Soc.* **76**, 3047.
 Doty, P. & Steiner, R. F. (1952). *J. chem. Phys.* **20**, 85.
 Felix, K., Fischer, H., Krekels, A. & Mohr, R. (1951) *Hoppe-Seyl. Z.* **287**, 224; *ibid.* **289**, 10.
 Franklin, R. E. & Gosling, R. G. (1953). *Nature, Lond.*, **171**, 740; **172**, 153.
 Frick, G. (1954). *Biochim. biophys. Acta*, **13**, 374.
 Greenstein, J. P. & Jenrette, W. V. (1941). *Cold Spr. Harb. Symp. quant. Biol.* **9**, 236.
 Gulland, J. M., Jordan, D. O. & Taylor, H. F. W. (1947). *J. Chem. Soc.* p. 1131.
 Kirkwood, J. G. & Goldberg, R. J. (1950). *J. chem. Phys.* **18**, 54.
 Neurath, H., Greenstein, J. P., Putnam, F. W. & Erickson, J. O. (1944). *Chem. Rev.* **34**, 157.
 Norman, N. & Simmons, N. S. (1953). *Atomic Energy Commission Project Report*, no. UCLA 280.
 Northrop, R. G., Nutter, R. L. & Sinsheimer, R. L. (1953). *J. Amer. Chem. Soc.* **75**, 5134.
 Overend, W. G., Stacey, M. & Webb, M. (1951). *J. Chem. Soc.* p. 12450.
 Peacocke, A. R. & Lea, W. E. (1953). *Research, Lond.*, **6**, 159.
 Peterlin, A. (1952). *J. Polymer Sci.* **10**, 425.
 Pouyet, J. & Sadron, Ch. (1954). *Trans. Faraday Soc.* **50**, 303.
 Schwander, H. & Cerf, R. (1949). *Helv. chim. acta*, **32**, 2356.
 Stacey, K. A. & Alexander, P. (1954a). *Ric. Scientifica* (in the Press).
 Stacey, K. A. & Alexander, P. (1954b). *Trans. Faraday Soc.* **50**, 303.
 Stamm, R. F. (1950). *J. opt. Soc. Amer.* **40**, 788.
 Steinhardt, J. (1938). *J. biol. Chem.* **123**, 543.
 Wilkins, M. H. F., Stokes, A. R. & Wilson, H. R. (1953). *Nature, Lond.*, **171**, 739.
 Zimm, B. H. (1948). *J. chem. Phys.* **16**, 1099.

Estimation of Monosaccharides by the Orcinol-Sulphuric Acid Reaction

By J. BRÜCKNER

Chemical Section of the Department of Pathology, Medical School, University of Otago, Dunedin, New Zealand

(Received 1 October 1954)

Methods of estimating carbohydrates, based on the colour reaction with orcinol and sulphuric acid, have been described by Tillmans & Philippi (1929), Sørensen & Haugaard (1933) and Brückner (1943). This paper describes an improved method based on the same principle.

MATERIALS AND METHOD

Orcinol reagent. Orcinol (0.4 g.) was dissolved in 10 ml. of water. The orcinol and its solution should be colourless. If this is not the case, the orcinol must be repeatedly crystallized from water and/or CHCl_3 until a colourless product is obtained. Freshly prepared solutions should be used.

Sulphuric acid reagent (31.2N). This was prepared by mixing 800 ml. of analytical grade H_2SO_4 (sp.gr. 1.84) with 200 ml. of water and adjusting the normality of the solution.

Carbohydrates. Materials (whenever possible analytical reagents or C.P. grade reagents) were obtained from the following sources: D-glucose and D-mannose from British

Drug Houses Ltd.; D-fructose from Schering-Kahlbaum; D-galactose from Hoffman La Roche; L-sorbose, L-rhamnose, D-ribose, D-xylose and D-lyxose from the Pfanstiehl Chemical Co.; D-arabinose from Nutritional Biochemicals Inc. The carbohydrates were dried to constant weight in a vacuum desiccator over H_2SO_4 and NaOH pellets and used in 0.01% (w/v) solutions in water. D-Xylulose was prepared and kindly supplied by Mr R. G. Kulka, Department of Biochemistry, University of Otago. It was used in a solution of approximately 0.01% in water.

Test tubes. It is advisable to use uniformly made test tubes, approximately 17 cm. long, prepared from glass tubing of approximately 0.7 cm. internal diam. Thin-walled test tubes (glass thickness about 0.5–0.6 mm.) are preferred.

General method

The carbohydrate solution (1.0 ml.) and 1.0 ml. of orcinol reagent are mixed in a 100 ml. round flask. The flask is cooled in cold water and 8 ml. of sulphuric acid reagent added, drop by drop, while the flask is steadily rotated in

cold water. Addition of the acid should take about 1.5 min. and heating of the mixture is avoided. About 5–6 ml. of the mixture are then transferred to the test tubes described above, covered with small glass marbles and immediately immersed for exactly 50 sec. in a vigorously boiling water bath. They are then cooled immediately in cold tap water for 3 min. The colour is measured as soon as possible in a spectrophotometer or in a colorimeter fitted with suitable spectrum filters, against a blank containing 1.0 ml. of water in place of the carbohydrate solution and treated as described above. Cuvettes having a 1.0 cm. light path are used.

Adequate curves for an absorption spectrum can be obtained by taking readings at 415, 430, 455, 475, 500, 520, 540, 560, 600 and 625 $m\mu$., the readings being commenced from the blue end of the spectrum. Corresponding spectrum filters may be used for instruments other than spectrophotometers.

Unless stated otherwise, the above conditions, which are recommended for a standard routine method, have been used throughout this work.

RESULTS

The colour reactions of carbohydrates with orcinol and sulphuric acid depend on several factors and vary considerably with the experimental conditions. The colour depends mainly on time of heating, but its intensity depends on the concentration of sulphuric acid and on the temperature during the period of heating. The influence of these factors was investigated in order to obtain optimum conditions for the reaction.

Influence of sulphuric acid concentration

The effect of varying the concentration of sulphuric acid in the reaction mixture was studied with a solution of galactose. The time of heating was 1 min., followed by cooling in cold water. Colours were measured in a Spekker absorptiometer with Ilford spectrum filter no. 604 (500–540 $m\mu$.) against an orcinol-sulphuric acid blank.

The intensity of colour increased with increasing sulphuric acid concentration up to 32.4N, at which concentration maximum colour developed. No colour developed when the concentration of the added sulphuric acid was below 21N. As a convenient concentration 31.2N acid was selected for the experiments. This gives almost maximum colour intensity.

Influence of temperature

Reaction mixtures containing galactose were heated in a water bath to different temperatures for 1 min. Colours were measured in a Spekker absorptiometer with Ilford spectrum filter no. 604 against an orcinol-sulphuric acid blank.

No colour reaction developed when the temperature of the water bath was below 74°. At 100° the

colour was of satisfactory, although not of maximum, intensity. Accordingly, a boiling-water bath was chosen for the standard method, since it ensures an easily reproducible temperature of reasonable constancy.

Influence of time of heating

The effect of time of heating on the development of colour in the reactions with galactose, fructose and glucose was also studied (Figs. 1–4).

On heating for 30 sec. fructose shows a yellow colour of considerable intensity. The absorption maximum at 475 $m\mu$. gradually fades on continued heating and shifts to 420 $m\mu$. and the colour changes to brownish yellow. Glucose and galactose, on the other hand, do not show any appreciable colour on such brief heating. Cherry-red and bluish red colours develop in glucose and galactose solutions on continued heating. Both colours change finally to a brownish yellow, similar to that given by fructose. The heating period of 50 sec. has been chosen for a standard method, because maximum differences between sugars in the colour reactions are produced in this time.

Influence of carbohydrate concentration

It is well known that the colour reactions of carbohydrates with mono- or poly-hydric phenols and sulphuric acid do not strictly follow Beer's law. The relationship of colour development to sugar concentration was investigated with glucose, fructose, galactose and mannose solutions under the standard conditions. The extinction values were measured at suitable maxima in a Unicam SP. 600 spectrophotometer.

Beer's law appeared to be obeyed satisfactorily, since the colours were proportional to concentrations up to 0.1 mg. monosaccharide/ml. Extinction values measured at this concentration were: glucose (475 $m\mu$.), 0.221; fructose (475 $m\mu$.), 1.445; galactose (520 $m\mu$.), 0.492; mannose (520 $m\mu$.), 0.424.

Absorption spectra

The absorption spectra of the colours developed with glucose, fructose, galactose, mannose, sorbose and arabinose are illustrated in Fig. 2. Data of spectral absorptions for the colour reactions given by ribose, xylose, lyxose, rhamnose and xylulose are given in Table 1.

Absorption spectra of the three aldohexoses show differences by which they may be distinguished from one another and from the ketohexoses. The pentoses investigated possess an absorption maximum at 560 $m\mu$., which distinguishes them from hexoses.

Fig. 1. Absorption spectra of carbohydrate solutions (0.1 mg./ml.) in the orcinol- H_2SO_4 reaction on 30 sec. heating at 100° . Colours measured in Unicam SP. 600 spectrophotometer against an orcinol- H_2SO_4 blank. ○, Glucose; ●, galactose; +, fructose.

Fig. 2. Absorption spectra of carbohydrate solutions (0.1 mg./ml.) in the orcinol- H_2SO_4 reaction on 50 sec. heating at 100° . Colours measured in Beckman DU spectrophotometer against an orcinol- H_2SO_4 blank. ▲, Sorbose; x, fructose; ○, glucose; +, galactose, ●, mannose; □, arabinose. Right-hand ordinate refers to sorbose and fructose only.

Fig. 3. Absorption spectra of carbohydrate solutions (0.1 mg./ml.) in the orcinol- H_2SO_4 reaction on 2 min. heating at 100° . Colours were measured in a Unicam SP. 600 spectrophotometer against an orcinol- H_2SO_4 blank. ○, Glucose; ●, galactose; +, fructose.

Fig. 4. Absorption spectra of carbohydrate solutions (0.1 mg./ml.) in the orcinol- H_2SO_4 reaction on 15 min. heating at 100° . Colours were measured in a Unicam SP. 600 spectrophotometer against an orcinol- H_2SO_4 blank. ○, Glucose; ●, galactose; +, fructose.

Orcinol-sulphuric acid reaction of individual monosaccharides

It follows from the above that the shape of the absorption curves will permit a certain characterization of the monosaccharides studied, when only one sugar is present in solution. Aldopentoses have similar absorption spectra and, therefore, cannot be distinguished with the test described. Another

An additional quotient, however, $E_{520\text{ m}\mu}/E_{560\text{ m}\mu}$, may be more suitable for the characterization of pentoses (0.77–0.82 for aldopentoses, 0.94 for xylulose, 1.19–1.29 for aldohexoses, 2.59 for sorbose and 3.76 for fructose).

Individual monosaccharides may be quantitatively estimated with the orcinol-sulphuric acid reaction by determining the extinctions at the maxima given in Fig. 2 and Table 1.

Table 1. *Extinction values of colour reactions of carbohydrates with orcinol-H₂SO₄ reagent*

Carbohydrate solutions (0.1 mg./ml.) were heated with the reagents for 50 sec. at 100°. Extinction values were measured in a Beckman DU spectrophotometer against an orcinol-H₂SO₄ blank. The colour reactions of the carbohydrates investigated do not show appreciable absorption in the orange or red part of the spectrum, except xylulose, which had a third maximum at 625 m μ . ($E=0.264$).

Sugar	Wavelength (m μ)								
	415	430	455	475	500	520	540	560	600
	Extinction values								
Ribose	0.822	0.760	0.406	0.318	0.348	0.375	0.419	0.476	0.106
Xylose	0.770	0.726	0.428	0.334	0.352	0.378	0.412	0.462	0.109
Lyxose	0.755	0.690	0.368	0.278	0.316	0.352	0.402	0.458	0.098
Rhamnose	0.720	0.682	0.425	0.332	0.358	0.379	0.416	0.469	0.111
Xylulose	0.657	0.899	1.135	0.813	0.396	0.276	0.260	0.294	0.200

Table 2. *Quotient values for colour reactions of carbohydrates with orcinol-H₂SO₄ reagent*

Sugar	$E_{475\text{ m}\mu}/E_{540\text{ m}\mu}$	$E_{415\text{ m}\mu}/E_{520\text{ m}\mu}$	$E_{415\text{ m}\mu}/E_{475\text{ m}\mu}$
Glucose	1.59	1.29	0.93
Fructose	10.92	1.68	0.37
Galactose	0.76	0.69	1.00
Mannose	0.89	1.37	1.65
Sorbose	8.31	1.65	0.40
Arabinose	0.71	1.65	2.11
Ribose	0.76	2.19	2.58
Xylose	0.81	2.04	2.30
Lyxose	0.69	2.14	2.72
Rhamnose	0.80	1.90	2.17
Xylulose	3.13	2.38	0.81

similar limitation occurs in the case of fructose and sorbose. Here, however, the specific extinction coefficients measured at 475 m μ . (i.e. λ_{max}) show a great difference for the two sugars.

Alternatively, the studied monosaccharides may be conveniently characterized by determining the ratios of the extinctions measured at wavelengths where maxima and minima and/or identical extinctions occur. Values for the quotients $E_{475\text{ m}\mu}/E_{540\text{ m}\mu}$, $E_{415\text{ m}\mu}/E_{520\text{ m}\mu}$ and $E_{415\text{ m}\mu}/E_{475\text{ m}\mu}$ are listed in Table 2 (the value for $E_{430\text{ m}\mu}/E_{520\text{ m}\mu}$ may be used instead of $E_{415\text{ m}\mu}/E_{520\text{ m}\mu}$).

The first quotient of this series is suitable for characterizing ketoses since the values are much higher than those given by any of the aldoses. The second quotient may be used for the characterization of galactose. Its value (0.69) for galactose is much lower than that found for any other sugar.

Estimation of monosaccharides in mixtures

This may be done by calculation, using Hüfner's method (see Heilmeyer, 1943) of calculating the partition of two pigments mixed in solution, provided individual absorption spectra of the two coloured components are suitable for such a calculation.

Alternatively, it may be more convenient to prepare a graph in which values of the appropriate quotient obtained for two monosaccharides in a mixture are plotted against the corresponding proportions of the same two monosaccharides in the mixture. If the extinction coefficients are measured on solutions containing each of the two monosaccharides in identical concentrations, the following formula may be used for constructing the curve:

Quotient value of an arbitrary mixture

$$= \frac{(K_1 \times M) + (k_1 \times m)}{(K_2 \times M) + (k_2 \times m)}$$

(K_1 and K_2 , k_1 and k_2 are the extinction coefficients of the two monosaccharides at two different wavelengths; M and m are the percentage proportions of the two monosaccharides in the arbitrary mixture). Experimentally the curve may be prepared by measuring the corresponding quotient values for mixed solutions of the two monosaccharides in varying proportions. With the aid of such a curve (Fig. 5) the proportion of glucose and fructose in an unknown mixture can be determined by estimating the extinctions at 475 and 540 m μ . and calculating the ratio (see also Brückner, 1951).

The test may also be performed by comparing the absorption curve obtained on an unknown solution with that of known mixtures of monosaccharides, since calculations for the estimation of three (or more) monosaccharides occurring together are usually too complicated for practical use. With the exception of glucose the monosaccharides investigated possess characteristic absorption maxima by which in most cases they may be recognized in mixtures. Similarly, an elevation or a depression of the values given in Table 2 for the quotients may indicate the presence of a particular monosaccharide.

Fig. 5. Curve for the estimation of the proportions of glucose and fructose in mixtures of the two sugars. The curve has been calculated from values of $E_{475 \text{ m}\mu}$ and $E_{540 \text{ m}\mu}$ of the two sugars. ●, Experimental values for mixtures of the two sugars.

DISCUSSION

The development of colours in reactions between carbohydrates and mono- or poly-hydric phenols heated in the presence of strong mineral acids is usually attributed to the formation of furfurals (from pentoses), and hydroxymethylfurfurals (from hexoses) which condense with phenolic reagents to give coloured products. This view originated with Tollens in his classical investigations on the detection and estimation of carbohydrates. A similar explanation has been offered for the development of colour in Molisch's α -naphthol-sulphuric acid test (Udránszky, 1888, 1910) and the anthrone-

sulphuric acid reaction (Kapuscinski & Zak, 1953).

When the orcinol-sulphuric acid test is applied to carbohydrates under the conditions described in this paper various colours, ranging from yellow to bluish red, appear in the initial stage of the reaction. This suggested that several decomposition products of carbohydrates might be formed in varying proportions. The suggestion is consistent with the findings of Love (1953), who studied the ultraviolet spectra of carbohydrate solutions which had been heated with sulphuric acid. The spectral data indicated the formation of several decomposition products. As a working hypothesis it may be assumed that these decomposition products of unknown structure react with orcinol in the initial stage of the test, but after longer heating (15 min.) the uniform colour developed with all the carbohydrates is due predominantly to condensation of orcinol with furfurals. This assumption may explain why the test shows considerable flexibility on altering the experimental conditions.

The use of 31.2N sulphuric acid and a heating time of 50 sec. have been chosen for the standard procedure, because under these conditions satisfactory extinctions with maximum differences in the spectra develop at reasonably low concentrations of all the monosaccharides studied. Other conditions may be appropriate for special purposes. Thus, heating the mixture for 30 sec. produces a considerable difference in the intensity of the spectral absorptions obtained with fructose on the one hand and glucose and galactose on the other (Fig. 1).

Although aldopentoses have similar absorption spectra in the orcinol-sulphuric acid reaction, xylulose and the hexoses studied show characteristic differences. In most cases the differences are large enough to fulfil requirements for the spectrochemical analysis of a two-component mixture. When unfavourable combinations are encountered yeast may provide a suitable means of removing the fermentable component.

SUMMARY

1. The orcinol-sulphuric acid reaction for carbohydrates has been studied in order to determine optimum conditions in respect of sulphuric acid concentration, and time and temperature of heating. A standard procedure is recommended for the characterization of individual monosaccharides and for their estimation, occurring in solution singly or in binary mixtures.

2. Data of spectral absorption of the reaction products in the visible range and methods for calculation are given.

The author wishes to thank Professor N. L. Edson for his kindly advice and assistance in preparation of this paper.

REFERENCES

- Brückner, J. (1943). *Hoppe-Seyl. Z.* **277**, 181.
 Brückner, J. (1951). *Helv. physiol. acta*, **9**, 259.
 Heilmeyer, L. (1943). *Spectrophotometry in Medicine*.
 English ed. London: Adam Hilger.
- Kapuscinski, V. & Zak, B. (1953). *Amer. J. clin. Path.* **23**, 784.
 Love, R. M. (1953). *Biochem. J.* **55**, 126.
 Sørensen, M. & Haugaard, G. (1933). *Biochem. Z.* **260**, 247.
 Tillmans, J. & Philippi, K. (1929). *Biochem. Z.* **215**, 36.
 Udránszky, L. (1888). *Hoppe-Seyl. Z.* **12**, 355.
 Udránszky, L. (1910). *Hoppe-Seyl. Z.* **68**, 88.

Paper Electrophoresis as a Quantitative Method

SERUM PROTEINS

BY W. P. JENCKS, MERA R. JETTON AND E. L. DURRUM*

*Department of Pharmacology, Army Medical Service Graduate School,
 Walter Reed Army Medical Center, Washington, D.C.*

(Received 25 October 1954)

A number of workers have recently questioned the validity of paper electrophoresis as a method for the separation and quantitative estimation of serum proteins. The four main problems are: (1) albumin 'tailing' due to irreversible adsorption of albumin on paper during its migration (Kunkel & Tiselius, 1951; Merklen & Masseyeff, 1952; Sommerfelt, 1953); (2) variation in dye binding capacity with the amount and area of application of protein (Franglen & Martin, 1954; Hardwicke, 1954); (3) fading and incomplete elution of the commonly used dyes bromophenol blue and amidoschwarz 10B (Flynn & de Mayo, 1951; Pezold & Peiser, 1953; Franglen & Martin, 1954); and (4) the lack of a linear relationship of dye concentration to photocell response in the direct scanning of dye on paper (Van Os, 1953; Crook, Harris, Hassan & Warren, 1954). In addition, there has been widespread disagreement as to the magnitude and usefulness of 'correction factors' for the conversion of the results of paper electrophoresis expressed in terms of dye binding to those of moving boundary electrophoresis expressed in terms of refractive index increment.

A study of these and some related problems is presented here. On the basis of this work, a detailed procedure for the paper electrophoretic analysis of serum proteins has been developed.

MATERIALS AND METHODS

Paper electrophoresis of serum. This was carried out on Whatman 3MM paper strips 2.9 × 30.0 cm. in a hanging strip type cell (Williams, Pickels & Durrum, 1955). The paper was saturated with barbital buffer pH 8.6 ionic strength 0.075 and allowed to drain in position in the cell for 15 min. Serum (0.01 ml.) was then applied from a micropipette to the apex of the strip directly over the silicone-coated glass

supporting rod. Separation was carried out with a constant current of 8 ma for 8 strips (90–100 v, about 3–3.5 v/cm.) for 16 hr. at room temperature, avoiding any marked inequalities of temperature from drafts, sunlight, etc. The strips were then removed from contact with the feed wick, spread flat, and dried in an oven at 110–120° for 30 min.

Staining. The strips were transferred to a staining rack which holds them apart, and placed in an aqueous dye bath containing 0.01% bromophenol blue, 5% acetic acid (v/v) and 5% ZnSO₄ · 7H₂O (w/v) for 16 hr. at room temperature. They were rinsed 3 times for 5, 5 and 10 min. in 2% acetic acid (v/v) without agitation, and then for 2 min. in a bath of 2% sodium acetate (CH₃COONa, 3H₂O, w/v) and 10% acetic acid (v/v). After being blotted gently on clean filter paper (S & S 470), the strips were transferred to a drying rack which touches them only on the ends and dried in the oven for 8–10 min. at 110–120°. A typical pattern is shown in Fig. 1.

Fig. 1. A typical paper electrophoretic pattern of serum proteins prepared by the technique described.

Elution. The strips were cut into the desired segments and eluted in 0.01 N-NaOH for 30 min. with occasional agitation, thoroughly mixed, and read within 1 hr. in a Beckman model DU spectrophotometer at 590 mμ. in 1 cm. cuvettes. The monochromator was calibrated with a mercury lamp.

Scanning. Dyed strips were scanned with an automatic recording and integrating scanner (Durrum & Gilford, 1954) after the strips had been 'cleared' by dipping in liquid petrolatum (mineral oil), allowed to drain for 2–18 hr., and blotted. After scanning, the petrolatum was removed with ether.

Protein was determined by the biuret method (Gornall, Bardawill & David, 1949) or by micro-Kjeldahl determination of nitrogen.

* Present address: Department of Pharmacology, Stanford Medical School, San Francisco 15, California.