linked to the ϵ -amino group. However, certain properties of a substance containing lysine and aspartic acid raise the question whether its structure is more complicated than that of a simple dipeptide.

5. One of the aspartic acid residues in bacitracin A is readily liberated by hydrolysis in the free state. The second residue is linked to lysine by a bond which is unusually resistant to hydrolysis. A fraction which contained mainly the first aspartic acid was separated from a fraction which contained mainly the second. In one experiment the two fractions were subjected separately to the action of D-aspartic acid oxidase. In another the aspartic acid in each fraction was isolated and its specific rotation measured. The results indicate that the easily liberated aspartic acid is derived from a residue with the D-configuration and that the aspartic acid residue next to lysine has the L-configuration.

We wish to thank Dr F. Sanger, F.R.S., for a sample of α -DNP-lysine, and the Commercial Solvents Corporation, Terre Haute, Indiana, U.S.A., for generous gifts of bacitracin. We are grateful to Miss M. Read and Mrs A. Talboys for expert technical assistance. One of us (I.M.L.) is indebted to the Medical Research Council for a Scholarship.

REFERENCES

- Barry, G. T., Gregory, J. D. & Craig, L. C. (1948). *J. biol. Chem.* **175**, 485.
- Blackburn, S. & Lowther, A. G. (1951). *Biochem. J.* **48**, 126.
- Craig, L. C., Gregory, J. D. & Hausmann, W. (1950). *Analyt. Chem.* **22**, 1462.
- Craig, L. C., Hausmann, W. & Weisiger, J. R. (1952). *J. biol. Chem.* **199**, 865.
- Craig, L. C., Hausmann, W. & Weisiger, J. R. (1954). *J. Amer. chem. Soc.* **76**, 2839.
- Fischer, E. (1899). *Ber. dtsh. chem. Ges.* **32**, 2451.
- Hausmann, W., Weisiger, J. R. & Craig, L. C. (1955). *J. Amer. chem. Soc.* **77**, 723.
- Hirs, C. H. W., Moore, S. & Stein, W. H. (1954). *J. Amer. chem. Soc.* **76**, 6063.
- Keilin, D. & Hartree, E. F. (1936). *Proc. Roy. Soc. B*, **119**, 141.
- Lequesne, W. J. & Young, G. T. (1952). *J. chem. Soc.* p. 24.
- Lockhart, I. M. & Abraham, E. P. (1954a). *Biochem. J.* **58**, 633.
- Lockhart, I. M. & Abraham, E. P. (1954b). *Biochem. J.* **58**, xlvii.
- Lockhart, I. M., Abraham, E. P. & Newton, G. G. F. (1955). *Biochem. J.* **61**, 534.
- Moore, S. & Stein, W. H. (1948). *J. biol. Chem.* **176**, 367.
- Moore, S. & Stein, W. H. (1951). *J. biol. Chem.* **192**, 663.
- Moore, S. & Stein, W. H. (1954). *J. biol. Chem.* **211**, 907.
- Newton, G. G. F. & Abraham, E. P. (1953). *Biochem. J.* **53**, 604.
- Newton, G. G. F. & Abraham, E. P. (1954). *Biochem. J.* **58**, 103.
- Sanger, F. & Thompson, E. O. P. (1953). *Biochem. J.* **53**, 353.
- Still, J. L., Buell, H. V., Knox, W. E. & Green, D. E. (1949). *J. biol. Chem.* **179**, 831.
- Synge, R. L. M. (1951). *Biochem. J.* **49**, 642.
- Weisiger, J. R., Hausmann, W. & Craig, L. C. (1955). *J. Amer. chem. Soc.* **77**, 731.
- Woiwod, A. J. (1949). *J. gen. Microbiol.* **3**, 312.
- Work, E. (1949). *Biochim. biophys. Acta*, **3**, 400.

Isolation of Cephalosporin C, a Penicillin-like Antibiotic Containing D- α -Amino adipic Acid

BY G. G. F. NEWTON* AND E. P. ABRAHAM
Sir William Dunn School of Pathology, University of Oxford

(Received 26 September 1955)

Abraham, Newton & Hale (1954) described the purification of a hydrophilic antibiotic, cephalosporin N, which was produced by a species of *Cephalosporium* (I.M.I. 49137). Cephalosporin N yielded D- α -amino adipic acid on hydrolysis, and the results of degradative experiments showed that it was probably (D-4-amino-4-carboxy-n-butyl)-penicillin (Newton & Abraham, 1954).

Crude preparations of cephalosporin N were reported to contain a second antibiotic named cephalosporin C (Newton & Abraham, 1955). The

latter contained sulphur and gave D- α -amino adipic acid on hydrolysis. Unlike cephalosporin N, it was stable in aqueous solution at pH 3 and showed an absorption band in ultraviolet light with a maximum at 260 m μ . This paper describes the isolation of cephalosporin C, in the form of a crystalline sodium salt, and gives some of its properties.

Cephalosporin C was first encountered as a distinct entity during the purification of crude cephalosporin N penillic acid by chromatography on an anion-exchange resin, when it followed the penillic acid from the column as a well-defined band. After isolation it was found to show

* Member of the Scientific Staff, Medical Research Council.

anti-bacterial activity. Cephalosporin C was not formed from cephalosporin N during the conversion of the latter into its penillic acid at pH 3, because no cephalosporin C was present in the penillic acid prepared from highly purified cephalosporin N (Newton & Abraham, 1954). The possibility that it was formed at pH 3 from an impurity in crude cephalosporin N was later eliminated by isolating it without recourse to the penillic acid rearrangement. Many of the physical properties of cephalosporin C are very similar to those of cephalosporin N, and the separation of these two compounds proved more difficult than the separation of the former from cephalosporin N penillic acid. However, a partial resolution was achieved in a manner which left little doubt that cephalosporin C was present as such in the culture fluid of the *Cephalosporium*.

An account of some of the biological properties of cephalosporin C has been given by Florey (1955). The substance was remarkably non-toxic to the mouse, a dose of 100 mg. given intravenously having no noticeable ill effect. It was able to afford complete protection to mice infected with a strain of haemolytic streptococcus which killed untreated animals within 12 hr.

RESULTS

Of the preparations of cephalosporin N examined, all except those that had been finally purified by countercurrent distribution (Abraham *et al.* 1954) were found to contain cephalosporin C. In the more active preparations, cephalosporin C was the main contaminant. It was largely responsible for the survival of some antibacterial activity when the cephalosporin N was converted into the corresponding penillic acid by keeping the preparations at pH 3 in aqueous solution. The amount of cephalosporin C in the preparations varied from batch to batch, but in most cases it was much smaller than the amount of cephalosporin N.

Separation of cephalosporin C from cephalosporin N penillic acid

Cephalosporin C and cephalosporin N penillic acid were readily separated by chromatography on a column of Amberlite IR 4B or by countercurrent distribution in a phenol-carbon tetrachloride-water system containing acetic acid.

Elution from Amberlite IR 4B. Fig. 1 shows the elution curves obtained when two preparations of cephalosporin N penillic acid were chromatographed on a column of Amberlite IR 4B in ammonium acetate buffer, pH 5.0. The top curve was obtained with a preparation of the penillic acid from partly purified cephalosporin N (51 u./mg.). The material in the major peak was cephalosporin N penillic acid

and that in the minor peak was cephalosporin C. The crystalline sodium salt of cephalosporin C was isolated from the contents of fractions 92–112. The bottom curve was obtained with a preparation of the penillic acid from a sample of cephalosporin N (68 u./mg.) that had been purified by countercurrent distribution in the manner described by Abraham *et al.* (1954).

Fig. 1. Separation of cephalosporin C and cephalosporin N penillic acid on a column (1.7 cm. $\times$ 20 cm.) of Amberlite IR 4B. Elution with 0.2M ammonium acetate, pH 5.0, 3 ml. fractions being collected. Top curve, crude penillic acid from partly purified cephalosporin N. Bottom curve, penillic acid from highly purified cephalosporin N.

Complete resolution of cephalosporin N penillic acid and cephalosporin C was also obtained on a column of Amberlite IR 4B when pyridine acetate, pH 5.0, was used as a buffer. This buffer was removed more easily than ammonium acetate from the products eluted from the column.

Countercurrent distribution between solvents. Fig. 2 shows the results obtained when a preparation of crude cephalosporin N penillic acid was subjected to ninety-seven transfers in a 100-tube machine, a solvent system composed of phenol, carbon tetrachloride and aqueous acetic acid being used. The distribution curve represents the ninhydrin colour densities given by samples from the upper (aqueous) layers. The material used in this experiment contained a considerable amount of substances other than cephalosporin N penillic acid and cephalosporin C. Cephalosporin N penillic acid was located in tubes 48–75 and cephalosporin C in tubes 20–50.

After the analysis shown in Fig. 2, the distribution was continued until 300 transfers had been completed. All the cephalosporin C was present in

fractions withdrawn from the machine between transfers 198 and 292. These fractions yielded crystalline cephalosporin C sodium salt.

Separation of cephalosporin C from cephalosporin N

Elution from Amberlite IR 4B. The recovery of active cephalosporin N after chromatography on a buffered ion-exchange resin was complicated by the instability of the substance at a pH lower than 5. Desalting by conventional methods, including those depending on the sublimation of volatile

Fig. 3 shows an elution curve obtained when a partly purified preparation of cephalosporin N (40 u./mg.) was chromatographed on a column of Amberlite IR 4B (20 cm. long) in pyridine sulphate buffer, pH 5.0, at 5°. The first small peak contained unidentified material without antibacterial activity. The major peak was asymmetrical. Material from its leading edge contained cephalosporin N virtually free from cephalosporin C. Material from its trailing edge contained a progressively large proportion of cephalosporin C and smaller proportion of cephalosporin N. Fractions 18–24 combined yielded a preparation of cephalosporin N barium salt with an activity of 50 u./mg. Fractions

Fig. 2. Countercurrent distribution of crude cephalosporin N penillic acid in a phenol-carbon tetrachloride-water system containing acetic acid. The curve shows the fundamental series after ninety-seven transfers. The upper (aqueous) phase only was analysed.

buffers, was unsuccessful. This difficulty was overcome by using pyridine sulphate as a buffer. Addition of the appropriate amount of barium hydroxide to a solution of either cephalosporin N or cephalosporin C in the buffer removed the sulphate ions and left the barium salt of the antibiotic in aqueous pyridine; the pyridine could then be removed *in vacuo* without loss of antibacterial activity. When mixtures of cephalosporin N and cephalosporin C were obtained, the maximum amount of the latter in the product could be estimated by measuring the optical density at 260 mμ.

Fig. 3. Partial separation of cephalosporin C and cephalosporin N on a column (1.9 cm. $\times$ 20 cm.) of Amberlite IR 4B. Elution with pyridine sulphate, pH 5.0, 3 ml. fractions being collected. ●, Ninhydrin colour values; x, antibacterial activity. Cephalosporin N is concentrated in the leading edge of the main peak, and cephalosporin C in the trailing edge.

30–45 yielded a crude barium salt of cephalosporin C still mixed with some cephalosporin N. This product was converted into its sodium salt. On concentrating an aqueous solution of the latter to a syrup the sodium salt of cephalosporin C separated as a crystalline mass.

In a similar experiment with a column 130 cm. long the trailing edge of the main peak was prolonged, but complete resolution of cephalosporin N and cephalosporin C was not obtained. This experiment yielded a preparation of the barium salt of cephalosporin N with an activity of 60 u./mg. and also a fraction from which cephalosporin C was obtained as a crystalline sodium salt.

Separation on paper chromatograms. On buffered paper chromatograms developed with methanol, which were used by Olson, Jennings & Junek (1953) for studying synnematin, cephalosporin N ran significantly faster than cephalosporin C. Thus

on paper buffered with sodium citrate, pH 5.5, cephalosporin N showed an R_f value of 0.44 and cephalosporin C an R_f value of 0.34. When a mixture of the two compounds was run on the chromatogram for 48 hr., the solvent being allowed to drip from the bottom of the paper, two clearly resolved spots were obtained on spraying with ninhydrin. Resolution that was nearly complete was obtained on paper buffered with sodium phosphate at pH 6.0. Fig. 4 shows the result of an experiment in which cephalosporin N (25 μ g.), cephalosporin C (100 μ g.) and a mixture of the two were run on a chromatogram under these conditions

Fig. 4. Zones of inhibition on a plate seeded with *Salmonella typhi*, showing the separation of cephalosporin C and cephalosporin N on Whatman no. 1 paper buffered with sodium phosphate, pH 6.0. The chromatogram was developed with methanol. 1, Cephalosporin C; 2, cephalosporin C and cephalosporin N; 3, cephalosporin N.

for 48 hr. at 4°. The paper was placed for 30 min. on a plate of agar seeded with a strain of *Salmonella typhi*, and the plate then incubated overnight. The positions of the antibiotics are marked by clear zones in an opaque area of bacterial growth.

Most of the crude preparations of cephalosporin N that were tested on paper chromatograms gave a spot in the position of cephalosporin C, as well as one in the position of cephalosporin N, when the paper was sprayed with ninhydrin. With the preparations that were richest in cephalosporin C the latter could also be distinguished from cephalo-

sporin N on the chromatograms by the distribution of antibacterial activity. However, the fact that the activity/mg. of cephalosporin C was only about one-tenth of that of cephalosporin N made this impossible with other preparations in which cephalosporin C was present in lower concentration.

When subjected to ionophoresis on paper in collidine-acetate buffer, pH 7, in the manner described by Newton & Abraham (1954), cephalosporin C migrated towards the anode at almost the same rate as cephalosporin N. No separation of the compounds could be obtained under these conditions.

Fig. 5. Photomicrograph ($\times 132$) of cephalosporin C sodium salt, crystallized from aqueous *n*-propanol. The crystals were photographed between crossed nicol prisms.

Some properties of cephalosporin C

When preparations of the sodium salt of cephalosporin C were concentrated, in aqueous solution, to a syrup, crystallization began readily and continued until the preparation contained a solid mass of monoclinic crystals. Fig. 5 shows a product obtained by recrystallization from aqueous *n*-propanol. An X-ray crystallographic analysis was kindly carried out by Dr Dorothy Hodgkin and Mrs D. Mackay, who reported as follows. Unit cell dimensions: $a = 13.12 \text{ \AA}$; $b = 4.97 \text{ \AA}$; $c = 17.82 \text{ \AA}$; $\beta = 106^\circ, 30$. Space group, $P2_1$; number of molecules in the unit cell, 2; density, 1.38. The molecular weight calculated from these figures is 470 ± 15 .

On drying *in vacuo* at room temperature crystalline cephalosporin C sodium salt, $[\alpha]_D^{20} +103^\circ$, lost 7.6% of its weight. This weight was rapidly regained on exposure to the laboratory air. The results of elementary analysis indicated that a possible formula for the hydrated product was $C_{18}H_{20}O_8N_3SNa \cdot 2H_2O$. Analytical figures for the amorphous free acid, obtained from the sodium salt by ionophoresis in a four-compartment cell (Synge, 1951), were consistent with this formula.

Cephalosporin C gave a positive ninhydrin reaction and a negative nitroprusside reaction. Electrometric titration indicated that it contained two acidic groups, with pK values of 3.1 and <2.6 respectively, and a basic group with a pK value of 9.8. The equivalent weight, calculated from the span of the titration curve for the basic group, was 480 ± 15 . In these properties cephalosporin C behaved as a monoaminodicarboxylic acid closely resembling cephalosporin N.

Fig. 6. Ultraviolet absorption spectrum of cephalosporin C sodium salt in water.

The ultraviolet absorption spectrum of cephalosporin C sodium salt in water ($\lambda_{\max}$ 260 $m\mu$, $\epsilon_{\max}$ 9000) is shown in Fig. 6. The infrared spectrum showed bands at the following wavelengths: 2.94, 3.06, 5.61, 5.77, 6.05, 6.29, 6.57, 7.17 and 7.36 μ . The bands at 6.05 and 6.57 μ . could be attributed to the C=O of a monosubstituted amide, the band at 6.29 μ . to the carboxylate ion, the bands at 7.17 and 7.36 μ . to an *isopropyl* group, and the band at 5.77 μ . to an ester or lactone grouping. The band at 5.61 μ . is of particular interest because a band in this region is shown by cephalosporin N and the common penicillins, and has been attributed in the latter to the C=O of the β -lactam-thiazolidine ring system (Thompson, Brattain, Randall & Rasmussen, 1949).

When tested against a strain of *Salm. typhi* and against a strain of *Staphylococcus aureus* which did not produce penicillinase, cephalosporin C sodium salt showed an activity of 8–10 u./mg., the unit being that used by Abraham *et al.* (1954) for assaying cephalosporin N. Against these organisms cephalosporin C was only about one-tenth as active as cephalosporin N, but the two compounds showed similar activities against a strain of *Escherichia coli*.

Cephalosporin C was relatively stable to acid, losing no detectable activity on standing in 0.1N-hydrochloric acid at room temperature for 4 hr. It was also more stable than cephalosporin N in the presence of heavy-metal ions, such as those of copper, zinc and mercury. It was rapidly inactivated, however, in alkaline solution above pH 11. Unlike cephalosporin N, it rapidly gave a heavy black precipitate when treated with sodium plumbite in 50% (w/w) sodium hydroxide solution at room temperature.

The sensitivity of cephalosporin C to four different preparations of penicillinase was compared with that of cephalosporin N under similar conditions. Table 1 shows the amount of each enzyme preparation required to destroy half the activity of a solution of cephalosporin C, relative to the amount required to destroy half the activity of a solution containing the same weight

Table 1. Relative amounts of different preparations of penicillinase required to inactivate equal weights of cephalosporin C and cephalosporin N under similar conditions

For details see Experimental section.

Penicillinase		Relative amounts required for 50% inactivation (amount for cephalosporin C/amount for cephalosporin N)
Source	Nature	
<i>B. subtilis</i> strain 749	Crude	> 100
<i>B. cereus</i> (NRRL 569)	Crude	5
<i>B. cereus</i> (NRRL 569)	Purified	30
<i>B. cereus</i> 5/B	Crystalline	10 000

of cephalosporin N. A crude preparation of penicillinase from a strain of *Bacillus subtilis* had no effect on cephalosporin C in the highest concentration at which it was tested. A crude preparation of penicillinase from a strain of *Bacillus cereus* was active against cephalosporin C, but considerably less active than it was against cephalosporin N. The relative activity against cephalosporin C of a purified preparation of penicillinase from the same source was smaller still, and that of crystalline penicillinase from another strain of *B. cereus* (Pollock, Torriani & Tridgell, 1956) was very small indeed. These results are most easily explained by the assumption that the major component, at least, of the penicillinase produced by *B. subtilis* and *B. cereus* has no significant activity against cephalosporin C, but that the last-named is inactivated by a separate enzyme, C-cephalosporinase. However, they leave open the question whether this cephalosporinase has any activity against the penicillins.

EXPERIMENTAL

General methods

Antibacterial activity was assayed by the method used by Abraham *et al.* (1954), with *Salm. typhi* as the test organism and cephalosporin N as a standard. Photometric ninhydrin determinations were made by the method of Moore & Stein (1948). Electrometric titrations were carried out at 20° in a manner described by Newton & Abraham (1953). Infrared spectra of samples in paraffin paste were kindly measured by Dr F. B. Strauss, using a Perkin-Elmer double-beam photometer, model 21.

Paper chromatography

Paper chromatograms were run on Whatman no. 1 paper which had been treated with 0.05 M sodium citrate buffer, pH 5.5, or with M sodium phosphate buffer, pH 6. Before use the paper was dipped in the buffer solution and dried in air at 37°. Usually about 100 µg. of material was applied to the paper. The chromatograms were developed with methanol which had been dried over CaO and distilled.

Chromatography on Amberlite resin

Resin. The weakly basic anion-exchange resin Amberlite IR 4B (150–300 mesh) was used in these experiments. In each case the resin was first brought to equilibrium with the buffer solution used for elution (cf. Hirs, Moore & Stein, 1952).

Buffer solutions. The following buffers were used: (1) ammonium acetate, pH 5.0 (0.2M), prepared as described by Hirs *et al.* (1952); (2) pyridine acetate, prepared by adding N pyridine (about 250 ml.) to 2N acetic acid (150 ml.) until the pH rose to 5.0, and diluting the solution to 1 l. with water; (3) pyridine sulphate, prepared by adding N pyridine (about 450 ml.) to 150 ml. of 2N-H₂SO₄ until the pH rose to 5.0 and diluting the solution to 1 l. with water.

Separation of cephalosporin C from crude cephalosporin N penicillin acid. (a) Crude cephalosporin N penicillin acid

(98 mg.) was dissolved in 5 ml. of ammonium acetate buffer and the solution added to a column (1.7 cm. × 20 cm.) of Amberlite resin. Elution was carried out with ammonium acetate buffer, the flow rate being adjusted so that 3 ml. fractions were collected every 15 min. Samples taken from odd-numbered fractions were diluted 10 times with water and the optical densities of the resulting solutions were measured at 240 mµ. (Fig. 1). Fractions 92–112 were combined, concentrated to a syrup in a rotating evaporator (Craig, Gregory & Hausmann, 1950), and finally dried *in vacuo* in a desiccator containing P₂O₅ and KOH. The solid was dissolved in water (25 ml.) and the solution placed in the specimen compartment of a four-compartment cell (Synge, 1951), fitted with a cooling coil carrying aqueous ethanol at -5°. Ionophoresis was continued until the current had fallen to 25 ma at 500 v. Cephalosporin C was recovered from the acetic acid compartment of the cell by freeze-drying. The product, which still retained some acetic acid, was dissolved in water (0.5 ml.) and precipitated with acetone (10 ml.). The sticky precipitate was separated by centrifuging and stirred with dry acetone until it formed a powder (7.7 mg.). The powder was dissolved in water (2 ml.), the pH adjusted to 6.5 by the addition of 0.1N-NaOH (0.17 ml.), and the resulting solution concentrated to a syrup. Within a few hours the syrup changed to a mass of crystals.

(b) Cephalosporin C was isolated in a similar manner in an experiment in which pyridine acetate was used instead of ammonium acetate. In this case the elution curve was obtained by measuring ninhydrin colour densities, using 0.2 ml. samples of alternate fractions from the column. Ionophoresis in the four-compartment cell was unnecessary for the complete removal of the buffer at room temperature. The fractions containing cephalosporin C were evaporated *in vacuo* and the product precipitated from a small volume of water by acetone. The precipitate was dissolved in water, the pH of the solution brought to 7.5 with NaOH, and the solution of the resulting sodium salt then concentrated *in vacuo*. On standing, the resulting syrup changed to a mass of crystals.

Separation of cephalosporin C from cephalosporin N. Partly purified cephalosporin N (600 mg., 41 u./mg.) was dissolved in 4 ml. of pyridine sulphate buffer and added to a column (1.9 cm. × 20 cm.) of Amberlite resin. The column was surrounded by a jacket carrying running water at 5°. Elution was carried out with pyridine sulphate buffer, 3 ml. fractions being collected every 20 min. Each fraction was collected in a tube containing 6 drops of pyridine, whereby its final pH was raised to 6. Photometric ninhydrin determinations were made on 0.1 ml. samples of alternate fractions, and samples from a few selected fractions were withdrawn for measurement of antibacterial activity (Fig. 3). Fractions 18–24 were combined and 0.3N-Ba(OH)₂ was added until the pH was 7.6. The precipitate was removed by centrifuging and the supernatant freeze-dried to yield a preparation of the barium salt of cephalosporin N (195 mg.) as a white powder. Its activity was 50 u./mg. Traces of pyridine still remaining in this product were removed by dissolving it in water (2 ml.), adjusting the pH to 7.2, and precipitating the product by the addition of 40 ml. of acetone.

Fractions 30–45 were combined and treated in a similar manner. They yielded 44 mg. of a crude barium salt of cephalosporin C. This product was dissolved in water and

a solution of Na_2SO_4 (10 mg./ml.) added in an amount just sufficient to precipitate the Ba^{2+} as BaSO_4 . After centrifuging, the supernatant solution, containing the sodium salt of cephalosporin C, was concentrated *in vacuo* to a syrup. On standing the latter changed to a crystalline mass.

Countercurrent distribution

Solvent system. This was made by mixing phenol saturated with water (2 l.), CCl_4 (0.8 l.), water saturated with phenol (9 l.), and glacial acetic acid (0.2 l.).

Separation of cephalosporin C from crude cephalosporin N penillic acid. Crude cephalosporin N penillic acid (40 g.) was dissolved in 75 ml. of the upper layer and 60 ml. of the lower layer of the solvent system. The pH of the upper layer was adjusted to 3.1 with 11 N-HCl (6.5 ml.). The solutions were then added to tubes 0, 1 and 2 of an all-glass, automatic, countercurrent-distribution machine, the fixed vol. of the lower layer in each tube being 20 ml. and the vol. of the mobile upper layer 27 ml. The machine was first set to complete 97 transfers, by means of the fundamental procedure (Craig & Craig, 1950). Samples (4.8 $\mu\text{l.}$) from the upper layer of every fourth tube were analysed by the photometric ninhydrin procedure (Fig. 2). The machine was then set to carry out a further 200 transfers, the upper layers being withdrawn from tube 100. The fractions withdrawn during transfers 99–164 were not examined further. Photometric ninhydrin determinations were made on samples (20 $\mu\text{l.}$) of alternate fractions withdrawn during transfers 165–300. Antibacterial activities were measured on samples (1.0 ml.) of every sixth fraction withdrawn during transfers 182–296. For this purpose the samples were diluted with 1 vol. of 0.2 M- Na_2POH_4 and the resulting solutions extracted twice with 1 vol. of CCl_4 to remove phenol. The final pH of the solutions was 6.8. The ninhydrin and activity determinations indicated that cephalosporin C was located in the fractions withdrawn during transfers 198–292.

Recovery of cephalosporin C. The pooled withdrawn fractions containing cephalosporin C were evaporated to about 250 ml. in a rotating evaporator (Craig *et al.* 1950) and the concentrated solution was extracted first with 2 vol. of CCl_4 and then three times with 1 vol. of benzene. The pH of the aqueous solution was then adjusted to 3.3 by the addition of pyridine, the solution concentrated to 50 ml. in the rotating evaporator and the concentrate freeze-dried. The resulting cephalosporin C was dissolved in a small volume of water and precipitated by the addition of 20 vol. of acetone. The precipitate was washed with acetone and dissolved in water, and the pH of the solution brought to 7.5 by addition of 2 N-NaOH. On concentrating the solution to a syrup, cephalosporin C separated as the crystalline sodium salt. The total yield was 800 mg.

Isolation and properties of cephalosporin C

Concentration of solutions of the sodium salt of cephalosporin C gave a syrup which crystallized on standing. The crystalline mass was stirred with a small amount of 70% (v/v) ethanol, which dissolved amorphous material, and the crystals were removed by filtration. The product was recrystallized by dissolving it in a small volume of water, slowly adding *n*-propanol until crystallization began, and then allowing the mixture to stand in the cold room. The air-dried crystalline sodium salt had $[\alpha]_D^{20} + 103^\circ$ in water (c, 0.9). When dried *in vacuo* at room temperature (or at

40°) it lost weight (7.5%) which was rapidly regained on exposure to the laboratory air. Elementary analyses were made on the air-dried products. (Found: C, 40.0; H, 5.2; N, 8.9; S, 6.2; Na, 4.9. Calc. for $\text{C}_{16}\text{H}_{20}\text{O}_8\text{N}_3\text{SNa}\cdot 2\text{H}_2\text{O}$: C, 40.5; H, 5.1; N, 8.9; S, 6.7; Na, 4.9; $2\text{H}_2\text{O}$, 7.5%; mol.wt. 473.)

Amorphous cephalosporin C free acid was prepared by ionophoresis of the crystalline sodium salt in the four-compartment cell. The free acid, obtained by freeze-drying the contents of the acetic acid compartment, was precipitated from aqueous solution by addition of 20 vol. of acetone. The air-dried product lost weight (4.2%) on drying *in vacuo* at room temperature. Elementary analyses were made on the air-dried product. (Found: C, 44.4; H, 5.6; N, 9.8; S, 7.3. Calc. for $\text{C}_{16}\text{H}_{21}\text{O}_8\text{N}_3\text{S}\cdot \text{H}_2\text{O}$: C, 44.5; H, 5.3; N, 9.7; S, 7.4%.)

Stability to heavy-metal ions. Solutions containing 0.2 mg. of cephalosporin C sodium salt and 1 mg. of $\text{CuSO}_4\cdot 5\text{H}_2\text{O}$, $\text{Pb}(\text{NO}_3)_2$ or ZnSO_4 in 1 ml. of water were kept at 37° for 3 hr. Antibacterial assays showed that no detectable activity had been lost in the presence of Cu^{2+} and Zn^{2+} and that the loss was less than 50% in the presence of Pb^{2+} . Under similar conditions cephalosporin N was completely inactivated.

Effect of penicillinase preparations. Four different preparations of penicillinase were used: (1) a solution of crude penicillinase from *B. subtilis* strain 749 (cf. Manson, Pollock & Tridgell, 1954), kindly provided by The Distillers Co. (Biochemicals) Ltd., (2) a crude freeze-dried preparation of penicillinase from *B. cereus* strain 569; (3) a highly purified preparation of penicillinase from *B. cereus* strain 569 (Pollock & Torriani, 1953); (4) twice-crystallized penicillinase from *B. cereus* strain 5/B, a constitutive mutant from a strain of *B. cereus* which does not produce penicillinase (Pollock *et al.* 1956). The three preparations of penicillinase from *B. cereus* were kindly provided by Dr M. R. Pollock.

Solutions of cephalosporin C sodium salt (1 mg.) in 0.9 ml. of 0.6 M sodium phosphate buffer, pH 7.0, were mixed with 0.1 ml. samples of solutions of the penicillinase and the mixtures kept at 37° for 3 hr. Each preparation of penicillinase was used in a series of decreasing concentrations, made by serial dilution of a stock solution of appropriate strength. Solutions of cephalosporin N (1 mg.) were treated in the same way as those of cephalosporin C. After incubation, the solutions containing penicillinase and control solutions to which no enzyme had been added were assayed against *Salm. typhi*. Under these conditions the degree of inactivation was normally reduced from more than 75% to less than 25% over a tenfold dilution of the enzyme solution. The dilution at which about 50% inactivation occurred was taken as the end point.

This method of measuring enzyme activities was a relatively crude one, since enzymic action continued during the period of assay. However, the method was simple and the results were sufficiently clear-cut to justify its use in the present case.

SUMMARY

1. Crude preparations of cephalosporin N, a hydrophilic penicillin, have been found to contain a second hydrophilic antibiotic. This substance, named cephalosporin C, has been isolated as a crystalline sodium salt.

2. Mixtures of cephalosporin C and cephalosporin N penillic acid have been completely resolved by chromatography on an anion-exchange resin and by countercurrent distribution in a phenol-water system containing acetic acid. Mixtures of cephalosporin C and cephalosporin N have been resolved on paper chromatograms and partly resolved on an ion-exchange resin.

3. Cephalosporin C has been provisionally assigned the molecular formula $C_{16}H_{21}O_8N_3S$. It behaves as a monoaminodicarboxylic acid. It shows an absorption maximum at $260\text{ m}\mu$, and its infrared spectrum contains a band at $5.61\text{ }\mu$.

4. Cephalosporin C shows similar activity to *Salm. typhi* and *Staph. aureus*. Against these organisms its activity is about 10% of that of cephalosporin N.

5. Cephalosporin C is stable in dilute acid but readily inactivated by dilute alkali. It is insensitive to some preparations of penicillinase, but not to others. A new enzyme, C-cephalosporinase, may therefore catalyse its inactivation.

We are grateful to the staff of the Medical Research Council's Antibiotic Research Station, Clevedon, for supplies of crude cephalosporin N and to Dr M. R. Pollock and The Distillers Co. (Biochemicals Ltd.) for gifts of penicillinase. We wish to thank Miss M. Read and Mr O. Boys for expert technical assistance.

REFERENCES

- Abraham, E. P., Newton, G. G. F. & Hale, C. W. (1954). *Biochem. J.* **58**, 94.
 Craig, L. C. & Craig, D. (1950). In *Technique of Organic Chemistry*, vol. 3, chap. 4. Ed. by Weissberger, A. New York: Interscience.
 Craig, L. C., Gregory, J. D. & Hausmann, W. (1950). *Analyt. Chem.* **22**, 1462.
 Florey, H. W. (1955). *Ann. intern. Med.* **43**, 480.
 Hirs, C. H. W., Moore, S. & Stein, W. H. (1952). *J. biol. Chem.* **195**, 669.
 Manson, E. E. D., Pollock, M. R. & Tridgell, E. J. (1954). *J. gen. Microbiol.* **11**, 493.
 Moore, S. & Stein, W. H. (1948). *J. biol. Chem.* **176**, 367.
 Newton, G. G. F. & Abraham, E. P. (1953). *Biochem. J.* **53**, 604.
 Newton, G. G. F. & Abraham, E. P. (1954). *Biochem. J.* **58**, 103.
 Newton, G. G. F. & Abraham, E. P. (1955). *Nature, Lond.*, **175**, 548.
 Olson, B. H., Jennings, J. C. & Juneke, A. J. (1953). *Science*, **117**, 76.
 Pollock, M. R. & Torriani, A. M. (1953). *C.R. Acad. Sci., Paris*, **237**, 276.
 Pollock, M. R., Torriani, A. M. & Tridgell, E. J. (1956). *Biochem. J.* **62**, 387.
 Syngé, R. L. M. (1951). *Biochem. J.* **49**, 642.
 Thompson, H. W., Brattain, R. R., Randall, H. M. & Rasmussen, R. S. (1949). In *The Chemistry of Penicillin*. p. 382. Princeton University Press.

Experiments on the Degradation of Cephalosporin C

BY E. P. ABRAHAM AND G. G. F. NEWTON*
Sir William Dunn School of Pathology, University of Oxford

(Received 26 September 1955)

The isolation and some of the properties of cephalosporin C have been described by Newton & Abraham (1956). This substance behaves as a monoaminodicarboxylic acid and has been given a provisional molecular formula which differs from that assigned to cephalosporin N (D-4-amino-4-carboxy-*n*-butylpenicillin) by only two carbon and two oxygen atoms. Moreover, the infrared spectrum of the substance shows a band at $5.61\text{ }\mu$, which is believed to be characteristic of the penicillins. These facts indicate that cephalosporin C is related to cephalosporin N, but other properties of cephalosporin C, such as its relative stability to acid and to crystalline penicillinase, suggest that it should not be regarded as a true penicillin. This paper gives an account of some of the changes which occur on hydrolysis of cephalosporin C and de-

scribes the isolation, from both cephalosporin C and the material formed from it by hydrogenolysis with Raney nickel, of degradation products showing a positive ninhydrin reaction.

Electrometric titration of cephalosporin C with 0.1N sodium hydroxide revealed the presence of a group with a pK of 9.8. At a pH above 11 the substance underwent rapid hydrolysis, with loss of activity and of the absorption maximum at $260\text{ m}\mu$. (Fig. 1). After standing at pH 12 for 2 hr., back titration showed that two new acidic groups had been formed and that buffering now occurred in the pH range of 4-6 (Fig. 2). The resulting solution gave a positive nitroprusside test, but the titration revealed no new group with a pK that seemed likely to be that of a simple thiol. The characteristic absorption band of cephalosporin C at $260\text{ m}\mu$ had disappeared.

* Member of the Scientific Staff, Medical Research Council.