The Metabolism of [1-¹⁴C]Glucose in an Enzyme System from *Propionibacterium*

By H. G. WOOD,* R. G. KULKA AND N. L. EDSON

Department of Biochemistry, Medical School, University of Otago, New Zealand

(Received 4 November 1955)

Wood, Stjernholm & Leaver (1955) have proposed that propionic acid fermentation may involve the simultaneous occurrence of the Embden-Meyerhof cleavage and a second, undetermined, cleavage. The suggestion was based on the distribution of ¹⁴C in the products obtained by fermenting [1-¹⁴C]-glucose, [2-¹⁴C]glucose, [6-¹⁴C]glucose and [3:4-¹⁴C]-glucose with cell suspensions. No matter which carbon atom of the glucose was labelled, there was considerable radioactivity in all the carbon atoms of the products. Although the widespread distribution of ¹⁴C can probably be accounted for by the Embden-Meyerhof cleavage and subsequent randomization of ¹⁴C via symmetrical C₄-dicarboxylic acids, and possibly via a symmetrical C₃ intermediate, the occurrence of a pathway other than the Embden-Meyerhof cleavage in whole cells is indicated by the following considerations: (1) The carbon dioxide formed from [1-¹⁴C]glucose had a higher activity than any of the carboxyl groups of the end products and a much greater activity than the carbon dioxide from [6-¹⁴C]glucose; (2) the acetate from [1-¹⁴C]glucose had a higher activity than that from [6-¹⁴C]glucose; (3) the peculiar distribution of ¹⁴C in the products from the other types of labelled glucose suggested the existence of an unknown metabolic pathway.

In order to obtain more detailed information about the mechanism it appeared desirable to have a cell-free enzyme preparation which could carry out the whole or part of the propionic acid fermentation. It has been found that such an enzyme system can be obtained by sonic disintegration of propionic acid bacteria. The preparations form propionate, succinate, acetate and carbon dioxide from [1-¹⁴C]glucose and the ¹⁴C is randomized into all positions of the products. The enzyme system of *Propionibacterium shermanii* is remarkable in that it promotes a net uptake of carbon dioxide when glucose is metabolized.

EXPERIMENTAL

The organisms were *Propionibacterium arabinosum* (34W) and *Pr. shermanii* (52W) used previously by Wood *et al.* (1955) and maintained in lactate-glucose stab cultures.

* Fulbright Visiting Professor. Permanent address: Department of Biochemistry, Western Reserve University, Cleveland, Ohio, U.S.A.

These strains were selected for investigation because they give quite different proportions of propionic acid, acetic acid and CO₂ in the fermentation of glucose (Wood & Werkman, 1936). The bacteria were grown for 3-5 days at 30° in 0.05M potassium phosphate buffered medium (pH 6.8), 1 l. of which contained 5.0 g. of glycerol, 5.0 g. of Difco yeast extract and added vitamins (1.0 mg. of calcium pantothenate, 1.0 mg. of thiamine hydrochloride, 0.01 mg. of biotin). For use in experiments they were transferred through 10 and 75 ml. of the glycerol medium and then cultivated in 2 l. conical flasks containing 1.5 l. of medium. The flasks were plugged with cotton wool. The cells were harvested by centrifuging and washed twice with 10 vol. of distilled water. The yields (wet wt.) were about 4 g. of *Pr. arabinosum* and 1 g. of *Pr. shermanii*/l. of medium.

Preparation of cell-free extracts. The wet cells were suspended in 1.5-5 times their weight of ice-cold 0.04-0.1M potassium phosphate buffer, pH 7.0. Batches (20-25 ml.) of the suspensions were treated for 1.5-2 hr. in a Raytheon 9 kc./sec. sonic disintegrator. The temperature of the suspension was maintained at 0-1° by placing the cup assembly in a cold room at 2-3° and circulating ice water through the cooling chamber at the rate of 1.8 l./min. The extracts were clarified by centrifuging for 0.5 hr. at 20 000 g in a refrigerated centrifuge operating at -4°. The brown supernatant solutions were quite clear but showed a pronounced Tyndall effect. When the original suspension contained sufficient phosphate buffer the pH of the extracts was 6.5-6.9. Extracts which were frozen and stored at -40° retained their activity for 8 weeks. There was about 50% loss of activity when they were freeze-dried.

Materials. [1-¹⁴C]Glucose was obtained from the National Bureau of Standards, Washington, D.C.; NaH¹⁴CO₃ was obtained from the Radiochemical Centre, Amersham, England. Adenosine triphosphate (ATP) was either the cryst. disodium salt (Sigma Chemical Co., St Louis, U.S.A.) or a solution of the sodium salt prepared from dibarium ATP (Pabst Laboratories, Milwaukee, U.S.A.) by ion exchange on Amberlite IRC 50 (sodium form). Hexose diphosphate (HDP) was used as a solution of the sodium salt prepared from the dibarium salt (Schwarz Laboratories, Inc., New York, U.S.A.) by removal of Ba on a Dowex 50 column (acid form) and by neutralization with NaOH. Diphosphopyridine nucleotide (DPN) was the chromatographically purified product of Schwarz Laboratories, Inc., New York. Sodium pyruvate was prepared by the neutralization with sodium bicarbonate of pyruvic acid which had been freshly distilled *in vacuo*.

Analytical methods

Carbon dioxide evolution during fermentation, and bound CO₂, were measured manometrically either in conventional Warburg vessels or in 200 ml. vessels. Net

utilization of CO_2 was measured by determining the bound CO_2 at zero time, the CO_2 liberated during fermentation, and the bound CO_2 at the end of the experiment. There was a large evolution of gas from NaHCO_3 when H_2SO_4 was tipped into the vessels, which were not shaken until most of the bound CO_2 had been liberated. When ^{14}C in the CO_2 was to be measured, the vessel was removed after the manometric determination of bound CO_2 , and alkali was added quickly to the centre well and the vessel was closed again in order to collect a sample of the CO_2 .

Glucose was determined colorimetrically by the methods of Somogyi (1945) and Nelson (1944).

Isolation of the products of fermentation. The acidified fermentation mixture was centrifuged and the protein precipitate washed with water. The combined supernatant and washings were extracted continuously with diethyl ether for 2 days. The water-ether mixture in the extraction flask was titrated with NaOH and then evaporated to dryness. The acids were separated on a Celite column (Swim & Krampitz, 1954). The succinic acid fraction was oxidized with acid permanganate to remove any lactic acid that might have been present, and re-extracted with ether and then sublimed at 115° under vacuum. The volatile-acid fractions were distilled in steam and titrated. Carrier acids were added when this was found to be necessary, on the basis of the titrations.

Degradations. Acetic and propionic acids were degraded by the Schmidt reaction (Phares, 1951). Formate was oxidized with mercuric oxide (Osburn, Wood & Werkman, 1933).

Succinate was converted into propionate and CO_2 with a suspension of *Micrococcus lactilyticus*. The bacteria were grown anaerobically at 37° under $\text{N}_2 + \text{CO}_2$ (95:5, v/v) in a medium containing 2.0% Difco peptone, 1.0% Difco yeast extract, 2.0% sodium lactate, 0.5% NaCl , 0.25%

K_2HPO_4 and 0.1 mg. % thiamine hydrochloride. The cells were harvested when they liberated gas vigorously (24 hr.) and were washed twice with 0.1M potassium phosphate buffer, pH 5.9. The succinate was fermented in a 200 ml. Warburg flask under nitrogen at 37° . The reaction mixture contained $\sim 0.03\text{M}$ succinate, 0.1M potassium phosphate buffer (pH 5.9), 0.005M- MgSO_4 , 0.01M- NaF and 0.7% (wet wt.) of cells. The gas evolution was observed manometrically until the rate decreased sharply, when the bound CO_2 was liberated by addition of H_2SO_4 from the side-bulb. A control fermentation with the cells in the absence of succinate gave a small amount of acetate but no propionate. The volatile acids were distilled in steam, and the propionic acid was separated on a Celite column and subsequently degraded by the Schmidt reaction.

The accuracy of each degradation was checked by comparing the sum of the activities in the degradation fractions with the total activity of the compound determined on the CO_2 after combustion with chromic acid (Van Slyke & Folch, 1940).

Measurement of radioactivity. All samples were plated on filter paper as barium carbonate and counted with an end-window Geiger-Müller counter. The counts were corrected for background and for self-absorption (to infinite thinness).

RESULTS

Manometric experiments

The enzyme systems from the two species of *Propionibacterium* had very different properties (Table 1). The extract of *Pr. arabinosum* required a combination of substrates (glucose, HDP and pyruvate), and also ATP and DPN, for maximal activity. Even with these additions (Table 1, no. 7)

Table 1. *Fermentation of glucose, hexose diphosphate and pyruvate by cell-free extracts of Propionibacterium*

Warburg vessels contained 1.60 ml. of medium, pH 6.9; gas, CO_2 ; temp. 30° . Final concentrations of constituents in the fermentation mixtures were: 0.05M glucose, 0.01M HDP, 0.02M pyruvate, 0.001M ATP, 0.0005M DPN, 0.12M- NaHCO_3 . The bacterial extract was added from the side bulb and readings began after 10 min. The extract in each vessel represented 0.16 g. of *Pr. arabinosum* and 0.13 g. of *Pr. shermanii* (both wet wt.). Vessels contained ATP and DPN unless omission is indicated.

Time interval (hr.) ...	Substrate added	$\mu\text{l. of CO}_2$ evolved/hr.*					
		<i>Pr. arabinosum</i> (34 W)			<i>Pr. shermanii</i> (52 W)		
		0-1	1.5-2.5	2.5-3.5	0-1	4-5	27-28
Vessel no.							
1	Glucose	-29	-10	-10	60	50	44
2	HDP	58	37	7	19	11	16
3	Pyruvate	-2	2	0	44	-5	4
4	Glucose, HDP	54	55	5	54	56	42
5	Glucose, pyruvate	8	7	-2	131	57	48
6	HDP, pyruvate	222	32	5	86	13	16
7	Glucose, HDP, pyruvate	668	490	172	162	74	47
8	Glucose, HDP, pyruvate (ATP and DPN omitted)	328	97	0	123	53	39
9	Glucose, HDP, pyruvate (DPN omitted)	605	207	19	162	63	35
10	Glucose (ATP and DPN omitted)	—	—	—	35	31	36
11	None	45	16	11	56	38	15

* The values are the averages of two tests, each with different extracts, and have been corrected for the value obtained in the absence of added substrate.

the activity decreased rapidly, and in 4–5 hr. there was no further formation of CO_2 or acid. In contrast the extract of *Pr. shermanii* fermented glucose at a constant rate in the absence of added ATP and DPN (Table 1, no. 10). The rate was somewhat higher when ATP and DPN were added (Table 1, no. 1), and the activity was greater with the combined substrates (Table 1, no. 7), but after 5 hr. the rate fell almost to that of glucose alone. The enzyme system of *Pr. shermanii* was remarkably stable, fermenting glucose at a constant rate for 30 hr. This extract metabolized pyruvate very actively as a single substrate, all the pyruvate disappearing in 6 hr. (negative nitroprusside test).

With *Pr. shermanii* the sum of the activities in the first hour on the individual substrates (glucose, pyruvate or HDP) was almost equal to the activity on the combination of the three. In the presence of single substrates the extract of *Pr. arabinosum* acted only on HDP, and no combination of two substrates gave as much fermentation as the three combined. When the concentrations of ATP and HDP were double those in Table 1 (no. 7), a higher rate of activity was maintained for a longer time. If more ATP was added to no. 7 when the fermentation rate had fallen to a low value, the activity increased again. Thus it seems probable that one

function of the HDP is to maintain a supply of ATP, the concentration of which limits the fermentation. Pyruvate may serve as a hydrogen acceptor: it is still present when the reaction stops (positive nitroprusside test) and doubling the amount of glucose does not increase the yield of CO_2 .

The extract of *Pr. shermanii* is remarkable in bringing about a net uptake of CO_2 when glucose is fermented. A net uptake of CO_2 in the propionic acid fermentation has been obtained before only with polyols such as glycerol, erythritol, ribitol or mannitol (Wood & Leaver, 1953; Wood & Werkman, 1940). Under conditions corresponding to those of Table 1 (no. 1), 91.1 μmoles of CO_2 were evolved during the fermentation and 78.6 μmoles of bound CO_2 were released on addition of acid at 30 hr. At zero time the bound CO_2 , determined by addition of acid, was 201.3 μmoles . Accordingly, the net uptake was 31.6 μmoles of CO_2 . Since 80 μmoles of glucose were added, 40 μmoles of CO_2 were utilized/100 μmoles of glucose. This substantial uptake of CO_2 is comparable with that previously obtained with glycerol (Wood & Werkman, 1938). The extract of *Pr. arabinosum* did not cause net utilization of CO_2 under the same conditions.

Table 2. *Products of fermentation of [1- ^{14}C]glucose by extracts of Propionibacterium*

Fermentations were conducted in 200 ml. Warburg vessels containing 32 ml. of reaction mixture with the bacterial extract initially in one side bulb and H_2SO_4 in the other: gas, CO_2 ; temp. 30°. Time, 33.5 hr. with *Pr. shermanii* and 6.5 hr. with *Pr. arabinosum*. The final concentrations of the constituents were: with *Pr. arabinosum*, 0.05M [1- ^{14}C]glucose (40300 counts/min./m-mole); 0.01M HDP; 0.02M pyruvate; 0.0019M ATP; 0.00025M DPN; 0.12M- NaHCO_3 ; bacterial extract corresponding to 2.6 g. (wet wt.) of cells; with *Pr. shermanii* an extract corresponding to 2.25 g. (wet wt.) of cells was mixed with the same constituents except HDP, pyruvate and DPN, which were omitted. To determine the initial bound CO_2 control vessels were set up with one-quarter the amount of the constituents, and the bacterial extract and H_2SO_4 were tipped simultaneously. The organic acids in the ether extracts from these control vessels were determined. The values ($\times 4$) for the organic acids in the control vessels were 0.069 m-mole of propionate and 0.043 m-mole of succinate for *Pr. shermanii* extract, and 0.044 m-mole of propionate and 0.011 m-mole of succinate for *Pr. arabinosum* extract. Formate and acetate values were not obtained. No correction has been made for the acids initially present in the extracts. *Pr. shermanii*: recovery of carbon in the products from the fermented glucose (1.27 m-moles) is 72%; oxidation/reduction = 1.28; added radioactivity accounted for, 90%; CO_2 liberated during fermentation (equivalent of acid formation less CO_2 fixed), 2.64 m-moles. *Pr. arabinosum*: added radioactivity accounted for, 85%; CO_2 liberated during fermentation (equivalent of acid formation plus CO_2 produced), 1.58 m-moles.

Product	<i>Pr. shermanii</i>					<i>Pr. arabinosum</i>			
	m-moles	Counts/ min./ m-mole	Total counts/ min.	Percentage added counts/min.	m-moles/ 100 m-moles of glucose fermented	m-moles	Counts/ min./ m-mole	Total counts/ min.	Percentage added counts/min.
Propionate	0.412	14 700	6 050	9.4	32.4	0.591	6 060	3 580	5.6
Acetate	0.434	10 800	4 670	7.3	34.2	0.434	2 460	1 070	1.7
Formate	0.072	2 180	157	0.2	5.7	0.160	195	31	0.05
Succinate	0.85	16 560	14 100	21.9	66.9	0.125	11 000	1 380	2.2
CO_2	-0.155	526*	6 310*	9.8	-12.2	0.40	183*	2 200*	3.4
Carbon in residue from ether extraction	9.54	2 780	26 500	41.2	—	11.4	4 080	46 500	72.2

* Approximately 12 m-moles of gaseous CO_2 , NaHCO_3 and dissolved CO_2 were estimated to be present in the vessels after fermentation.

The addition of sodium fluoride (0.03M final concentration) inhibited the fermentations conducted with *Pr. shermanii* and *Pr. arabinosum* under the conditions of Table 1, in nos. 1 and 7 respectively. Carbon dioxide evolution was almost completely inhibited from the beginning in the *Pr. shermanii* fermentation; and the inhibition was about 50 % with *Pr. arabinosum* in the first 35 min. (observations not continued).

Products from [1-¹⁴C]glucose and the distribution of ¹⁴C

Fermentations of [1-¹⁴C]glucose were carried out with extracts of *Pr. arabinosum* and *Pr. shermanii*, HDP and pyruvate being present in addition to glucose in the former extract. The conditions of the experiments, yields of products and the recovery of ¹⁴C are shown in Table 2. The yield of products per 100 m-moles of fermented glucose has been calculated for the *Pr. shermanii* fermentation in order to present the data in a form comparable with that of previous experiments. Recovery of the carbon of the fermented glucose was only 72 % and the oxidation/reduction balance was 1.28. Probably some product other than those determined was formed by the extract. The unfermented glucose (0.33 m-mole) was equivalent to 13 300 counts/min., but 26 500 counts/min. were found in the residue from ether extraction. The identity of this unknown product is being investigated. The most noteworthy point regarding the yield of products is the very large quantity of succinate (66.9 m-moles/100 m-moles of glucose), which is much greater than the amount (15 m-moles/100 m-moles of glucose) found by Wood & Werkman (1936) with proliferating cells of *Pr. shermanii*. The high yield of succinate in the enzymic fermentation accompanies the net utilization of CO₂ (12.2 m-moles/100 m-moles of glucose). A similar relationship was observed previously in glycerol fermentations (Wood & Werkman, 1938). It is to be noted also that 9.8 % of the added ¹⁴C was converted into CO₂, showing that the enzyme system, like the intact cells, converts C-1 of glucose into CO₂.

The extract of *Pr. arabinosum* converted much less of the ¹⁴C of glucose into propionate, acetate, succinate and CO₂ than the extract of *Pr. shermanii* (13.0 % compared with 48.6 %). Apparently a considerable portion of the products arose from HDP and pyruvate, and much of the glucose carbon remained in the residue after ether extraction (72 % of the ¹⁴C), possibly as phosphate esters.

The distribution of ¹⁴C in the products is shown in Table 3, in which data obtained by Wood *et al.* (1955) with resting cells have been included for comparison. It will be seen that the enzyme preparations randomize ¹⁴C from C-1 of glucose into all positions of the products. Any comparison

Table 3. *Distribution of ¹⁴C in the products of fermentation of [1-¹⁴C]glucose*

		Specific activity as percentage of the specific activity in C-1 of glucose											
		Propionate				Succinate				Acetate			
Organism	Formate	Com-bustion (C × 3)			Com-bustion (C × 4)			Com-bustion (C × 2)			Com-bustion (C × 2)		
		CH ₃ -	CH ₃ - CO ₂ H	Sum	CH ₃ -	CH ₃ - CO ₂ H ₂	Sum	CH ₃ -	CH ₃ - CO ₂ H	Sum	CH ₃ -	CH ₃ - CO ₂ H	Sum
<i>Pr. shermanii</i>	CO ₂	17.2	18.3	35.5	16.4	2.4	37.6	14.1	12.3	26.4	14.1	12.3	26.4
	Formate	5.4	7.7	13.1	10.2	3.3	27.0	4.7	2.7	7.4	4.7	2.7	7.4
<i>Pr. arabinosum</i>	CO ₂	13.2	10.9	24.1	12.7	9.1	31.8	9.5	2.1	11.6	9.5	2.1	11.6
	Formate	10.5	4.9	15.4	7.3	3.7	11.0	15.1	5.2	20.3	15.1	5.2	20.3
<i>*Pr. shermanii</i>	CO ₂	14.2	15.7	29.9	15.5	4.5	20.0	—	—	—	—	—	—
	Formate	—	—	—	—	—	—	—	—	—	—	—	—
<i>*Pr. arabinosum</i>	CO ₂	—	—	—	—	—	—	—	—	—	—	—	—
	Formate	—	—	—	—	—	—	—	—	—	—	—	—

* Data from Table 1 of Wood *et al.* (1955).

between experiments with extracts and intact cells must take cognizance of the differences in experimental conditions. In the experiment with extracts the gaseous phase was CO_2 , whereas with cells it was nitrogen and the CO_2 formed from glucose was absorbed in alkali. Thus the ^{14}C in the CO_2 arising from glucose was greatly diluted by $^{12}\text{CO}_2$ in the extracts, and the specific activity was low. This in turn caused a decrease in the specific activity in the carboxyl groups of the propionate and succinate since CO_2 fixation introduced carbon with low radioactivity. If this difference in the experiments is taken into account, the results with the extracts and the intact cells are quite similar. The radioactivity in the products formed by the *Pr. arabinosum* extract was diluted further by metabolites arising from HDP and pyruvate.

Formate showed considerable radioactivity in the experiment with *Pr. shermanii* extract. Formate is not usually a product in the propionic acid fermentation, but Wood & Leaver (1953) have noticed its formation, especially from polyols.

Fermentation of pyruvate and fixation of CO_2

Fermentation of pyruvate in the presence of radioactive bicarbonate was conducted with an extract of *Pr. shermanii* to determine the products and the extent of CO_2 fixation (Table 4). An odour of diacetyl was noticed and for this reason diacetyl and acetoin were determined together in a portion of the extract by oxidation with ferric chloride, distillation and estimation of the resulting diacetyl (Westerfield, 1945). Usually acetoin and diacetyl are not formed by the propionic acid bacteria, but

a considerable amount was produced with the extract. The other products were propionate, acetate, succinate and CO_2 . The CO_2 production was quite substantial in contrast to the net uptake of CO_2 which occurs in the presence of glucose (Table 2). Nevertheless, there was a large fixation of CO_2 since the propionate contained 14 300 counts/min./m-mole as compared to the CO_2 which had 24 200 counts/min./m-mole at the termination of the experiment. Although the propionate was not degraded it is reasonably certain that all the activity was in the carboxyl group (Wood & Leaver, 1953). This means that about half the carboxyl groups of the propionate must have originated from CO_2 .

DISCUSSION

The results demonstrate that it is possible to extract from *Pr. shermanii* (52W) a soluble-enzyme system, which contains all the enzymes necessary for conversion of glucose or pyruvate into propionate, succinate, acetate and carbon dioxide, and for the production of carbon dioxide from C-1 of glucose. The remarkably widespread distribution of ^{14}C into all the carbon atoms of the products formed by this enzyme system during the fermentation of $[1-^{14}\text{C}]$ glucose is comparable with that observed in the intact cells. The preparation contains a very active system for the fixation of carbon dioxide and for the production of succinate. For this reason it appears to be very suitable for a study of the complete propionic acid fermentation.

Although the extract of *Pr. arabinosum* (34W) has many similar properties, it is inactive with glucose as the sole substrate and demands, in addition, HDP and pyruvate. It appears to contain enzymes destroying the ATP that is needed in the fermentation; and it may also lack some enzymes required for glucose breakdown.

The formation of diacetyl is interesting because the propionic acid fermentation is not known to involve acetaldehyde. Since acetaldehyde is thought to be an intermediate of acetoin formation, it is possible that acetaldehyde or a derivative may participate in the propionic acid fermentation.

Table 4. *Fermentation products formed from pyruvate by an extract of Pr. shermanii (52W) in the presence of $\text{NaH}^{14}\text{CO}_3$*

The fermentation was carried out in a 200 ml. Warburg vessel containing 8 ml. of reaction mixture with the bacterial extract in one side bulb and H_2SO_4 in the other: gas, CO_2 ; temp. 30° , time, 20 hr. Final concentrations: 0.08M pyruvate, 0.0019M ATP, 0.12M- $\text{NaH}^{14}\text{CO}_3$. The extract represented 0.56 g. (wet wt.) of cells. Pyruvate added was 0.64 m-mole. Carbon recovery in the products was 93.7%; oxidation/reduction = 1.10; the CO_2 in the gaseous phase after fermentation contained 24 200 counts/min./m-mole. Propionate contained 14 300 counts/min./m-mole.

Product	Amount formed	
	m-mole	m-moles/ 100 m-moles of pyruvate
Propionate	0.138	21.6
Acetate	0.212	33.0
Formate	0.029	4.5
Succinate	0.052	8.1
CO_2	0.463	72.3
Acetoin + diacetyl	0.066	10.3

SUMMARY

1. Soluble-enzyme systems, free from particulate matter, have been extracted from *Propionibacterium shermanii* and *Propionibacterium arabinosum* by sonic disintegration of the cells.

2. The extracts ferment glucose or pyruvate. The fermentation of glucose by *Pr. shermanii* extract is unique in that there is a net uptake of carbon dioxide.

3. The products of fermentation of $[1-^{14}\text{C}]$ -glucose have been isolated and degraded. The enzyme system randomizes ^{14}C into all positions of the products. Carbon dioxide is produced from C-1 of glucose, and this may occur by some pathway additional to the Embden-Meyerhof cleavage.

REFERENCES

- Nelson, N. (1944). *J. biol. Chem.* **153**, 375.
 Osburn, O. L., Wood, H. G. & Werkman, C. H. (1933). *Industr. Engng Chem. (Anal.)*, **5**, 247.
 Phares, E. F. (1951). *Arch. Biochem. Biophys.* **33**, 173.
 Somogyi, M. (1945). *J. biol. Chem.* **160**, 61.
 Swim, H. E. & Krampitz, L. O. (1954). *J. Bact.* **67**, 419.
 Van Slyke, D. D. & Folch, J. (1940). *J. biol. Chem.* **136**, 509.
 Westerfield, W. W. (1945). *J. biol. Chem.* **161**, 495.
 Wood, H. G. & Leaver, F. W. (1953). *Biochim. biophys. Acta*, **12**, 207.
 Wood, H. G., Stjernholm, R. & Leaver, F. W. (1955). *J. Bact.* **70**, 510.
 Wood, H. G. & Werkman, C. H. (1936). *Biochem. J.* **30**, 618.
 Wood, H. G. & Werkman, C. H. (1938). *Biochem. J.* **32**, 1262.
 Wood, H. G. & Werkman, C. H. (1940). *Biochem. J.* **34**, 7.

The Influence of Acetamide on Citrate Accumulation after Fluoroacetate Poisoning

By S. GITTER*

Department of Biochemistry, University of Oxford

(Received 28 September 1955)

Up to 1949 only symptomatic therapy against fluoroacetate poisoning had been attempted, but no satisfactory results could be obtained by several workers. Sedatives and anticonvulsants gave no protection (Foss, 1948).

Bartlett & Barron (1947) and Saunders & Stacey (1948) considered the action of fluoroacetate to be a specific inhibition of acetate metabolism; this led Chenoweth, Scott & Seki (1949) to ascertain whether acetate would protect the animal against fluoroacetate, but little or no protection of the intact rabbit could be obtained by them even after very high doses of sodium acetate. Only by combining acetate with ethanol could Tourtelotte & Coon (1949) obtain protection in mice, and good results were also reported (Hutchens, Wagner, Podolsky & McMahon, 1949) by the use of ethanol alone. Chenoweth, Kandel, Johnson & Bennett (1951) then reported monoacetin as the most efficient protective agent against fluoroacetate poisoning.

Gitter, Blank & Bergmann (1953*a*) showed that acetamide protected white rats against fluoroacetate poisoning when given simultaneously or up to 24 hr. before the dose of fluoroacetate. Acetamide did not protect when given 30 min. or more after the administration of fluoroacetate, even before the convulsions had started.

The protective results obtained suggested that acetamide might belong to those two-carbon sub-

stances, which, according to Peters (1952*a*), penetrate into the mitochondria and are consequently able to interfere with the formation of the fluorotricarboxylic acid. The influence of acetamide on the accumulation of citrate after the administration of fluoroacetate has been studied. The effects of monoacetin (Chenoweth *et al.* 1951), propionamide and butyramide on citrate accumulation were also tested. A related problem was the low toxicity which we had observed for fluoroacetamide (Gitter, Blank & Bergmann, 1953*b*), and the effect of this substance on citrate accumulation has also been examined.

EXPERIMENTAL

Materials

Acetamide. Acetamide (B.D.H.) was recrystallized and an aqueous solution neutralized by addition of 0.1*N*-NaOH. The neutralized solution, containing 250 mg. of acetamide/ml., was injected subcutaneously to rat in doses of 125 mg./100 g. body wt.

Sodium fluoroacetate. This was prepared in aqueous solutions of 5 and 10 mg./ml.; 0.1 ml./100 g. body wt. was injected intraperitoneally.

Monoacetin. This was given undiluted by subcutaneous or intraperitoneal injection in a dose of 0.5 ml./100 g.

Propionamide and butyramide (a mixture of equal quantities of *n*- and *iso*-butyramide). These were injected in a similar manner and dosage as for acetamide (125 mg./100 g. body wt.).

White male rats, weighing 120–240 g., and female guinea pigs were used. The animals were killed by decapitation,

* Present address: The Rogoff Medical Research Institute, Beilinson Hospital, Petah Tiqva, Israel.