- Liébecq, C. & Peters, R. A. (1949). *Biochim. biophys. Acta*, **2**, 215.
- Lindenbaum, A., White, M. R. & Schubert, J. (1951). *J. biol. Chem.* **190**, 585.
- Lotspeich, W. D., Peters, R. A. & Wilson, T. H. (1951). *J. Physiol.* **115**, 24.
- Malloy, S. & Tepperman, J. (1953). *Fed. Proc.* **12**, 1143.
- Peters, R. A. (1952a). *Brit. med. J.* **2**, 1165.
- Peters, R. A. (1952b). *Proc. Roy. Soc. B*, **139**, 143.
- Peters, R. A., Wakelin, R. W., Buffa, P. & Thomas, L. C. (1953). *Proc. Roy. Soc. B*, **140**, 497.
- Peters, R. A. & Wilson, T. H. (1952). *Biochim. biophys. Acta*, **9**, 310.
- Potter, V. R., Busch, H. & Bothwell, J. (1951). *Proc. Soc. exp. Biol., N.Y.*, **76**, 38.
- Pucher, G. W., Sherman, C. C. & Vickery, H. C. (1936). *J. biol. Chem.* **113**, 235.
- Saunders, B. C. & Stacey, G. J. (1948). *J. chem. Soc.* p. 1773.
- Saunders, B. C. & Stacey, G. J. (1949). *J. chem. Soc.* p. 773.
- Tourtelotte, W. W. & Coon, J. M. (1949). *Fed. Proc.* **8**, 339.

Studies of the Copper Compounds in Aqueous Extracts of Herbage

By C. F. MILLS

The Rowett Research Institute, Bucksburn, Aberdeenshire

(Received 29 June 1955)

In a previous publication (Mills, 1955a) it has been shown that the physiological response of the copper-deficient rat may be used to compare the availability of copper in different dietary supplements. The results of this work indicated that the copper of an aqueous extract of freeze-dried herbage was more readily utilized for haemoglobin formation, melanogenesis, growth and liver-copper storage than an equivalent quantity of copper administered as cupric sulphate. The results of earlier work (Mills, 1954) suggested that much of the copper in herbage exists as organic complexes, but with the methods employed it was not possible to determine the contribution of ionic copper to the total copper content of the soluble components of herbage. The present investigation was designed to determine whether the water-soluble component of herbage which promotes rapid recovery of the copper-deficient rat contains copper mainly as ionic salts or as organic complexes which may or may not be broken down to liberate the free ion in the digestive tract.

EXPERIMENTAL

Electrodialysis of herbage extracts. A sample (20 g.) of milled, freeze-dried herbage was extracted with 200 ml. of glass-distilled water by stirring for 18 hr. at 37°. The mixture was filtered through muslin and the filtrate centrifuged at 8300 g for 1 hr. Before electrodialysis the supernatant was concentrated to 50 ml. by evaporation in a freeze drier. The apparatus described by Theorell & Åkeson (1943) was modified by substituting cellophan and filter-paper membranes for the parchment membranes used by these workers. This modification was necessary to prevent complex-formation between the Cu^{2+} ions of the dialysate and free amino and carboxyl groups of parchment protein. A further modification was the introduction of a fourth dialysis cell between the cell containing the sample

and the catholyte cell (Fig. 1). Dialysis cells were constructed of $\frac{1}{8}$ in. Perspex and were connected by $\frac{1}{8}$ in. diam. holes in which were placed the membranes interposed between $\frac{1}{8}$ in. rubber gaskets. Electrodes were of 22 s.w.g. Pt wire wound in $\frac{1}{8}$ in. diam. spirals. Cooling water was circulated through glass U tubes in each cell and the cell contents were stirred with a stream of N_2 . Power supplies were from a stabilized power unit of maximum output 100 mA at 250 v. Electrodialysis was carried out for about 9 hr., the current being restricted to 75 mA by a variable rheostat to avoid overheating the cell contents. After this period the current rapidly fell to about 3 mA, indicating that ion transport had virtually ceased. Determinations of pH on cell contents were carried out at intervals during electrodialysis. Cell contents were evaporated to dryness before Cu analysis.

Copper analyses. These were carried out by an adaptation of the method of Eden & Green (1940). The yellow Cu diethyldithiocarbamate complex was extracted into 2 ml. of amyl alcohol at 20° and the absorption at 432 m μ . was determined by a Beckman spectrophotometer.

Fig. 1. Four-compartment electrodialysis unit.

Table 1. *Electrodialysis of a concentrated aqueous extract of freeze-dried herbage*

Time (min.)	Cell 1 (anode)	Cell 2 (sample cell)	Cell 3	Cell 4 (cathode)	Current (mA)
	(i) pH of cell contents during electrodialysis				
10	3.2	4.3	5.8	5.8	7
25	3.0	4.1	6.2	8.1	15
60	2.0	3.4	8.5	9.0	54
120	<2	3.2	8.0	9.5	75
480	<2	2.8	8.0	>10	4
	(ii) Migration of copper during electrodialysis				
Copper distribution after electrodialysis ($\mu\text{g. of Cu in cell}$)	0.20	6.92	0.19	0.26	—
	Total Cu in sample electrodialysed = 7.42 $\mu\text{g.}$ Total Cu recovered = 7.57 $\mu\text{g.}$				

Preparation and use of radioactive copper. Radioactive Cu was received in the form of irradiated Specpure foil. A sample (0.04 g.) was dissolved in 1 ml. of 70% (w/w) HNO_3 (redist. A.R.), and the excess of acid neutralized with 2N-NaOH (A.R.) with vigorous stirring to prevent precipitation of $\text{Cu}(\text{OH})_2$. This solution was diluted and a portion containing 10 $\mu\text{g.}$ (^{64}Cu + carrier Cu) was added to the freeze-dried concentrate corresponding to 150 ml. of the original aqueous extract of herbage. After dilution to 50 ml. the product was electrodialysed for 9 hr. After wet-ashing with nitric, sulphuric and perchloric acids the ^{64}Cu activity of cell contents was determined by means of a 10 ml. capacity Mullard liquid counter with Dynatron scaler unit.

Uptake of copper by a cation-exchange resin from herbage extracts after in vitro peptic digestion. Samples (5 ml.) of an aqueous extract of herbage prepared as described above were incubated for 3 hr. at 37° with 0.005 g. of pepsin (B.D.H.) at various pH values after the addition of 2N-HCl. After cooling to 18°, 2 g. (wet weight) of Zeo-Karb 215 (Na activated) resin was added to each digest and the pH adjusted to its original value by the addition of further acid. After being allowed to stand with frequent stirring for 1 hr. the supernatant liquor was decanted off and the resin washed with 30 ml. of glass-distilled water. The supernatant and resin washings were pooled for Cu analysis. To study the uptake of the free Cu^{2+} ion by Zeo-Karb 215 (Na) under similar conditions of pH, 1.85 $\mu\text{g.}$ of Cu^{2+} as $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$ (A.R.) was added to 5 ml. samples of the catholyte solution obtained by the electrodialysis of an aqueous extract of herbage; these solutions were then subjected to resin treatment as described above.

pH determinations. All pH determinations were carried out with the glass electrode.

RESULTS

Electrodialysis of aqueous extracts of herbage. The results of a typical electrodialysis of a concentrated aqueous extract of herbage are presented in Table 1. Only 0.26 $\mu\text{g.}$ of the copper in the herbage preparation migrated through the toughened filter paper and cellophan membranes into the cathode compartment. A further 0.19 $\mu\text{g.}$ of the copper migrated towards the cathode, but was prevented by the cellophan membrane from entering the cathode

Table 2. *Partition of radioactive copper by electrodialysis of a herbage extract treated with 10 $\mu\text{g.}$ of ^{64}Cu*

	Cell pH	^{64}Cu content of cell ($\mu\text{g.}$)
Cell 1 (anode)	<2	0.06
Cell 2 (sample cell)	3.1	9.85
Cell 3	7.5	0.03
Cell 4 (cathode)	>10	0.04

Recovery of added ^{64}Cu , 99.8%.

compartment. Other ions freely migrated during electrodialysis, as was shown by the changes in pH of the contents of cathode and anode compartments. Separate experiments have shown that the free Cu^{2+} ion in the absence of herbage extract readily migrates to the cathode, the migration of 50 $\mu\text{g.}$ of copper being complete within 9 hr.

These results show that cationic copper is not readily liberated from the herbage extract, despite the low pH attained during the electrodialysis. Migration of copper into the anode compartment, presumably in the form of negatively charged complex ions, accounted for about 2.7% of the copper present in the original material. However, the substitution of the negatively charged cellophan membranes for the ampholytic parchment used by Theorell & Åkeson (1943) may prevent the efficient separation of complexes of weak negative charge from neutral materials. Thus it is probable that the material remaining in the sample cell after electrodialysis may contain some copper in the form of complexes of negative charge in addition to neutral complexes.

Effect of adding Cu^{2+} ions to an aqueous extract of herbage. The uptake of free Cu^{2+} ions by the water-soluble components of herbage was investigated by adding radioactive copper, as $^{64}\text{Cu}(\text{NO}_3)_2$, to a concentrated aqueous extract of herbage. The product was electrodialysed for 9 hr. The results are presented in Table 2. They demonstrate the avidity of the water-soluble components of herbage for ionic

copper. In the untreated herbage extract a considerable excess of free metal-binding ligands must be present, for, in this experiment, the copper content of the extract was more than doubled by the addition of the radioactive cupric ion, yet these ions were rapidly removed by complex formation and only 0.4% of the added copper migrated into the cathode cell.

Uptake of copper by Zeo-Karb 215 (Na) from in vitro peptic digests of the water-soluble components of herbage. In earlier work (Mills, 1955b) the uptake of copper by columns of the cation-exchange resins Zeo-Karb 215 and 226 (H^+ -activated forms) from the water-soluble components of herbage was examined. It was found that under moderately acid conditions (pH range 2.6–4.9) only a very small proportion of the total copper was retained by ion exchange with either resin.

In the present investigation the uptake of copper by Zeo-Karb 215 (Na) under physiologically acid conditions after *in vitro* incubation of an aqueous extract of herbage was examined. The results are presented in Fig. 2. It is seen that above pH 2.5 less than 10% of the copper in the herbage extract was retained by the resin (curve A). Under more acid conditions copper uptake increased, until at pH 0.8 68% was retained. The gradual decline in copper uptake as the pH was further reduced is apparently the result of competition at ion-exchange sites by the other freely soluble cations of the herbage extract, as is seen on studying the

uptake by the resin of Cu^{2+} ions added to the catholyte obtained from electrodialysis of a herbage extract. Thus above pH 1 more than 90% of the ionic copper is retained, but under more acid conditions the efficiency of the resin rapidly falls (curve B). In the absence of the other soluble cations of herbage, complete uptake of added Cu^{2+} ions is readily obtained even from N-HCl.

DISCUSSION

The transport of copper to the cathode of an electrodialysis cell does not show unequivocally that this element is in the form of the free Cu^{2+} ion. Cathode-migrating species in electrodialysis include the free ion and positively charged stable complexes. Both these components would freely penetrate the filter-paper membrane on the cathode side of the sample cell in the electrodialysis unit. It is possible that positively charged complexes of large molecular size, although migrating in the direction of the cathode, may not be able to diffuse through the cellophan membrane and into the cathode compartment. Thus in this experiment about 5% of the total copper migrated towards the cathode, but the components migrating into the cathode compartment accounted for only 2.5% of the copper. From these results there is evidence to suggest that in an aqueous extract of herbage a small quantity of copper is present as the free ion or as positively charged complexes, but the greater part of the copper is in the form of neutral or negatively charged complexes which do not readily liberate the free ion under moderately acid conditions.

These suggestions are supported by the results of studies with Zeo-Karb 215 (Na). The soluble organic complexes of herbage appear to be stable above pH 2.5. Although a small proportion of the total Cu was retained by the cation-exchange resin above this pH, the retention appeared to be independent of pH; this suggests that the loss was incurred through adsorption of material in the resin matrix rather than through removal of copper by ion exchange.

SUMMARY

1. Electrodialysis of aqueous extracts of freeze-dried herbage in a modified Theorell-Åkeson four-compartment cell liberated only a small proportion of the copper in the extracts as the free ion. The greater part of the copper was present in the form of neutral or anionic complexes.

2. The avidity of the complex-forming ligands of an aqueous extract of herbage for added Cu^{2+} ions was demonstrated by adding radioactive copper to the extract before electrodialysis.

Fig. 2. Uptake of Cu by Zeo-Karb 215 (Na) from an aqueous extract of herbage incubated *in vitro* + pepsin and from an electrodialysis catholyte of herbage containing added Cu^{2+} ions as a function of pH. Curve A, ϕ — ϕ , residual Cu content of herbage extract after resin treatment (three determinations to each point; spread indicated by vertical line). Curve B, $\times$ — $\times$, residual Cu content of catholyte + Cu^{2+} ions after resin treatment.

3. The organic complexes of copper in these extracts appeared to be stable in the presence of a cation-exchange resin after *in vitro* incubation at pH values above 2.5. Below pH 2.0 decomposition of complexes was appreciable.

I wish to thank Dr J. Duckworth for his interest and encouragement and Dr R. L. M. Synge for advice on electrodialysis techniques. Thanks are due to Miss Grace Murray and Mr A. Forbes for skilled technical assistance.

REFERENCES

- Eden, A. & Green, H. H. (1940). *Biochem. J.* **34**, 1202.
 Mills, C. F. (1954). *Biochem. J.* **57**, 603.
 Mills, C. F. (1955a). *Brit. J. Nutr.* **9**, 398.
 Mills, C. F. (1955b). Ph.D. Thesis: University of London.
 Theorell, H. & Åkeson, A. (1943). *Ark. Kemi Min. Geol. A*, **16**, no. 8.

The Dietary Availability of Copper in the Form of Naturally Occurring Organic Complexes

By C. F. MILLS

The Rowett Research Institute, Bucksburn, Aberdeenshire

(Received 29 June 1955)

In the preceding paper (Mills, 1956) it was shown that the greater part of the copper in an aqueous extract of herbage was in the form of complexes which appear to be stable above pH 2.5. Earlier it had been found that such extracts promoted a more rapid recovery of the copper-deficient rat than a corresponding quantity of copper administered as an inorganic supplement (Mills, 1955).

The object of the present work was to investigate the availability of copper in herbage extracts treated with a cation-exchange resin to remove traces of ionic copper and copper from complexes of low stability and to determine whether the rapid response to such preparations in earlier biological trials was due to either: (1) the efficient utilization of any ionic copper present in the original extract or liberated from complexes during gastric digestion, or (2) the utilization of stable organic complexes of copper.

The use of the copper-deficient rat to determine the biological availability of dietary copper has been described elsewhere (Mills, 1955). The technique consists in depleting the weanling black-hooded rat of liver-copper reserves on a copper-deficient basal diet supplemented with vitamins and with a purified-iron salt. As the animal becomes copper-deficient a marked anaemia develops, with severe depigmentation of hair and failure of growth. The effect of a dietary supplement containing copper on haematopoiesis, melanin-pigment regeneration, growth and liver-copper storage is taken as an index of the biological availability of copper in the test material.

EXPERIMENTAL

Preparation of 'de-ionized' herbage extract. [The term 'de-ionized' herbage extract is in this paper used to indicate a herbage extract treated with Zeo-Karb 215 (H) to remove

ionic copper.] Milled, freeze-dried herbage (200 g.) was stirred vigorously for 18 hr. with 2 l. of glass-distilled water at 37°. Coarse solid material was filtered off with muslin and the filtrate centrifuged at 1500 g for 1 hr. at -0°. A 300 mm. x 18 mm. column of Zeo-Karb 215 was activated with 4 vol. of 2N-HCl (A.R.) and washed with glass-distilled water until the eluate was chloride-free. The centrifuged herbage extract was percolated through the cation-exchange resin at the rate of 5 ml./minute. The first samples of eluate were strongly acid (pH < 1), but after about 45 ml. of eluate had been run off the pH rose rapidly to 3.4 and the collection of a bulk sample was commenced. This extract was freeze-dried and then reconstituted with glass-distilled water so that 0.5 ml. of concentrate contained 3.0 µg. of Cu. This concentrate was stored under N₂ at +1°.

Production of copper-deficient rats. Rats of the Rowett Institute strain of black-hooded Lister were used for this work. (i) Pre-weaning treatment: fresh whole milk was gathered into acid-washed glass vessels and was supplemented with the following vitamins (mg./l.) before feeding *ad lib.* to dam and litter: thiamine 0.3, riboflavin 0.4, Ca pantothenate 2.0, folic acid 0.2, nicotinic acid 0.2, *p*-aminobenzoic acid 0.2 (added in aqueous solution); vitamin A (alcohol) 0.8, α -tocopherol acetate 5.0 (added in arachis oil). This supplemented milk was shaken vigorously before use to ensure uniform dispersion of arachis oil containing fat-soluble vitamins. The dam was removed to a separate cage for three 1 hr. periods daily for the feeding of a solid diet. (ii) Post-weaning depletion diet: a solid basal diet of the following composition was fed at the rate of 6 g./rat/day for the first two weeks after weaning and 10 g./rat/day thereafter: roller-dried full-cream milk (low Cu batch) 95%, sucrose (A.R.) 5%. To this the following water-soluble vitamins (mg./kg.) were added: thiamine 5.0, riboflavin 10.0, pyridoxine 5.0, nicotinic acid 20.0, Ca pantothenate 50.0, *p*-aminobenzoic acid 10.0, folic acid 11.0, inositol 400.0. The copper content of this diet was 0.18 µg./g. of D.M. A purified FeCl₃ supplement was administered orally to give 0.5 mg. of Fe²⁺/rat/day. The following fat-soluble vitamins were administered orally in 0.2 ml. of arachis oil: vitamin A (alcohol), 37 µg./rat/day;