following results were obtained. Rugulosin concentrations, 1:50 000, no growth; 1:100 000, partial inhibition, growth retarded and very thin; 1:200 000, partial inhibition; 1:400 000, normal growth.

SUMMARY

1. Two crystalline colouring matters—one red and the other yellow—have been isolated from the mycelium of laboratory cultures of a number of strains of *Penicillium rugulosum* Thom and from one strain of *P. wortmanni* Klöcker.

2. The red colouring matter is skyrin, $C_{30}H_{18}O_{10}$, previously reported from *P. islandicum* Sopp.

3. The yellow colouring matter, which has not been described previously and has been named rugulosin, $C_{30}H_{22}O_{10}$, forms yellow prisms, m.p. about 293° (decomp.) which are strongly dextro-rotatory. It has a markedly specific antibacterial activity and is moderately antifungal.

4. Good yields of chrysophanol (4:5-dihydroxy-2-methylanthraquinone) and emodin (4:5:7-trihydroxy-2-methylanthraquinone) have been obtained by degradation of rugulosin.

5. Treatment of rugulosin with concentrated sulphuric acid at 140° gives aurantio-rugulosin, $C_{30}H_{18}O_8$, which is believed to be a substituted dianthraquinone.

6. The structural relationship of rugulosin and rubroskyrin, $C_{30}H_{22}O_{12}$ from *P. islandicum*, is discussed as well as that of aurantio-rugulosin and iridoskyrin, $C_{30}H_{18}O_{10}$, from *P. islandicum*.

7. It is believed that rugulosin and rubroskyrin belong to a new class of colouring matters which, while not anthraquinones themselves, readily give rise to well known polyhydroxyanthraquinones.

REFERENCES

- Adkins, H. & Connor, R. (1931). *J. Amer. chem. Soc.* **53**, 1091.
 Dacre, J. C. (1950). The Biochemistry of *Penicillium rugulosum* Thom and certain other species of moulds, p. 1. Ph.D. Thesis, University of London.
 Howard, B. H. & Raistrick, H. (1949). *Biochem. J.* **44**, 227.
 Howard, B. H. & Raistrick, H. (1950). *Biochem. J.* **46**, 49.
 Howard, B. H. & Raistrick, H. (1954a). *Biochem. J.* **56**, 56.
 Howard, B. H. & Raistrick, H. (1954b). *Biochem. J.* **57**, 212.
 Kinney, C. R. & Langlois, D. P. (1931). *J. Amer. chem. Soc.* **53**, 2189.
 Peterson, V. L. & West, E. S. (1927). *J. biol. Chem.* **74**, 379.
 Raistrick, H. (1949). *Proc. Roy. Soc. A*, **199**, 158.
 Raper, K. B. & Thom, C. (1949). *A Manual of the Penicillia*, p. 584. Baltimore: Williams and Wilkins.
 Shibata, S., Tanaka, O., Chihara, G. & Mitsuhashi, H. (1953). *Pharm. Bull. Japan*, **1**, 302.

The Reactivity of the Cystine Linkages in Wool towards Reducing Agents

By A. J. FARNWORTH*

Textile Research and Testing Department, Gordon Institute of Technology, Geelong, Victoria, Australia

(Received 24 December 1954)

Of all the various reactive side linkages in wool, none has been studied more extensively than the cystine linkage. This is probably accounted for by the fact that the keratins contain a much higher proportion of this amino acid than other proteins. Possibly because of this there has been a tendency to explain any anomalous physical or chemical behaviour of wool in terms of cystine.

In a very comprehensive series of studies of the reactivity of the cystine linkages in wool, Phillips and his colleagues have examined the behaviour of untreated and modified wools towards sulphites, alkalis, formaldehyde and thioglycolic acid (Elsworth & Phillips, 1938, 1941; Middlebrook & Phillips, 1942a, b, 1947; Cuthbertson & Phillips, 1945; Lindley & Phillips, 1945, 1947; Carter,

Middlebrook & Phillips, 1946; Phillips, 1946.) As a result of this work, the combined cystine of wool has been divided into two fractions of differing reactivity, each fraction being further divided into two subfractions. For convenience, the more reactive fraction has been termed (*A* + *B*) and the less reactive portion (*C* + *D*). By definition, that portion of cystine which reacts with sodium bisulphite solutions to give thiol groups on subsequent hydrolysis is known as fraction (*A* + *B*). Most of this fraction reacts in the cold. When rinsing with water is interposed between bisulphiting and hydrolysis, the residual thiol-S determined in the hydrolysate gives a measure of subfraction *B*, subfraction *A* being obtained by difference.

Fraction (*C* + *D*) is defined as that portion of the cystine which does not give cysteine and S-cysteine

* Present address: C.S.I.R.O. Wool Textile Research Laboratory, Belmont, Geelong, Victoria, Australia.

sulphonate side chains when wool is treated with sodium bisulphite. Subfraction *D* gives combined α -amino acrylic acid very slowly at room temperatures but more rapidly at higher temperatures.

The differences between subfractions *A* and *B* and between subfractions *C* and *D* are not nearly so marked as those between fractions (*A* + *B*) and (*C* + *D*). Similar behaviour has been found to exist with the cystine linkages of proteins other than wool (Lindley, 1948).

Phillips and his colleagues have suggested and rejected several possible explanations of this behaviour, and eventually came to the conclusion that the reactivities of the subfractions are influenced by the nature of the amino acid in the near vicinity of the cystine in the protein chain. Astbury (1942) has deduced that the number of polar and non-polar side chains in wool are approximately equal and, in view of the roughly equivalent proportions of fractions (*A* + *B*) and (*C* + *D*), it has been suggested that fraction (*A* + *B*) is associated with a polar side-chain environment and (*C* + *D*) with non-polar side chains. Concerning the difference between subfractions *A* and *B*, Phillips and his co-workers have suggested that ionizable carboxyl groups may predominate in the environment of subfraction *A*, whilst *B* may be in the vicinity of acid amide side chains. Supporting evidence for this is given by the behaviour of methylated wool which, on treatment with cold solutions of sodium bisulphite, showed that subfraction *B* increased at the expense of subfraction *A* with increasing degree of esterification (Lindley & Phillips, 1947).

More recently, Alexander and his co-workers have investigated the reaction of oxidizing agents with wool and have shown that both the cystine linkages and the tyrosine side chains show varying reactivity towards chlorine, hypochlorous acid, peracetic acid, acid permanganate and alkaline hypochlorite (Alexander, Hudson & Fox, 1950; Alexander & Gough, 1951). They showed that the cystine and tyrosine could be divided into two fractions. In the case of cystine, however, these fractions are not related to those of Phillips. Alexander considered that the different reactivity could not be ascribed to differences in morphological factors and has attempted to explain the results of their work in terms of a two-phase structural system for wool. One phase is considered to be a high molecular weight micellar arrangement of low chemical reactivity. The other phase is a low molecular weight cementing material of high reactivity situated between the micelles.

Schuringa, Schooneveldt & Konings (1951) examined the action of sodium bisulphite and potassium cyanide on wool and concluded that accessibility of the cystine groups was partly responsible for the varying reactivity.

In this present paper the reaction of wool with reducing agents has been further studied in an attempt to find evidence for the influence of polar side chains on the reactivity of the cystine linkages. Two methods of approach have been used: (a) the behaviour of wool towards sodium bisulphite after modification of the cystine linkages themselves; (b) the reactivity of the cystine linkages with various reagents after modification of the polar side chains.

EXPERIMENTAL METHODS

Materials. The wools used were a 64's quality merino and a 58-60's quality crossbred. The tip ends were removed and the root portions cleaned by extraction with ether followed by a large number of changes of distilled water. They were then allowed to condition at 65% relative humidity and 21°.

Treatment with sodium bisulphite solutions. In general the methods of Phillips *et al.* were used (Elsworth & Phillips, 1938, 1941; Middlebrook & Phillips, 1942*a, b*; Carter *et al.* 1946). Treatments were carried out using cold and boiling solutions. In the former case, wool (0.2 g.) was treated with 100 times its weight of 30% (w/v) NaHSO₃ for 18 hr. at 21°, while in the latter, boiling was carried out for 30 min. When fraction (*A* + *B*) was being estimated, the wool was removed, squeezed and hydrolysed in 5*N*-HCl for 4 hr. For determining cystine subfraction *B* (water stable), the bisulphite-treated wool was rinsed in three changes of distilled water (200 ml.) for 10 min. in each change, and then immersed overnight in 2 l. distilled water. After squeezing, it was hydrolysed as above.

Reduction with thioglycollic acid. Wool was reduced by treatment at 35° for 18 hr. with 50 times its weight of 0.5*M* thioglycollic acid brought to pH 4.8 with NaOH (Patterson, Geiger, Mizell & Harris, 1941). The wool was then rinsed in water, ethanol and finally water.

Alkylation of reduced wool. Wool, reduced as above was alkylated with methyl iodide or ethyl iodide by treatment at 35° for 18 hr. with 100 times its weight of a phosphate buffer (0.2*M*-Na₂HPO₄-KH₂PO₄) at pH 7.5, to which had been added methyl iodide (0.12 ml./g. wool) or ethyl iodide (0.16 ml./g. wool). It was then thoroughly rinsed with distilled water.

Cross-linking of reduced wool with alkylene dihalides. Wool, reduced as above, was treated with methylene diiodide (0.16 ml./g. wool) and ethylene dibromide (0.20 ml./g. wool) under the same conditions as those used in alkylation.

Treatment with phenyl isocyanate. Wool, dried in an oven at 105° for 4 hr. and then for 24 hr. over phosphorus pentoxide *in vacuo*, was treated with phenyl isocyanate (2 ml./g. original wool) in anhydrous pyridine (18 ml./g. original wool) for 72 hr. at 70°. The reaction was terminated by addition of water. The wool was then rinsed with ethanol, extracted with ether and rinsed thoroughly in distilled water (Farnworth, 1955).

Acetylation. Wool was boiled in acetic anhydride under reflux for 30 min. followed by prolonged rinsing with water to remove excess reagent.

Treatment with alkali. Wool was treated with 0.1*N*-NaOH (1 l./g. wool) at 22° for 24 hr. followed by prolonged rinsing with distilled water.

Thiol and disulphide estimations. Analyses for thiol-S and disulphide-S were carried out on the hydrolysate by the method of Shinohara (1935*a-c*). The results are expressed as a percentage sulphur on the weight of the oven-dry wool.

RESULTS

Reaction of untreated wool with sodium bisulphite solution. In order to confirm the existence and assess the proportions of the cystine fractions of Phillips in the wool being used, both wools were treated with cold and boiling solutions of sodium bisulphite, with and without subsequent rinsing before hydrolysis.

Table 1 shows the results obtained from analyses of wools before and after treatment with bisulphite and rinsing in various ways.

fraction ($C + D$) is 1.25 % and subfraction C is 0.56 %. These values, together with their relationship to one another and to the total disulphide S, are shown in Table 2.

It is seen that there are slight variations between the various values for the two wools, and also from the 64's Cape merino wool used by Phillips and his colleagues, the figures for which are included for comparison. These latter figures were obtained from the analytical data of Middlebrook & Phillips (1942*b*).

Apart from these slight differences, the only major discrepancy between the various sets of figures is in the relative magnitude and ratio of subfractions A and B . In the present work the ratio A/B is very nearly 3, as against a ratio of 1 for the

Table 1. *The thiol-S and disulphide-S contents of bisulphite-treated wools*

Values are expressed as percentage of dry weight of wool.

Wool	Treatment	Thiol-S	Disulphide-S	Thiol + disulphide-S
58-60's	None	0.06	2.97	3.03
	Cold bisulphite, unrinsed	1.78	0.94	2.72
	Cold bisulphite, rinsed	0.47	2.35	2.82
	Boiling bisulphite, unrinsed	1.81	0.57	2.38
	Boiling bisulphite, rinsed	0.57	1.90	2.47
64's	None	0.11	2.99	3.10
	Cold bisulphite, unrinsed	1.85	0.96	2.81
	Cold bisulphite, rinsed	0.49	2.49	2.98
	Boiling bisulphite, unrinsed	1.93	0.56	2.49
	Boiling bisulphite, rinsed	0.51	2.16	2.67

Table 2. *The relative proportions and relationship of the various cystine subfractions in 64's and 58-60's wool*

For definition of subfractions see text. S values are expressed as percentage of dry wt. of wool.

	58-60's wool	64's wool	64's Cape wool used by Phillips
Fraction ($A + B$)-S	1.78	1.85	1.86
($A + B$)/total disulphide-S	0.59	0.60	0.57
Subfraction A -S (by difference)	1.31	1.36	0.94
Subfraction B -S	0.47	0.49	0.92
A/B	2.8	2.8	1.0
Fraction ($C + D$)-S (by difference)	1.25	1.25	1.39
($C + D$)/total disulphide-S	0.41	0.40	0.43
Subfraction C -S	0.57	0.56	0.67
Subfraction D -S (by difference)	0.68	0.69	0.72
C/D	0.8	0.8	0.9

In calculating the values for the various fractions and subfractions, the more direct method of Lindley (1948) has been used in preference to that of Carter *et al.* (1946), in which the non-disulphide sulphur initially present in the wool is included with fraction ($C + D$), and various other assumptions are made. Such calculations show that for the 58-60's wool, fraction ($A + B$) amounts to 1.78 % S, of which 0.47 % is in subfraction B . Fraction ($C + D$), obtained by difference, is 1.25 % S, of which 0.57 % is in subfraction C . In the case of the 64's wool, fraction ($A + B$) is 1.85 %, subfraction B is 0.49 %,

wool used by Phillips and his co-workers. These workers have pointed out that the difference between A and B is not nearly so marked as that between ($A + B$) and ($C + D$), and the wide discrepancy between their results for the ratio A/B and those of the present work also seems to suggest that there is no clear-cut division between these two subfractions. In fact, it is felt that in the case of the reaction under consideration—namely, that with sodium bisulphite solutions—the difference does not actually exist, because if the rinsing is carried out with running distilled water for 48 hr.,

the residual thiol-S is further reduced to 0.26 % in the case of the 58–60's wool. This gives a value of 0.26 % for subfraction B, 1.52 % for subfraction A and 5.8 for the ratio A/B . Carter *et al.* (1946) showed that the relative magnitudes of A and B vary considerably with the pH and composition of the rinsing solution.

The effect of modification of the cystine linkages on the reaction of wool with sodium bisulphite and thioglycollic acid. Middlebrook & Phillips (1942b) showed that molar solutions of thioglycollic acid at pH 4.5 and 35° reduced the cystine linkages of wool, and that subsequent treatment with cold NaHSO_3 solutions produced no further change. Patterson *et al.* (1941) had shown that increasing the concentration of thioglycollic acid to 2M at pH 4.5 gave very little more reduction, and Phillips and his colleagues came to the conclusion that reduction with thioglycollic acid under these conditions reached a limiting value and was confined to fraction (A + B).

In order to investigate these claims fully, wool has been reduced with thioglycollic acid solutions and then subjected to treatment with NaHSO_3 and also to further reduction with thioglycollic acid. In addition, after the initial reduction, the thiol groups have been alkylated with alkyl halides or cross-linked with alkylene dihalides, and the wool further treated with NaHSO_3 or thioglycollic acid. The results obtained for the thiol-S and disulphide-S of these various wools are shown in Tables 3 and 4.

In calculating the values for fraction (A + B) it has been necessary to make certain corrections in some cases, to allow for the free thiol-S present in the wool after alkylating or cross-linking and before the treatment with bisulphite or the second reduction with thioglycollic acid. Furthermore, during the alkylation or cross-linking procedure, in some cases part of the thiol-S reverts to disulphide-S. This would no doubt be re-converted to thiol-S during subsequent reduction with bisulphite or thioglycollic acid and would lead to an incorrect figure

Table 3. *The thiol-S and disulphide-S of wool treated with bisulphite after modification of the cystine linkages (58–60's wool)*

For details of calculations see text. S values refer to dry wool.

Treatment of wool	Thiol-S (%)	Disulphide-S (%)	Thiol + disulphide-S (%)	Fraction (A + B)-S (%)	(A + B)/total disulphide-S
None	0.06	2.97	3.03	1.78	0.59
Reduced with thioglycollic acid	1.75	1.08	2.83	1.75	0.58
Reduced with thioglycollic acid followed by cold NaHSO_3	2.03	0.76	2.79	2.03	0.67
Reduced with thioglycollic acid followed by boiling NaHSO_3	2.04	0.39	2.43	2.04	0.67
Reduced with thioglycollic acid followed by methylation	0.08	1.17	1.25	—	—
Reduced and methylated followed by cold NaHSO_3	0.61	0.74	1.35	2.19	0.72
Reduced and methylated followed by boiling NaHSO_3	0.99	0.21	1.20	2.59	0.85
Reduced with thioglycollic acid followed by ethylation	0.13	1.37	1.50	—	—
Reduced and ethylated followed by cold NaHSO_3	0.81	0.81	1.62	2.14	0.71
Reduced and ethylated followed by boiling NaHSO_3	1.14	0.35	1.48	2.47	0.82
Reduced with thioglycollic acid followed by cross-linking with methylene diiodide	0.71	1.50	2.21	—	—
Reduced and cross-linked followed by cold NaHSO_3	1.06	0.94	2.10	1.68	0.55
Reduced and cross-linked followed by boiling NaHSO_3	1.37	0.61	1.98	1.99	0.66
Reduced with thioglycollic acid followed by cross-linking with ethylene dibromide	0.16	1.36	1.52	—	—
Reduced and cross-linked followed by cold NaHSO_3	0.69	0.76	1.45	2.00	0.66
Reduced and cross-linked followed by boiling NaHSO_3	0.83	0.60	1.43	2.14	0.71

for 'new' thiol-S unless allowance was made for it. Thus, in Table 3, analysis after cold bisulphite solution has reacted with reduced and methylated wool shows a thiol-S content of 0.61%, of which 0.08% was present after methylation and before treating with bisulphite. Hence the bisulphite treatment has produced an additional 0.53% of thiol-S. However, during methylation it is seen that the disulphide-S rose from 1.08 to 1.17%, indicating that 0.09% thiol-S has been reoxidized to disulphide-S, i.e. 0.09% of fraction (A + B) has reverted to disulphide-S. It is reasonable to assume

that this disulphide-S would once again be reduced to thiol-S during the bisulphite treatment, so that, of the 0.53% additional thiol-S thus produced, 0.09% has already been included in the original (A + B) fraction (1.75%) shown after the initial reduction with thioglycollic acid. Hence the total thiol-S after the complete reduction-methylation-bisulphiting sequence is given by the thiol-S after the initial reduction (1.75%) plus the new thiol-S arising from the action of bisulphite on the disulphide bonds still intact after the initial reduction (0.53 - 0.09 = 0.44%), a total of 2.19%. Fraction (A + B)

Table 4. *The thiol-S and disulphide-S contents of wools reduced with thioglycollic acid after modification of the cystine linkages*

For details of calculations see text. S values refer to dry wool.

Treatment	Thiol-S (%)	Disulphide-S (%)	Thiol + disulphide-S (%)	Fraction (A + B)-S (%)	(A + B)/total disulphide-S
64's wool					
None	0.11	2.99	3.10	1.61	0.52
Reduced with thioglycollic acid	1.75	1.08	2.83	1.75	0.56
Reduced with thioglycollic acid followed by a second reduction with thioglycollic acid	1.86	0.98	2.84	1.86	0.60
Reduced with thioglycollic acid followed by cross-linking with ethylene dibromide	0.14	1.06	1.20	—	—
Reduced and cross-linked followed by a second reduction with thioglycollic acid	0.42	0.82	1.24	2.03	0.65
58-60's wool					
None	0.06	2.97	3.03	1.78	0.59
Reduced with thioglycollic acid	1.66	1.17	2.83	1.66	0.55
Reduced with thioglycollic acid followed by a second reduction with thioglycollic acid	1.89	0.96	2.85	1.89	0.62
Reduced with thioglycollic acid followed by cross-linking with methylene diiodide	0.50	1.38	1.88	—	—
Reduced and cross-linked followed by a second reduction with thioglycollic acid	0.93	0.94	1.87	1.88	0.62
Reduced with thioglycollic acid followed by methylation	0.03	1.13	1.16	—	—
Reduced and methylated followed by a second reduction with thioglycollic acid	0.50	0.50	1.00	2.13	0.70
Reduced with thioglycollic acid followed by ethylation	0.10	1.08	1.18	—	—
Reduced and ethylated followed by a second reduction with thioglycollic acid	0.61	0.60	1.21	2.17	0.72
Reduced with thioglycollic acid	1.75	1.08	2.83	1.75	0.58
Reduced with thioglycollic acid followed by methylation	0.08	1.17	1.25	1.76	0.58
Reduced and methylated followed by a second reduction with thioglycollic acid and a second methylation	0.03	0.77	0.80	2.03	0.67
Three successive alternate reductions and methylations	0.04	0.45	0.49	2.34	0.77
Four successive alternate reductions and methylations	Nil	0.42	0.42	2.41	0.80
Five successive alternate reductions and methylations	Nil	0.31	0.31	2.52	0.83

after the complete sequence is therefore 2.19. Whilst such corrections are not of any great significance in this case, they do assume much greater significance with other treatments. Thus, during cross-linking with methylene diiodide (Table 3) 0.42 % of fraction ($A + B$) thiol-S apparently reverts to disulphide-S, and, in addition, the wool still contains 0.71 % free thiol-S before the treatment with bisulphite is commenced. Hence the thiol-S produced by boiling bisulphite solution additional to that formed during the initial reduction with thioglycollic acid is not 1.37 % as would appear at first glance, but $1.37 - (0.42 + 0.71) = 0.24$ %. Hence the total thiol-S ($A + B$) after the complete reduction-cross linking-bisulphiting sequence is $1.75 + 0.24 = 1.99$ %.

The effect of modification of the polar side chains on the reaction of wool with sodium bisulphite, thioglycollic acid and sodium hydroxide. In view of the suggestion that fraction ($A + B$) of the cystine is in the vicinity of polar side chains which confer enhanced reactivity on these linkages, it would appear that conversion of the polar side chains into less polar or non-polar groups should lead to a decreased reactivity of the ($A + B$) cystine. There is some evidence in favour of this, as Lindley & Phillips (1947) found that the magnitude of fraction

($A + B$) was decreased by deamination and also by acetylation with boiling acetic anhydride.

To investigate this matter further an alternative means of blocking polar side chains was desired, preferably one which did not introduce any coloration of the wool, as any such coloration would lead to interference with the colorimetric method used for analysis of thiol-S and disulphide-S. Such interference resulted when the amino side chains were blocked by treatment with 1-fluoro-2:4-dinitrobenzene, using the conditions recommended by Sanger (1945).

Phenyl isocyanate has been made to react with wool under anhydrous conditions, reaction occurring with virtually all the polar side chains. Furthermore, it has been found that acetylation with boiling acetic anhydride also leads to extensive reaction with the polar side chains (Farnworth, 1955). Hence, wools modified by treatment with phenyl isocyanate and also by acetylation have been examined as to their behaviour towards sodium bisulphite and thioglycollic acid. The effect of sodium hydroxide solution on such modified wools has also been examined, as the cystine linkages in wool are very susceptible to attack by alkali.

The results of these experiments are given in Table 5.

Table 5. *The thiol-S and disulphide-S contents of wools treated with sodium bisulphite, thioglycollic acid and sodium hydroxide after modification of the polar side chains (64's wool)*

S values refer to percentage of dry wt. of wool.

Treatment	Thiol-S (%)	Disulphide-S (%)	Thiol + disulphide-S (%)	Fraction ($A + B$)-S (%)		(A + B)/total disulphide-S	
None	0.11	2.99	3.10	—		—	
Cold bisulphite solution	1.85	0.96	2.81	1.85		0.60	
Boiling bisulphite solution	1.93	0.56	2.49	1.93		0.62	
Thioglycollic acid	1.75	1.08	2.83	1.75		0.56	
0.1N-NaOH (1 l./g. wool) 22°, 24 hr.	Nil	1.10	1.10	—		—	
Acetylated	0.13	2.59	2.72	—		—	
				Max.	Min.	Max.	Min.
Acetylated followed by cold bisulphite solution	1.17	1.62	2.79	1.57	1.17	0.51	0.38
Acetylated followed by boiling bisulphite solution	1.26	1.40	2.66	1.66	1.26	0.54	0.41
Phenyl isocyanate	Nil	2.79	2.79	—	—	—	—
Phenyl isocyanate followed by cold bisulphite solution	0.07	2.77	2.84	0.27	0.07	0.09	0.02
Phenyl isocyanate followed by boiling bisulphite solution	0.49	2.22	2.71	0.69	0.49	0.22	0.16
Phenyl isocyanate followed by thioglycollic acid	0.08	2.72	2.80	0.28	0.08	0.09	0.03
Phenyl isocyanate followed by 0.1N-NaOH (1 l./g. wool) 22°, 24 hr.	Nil	2.79	2.79	—	—	—	—

It will be noticed that a maximum and minimum value is given for fraction ($A + B$) after reduction of the acetylated and phenyl isocyanate-treated wool. This has been done because there is a slight loss of disulphide-S during the acetylation (0.4%) and phenyl isocyanate (0.2%) treatments. Most probably the cystine linkages degraded during these treatments would be from the ($A + B$) fraction, and if this assumption is made, then the amount of disulphide-S so lost must be added to the thiol-S produced during the subsequent reduction with NaHSO_3 or thioglycollic acid, and the maximum values shown in the table have been obtained in this way. On the other hand, if none of the degraded cystine belongs to the ($A + B$) fraction, then the thiol-S contained in the wool after treatment with NaHSO_3 and thioglycollic acid would represent the true ($A + B$) fraction. The minimum values shown in the table have been calculated on this basis.

The reformation of cystine from reduced cystine linkages. Under certain circumstances it has been found possible to induce reduced cystine linkages to revert to cystine. In fact, the original subdivision of fraction ($A + B$) into subfractions A and B was based on the lability to water of the thiol and sodium cysteine sulphonate groups formed during the bisulphiting of wool (Carter *et al.* 1946), as rinsing with water after bisulphiting led to a loss of thiol-S and an increase in disulphide-S in the hydrolysate. These workers have shown that reversion to cystine is at a minimum with acidic rinsing solutions but increases with increasing pH, maximum reversion occurring in an alkaline medium. Speakman (1936) showed that some of the thiol groups produced by reduction of wool with thioglycollic acid could be reoxidized to cystine linkages by means of hydrogen peroxide.

Two other observations worthy of note have been made in connexion with the reversion of broken cystine linkages. (a) When wool reduced with thioglycollic acid is treated with cold NaHSO_3 solution and then rinsed with water according to the conditions outlined in the experimental section, there is a decrease in thiol-S of approximately 0.5%, the disulphide-S increasing accordingly. If boiling NaHSO_3 solution is used followed by rinsing with water, the thiol-S decreases by 1.4%, there being a corresponding increase in the disulphide-S. In this latter case, the thiol-S is decreased from 1.75% in the initially reduced wool to 0.41% in the bisulphited and rinsed wool. (b) When wool reduced with thioglycollic acid is heated in anhydrous pyridine at 70° for 72 hr., the thiol-S is decreased to less than 15% of its initial value, the disulphide-S showing a quantitatively corresponding increase. Apparently, heating under these conditions leads to re-oxidation of the major portion of the thiol groups.

DISCUSSION

Throughout the discussion it has been found convenient to refer to the ($A + B$)-S as a proportion of the total disulphide-S present in the original wool.

The results of the present work are in agreement with the very large amount of evidence put forward by Phillips and his colleagues for the variation in reactivity of the cystine in wool. However, the aim of this work has been to attempt to find convincing evidence for the influence of adjacent amino acids in the protein chain. The bulk of the evidence appears to be against the idea of some cystine linkages being activated by an environment of polar side chains. Thus in Table 4 it is seen that a second successive reduction with thioglycollic acid gives a slight increase in thiol-S, the ($A + B$)/total disulphide-S ratio increasing to about 0.6. However, when the thiol groups are methylated or ethylated prior to a second reduction with thioglycollic acid, there is a more marked secondary reduction, the ($A + B$)/total disulphide-S ratio increasing to 0.7; if repeated successive reduction and methylation is carried out, the ratio rises to 0.83 after five such cycles. At this stage, only 0.31% disulphide-S remains unchanged. Similar behaviour has been observed by Geiger, Kobayashi & Harris (1942). In the original division of the cystine into its various fractions on the basis of the reaction with sodium bisulphite, the 'inert' subfraction C amounted to 0.56%. Hence virtually all the cystine in wool, including the chemically 'inert' portion, can be induced to react with thioglycollic acid under conditions which normally lead to reaction with only slightly more than 50% of the total disulphide-S, provided that the thiol groups are progressively alkylated. A possible explanation could be that a state of equilibrium is reached between a reversible oxidation-reduction system in solution and a similar system in the wool. The equilibrium point would be altered either by removal of the disulphide formed from thioglycollic acid by renewal of the reduction liquor, or by alkylation of the thiol groups in the wool. An alternative explanation is that breakdown of some of the disulphide groups renders the remaining groups more accessible to the reagent, allowing their reduction to proceed. The results of Table 4 seem to point to the second explanation as being more probable, as cross-linking of the thiol groups with a methylene group before the second treatment with thioglycollic acid prevents any additional reduction from taking place. When the cross-linking group is increased in size, some additional reduction does take place, but not to the same extent as when alkylation is used to block the thiol groups. Cross-linking by means of alkylene dihalides would be as effective as alkylation in upsetting an oxidation-reduction system.

Similar but far more convincing evidence for this behaviour is obtained from the results in Table 3. When wool in which the cystine linkages of fraction ($A+B$) have been converted into thiol groups by reduction with thioglycollic acid is treated with bisulphite solutions, there is a slight increase in the ratio ($A+B$)/total disulphide-S. However, when the thiol groups are alkylated before treating with bisulphite, there is a marked increase in the ratio following treatment with cold bisulphite solutions and a very considerable increase when treated with boiling bisulphite solutions. Thus when the thiol groups after thioglycollic reduction are methylated, boiling bisulphite increases the ($A+B$)/total disulphide-S ratio from 0.58 to 0.85. The residual unchanged disulphide-S after such treatment is only 0.21 %, a value considerably lower than the 0.57 % 'inert' subfraction C of the original subdivision. Furthermore, the thioglycollic reduction-alkylation-bisulphite procedure leads to the formation of only one reaction product, namely thiol and cysteine sulphonate groups, the latter being converted into thiol groups on hydrolysis prior to analysis. The characteristic properties of fraction ($C+D$) on which the original division was based were that it was less reactive than fraction ($A+B$) and followed a different reaction mechanism, giving rise to combined α -aminoacrylic acid with complete loss of sulphur when treated with boiling bisulphite solutions. Hence, the present results are in direct contrast with this, all the subfractions having reacted according to one mechanism to give thiol groups, irrespective of whether the bisulphite solutions used are cold or boiling.

When the thiol groups resulting from the initial reduction are cross-linked through a methylene group, subsequent bisulphiting is ineffective except in boiling solution, and even under these circumstances the extent to which reaction occurs is not very great. When a longer cross-linkage is inserted by using ethylene dibromide, reaction with bisulphite occurs to a degree intermediate between alkylation, which leaves the cystine linkages permanently broken, and cross-linking through the short methylene bridge. It is difficult to explain these results other than on the grounds of increased accessibility of the ($C+D$) disulphide-S after rupture of the ($A+B$) cystine linkages.

It appears, therefore, that the various subfractions of the cystine in wool are not as clearly defined as was considered to be the case. In fact, by suitable choice of conditions it is possible to induce virtually all the cystine to react according to one reaction mechanism.

When the polar side chains are modified by acetylation or treatment with phenyl isocyanate, an overall decrease in cystine reactivity is observed (Table 5). Thus, for the reaction with cold bisulphite

solutions, the ratio ($A+B$)/total disulphide-S is decreased from 1.85 for untreated wool to 1.57 for acetylated wool and 0.27 for phenyl isocyanate-treated wool, even when using the maximum figures for fraction ($A+B$). If the minimum figures are considered, the decreases in ($A+B$) are much greater. Similar decreases in the ratio ($A+B$)/total disulphide-S are observed when boiling bisulphite solution or thioglycollic acid is used. Since acetic anhydride and phenyl isocyanate each react extensively with the polar side chains in wool it might be expected that, if the effect on cystine reactivity was due to the conversion of the polar environment of fraction ($A+B$) into a less polar one, both reagents may behave similarly, and result in a similar decrease in ($A+B$) and an increase in ($C+D$). However, whereas acetylation leads to a marked decrease in the ratio ($A+B$)/total disulphide-S, treatment with phenyl isocyanate almost completely destroys the reactivity of the cystine linkages in wool. That this behaviour is not confined to the effect of reducing agents is shown by the experiment with sodium hydroxide. Under the conditions used, some 65 % of the disulphide-S in untreated wool is decomposed by sodium hydroxide, whereas with phenyl isocyanate-treated wool there is no change in the disulphide-S content.

The original subdivision of fraction ($A+B$) was based on the ability to water of the thiol and cysteine sulphonate groups resulting from treatment with cold bisulphite solutions, although it was pointed out that the relative magnitude of these subfractions was influenced by the conditions used in rinsing. A more definite differentiation between A and B was given by the reaction of thioglycollic-reduced wool with methylene diiodide and formaldehyde. The former reagent was found to react with only subfraction A to give combined djenkolic acid, whilst formaldehyde reacted with subfraction A to give combined djenkolic acid and also with subfraction B to give combined thiazolidine-4-carboxylic acid (Middlebrook & Phillips, 1947). It was suggested that the cystine linkages of subfraction A which are capable of undergoing reversion after preliminary rupture with bisulphite, and of being recombined through a methylene group after rupture with thioglycollic acid, are in the vicinity of salt linkages, the effect of which is to prevent separation of the two sulphur atoms after rupture of the linkage. The linkages incapable of being reunited once broken (subfraction B) were suggested as being in the vicinity of acid amide groups which are incapable of forming salt linkages. It is difficult to reconcile this view with the almost complete reversion to intact cystine linkages of the thiol groups produced by reduction with thioglycollic acid on subsequently heating the reduced wool in pyridine. It may be argued that separation of the

sulphur atoms only occurs by virtue of the swelling associated with an aqueous medium. However, pyridine is also capable of swelling wool fibres to a reasonable extent.

The preceding results and discussion do not lend very much support to the suggestion that some of the cystine linkages in wool are activated by an environment of polar side chains, and alternative ideas have been considered. The recent discovery of Horio & Kondo (1953) and Mercer (1953) that the cortex of wool fibres is composed of two different proteins, one of which is considerably more resistant to chemical attack than the other, offered a possible explanation, but Fraser, Lindley & Rogers (1954) have shown that the more resistant portion is much richer in disulphide linkages. Mercer, Golden & Jeffries (1954) have suggested that the cystine in the more reactive orthocortex behaves in much the same manner as is associated with fraction (*A* + *B*) and that there is about twice as much cystine in the less reactive paracortex as in the orthocortex. These ideas are not in keeping with the approximate equivalence of fractions (*A* + *B*) and (*C* + *D*) as assessed on the initial reaction with reducing agents.

The results, however, do point fairly consistently to accessibility of cystine linkages to the reagent as being of some importance in deciding the degree, and to a lesser extent the mechanism, of a particular reaction. Lindley (1948) has shown that the cystine of some non-fibrous proteins, including soluble proteins, shows a similar variation in reactivity and has suggested that this is not in keeping with a difference in accessibility as being the cause of the variation. This may be the case if accessibility is interpreted in terms of crystalline and amorphous regions of a fibrous protein. However the experiments using acetylated and phenyl isocyanate-treated wools seem to point to steric factors as a possible cause of the variation in reactivity. An overall decrease in reactivity is brought about by the substitution of an acetyl group on the ends of polar side chains, and when the size of this substituent group is increased to that of a phenyl group, the reactivity is almost completely destroyed.

It may be that in the untreated wool some of the cystine linkages are protected by an environment of bulky side chains offering steric hindrance to the approach of other molecules. Consden & Gordon (1950), by means of chromatographic studies of partial hydrolysates of wool, have isolated dipeptides of cystine with most of the other amino acids in wool and concluded that most of the mono amino-mono carboxylic acids are combined both through their amino and carboxyl groups to residues of cystine. Consden, Gordon & Martin (1949) found some peptides of cystine with dicarboxylic amino acids in partial hydrolysates of wool. Hence there

is considerable evidence for cystine being in close proximity to amino acid residues having both long and short side chains.

Whilst it is possible to ascribe the variation in degree of reactivity of the cystine linkages to steric hindrance by bulky side chains, this does not explain the different reaction mechanism and ultimate products which do undoubtedly arise in some instances as shown by Phillips and his colleagues. If the different products arise solely from the more drastic conditions necessary to induce the less reactive portion to undergo reaction, there does not seem to be any convincing reason why all the cystine should not then undergo the alternative reaction. For example, if cold bisulphite solutions react readily with part of the cystine to give thiol and cysteine sulphonate groups, but boiling solutions, which are necessary to reach the remainder of the cystine, lead to the formation of combined α -amino-acrylic acid, then it might reasonably be expected that all the cystine would give combined α -amino-acrylic acid under these conditions, if accessibility were the only factor governing the reactivity. It must be concluded, therefore, that accessibility is only one factor.

Recent work on oxytocin, vasopressin and insulin has shown that some of the cystine is in the form of intramolecular linkages with the two half-cystine residues in a 1:6 relationship along the chain.

Some fundamental structural difference such as this would be more in keeping with the observed differences in reaction mechanism and products than the influence of side chains. Furthermore, it is probable that such laterally situated disulphide bonds would be less protected by bulky side chains than would the conventional cross-linkages. However, there is no convincing evidence for, or against, the existence of this type of structure in wool.

SUMMARY

1. The reactivity of the cystine linkages in wool towards reducing agents has been investigated before and after modification of the wool in various ways.
2. Reduction and alkylation of part of the cystine (the (*A* + *B*) fraction) allows the remainder (the (*C* + *D*) fraction) to react with sodium bisulphite and thioglycolic acid under conditions where it is normally inert. The reaction products are then the same as normally obtained with fraction (*A* + *B*).
3. In contrast with this, reduction followed by cross-linking of the thiol groups from fraction

($A + B$) through a methylene bridge does not allow fraction ($C + D$) to show this induced reactivity.

4. Acetylation of polar side chains in wool leads to a general reduction in the reactivity of cystine.

5. Treatment of wool with phenyl isocyanate leads to blocking of all the polar side chains without affecting the cystine. Such treatment almost completely inhibits subsequent reactivity of the cystine linkages.

6. It is suggested that the degree of reactivity of cystine linkages is largely governed by their accessibility to a reagent, close proximity of bulky side chains reducing reactivity through steric hindrance.

7. Other factors such as varying structural arrangement may be responsible for the variation in the nature of the products resulting from a particular reaction.

Grateful acknowledgement is made to the administrators of the Wool Levy Research Trust Fund, under whose grant this work was carried out, to Miss Joan Hill for assistance with experimental work, and to officers of the Wool Textile Research Laboratories, C.S.I.R.O., for helpful criticism.

REFERENCES

- Alexander, P. & Gough, D. (1951). *Biochem. J.* **48**, 504.
 Alexander, P., Hudson, R. F. & Fox, M. (1950). *Biochem. J.* **46**, 27.
 Astbury, W. T. (1942). *J. chem. Soc.* p. 337.
 Carter, E. G. H., Middlebrook, W. R. & Phillips, H. (1946). *J. Soc. Dy. Col.* **62**, 203.
 Consden, R. & Gordon, A. H. (1950). *Biochem. J.* **46**, 8.
 Consden, R., Gordon, A. H. & Martin, A. J. P. (1949). *Biochem. J.* **44**, 548.
 Cuthbertson, W. R. & Phillips, H. (1945). *Biochem. J.* **39**, 7.
 Elsworth, F. F. & Phillips, H. (1938). *Biochem. J.* **32**, 837.
 Elsworth, F. F. & Phillips, H. (1941). *Biochem. J.* **35**, 135.
 Farnworth, A. J. (1955). *Biochem. J.* **59**, 529.
 Fraser, R. D. B., Lindley, H. & Rogers, G. E. (1954). *Biochem. biophys. Acta*, **13**, 295.
 Geiger, W. B., Kobayashi, F. F. & Harris, M. (1942). *Industr. Engng Chem.* **34**, 1398.
 Horio, M. & Kondo, T. (1953). *Text. Res. (J.)* **23**, 373.
 Lindley, H. (1948). *Biochem. J.* **42**, 481.
 Lindley, H. & Phillips, H. (1945). *Biochem. J.* **39**, 17.
 Lindley, H. & Phillips, H. (1947). *Biochem. J.* **41**, 34.
 Mercer, E. H. (1953). *Tex. Res. (J.)* **23**, 388.
 Mercer, E. H., Golden, R. L. & Jeffries, E. B. (1954). *Text. Res. (J.)* **24**, 615.
 Middlebrook, W. R. & Phillips, H. (1942*a*). *Biochem. J.* **36**, 294.
 Middlebrook, W. R. & Phillips, H. (1942*b*). *Biochem. J.* **36**, 428.
 Middlebrook, W. R. & Phillips, H. (1947). *Biochem. J.* **41**, 218.
 Patterson, W. I., Geiger, W. B., Mizell, L. R. & Harris, M. (1941). *Bur. Stand. J. Res., Wash.*, **27**, 89.
 Phillips, H. (1946). *Fibrous Proteins Symposium*, p. 39. Society Dyers and Colourists, Bradford.
 Sanger, F. (1945). *Biochem. J.* **39**, 507.
 Schuringa, G. J., Schooneveldt, C. & Konings, T. (1951). *Text. Res. (J.)* **21**, 281.
 Shinohara, K. (1935*a*). *J. biol. Chem.* **109**, 665.
 Shinohara, K. (1935*b*). *J. biol. Chem.* **112**, 671.
 Shinohara, K. (1935*c*). *J. biol. Chem.* **112**, 683.
 Speakman, J. B. (1936). *J. Soc. Dy. Col.* **52**, 423.

Selective Inhibition of the Liberation of Extracellular Enzymes and Protein in Cultures of *Staphylococcus aureus*

BY H. J. ROGERS AND P. C. SPENSLEY

National Institute for Medical Research, Mill Hill, London, N.W. 7

(Received 27 September 1954)

Most substances, such as amino acid analogues, which inhibit protein formation by bacteria, do so in a non-specific manner. That is, the formation of all protein ceases and growth stops when they are added to the culture medium. Any substance which acts upon the formation of a single protein or group of proteins is likely to be a valuable tool in studying biosynthetic processes. The present work describes the effect of some negatively charged polymers upon cultures of staphylococci. The liberation into the culture fluid of a group of extracellular enzymic proteins is inhibited, but growth of the organisms is not impaired. These extracellular enzymes are not essential for growth under the conditions chosen. Recently, considerable attention has been paid to

the ability of synthetic charged polymers, 'macroanions' and 'macroocations', to inhibit the action of formed enzymes (cf. Spensley & Rogers, 1954, for some references). These studies have not, however, been extended to the effects of macro-ions upon living cells. An exception is the study by Katchalski, Bichowski-Slonmizki & Volcani (1953) of the growth inhibition of *Escherichia coli* and *Staphylococcus aureus* by the macroocations polylysine and polyarginine. Polyglutamic and polyaspartic acids, two macroanions examined by these authors, did not inhibit growth.

The ability of some macroanions to inhibit the liberation of at least two extracellular enzymes in cultures of staphylococci is proved in the work