REFERENCES

- Danishefsky, I. & Eiber, H. B. (1959). *Arch. Biochem. Biophys.* **85**, 53.
- Dodgson, K. S. (1961). *Biochem. J.* **78**, 324.
- Dodgson, K. S. & Lloyd, A. G. (1961). *Biochem. J.* **78**, 319.
- Dodgson, K. S., Powell, G. M. & Tudball, N. (1960). *Biochem. biophys. Res. Commun.* **2**, 130.
- Dodgson, K. S. & Spencer, B. (1953). *Biochem. J.* **55**, 436.
- Dohlman, C. H. (1956). *Acta physiol. scand.* **37**, 220.
- Dziewiatkowski, D. D. (1956). *J. biol. Chem.* **223**, 239.
- Gardell, S. (1955). *Acta chem. scand.* **9**, 1035.
- Gregory, J. D. & Robbins, P. W. (1960). *Annu. Rev. Biochem.* **29**, 347.
- Hirano, S., Hoffman, P. & Meyer, K. (1960). *Fed. Proc.* **19**, 146.
- Huseman, E., Hoffman, E. G., Lotterle, R. & Widersheim, M. (1952). *Experientia*, **8**, 153.
- Lloyd, A. G. (1959a). *Nature, Lond.*, **183**, 109.
- Lloyd, A. G. (1959b). *Biochem. J.* **72**, 133.
- Lloyd, A. G. (1959c). *Rep. Emp. Rheum. Council.* **22**, 42.
- Lloyd, A. G. (1960). *Biochem. J.* **75**, 478.
- Lloyd, A. G. (1961). *Nature, Lond.*, **190**, 914.
- Lloyd, A. G. & Dodgson, K. S. (1959). *Nature, Lond.*, **184**, 548.
- Lloyd, A. G. & Dodgson, K. S. (1961). *Biochim. biophys. Acta*, **46**, 116.
- Lloyd, A. G., Dodgson, K. S., Price, R. G. & Rose, F. A. (1961). *Biochim. biophys. Acta*, **46**, 108.
- Lutwak-Mann, C. (1941). *Biochem. J.* **35**, 610.
- Meyer, K. (1960). In *Molecular Biology*, p. 69. London: Academic Press Inc.
- Meyer, K., Hoffman, P. & Linker, A. (1958). *Science*, **128**, 896.
- Meyer, K., Linker, A., Davidson, E. A. & Weissmann, B. (1953). *J. biol. Chem.* **205**, 611.
- Morrow, P. E., Hodge, H. C., Neumann, W. F., Maynard, E. A., Blanchet, H. K., Fasset, D. W. & Birk, R. E. (1952). *J. Pharmacol.* **105**, 273.
- Partridge, S. M. (1948). *Biochem. J.* **42**, 238.
- Peacock, A. C. & Harris, R. S. (1950). *Arch. Biochem.* **27**, 198.
- Ricketts, C. R., Walton, K. W. & Saddington, S. M. (1954). *Biochem. J.* **58**, 532.
- Roseman, S. (1959). *Annu. Rev. Biochem.* **28**, 545.
- Roseman, S. & Ludowieg, J. (1954). *J. Amer. chem. Soc.* **76**, 301.
- Rosen, O., Hoffman, P. & Meyer, K. (1960). *Fed. Proc.* **19**, 147.
- Tanko, B. (1932). *Biochem. Z.* **247**, 486.
- Walkenstein, S. S. & Knebel, C. M. (1957). *Analyt. Chem.* **29**, 1516.
- Wolfrom, M. L., Montgomery, R., Karabinos, J. V. & Rathgeb, P. (1950). *J. Amer. chem. Soc.* **72**, 5796.
- Young, L., Edson, M. & McCarter, J. A. (1949). *Biochem. J.* **44**, 179.

Biochem. J. (1961) **80**, 578

The Isolation of the α -Toxin of *Clostridium welchii* Type A by Zone Electrophoresis in Vertical Columns

By J. STEPHEN

The Rowett Research Institute, Bucksburn, Aberdeen

(Received 16 January 1961)

The α -toxin of *Clostridium welchii* type A is believed to be the most important single factor involved in clostridial toxæmia (van Heyningen, 1950, p. 25). It has been extensively studied biochemically and serologically, as a result of which it has been recognized as a lecithinase capable of splitting phosphorylcholine from lecithin (Macfarlane & Knight, 1941). Such enzymic activity is believed sufficient (Macfarlane, 1950) to explain the haemolytic, necrotizing and lethal activities assigned to it by Glenny, Barr, Llewellyn-Jones, Dalling & Ross (1933). Previous attempts have been made by van Heyningen (1941a, b), Adams & Hendee (1945), van Heyningen & Bidwell (1948) and Roth & Pillemer (1953) to isolate it from the other *C. welchii* antigens such as hyaluronidase, collagenase and θ -toxin (a reversibly oxidizable

haemolysin), with which it usually occurs in culture filtrates. The techniques employed by these workers were mainly fractional precipitation with salt and organic solvents.

A series of experiments was planned in which the newer techniques applicable to such a problem were to be applied, and the first attempt was by zone electrophoresis in vertical columns. At the conclusion of the experimental stage of this part of the work, a paper was published by Habermann (1959), who described the results of a similar approach to the same problem. He found that the α -toxin could be purified to a considerable degree by a combination of either fractional precipitation and chromatographic techniques or electrophoretic and chromatographic techniques. The yield from the chromatographic stage was high but that from

the electrophoretic stage was relatively poor, and the α -toxin was not quantitatively resolved from θ -toxin by the latter.

MATERIALS AND METHODS

Organisms. *Clostridium welchii* strain CN.1491 was used throughout this work and was kindly given by the Wellcome Research Laboratories. Stock cultures were kept in Robertson's alkaline egg medium (Levine & Schoenlein, 1930). Fresh cultures were prepared about every 3 months and stored at 4°.

Culture media

Cooked-meat broth. Ox muscle was substituted for heart in the preparation described by Mackie & McCartney (1948), otherwise the medium was the same.

Medium for production of α -toxin. Peptone (Evans Medical Supplies Ltd.) and Bacto-Casitone (Difco Laboratories Inc., Detroit, Mich., U.S.A.) were each added to 3 l. of tap water to a final concentration of 1% (w/v). The pH was adjusted to 8 and the solution steamed for 30 min., after which it was filtered. N-Hydrochloric acid was then added to bring the pH to 7.8. Meat, with superficial fat removed, was minced, treated with 0.05N-NaOH (1 l./kg. of minced meat) for 20 min. at just below boiling point and squeezed dry. The meat particles (200 g./l.) were added to the peptone-Casitone solution and the whole was autoclaved at 15 lb./in.² for 15 min. An aqueous solution of glucose (50%, w/v) was autoclaved at 15 lb./in.² for 15 min. and, immediately before inoculation with the culture, was added to the medium to a final concentration of 0.6% (v/v).

Medium for production of θ -toxin. The medium used was that described by Brooks, Sterne & Warrack (1957).

Enzyme and toxin assays

α -Toxin. α -Toxin was assayed as lecithinase by a method based on that described by van Heyningen (1941 a), in which the turbidity produced in egg-yolk solutions is measured. This technique is not completely specific, other enzymes being known which will evoke the same phenomenon. However, it is assumed (cf. Macfarlane & Knight, 1941; Kushner, 1957) that if this reaction is specifically neutralized by appropriate antisera it can be used for the assay of the lecithinase produced by *C. welchii* type A. This condition was fulfilled with preparation A (see Results section). A standard toxin was prepared by dialysing a culture filtrate against glycerol for 72 hr. (Macfarlane & Knight, 1941) and examined by Mrs I. Batty of the Wellcome Research Laboratories and found to contain 14.95 lecithovitellin (LV) units/ml. (1 LV unit is that amount of α -toxin which in the presence of 1 i.u. of antitoxin will just cause turbidity in an egg-yolk solution: see van Heyningen, 1950, p. 27); this preparation remained stable for 2½ years. The egg-yolk broth was made by suspending 10% (v/v) broken yolks of eggs in papain-digest broth which included peptone 1% (w/v) and NaCl 0.5% (w/v). After centrifuging this was Seitz-filtered. The composition of the borate buffer diluent was as given by Howie, Duncan & Mackie (1953), but the pH was adjusted to 7.4 instead of 8.0-8.4.

Volumes of 5 ml. of the egg-yolk broth in borate buffer (15%, v/v) were dispensed into standard EEL (Evans

Electroselenium Ltd., Harlow, Essex) nephelometer tubes. Dilutions of preparation A, in the range 0.09-0.33 LV unit/ml., were made in steps of 0.03 LV unit/ml. and 1.0 ml. was added to each assay tube. The tube contents were mixed rapidly by inversion and placed at 1 min. intervals in a water bath at 37° and incubated for 15 min., after which they were withdrawn at 1 min. intervals, wiped dry and the turbidity was read on the nephelometer. Each assay was done in duplicate and the results were used to construct a standard curve. This procedure was repeated before each assay because of the variability of the egg-yolk solutions.

Proteolytic activity. Proteolytic activity of our preparations was assayed with azocoll as substrate (Oakley, Warrack & van Heyningen, 1946). The method of assay was based on that described by Bidwell (1950). Azocoll was prepared according to Oakley *et al.* (1946) and sieved through an 80-mesh sieve. The diluent used throughout was NaCl, 0.85%, to which was added tricresol ether, 0.5%.

The test system was calibrated in terms of arbitrary units; preparation A, which was rich in 'azocoll-splitting' activity, was used as standard and said to contain 1.0 unit/0.0135 mg.

Solutions containing from 1 to 10 azocoll units/ml. were made and 1 ml. portions were pipetted into EEL tubes (8 mm. diam., suitable for EEL colorimeter), each of which contained 20 mg. of azocoll, and incubated overnight at 37° after mixing. Residual azocoll was centrifuged and the extinction of the supernatant measured in the colorimeter. One batch of azocoll was sufficient for the whole of this work and did not require repeated standardization.

Hyaluronidase. Hyaluronidase was assayed by the turbidimetric assay described by Dorfmann & Ott (1948) with serum albumin (Armour Laboratories, fraction V).

θ -Toxin. The reversibly oxidizable haemolysin (θ -toxin) was assayed as an antigen against specific antibody. The procedure of Brooks *et al.* (1957) was followed.

Glycosidases. α -Mannosidase, β -galactosidase, β -N-acetylglucosaminidase and β -glucuronidase were assayed by the methods described by Conchie, Findlay & Levy (1959).

Ultrafiltration

The high-pressure ultrafilter described by Bullen & Batty (1956) was used for the concentration of our culture filtrates. Membranes were cast from 8% collodion in acetic acid (Necol; Imperial Chemical Industries Ltd.). Full details of membrane production and the manipulation of the apparatus are given by Bullen, Thurlbourn & Brown (1954).

Small volumes, e.g. eluate fractions, were conveniently concentrated by the 'polyethylene glycol' technique described by Kohn (1959).

Zone electrophoresis in columns

The apparatus was a development of the one described by Mould & Stephen (1958). Attempts to cool packed columns centrally in order to increase the applied field have been made by introducing a cold-finger type of arrangement (Porath, 1957). The disadvantage of such a design is in the local evolution of heat at the porous disk at the bottom of the column.

The following is a description of the construction of an annulus column (i.e. one in which the refrigerant goes right

through the porous disk). The essential part of the apparatus is shown in Fig. 1.

A is a column, 65 cm. long and 4 cm. diam., with a B34 standard ground-glass socket at the top and a B34 standard ground-glass cone at the bottom. A central tube *B*, 1 cm. diam., emerges from the side of the column just below the socket at the top and terminates at the lower end in a B10 socket *C*, flush with the lower edge of the B34 cone. The porous disk *D* is formed *in situ* from an Araldite resin and a 'filler' (all materials supplied by Aero Research, Duxford, Cambs.). The composition of the mixture was Araldite 103 (10 g.), Hardener 951 (1 g.) and Silica flour (15 g.). This was allowed to set overnight at room temperature, after which a series of holes was drilled in the plug. The anode chamber consists of a three-necked 5 l. vessel, into the bottom of which is sealed a tube *E* of 1 cm. diam., terminating in a B10 cone. This is designed so that, with the column *in situ*, the B10 cone of *E* engages the B10 socket of *B*. To obviate glass-blowing difficulties, the seal *F* in the bottom of the flask is made with the same Araldite mixture as for the plug. The rest of the apparatus was exactly as described by Mould & Stephen (1958). The effective field across the column is about 80% of the applied p.d.

Cellulose. Ethanolyzed cellulose was prepared according to Flodin & Kupke (1956). Our experience showed that the best source of cellulose was Regaid cotton wool (Boots),

although variations in the suitability of the ethanolyzed product were detected from batch to batch.

Buffers. Borate buffer, pH 7.1, *I* 0.025, which contained 0.025 M-NaCl, was used since at this pH the α -toxin is stable (Gale & van Heyningen, 1942) and in borate buffer can be conveniently assayed as leicithinase.

Packing columns. Ethanolyzed cellulose, 200 g., was suspended in borate buffer, pH 7.1, *I* 0.025. The column was packed to a depth of 55 cm. by allowing this slurry to filter into the column at a controlled rate, from a height of 12 ft. A filter-paper disk had been previously placed on top of the porous disk, and one was finally placed on top of the packing to prevent its being disturbed. The suitability of the packed column was always tested by passing through it a sample of the material to be fractionated, the eluate being examined for activity recovery and any evidence of adsorption.

Immunoelectrophoresis

This was largely carried out according to Grabar & Williams (1953, 1955).

Glass plates, 30 cm. $\times$ 11 cm., were coated with New Zealand agar, 1% (w/v) in borate buffer, pH 7.1, *I* 0.05, to a depth of 3 mm. The agar was cut into a convenient number of strips and inlays were cut with a cork borer of suitable diameter. The positions of the inlays were determined by the electrophoretic properties of the substances under study, due allowance being made for the electro-endosmotic streaming through the agar gel. The sample was made up in buffer to twice the concentration finally desired and diluted with an equal volume of agar, 2% (w/v) in buffer. This was introduced into the inlay. Paper wicks connected the ends of the gels to the buffer in the electrode chambers.

The plates were usually clamped to a refrigerated copper plate whose temperature was maintained at 1°. A voltage gradient of 2.6 v/cm. was applied, which produced currents of approx. 46 ma.

The electrode chambers were perfused continuously with fresh buffer to maintain a stable pH with borate buffer, pH 7.1. At the end of the electrophoretic run, antiserum, previously dialysed against the same buffer, was added to the longitudinal ditches of the gel, which was then usually incubated for 48 hr. at 37°. The gels were then washed in a solution of NaN_3 , 0.1%, and NaCl, 0.85%, for approx. 3 days with frequent changes of the wash solution. The precipitin lines were then stained with azocarmine G (G. T. Gurr Ltd., London, S.W. 6).

Other chemical and biological estimations

Nitrogen. The nitrogen content of specific samples was estimated by the micro-Kjeldahl technique.

Protein. The method of Lowry, Rosebrough, Farr & Randall (1951) was used to estimate the protein in fractions from the columns. Standards were not prepared as only the relative distribution of protein in the fractions was examined. Readings were taken at λ 500 m μ .

Measurements of pH. All pH measurements were carried out with a Pye pH meter.

LD₅₀ Assays. Mice (23–25 g.; T.O. strain, Mill Hill 1956) were used. The mice were injected intravenously with a total volume of 0.3 ml. of toxin dilution. Deaths were recorded at 24, 48 and 72 hr. Very few deaths were observed after 48 hr. and none after 72 hr., which was taken

Fig. 1. Annulus columns. For description see text.

as the maximum time limit for recording death from toxic effects. Adequate serum and diluent controls were included with the experimental groups.

Minimum haemolytic dose (MHD). The haemolytic potency of the lecithinase in preparation A was estimated as the number of MHD/LV unit. Doubling dilutions were made from a solution containing 1 LV unit/ml. in a saline diluent containing 5 mm-CaCl₂. A suspension of washed, sheep red blood cells, 6% in 0.85% NaCl soln., was made, 0.5 ml. portions being dispensed into each tube. The tubes were incubated for 1 hr. at 37°, and then cooled and centrifuged. The last tube to contain trace haemolysis was taken as the end point. (The content of θ -toxin in preparation A, produced in the presence of solid meat particles, was so low as to be negligible at this dilution.)

RESULTS

Production of α -toxin

Cooked-meat broth (20 ml.) was inoculated with 2 ml. of stock culture and incubated overnight at 37°. This culture was subcultured twice in cooked-meat broth (20 ml.) and once in the peptone medium (200 ml.) used for toxin production. Each subculture was grown for approx. 2 hr. (or until active gas production was observed) at 37°. The last subculture was used as inoculum for the 3 l. batch of peptone medium, which was then incubated for 3½ hr. at 37°. The culture was clarified through paper pulp and sterilized through a Carlson E.K. pad. The filtrate thus obtained contained 3.5 LV units/ml. and was concentrated by ultrafiltration.

Viable bacterial counts (Bullen & Scarisbrick, 1957) showed that most of the α -toxin was produced during active cell division, in confirmation of the earlier observation of Gale & van Heyningen (1942). Under normal conditions of culturing the pH dropped from an initial pH of 7.8 to about 6. Addition of bicarbonate buffer was shown to have no appreciable effect on the yield of α -toxin.

Concentration of Seitz filtrates

Potent crude material was obtained by high-pressure ultrafiltration with reasonable success. Filtrates from cultures grown in media containing peptone, 1% (w/v), could be concentrated almost to dryness. The viscous concentrate was washed off with a NaCl solution (0.1%), the washings being pooled, centrifuged if necessary until the supernatant was absolutely clear, and freeze-dried. Preparation A was the product obtained from four 3 l. batches of culture, the filtrates of which were separately concentrated and then pooled and freeze-dried. Preparation B was produced in a manner similar to that for preparation A, except that the culture was grown in a 3 l. batch of the medium described for θ -toxin production.

α -Toxin production was directly proportional to the peptone concentration of the medium over the range 1–3% (w/v). However, the culture filtrates from media containing peptone of initial concentration > 1% (w/v) could not be concentrated to the same degree as those from media containing $\leq$ 1% (w/v) peptone.

Zone electrophoresis

Preparation A (500 mg. equivalent to 500 LV units of α -toxin, 15 000 i.u. of hyaluronidase, 37 000 azocoll units of proteolytic activity) were dissolved in 2 ml. of the same buffer as was used for packing and run into the column to a depth of 3 cm. The column was then fitted into the rest of the apparatus, which was filled with buffer. A potential of 1000 v was applied, giving an approximate voltage gradient of 16 v/cm. across the packed column and producing a current of 78 ma. After 18 hr. the column was removed and eluted with the same buffer, 5 ml. fractions being collected. Each fraction was analysed for protein, α -toxin (as lecithinase), hyaluronidase and proteolytic activity. The result is shown in Fig. 2. The proteolytic activity appeared to be distributed between four components, and the hyaluronidase was recovered as a broad asymmetric peak. The α -toxin was recovered in 90–95% yield, the proteolytic activity in 95% and the hyaluronidase in 70–75% yield. The hyaluronidase and the proteolytic activities were clearly not separated. Immuno-electrophoresis showed the α -toxin to be contaminated with both faster- and slower-moving antigens.

A second experiment was done with preparation A, the duration of the run being 40 hr., and at the half-way stage the buffer was changed in the side

Fig. 2. Zone electrophoresis: 18 hr. run on preparation A, showing separation of α -toxin from the proteolytic and hyaluronidase activities. ○, α -Toxin; □, proteolytic activity; ■, hyaluronidase; ●, protein. Activities are plotted as percentages of their respective maxima.

arm and the cathode vessel. Under these conditions the hyaluronidase and proteolytic activities were run right off the column and the α -toxin was recovered in 85–90 % yield. Immuno-electrophoresis showed the isolated toxin to be a single precipitin system (see below). The results of the 18 and 40 hr. experiments show that the percentage recovery of the active toxin decreases with the duration of the experiment; this confirmed the observations of Habermann (1959). The material was not run again on the column as this would have resulted in a further substantial loss of activity.

Table 1. *Properties of α -toxin before and after fractionation*

Activity of the preparations is expressed in terms of leicthovittellin (LV) units, lethal dose (LD_{50}) and minimal haemolytic dose (MHD).

LV units/mg. of N	LD_{50} /LV unit	MHD/LV unit
Crude toxin (preparation A)		
7.8	55	16–32
Fractionated toxin		
50.6	55	16–32

Fig. 3. Zone electrophoresis: 40 hr. run on preparation B, showing separation of α -toxin from θ -toxin. $\circ$, α -Toxin; $\bullet$, θ -toxin. Activities are plotted as percentages of their respective maxima.

Fig. 4. Diagrammatic representation of the result obtained with preparation A with egg yolk and antiserum K680.

The properties of the isolated toxin are listed in Table 1.

An overall purification of approximately sixfold was achieved when expressed on a unit nitrogen basis. The lecithinase, lethal and haemolytic properties of α -toxin were not separated by electrophoresis.

A third experiment was done with preparation B to determine whether or not the α - and θ -toxins were electrophoretically separable. The equivalent of 500 LV units of α -toxin and 1300 units of θ -toxin were dissolved in 2 ml. of borate buffer and the experiment was conducted in exactly the same manner as described for the second experiment; the result is shown in Fig. 3. Under these conditions the α -toxin is more anodic than the θ -toxin and these two substances are clearly separable. The yield of θ -toxin was variable, 50–70 %, and, unaccountably, the yield of α -toxin was considerably lower, 50–60 %, from this preparation than that repeatedly obtained from preparation A. The fractions which contained the α - and θ -toxins respectively were shown by immuno-electrophoresis to be complex mixtures of antigens.

Immuno-electrophoresis

A borate gel with two strips was prepared. Two inlays were cut in the gel, one in each strip. One was near the cathode and one, in the other strip, was near the anode end, and into these preparation A (12 mg.) was introduced. At the end of the run (i.e. after 18 hr.) antiserum K680 (*Clostridium* Diagnostic Serum, *C. welchii* type A, produced by Wellcome Research Laboratories) was added to the two outer ditches and the egg-yolk solution used for lecithinase assays added to the centre ditch. After incubation nine precipitin lines developed in the gel, and an intense turbidity developed in the centre ditch which later spread into the gel itself. This was rather difficult to photograph and a diagrammatic representation is given in Fig. 4. The line opposite the zone of turbidity was assumed to be the one produced by the α -toxin and its corresponding antitoxin. This line is also indicated by an arrow in Plate 1 (a), which shows the result of repeating the experiment without the egg-yolk treatment. Another experiment was done with a third strip in the gel, in

EXPLANATION OF PLATE 1

(a) Immuno-electrophoresis of preparation A. The precipitin line (marked with an arrow) is the one opposite which the egg-yolk turbidity developed.

(b) Immuno-electrophoresis of preparation B.

(c) Immuno-electrophoresis of preparation A. Recovered on elution through a column packed with ethanolyzed cellulose.

(d) Immuno-electrophoresis of isolated α -toxin.

(a)

(b)

(c)

(d)

which was placed (near the anode end) the α -toxin isolated from the 40 hr. zone-electrophoresis run, previously described, with preparation A. A precipitin line appeared [Plate 1 (*d*)], whose position indicated a substance with the same mobility as that which produced the precipitin line shown by the arrow in Plate 1 (*a*); no other lines were observed. These experiments were repeated with 18 different *C. welchii* antisera, some of which revealed the same number of lines in preparation A as did serum K 680, but none revealed more.

By the use of the neutral indicator *N*-2:4-dinitrophenylethanolamine it was shown that the α -toxin was anionic, migration towards the cathode being due to the considerable electroendosmosis which occurred through these agar gels. It was also demonstrated, by repeating the experiment with dinitrophenylethanolamine in a phosphate-buffered agar (pH 7.1, *I* 0.05) that the effect of boric acid on the mobility of proteins having a carbohydrate moiety (Garbers & Joubert, 1958) was not operative with the α -toxin.

Preparation B, originally prepared for θ -toxin, was shown to contain at least 25 antigens when examined against serum K 680 [Plate 1 (*b*)]. The abundance of precipitin lines produced by these preparations prompted the search for other specifically identifiable substances. Preparation A was examined for α -mannosidase, β -galactosidase, β -*N*-acetylglucosaminidase and β -glucuronidase, of which the two last-named are usually associated with hyaluronidase and the first two with the second two in mammalian tissues. None of these enzymes was detected. Preliminary histochemical tests (Lillie, 1954) also failed to detect deoxyribonuclease, which is known to be produced by some *C. welchii* type A strains (Oakley & Warrack, 1953).

A buffer solution (2 ml.) containing 100 mg. of preparation A was then passed through a column, packed as for the electrophoresis runs but without application of the electric field. The eluate was concentrated by polyethylene glycol and examined immunoelectrophoretically; the result is shown in Plate 1 (*c*) and is seen to be similar to (*a*), obtained with the original untreated preparation A.

Both toxigenic media used for the production of preparations A and B were concentrated approximately tenfold and examined against serum K 680, no precipitin lines being developed. Preparation A was also tested against a serum from a rabbit immunized with the toxigenic peptone medium and again no result was obtained.

DISCUSSION

The method of concentration of the Seitz filtrates was regarded as an important aspect of this work, as the obtaining of crude concentrates

rich in α -toxin has been cited (Roth & Pillemer, 1953) as one of the major hindrances in purification studies. The crude ultrafiltered concentrates could be obtained within 24 hr. of the start of culture operations without their being subjected to precipitation with salts or organic solvents; a stage yield of 95% was obtained.

It is possible to obtain the α -toxin of *C. welchii* type A free from hyaluronidase, θ -toxin and the proteolytic activities, which degrade the crude substrate azocoll, by zone electrophoresis.

The α -toxin isolated from a 40 hr. run with preparation A as starting material was shown by immunoelectrophoresis to have been separated from the other eight antigens of bacterial origin that were originally present. Nevertheless the degree of purification achieved was only sixfold. The activity of the final material is very low compared with that of the product obtained by Roth & Pillemer (1953), which although of undefined antigenic purity, was 35 times as active. Failure to demonstrate precipitin lines with concentrated culture medium against antiserum K 680 would suggest that the low specific activity of the isolated toxin was due to the presence of non-antigenic residues from the medium, which were retained by the ultrafiltration membranes used in the concentration of the culture filtrates. Alternatively, the low activity could mean that there were present bacterial antigens to which there were no antibodies present in the 18 *C. welchii* antisera examined, but in view of the extremely varied composition of these antisera this is regarded as unlikely.

Oakley & Warrack (1953) listed type A strains of *C. welchii* as being capable of producing at least five serologically distinct antigens. The results reported in this paper show that culture filtrates from peptone-based meat-containing media may contain at least nine different antigens and those from muscle-digest meatless media 25 different antigens.

Examination of crude material passed through the column without application of the electrical field showed that passage through the packed column and the method for concentrating the eluate did not alter the serological properties of the mixture studied. This was considered important in respect of the analysis of the isolated α -toxin as it shows that the processes involved in fractionating and recovering the active material do not result in the denaturation of antigenic contaminants.

SUMMARY

1. Culture filtrates of *Clostridium welchii* type A, concentrated by high-pressure ultrafiltration, have been examined by zone electrophoresis and immunoelectrophoresis.

2. Zone electrophoresis was carried out with ethanolyzed cellulose as the stabilizing medium in columns of a novel design. The columns were internally as well as externally cooled; a description of their construction is given. With such an apparatus, the α -toxin of *C. welchii* type A was quantitatively separated from the hyaluronidase, proteolytic and θ -toxin activities. The lecithinase, haemolytic and lethal activities were electrophoretically inseparable.

3. Immunoelectrophoresis was carried out on the toxin preparations. At least nine different antigens were found in the one crude preparation and at least 25 in another. The α -toxin obtained from the former preparation by zone electrophoresis was shown to be free from the other eight antigens but was of low specific activity.

I wish to thank Dr J. J. Bullen, Head of Pathology Department, Rowett Research Institute, for his enthusiastic support; Dr D. L. Mould (now at Animal Diseases Research Association, Moredun Institute, Edinburgh) for initial guidance in the electrophoretic procedures and his stimulating interest throughout; Mr B. F. Fell of the Pathology Department of the Rowett Research Institute, for the deoxyribonuclease histochemical examination; Dr J. Conchie of the Enzymology Department of the Rowett Research Institute, for the α -mannosidase, β -galactosidase, β -N-acetylglucosaminidase and β -glucuronidase examinations; Mrs I. Batty of the Wellcome Research Laboratories for the standard α -toxin assay; the Wellcome Research Laboratories for the gift of valuable antisera and culture CN.1491; Miss Jean McBain, Mr W. G. Gordon and Mr G. M. Bruce for technical assistance, and Mr I. Robb and Mr R. Middleton for the photography.

REFERENCES

- Adams, M. H. & Hendee, E. D. (1945). *J. Immunol.* **51**, 249.
 Bidwell, E. (1950). *Biochem. J.* **46**, 589.
 Brooks, M. E., Sterne, M. & Warrack, G. H. (1957). *J. Path. Bact.* **74**, 185.
 Bullen, J. J. & Batty, I. (1956). *J. Path. Bact.* **71**, 311.
 Bullen, J. J. & Scarisbrick, R. (1957). *J. Path. Bact.* **73**, 495.
 Bullen, J. J., Thurlbourn, C. & Brown, P. (1954). *Nature, Lond.*, **173**, 254.
 Conchie, J., Findlay, J. & Levvy, G. A. (1959). *Biochem. J.* **71**, 318.
 Dorfman, A. & Ott, M. L. (1948). *J. biol. Chem.* **172**, 367.
 Flodin, P. & Kupke, D. W. (1956). *Biochim. biophys. Acta*, **21**, 368.
 Gale, E. F. & van Heyningen, W. E. (1942). *Biochem. J.* **36**, 624.
 Garbers, C. F. & Joubert, F. J. (1958). *Nature, Lond.*, **182**, 530.
 Glenny, A. T., Barr, M., Llewellyn-Jones, M., Dalling, T. & Ross, H. E. (1933). *J. Path. Bact.* **37**, 53.
 Grabar, P. & Williams, C. A. (1953). *Biochim. biophys. Acta*, **10**, 193.
 Grabar, P. & Williams, C. A. (1955). *Biochim. biophys. Acta*, **17**, 67.
 Habermann, E. (1959). *Arch. exp. Path. Pharmacol.* **235**, 513.
 Howie, J. H., Duncan, I. B. R. & Mackie, L. M. (1953). *Lancet*, ii, 1018.
 Kohn, J. (1959). *Nature, Lond.*, **183**, 1055.
 Kushner, D. J. (1957). *J. Bact.* **78**, 297.
 Levine, M. & Schoenlein, H. W. (1930). *A Compilation of Culture Media for the Cultivation of Micro-organisms*, p. 292. Baltimore: The Williams and Wilkins Co.
 Lillie, R. D. (1954). *Histopathologic Technic and Practical Histochemistry*. New York: Blackiston Co. Inc.
 Lowry, O. H., Rosebrough, N. J., Farr, A. L. & Randall, R. J. (1951). *J. biol. Chem.* **193**, 265.
 Macfarlane, M. G. (1950). *Biochem. J.* **47**, 270.
 Macfarlane, M. G. & Knight, B. C. J. G. (1941). *Biochem. J.* **35**, 884.
 Mackie, J. T. & McCartney, J. E. (1948). *Handbook of Practical Bacteriology*, 8th ed., p. 185. Edinburgh: E. and S. Livingstone Ltd.
 Mould, D. L. & Stephen, J. (1958). *Biochem. J.* **69**, 44P.
 Oakley, C. L. & Warrack, G. H. (1953). *J. Hyg., Camb.*, **51**, 102.
 Oakley, C. L., Warrack, G. H. & van Heyningen, W. E. (1946). *J. Path. Bact.* **58**, 229.
 Porath, J. (1957). *Ark. Kemi*, **11**, 161.
 Roth, F. B. & Pillemer, L. (1953). *J. Immunol.* **70**, 533.
 van Heyningen, W. E. (1941a). *Biochem. J.* **35**, 1246.
 van Heyningen, W. E. (1941b). *Biochem. J.* **35**, 1257.
 van Heyningen, W. E. (1950). *Bacterial Toxins*. Oxford: Blackwell Scientific Publications.
 van Heyningen, W. E. & Bidwell, E. (1948). *Biochem. J.* **42**, 130.