about pathways of normal metabolism (O'Brien & Peters, 1958b). It is gratifying that the investigation of the metabolism of 2-fluoroethanol should show the presence of a second alcohol dehydrogenase in liver.

SUMMARY

1. 2-Fluoroethanol, known to be toxic to rats, causes a considerable rise in tissue concentrations of citrate.

2. Guinea-pig-kidney-particle preparations are unaffected by 2-fluoroethanol, which, however, is converted by incubation with liver slices or by coupled oxidation by catalase into a compound capable of inducing citrate in the particle preparations.

3. Rat-liver slices oxidize 2-fluoroethanol to fluoroacetate, identified as the methyl ester by gas-liquid chromatography.

4. 2-Fluoroethanol is not a substrate for crystalline horse-liver alcohol dehydrogenase, and competitively inhibits the oxidation of ethanol by this enzyme.

5. A second liver alcohol dehydrogenase of different specificity readily oxidizes 2-fluoroethanol with nicotinamide-adenine dinucleotide, and to a lesser extent nicotinamide-adenine dinucleotide phosphate, as coenzymes.

6. Fluoroacetaldehyde is oxidized one and a half times as fast as is acetaldehyde by liver aldehyde dehydrogenase.

The author is deeply indebted to Sir Rudolph A. Peters, F.R.S., for his interest and advice. Thanks are due to Miss P. A. Ivall and Mrs M. A. Davies for skilled technical assistance and to the Wellcome Trust for a Research Fellowship during part of this work.

REFERENCES

- Bartlett, G. R. (1952). *J. Pharmacol.* **106**, 464.
 Bonnichsen, R. K. & Wassen, A. M. (1948). *Arch. Biochem.* **18**, 361.
 Bruce, H. M. & Parkes, A. S. (1949). *J. Hyg., Camb.*, **47**, 201.
 Dalziel, K. (1958). *Acta chem. scand.* **12**, 459.
 Keilin, D. & Hartree, E. F. (1945). *Biochem. J.* **39**, 293.
 Liébecq, C. & Peters, R. A. (1949). *Biochim. biophys. Acta*, **3**, 215.
 Lundsgaard, E. (1938). *Johns Hopk. Hosp. Bull.* **63**, 15.
 McArdle, B. (1955). *Biochem. J.* **60**, 647.
 McCombie, H. & Saunders, B. C. (1946). *Nature, Lond.*, **158**, 382.
 Mirsky, I. A. & Nelson, N. (1939). *Amer. J. Physiol.* **126**, 587.
 O'Brien, R. D. & Peters, R. A. (1958a). *Biochem. Pharmacol.* **1**, 3.
 O'Brien, R. D. & Peters, R. A. (1958b). *Biochem. J.* **70**, 188.
 Pattison, F. L. M., Howell, W. C., McNamara, A. J., Schneider, J. C. & Walker, J. F. (1956). *J. org. Chem.* **21**, 739.
 Peters, R. A. (1952). *Proc. Roy. Soc. B*, **139**, 143.
 Peters, R. A. & Wakelin, R. W. (1957). *Biochem. J.* **67**, 280.
 Racke, E. (1949). *J. biol. Chem.* **177**, 883.
 Saunders, B. C., Stacy, G. J. & Wilding, I. G. E. (1949). *J. chem. Soc.* p. 773.
 Schrader, G. (1935). *I. G. Leverkusen German Patent J.* 52803, 11 July.
 Singer, T. P. & Kearney, E. B. (1957). *Meth. biochem. Anal.* **4**, 307.
 Swarts, F. (1914). *Bull. Acad. Belg. Cl. Sci.* p. 7.
 Theorell, N. & Bonnichsen, R. (1951). *Acta chem. scand.* **5**, 1105.
 Umbreit, W. W., Burris, R. M. & Stauffer, J. F. (1957). *Manometric Techniques*, p. 149. Minneapolis: Burgess Publishing Co.

Biochem. J. (1962) **82**, 134

The Effect of Dinitrophenol and Amytal on the Reduction of Acetoacetate in the Presence of Succinate

By H. A. KREBS AND L. V. EGGLESTON

Medical Research Council Unit for Research in Cell Metabolism, Department of Biochemistry, University of Oxford

(Received 21 July 1961)

Chance & Hollunger (1957, 1960, 1961), Chance & Hagihara (1960), Chance (1961a, b), Chance & Fugmann (1961), Klingenberg, Slenczka & Ritt (1959), Azzone, Ernster & Klingenberg (1960), Azzone & Ernster (1961a, b), and Klingenberg & Schollmeyer (1960, 1961) have put forward the view that the reduction of diphosphopyridine

nucleotide observed on addition of succinate to mitochondria involves a reversal of oxidative phosphorylation. Evidence in favour of another mechanism, a competitive inhibition of the oxidation of reduced diphosphopyridine nucleotide by succinate, was put forward by Birt & Bartley (1960), and by Kulka, Krebs & Eggleston (1961)

and Krebs, Eggleston & d'Alessandro (1961), who investigated the effect of succinate on the diphosphopyridine nucleotide-dependent reduction of acetoacetate to β -hydroxybutyrate in tissue homogenates.

Further evidence against a major role of a reversal of oxidative phosphorylation is provided by experiments reported in this paper. It is shown that under suitable conditions high concentrations of 2,4-dinitrophenol (0.5 mM) plus Amytal (1.25 mM), which strongly inhibit oxidative phosphorylation, do not necessarily inhibit the reduction of acetoacetate (i.e. diphosphopyridine nucleotide) in the presence of succinate.

EXPERIMENTAL

Methods. The materials and procedures were as described by Krebs *et al.* (1961). The temperature of incubation was 20°, unless otherwise stated, because the rapid reaction rates at body temperature made the accurate measurements of rates in some experiments difficult.

The adenosine phosphates and inorganic phosphate were separated by paper electrophoresis on Whatman no. 1 paper, 65 cm. $\times$ 10 cm. Samples of trichloroacetic acid extracts (usually 50 μ l.) were dried on spots (1 cm. wide) on the starting line 10 cm. from one end of the paper. Drying was hastened with cold air from a hair drier. The Locarte electrophoresis apparatus (The Locarte Co., 24 Emperor's Gate, London, S.W. 7) was used. The runs were carried out in 0.1M-sodium citrate buffer, pH 3.5, for 50 min., at 8 kv and about 40 ma (see also Webster, 1953; Wade & Morgan, 1954). After drying, the papers were examined under u.v. light and absorbing areas due to purine derivatives were ringed with pencil. The papers were then sprayed with the acid molybdate reagent of Hanes & Isherwood (1949) and phosphate was determined in the spots by the method of Berenblum & Chain (1938) as modified by Bartley (1953). The average distances travelled

by the compounds were (cm.): adenosine, inosine, adenine -2 to +1.5; adenosine monophosphate, +5; inosine monophosphate, +16.5; adenosine diphosphate, +19.5; adenosine triphosphate, +27; inorganic phosphate, +31.

RESULTS

Effects of 2,4-dinitrophenol and Amytal on the metabolism of succinate and acetoacetate. Rat-liver homogenate was incubated with acetoacetate and succinate, and the effects of Amytal and dinitrophenol, separately and jointly, on the metabolic changes were measured (Table 1). Dinitrophenol alone caused a 37% stimulation of the oxygen uptake at 0.05 mM and a 15% inhibition at 0.5 mM. Large amounts of succinate were removed and converted mainly into fumarate and malate. The increased oxygen uptake at 0.05 mM-dinitrophenol was not accompanied by an increased succinate removal, and at 0.5 mM-dinitrophenol the inhibition of succinate removal was much greater (43%) than inhibition of the oxygen uptake (15%). The differential effect of dinitrophenol on the oxygen and succinate removal shows itself clearly by the change of the ratio (succinate used)/(oxygen used) shown in the last horizontal line of Table 1. Dinitrophenol decreases this ratio. The fall of the ratio indicates that less of the oxygen uptake was due to the oxidation of succinate to fumarate and a greater proportion to the oxidation of other substrates such as the fumarate formed from succinate. Both concentrations of dinitrophenol greatly decreased the rate of acetoacetate reduction. It was already reported by Kulka *et al.* (1961) and Krebs *et al.* (1961) that dinitrophenol had such effects whenever it accelerated the rate of respiration. What may at first be unexpected is the

Table 1. *Effects of 2,4-dinitrophenol and Amytal on the metabolism of rat-liver homogenates in the presence of succinate and acetoacetate*

A 10% homogenate was incubated at 30° for 30 min. in an atmosphere of O ₂ . To all vessels 19.5 μ moles of acetoacetate and 80 μ moles of succinate were added. The data refer to 4 ml.						
Vessel no.	1	2	3	4	5	6
Further additions (final concn.)		Amytal (1.25 mM)	Dinitro- phenol (0.05 mM)	Amytal (1.25 mM) + dinitro- phenol (0.05 mM)	Dinitro- phenol (0.5 mM)	Amytal (1.25 mM) + dinitro- phenol (0.5 mM)
Metabolic changes:						
O ₂ (μ moles)	-38.3	-38.8	-52.0	-44.3	-32.6	-41.2
Acetoacetate (μ moles)	-10.2	-9.4	-3.1	-12.3	-0.2	-7.3
β -Hydroxybutyrate (μ moles)	+9.5	+9.3	+2.9	+11.6	+1.7	+9.2
Succinate (μ moles)	-71.8	-70.6	-72.7	-77.2	-41.2	-72.5
Fumarate + malate (μ moles)	+57.5	+54.3	+55.5	+58.9	+23.7	+51.3
Pyruvate (μ moles)	+1.1	+1.6	+0.5	+0.1	+0.1	+0.1
α -Oxoglutarate (μ mole)	+0.4	+0.3	+0.5	+0.05	+0.2	+0.1
Citrate (μ moles)	+0.5	+0.7	+0.8	+0.5	+2.4	+1.7
Succinate used	1.87	1.82	1.40	1.75	1.26	1.76
O ₂ used						

inhibition of acetoacetate reduction at 0.5 mM-dinitrophenol when it did not accelerate but inhibited respiration. However, as 0.5 mM-dinitrophenol inhibited the oxidation of succinate a decreased reduction of acetoacetate is not surprising because the accelerating effect of succinate depends on the ready oxidation of this substrate.

Amytal alone had no major effects when succinate was present (as is well established), but in the presence of dinitrophenol Amytal had several effects which all represent reversals of the action of dinitrophenol: it abolished (1) the stimulation of respiration at 0.05 mM-dinitrophenol; (2) the inhibition of respiration at 0.5 mM-dinitrophenol; (3) the inhibition of the reduction of acetoacetate at both dinitrophenol concentrations; (4) the inhibition of succinate oxidation at 0.5 mM-dinitrophenol.

Effects of dinitrophenol and Amytal on oxidative phosphorylation. These results raise the question of whether, in the presence of succinate, Amytal also removes the uncoupling of oxidative phosphorylation by dinitrophenol. The effects of dinitrophenol and Amytal on oxidative phosphorylation were therefore tested under conditions which were close to those of the previous experiment. Adenosine triphosphate was added to liver homogenates in the presence of succinate and the changes in the concentrations of adenosine tri-, di- and mono-phosphate and inorganic phosphate on incubation were measured. Anaerobically adenosine triphosphate is rapidly lost in such homogenates but aerobically its concentration is maintained as long as oxidative phosphorylation is intact. Inhibition of oxidative phosphorylation therefore shows itself by the formation of inorganic phosphate, a loss of adenosine triphosphate and a gain in the products of dephosphorylation of adenosine triphosphate. The results of an experiment testing the effects of dinitrophenol and Amytal is shown in Table 2.

The added adenosine triphosphate contained some adenosine diphosphate (see column 0 min.) and the homogenate converted most of this into adenosine triphosphate within the first 15 min. After 15 min. there were no major changes in the concentrations of adenosine di- and tri-phosphate in the absence of inhibitors. Amytal did not affect the oxygen consumption or the adenosine tri- and di-phosphate concentrations, whereas dinitrophenol, as expected, caused a rapid fall of adenosine triphosphate during the first 30 min. After this the adenosine triphosphate concentration remained approximately constant at about 12% of the original value. The fall of adenosine triphosphate was accompanied by a rise of adenosine di- and mono-phosphate but much of the adenosine triphosphate loss was not accounted for by adenosine di- or mono-phosphate, and the forma-

tion of inorganic phosphate indicated that adenosine monophosphate underwent further dephosphorylation. The sum of adenosine tri-, di- and mono-phosphate and inorganic phosphate remained approximately constant.

Amytal had no effect on the changes in the concentrations of phosphate caused by dinitrophenol; thus it did not reverse the inhibition of oxidative phosphorylation.

The effects of dinitrophenol and Amytal on oxidative phosphorylation were also measured by a standard procedure which included the addition of fluoride, glucose and hexokinase to the medium (for details see Table 3). The rate of oxidative phosphorylation was determined by the disappearance of phosphate. In this test Amytal alone had no effect. In Expt. I dinitrophenol concentrations of 0.05 and 0.5 mM inhibited the phosphate removal by 73 and 86% respectively and these inhibitions were reduced by the addition of Amytal. The effect was slight (and possibly not significant) at 0.05 mM-dinitrophenol, but it was definite at 0.5 mM-dinitrophenol. In this experiment the P/O values were low because the supply of phosphate was insufficient for an incubation period of 60 min. In Expt. II where the total incubation time was 15 min., normal P/O values were obtained in the absence of dinitrophenol. The inhibitions were similar to those in Expt. I.

The two procedures thus give the clear-cut result that Amytal does not reverse the inhibition of oxidative phosphorylation by dinitrophenol. The degree of inhibition by 0.5 mM-dinitrophenol was approximately equal in both experiments: as judged by the amount of adenosine triphosphate remaining after 30 min. when a steady state of adenosine triphosphate had been reached the inhibition was about 88% in the experiment of Table 2, and it was 86% in Expt. I of Table 3 as judged by the phosphate uptake. The incompleteness of the inhibition can hardly be ascribed to the dinitrophenol-insensitive step because only one phosphorylating step, that associated with the reaction

is not affected by dinitrophenol, and this step contributed probably less than 3% to the total phosphorylating steps. This leads to the conclusion that the inhibition of the steps of oxidative phosphorylation which are sensitive to dinitrophenol is still incomplete at 0.5 mM-dinitrophenol.

Further observations on the effect of dinitrophenol on the removal of succinate. The inhibition of succinate oxidation by high concentrations of dinitrophenol is not dependent on the presence of acetoacetate, as shown in Table 4. Various concentrations of dinitrophenol were added to liver homo-

Table 2. Disappearance of ATP and ADP in the presence of dinitrophenol and Amytal

A 10% rat-liver homogenate was incubated at 20° in O₂. The data refer to 4 ml., which included 120 μmoles of succinate and 20 μmoles of 'ATP' containing some ADP (see zero values). Substrates and inhibitors were added from the side arm after 10 min. equilibration. The enzymic reactions were stopped by the addition of 0.5 ml. of 30% trichloroacetic acid. Supernatant (20 μl.) was used for the determination of inorganic phosphorus and 50 μl. for separation of nucleotides by paper electrophoresis. The zero values refer to an unincubated homogenate to which ATP had been added. Where added dinitrophenol was 0.5 mm and Amytal 1.25 mm.

Time of incubation (min.)		0	15			30			45		
Inhibitors added	...	None	None	Amytal	Dinitro-phenol + Amytal	None	Amytal	Dinitro-phenol + Amytal	None	Amytal	Dinitro-phenol + Amytal
O ₂ uptake (μmoles)	...	—	10.4	10.5	6.75	6.88	21.6	22.4	33.4	33.9	18.0
Inorganic phosphorus found (μmoles)	27.2	27.2	21.5	19.9	36.8	35.8	23.4	22.5	25.2	23.0	43.1
Inorganic phosphorus change (μmoles)	—	—	-5.7	-7.3	+9.6	+8.6	-3.8	-4.7	-2.0	-4.25	+15.9
ATP phosphorus found (μmoles)	19.5	19.5	24.0	22.2	7.50	8.74	19.7	24.0	22.2	21.6	3.04
ADP phosphorus found (μmoles)	4.51	4.51	2.90	6.35	2.78	2.06	7.35	3.36	3.87	2.61	7.30
Sum of ATP and ADP phosphorus found (μmoles)	24.0	24.0	26.9	28.5	10.3	10.8	27.1	27.4	26.1	24.2	10.3
Sum of ATP and ADP phosphorus change (μmoles)	—	—	+2.89	+4.54	-13.7	-13.2	+3.10	+3.40	+2.11	+0.20	-13.6
AMP phosphorus found	0	0	0	5.94	5.49	5.49	0	0	0	0	1.19
Total P recovered as ATP, ADP, AMP, P (μmoles)	51.2	51.2	48.4	48.4	53.0	52.1	50.5	49.9	51.3	47.2	54.6

Table 3. Effect of dinitrophenol and Amytal on oxidative phosphorylation in rat-liver homogenate

The homogenate (final concentration 10%) included NaF (0.04M), glucose (0.05M) and hexokinase (2 mg.). Succinate (80 μmoles), ATP (20 μmoles), Amytal and dinitrophenol were added from the side arm after 10 min. equilibration. Further incubation for 60 min. (Expt. I) or 15 min. (Expt. II). The gas space contained O₂; temperature, 20°. The reaction was stopped with 0.5 ml. of 30% trichloroacetic acid. Inorganic phosphate was determined in 0.1 ml. of the filtrate. Initial inorganic phosphate content: 18.4 μmoles. All data refer to 4 ml.

Cup no.	Inhibitors added (final concn.)	Expt. I				Expt. II			
		O ₂ uptake (μmoles)		Decrease in inorganic P in 60 min. (μmoles)	Inhibition of phosphate removal (%)	O ₂ uptake (15 min.) (μmoles)	Decrease in inorganic P (μmoles)	Inhibition of oxidative phosphorylation (%)	
		First 30 min.	Second 30 min.						P/O
1	None	7.36	5.34	-16.5	—	3.88	-14.28	1.84	—
2	Amytal (1.25 mm)	7.64	5.35	-17.2	0	4.50	-14.25	1.58	14
3	Dinitrophenol (0.05 mm)	5.80	3.22	-4.4	73	4.15	-4.80	0.58	69
4	Dinitrophenol (0.5 mm)	5.86	2.86	-2.3	86	4.42	-1.88	0.21	89
5	Dinitrophenol (0.05 mm) + Amytal (1.25 mm)	6.80	3.50	-5.3	68	5.18	-5.06	0.49	74
6	Dinitrophenol (0.5 mm) + Amytal (1.25 mm)	6.50	3.35	-4.8	71	4.78	-4.29	0.45	76

genates, with succinate as the sole added respiratory substrate. At lower concentrations (0.05 mM or less) dinitrophenol stimulated the rate of oxygen consumption and of succinate removal. The maximum stimulation of the oxygen consumption was at 0.05 mM when it amounted to 58 %, whereas the rate of succinate removal was increased by only 14 %. At the higher dinitrophenol concentrations (0.25 mM) the removal of succinate was inhibited and the inhibition was greater than that of the oxygen uptake. The gradual decrease of the ratio (succinate removed)/(oxygen used) shows that the conversion of succinate into fumarate, which was the main oxidative reaction in the absence of dinitrophenol, was increasingly replaced by other oxidative reactions as the dinitrophenol concentration rose.

In this experiment the initial concentration of succinate chosen was high in order to maintain a supply of substrate for at least 1 hr. In a similar experiment the concentration of succinate was reduced to one-quarter and the oxygen consumption was measured over the first 15 min. The stimulation of the oxygen consumption at low dinitrophenol concentration was less (23 compared with 59 %) but the inhibitions at high concentrations were of the same order. Thus the effect of

dinitrophenol was essentially independent of the substrate concentration.

Inhibition of the oxygen uptake by adenosine triphosphate. In the experiment shown in Table 2, in contrast with that of Table 1, Amytal did not stimulate the oxygen consumption in the presence of dinitrophenol. It was confirmed in other experiments (see, for example, Table 5) that the stimulation of respiration by Amytal in the presence of dinitrophenol is abolished by adenosine triphosphate. In other words, adenosine triphosphate inhibits the oxygen uptake in the presence of dinitrophenol plus Amytal, but does not do so in the absence of these substances. Adenosine diphosphate acted like adenosine triphosphate but adenosine monophosphate and creatine phosphate had no effect. Pyrophosphate slightly inhibited the stimulating effect of Amytal.

Joint effects of dinitrophenol and Amytal on the oxidation of substrates other than succinate. When succinate was replaced by other substrates, Amytal inhibited the oxygen uptake by about 70–80 %. Dinitrophenol at 0.5 mM raised the rate of oxygen consumption in the presence of citrate, glutamate and α -oxoglutarate but this stimulation was completely prevented by Amytal. This is in general agreement with the view that in the presence of

Table 4. *Effect of various dinitrophenol concentrations on the oxidation of succinate in rat-liver homogenates*

Data refer to 4 ml. of 10 % rat-liver homogenate which included 120 μ M-succinate. Succinate and dinitrophenol were added from the side arm after 10 min. equilibrations. The gas space contained O_2 ; temperature, 20°. Incubation was for 60 min. after mixing.

Concn. of dinitrophenol (mM)	Metabolic changes					Succinate removed O_2 used
	O_2 (μ moles)	Succinate (μ moles)	Fumarate + malate (μ moles)	Citrate (μ moles)	α -Oxo- glutarate (μ mole)	
0	-40.8	-79.3	+68.1	+0.56	+0.10	1.94
0.002	-47.2	-86.6	+75.6	+0.66	+0.25	1.83
0.01	-55.6	-95.5	+87.0	+1.2	+0.20	1.71
0.05	-64.9	-90.5	+75.0	+1.3	+0.13	1.40
0.25	-28.0	-30.5	+24.3	+1.6	+0.07	1.09

Table 5. *Effects of dinitrophenol and Amytal on the oxygen uptake and release of inorganic phosphate in the presence of ATP in rat-liver homogenates*

Homogenate (4 ml.; final concentration 10 %) was incubated at 20° in O_2 . All additions were mixed from the side arm after 10 min. equilibration. Further incubation was for 60 min. The reaction was stopped by the addition of 0.5 ml. of trichloroacetic acid (30 %).

Cup no.	Additions other than succinate (20 mM)	O_2 uptake (μ moles)	Increase in inorganic P due to ATP breakdown (μ moles)
1	None	38.8	—
2	Amytal (1.25 mM)	39.3	—
3	Dinitrophenol (0.5 mM)	23.5	—
4	Dinitrophenol (0.5 mM) + Amytal (1.25 mM)	35.4	—
5	ATP (2 mM)	38.0	+ 0.3
6	ATP (2 mM) + Amytal (1.25 mM)	37.4	+10.1
7	ATP (2 mM) + dinitrophenol (0.5 mM)	19.1	+22.8
8	ATP (2 mM) + dinitrophenol (0.5 mM) + Amytal (1.25 mM)	20.1	+22.9

Amytal electron transport from reduced diphosphopyridine nucleotide is no longer controlled by the concentration of adenosine diphosphate or phosphate, or both, but by the capacity of the electron carriers themselves which is decreased by Amytal. The stimulation of the oxidation of various substrates which occurs at lower dinitrophenol concentrations was also abolished by Amytal, except with succinate.

Effects of dinitrophenol and Amytal in mitochondria. The effects of dinitrophenol, Amytal and adenosine triphosphate were also investigated in rat-liver mitochondria prepared according to Aldridge (1957). When succinate was the sole added substrate, the results obtained were analogous to those with homogenates (Table 6). Dinitrophenol (0.5 mM) inhibited the oxygen uptake by 13% but the oxidation of succinate by 54%. Amytal alone slightly inhibited the uptake of oxygen and succinate without decreasing the formation of fumarate and malate, so that in the presence of Amytal almost the whole oxygen consumption was due to the conversion of succinate into fumarate. Dinitrophenol increased the rate of oxidation of succinate to fumarate by about 50% when Amytal was present. In other words dinitrophenol inhibited the oxidation of succinate when added alone but increased it in the presence of Amytal. The main effect of adenosine triphosphate was to prevent partially the stimulation by dinitrophenol of the oxygen uptake and succinate removal in the presence of Amytal. The experiment reported here in detail was carried out at 20°. At 40° the effects of dinitrophenol, Amytal and adenosine triphosphate were of the same type.

Dependence of the action of dinitrophenol on the treatment of the tissue preparation. The effect of dinitrophenol depended on the treatment of the respiring material. Deep-freezing and thawing a homogenate, or centrifuging the insoluble particles and washing them twice with a saline medium (see Table 6), caused a fall of about 40% in the rates of the oxygen uptake and the succinate removal

(Table 7). The respiration that remained was much less sensitive to 1 mM-dinitrophenol. In the fresh homogenate, dinitrophenol inhibited the oxygen uptake by 37%, and in the washed particles and in the deep-freeze-treated material by only 14 and 17% respectively. The removal of succinate was inhibited by 63% in the fresh homogenate and by 5 and 13% respectively in the washed particles and deep-freeze-treated homogenate. The differences in the effects of dinitrophenol are clearly expressed by the ratio (succinate removed)/(oxygen used). In fresh homogenates the ratio fell from 1.60 to 0.94 on addition of dinitrophenol. It was slightly increased in the washed particles (1.84–2.02) and in the deep-freeze-treated homogenate (1.48–1.55). As already shown (Table 6) mitochondria prepared according to Aldridge behave similarly to the fresh homogenate, but when mitochondria were washed with iso-osmotic potassium chloride solution and suspended in saline medium their response to dinitrophenol was like that of the washed particles or the deep-freeze-treated homogenate in Table 7.

When both succinate and acetoacetate were added to mitochondria the response differed from that of homogenates in respect of the reduction of acetoacetate (see Table 8). Both dinitrophenol and Amytal, when added individually, strongly inhibited the reduction but when added jointly the inhibition was relatively small.

The effect of Amytal on the reduction of acetoacetate depended on the concentration of succinate in the homogenate but not in mitochondrial suspensions (see Table 9). Amytal accelerated the removal (i.e. reduction) of acetoacetate in the absence of succinate, especially in the homogenate. In the mitochondria Amytal inhibited the reduction about 60% within a wide range of succinate concentrations. In the homogenate there was an 80% acceleration at 10 mM-succinate but higher concentrations caused increasing inhibitions.

Related observations. A number of related findings may be briefly mentioned. Dicoumarol showed the

Table 6. *Effects of dinitrophenol and Amytal on rat-liver mitochondria in the presence and absence of ATP*

The data refer to 4 ml. of suspension containing 80 μ moles of succinate and 30.3 mg. dry wt. of mitochondria. Incubation was at 20° for 60 min. in O ₂ . The final concentrations of dinitrophenol were 0.5 mM, of Amytal 1.25 mM and of ATP 1 mM. The mitochondria were prepared according to Aldridge (1957) in 0.3 M-sucrose. The mitochondria from 12 g. of liver from two rats were suspended in a total volume of 13 ml. of 0.3 M-sucrose and 1 ml. of this was used per cup. The saline medium used for dilution contained 0.154 M-KCl, 0.025 M-MgCl ₂ , 0.023 M-potassium phosphate buffer, pH 6.7, and neutralized 1.5 mM-ethylenediaminetetra-acetic acid.									
Other substances added	...	None	Dinitrophenol	Amytal	Dinitrophenol + Amytal	ATP	ATP + dinitrophenol	ATP + Amytal	ATP + dinitrophenol + Amytal
Metabolic changes									
O ₂ (μ moles)		-17.7	-15.4	-15.2	-23.1	-13.7	-12.9	-18.9	-18.5
Succinate (μ moles)		-40.3	-18.7	-32.2	-45.2	-35.5	-20.0	-35.6	-27.1
Fumarate + malate (μ moles)		+33.2	+20.3	+35.3	+44.8	+32.6	+17.5	+31.5	+27.8

Table 7. *Effects of dinitrophenol on the oxygen uptake and succinate removal in various rat-liver preparations*

Rat liver was homogenized with 6.5 vol. of saline medium. Part of the homogenate (fraction A) was used directly. Another (fraction B) was centrifuged at high speed for 7 min. at 0°; the sediment was washed twice with the saline medium and finally suspended in the original volume of saline. Fraction C was identical with A except that it was kept at -15° for 18 hr. and then thawed. Manometer cups containing 3 ml. of each fraction and 1 ml. additions including 80 μ moles of succinate. Dinitrophenol (1 mM final concentration) and succinate were added from the side arm after 10 min. equilibration. The gas space contained O₂; temperature, 20°. Incubation after mixing was for 30 min.

Fraction	...	...	...	...	A		B		C	
Concn. of dinitrophenol (mM)	...				0	1	0	1	0	1
Changes:										
O ₂ (μ moles)					-21.2	-13.4	-11.0	-9.5	-12.8	-10.6
Succinate (μ moles)					-34.0	-12.6	-20.3	-19.3	-19.0	-16.5
Inhibition by dinitrophenol:										
O ₂ uptake (%)					—	37	—	14	—	17
Succinate removal (%)					—	63	—	5	—	13
Succinate removed										
O ₂ used					1.60	0.94	1.84	2.02	1.48	1.55

Table 8. *Effects of dinitrophenol and Amytal on the metabolism of rat-liver mitochondria in the presence of succinate and acetoacetate*

Mitochondria were prepared according to Aldridge (see legend of Table 6). The data refer to 4 ml. of suspension containing 120 μ moles of succinate and 20 μ moles of acetoacetate. The final concentration of dinitrophenol was 0.5 mM and of Amytal 1.25 mM. Incubation was at 30° for 30 min. in O₂.

Inhibitors added	...	...	None	Dinitrophenol	Amytal	Dinitrophenol + Amytal
Metabolic changes:						
O ₂ (μ moles)			-33.1	-29.0	-42.8	-37.8
Succinate (μ moles)			-67.5	-44.6	-72.8	-63.5
Acetoacetate (μ moles)			-11.4	-0.5	-3.2	-7.0
β -Hydroxybutyrate (μ moles)			+8.3	+1.2	+2.6	+6.3

Table 9. *Effect of Amytal on acetoacetate reduction in rat-liver preparations at different succinate concentrations*

Each cup contained 4 ml. of 10% homogenate or mitochondria prepared as in Table 6, including 20 μ moles of acetoacetate. The dry weight of mitochondria was 22.5 mg./cup. Incubation was for 30 min. in O₂ at 30°.

		Metabolic changes					
		Homogenate			Mitochondria		
Final concn. of succinate (mM)	Amytal concn. (mM)	O ₂ uptake (μ moles)	Acetoacetate removal (μ moles)	Effect of Amytal on acetoacetate removal (%)	O ₂ uptake (μ moles)	Acetoacetate removal (μ moles)	Effect of Amytal on acetoacetate removal (%)
0	0	16.8	-0.16	—	0.74	-0.7	—
0	1.25	3.7	-5.23	—	0.83	-1.5	—
5	0	—	—	—	9.65	-5.4	—
5	1.25	—	—	—	11.7	-2.3	-57
10	0	30.6	-8.1	—	19.3	-6.1	—
10	1.25	28.3	-14.1	+80	21.4	-2.7	-56
15	0	36.2	-14.1	—	23.8	-5.4	—
15	1.25	36.3	-11.8	-17	26.1	-1.9	-65
20	0	41.3	-15.0	—	31.8	-9.4	—
20	1.25	40.1	-11.5	-23	37.8	-3.7	-61
30	0	37.2	-17.7	—	33.1	-11.4	—
30	1.25	39.1	-10.1	-42	42.5	-3.18	-72
40	0	39.8	-16.8	—	—	—	—
40	1.25	35.9	-7.9	-53	—	—	—

Table 10. *Effects of 2,4-dinitrophenol and Amytal on the metabolism of sheep-heart homogenate in the presence of succinate and acetoacetate*
A 10% homogenate was incubated at 30° for 30 min. in an atmosphere of O₂. To all vessels 23.1 μmoles of acetoacetate and 120 μmoles of succinate were added. The data refer to 4 ml.

Vessel no.	1	2	3	4	5	6	7	8
Further additions	...	...	...	...	...	...	...	...
	None	Amytal (1.25 mm)	Dinitro- phenol (0.05 mm)	Dinitro- phenol (0.5 mm)	Dinitro- phenol (1 mm)	Dinitro- phenol (0.05 mm) + Amytal (1.25 mm)	Dinitro- phenol (0.5 mm) + Amytal (1.25 mm)	Dinitro- phenol (1 mm) + Amytal (1.25 mm)
Metabolic changes:								
O ₂ (μmoles)	-46.0	-44.2	-42.6	-37.9	-35.2	-59.2	-43.1	-37.1
Acetoacetate (μmoles)	-9.6	-10.9	-4.3	-3.1	-2.5	-7.9	-5.5	-5.0
β-Hydroxybutyrate (μmoles)	+	+10.3	+3.5	+2.3	+3.1	+	+8.0	+7.1
Succinate (μmoles)	-63.0	-74.8	-42.3	-49.3	-43.1	-78.5	-57.8	-57.0
Fumarate + malate (μmoles)	+52.0	+60.7	+27.2	+29.9	+29.5	+68.0	+49.1	+43.0

same effects in homogenates as dinitrophenol in the presence of succinate, acetoacetate and Amytal: 0.5 mm-dicoumarol inhibited the oxygen uptake much more than the removal of succinate, and Amytal abolished these inhibitions. Some experiments were repeated with suspensions of minced pigeon-breast muscle and sheep-heart homogenate. These tissues gave responses to dinitrophenol and to Amytal, and to mixtures of these, which were similar to those of rat liver. An example is given in Table 10.

In the presence of succinate the inhibition of respiration by dinitrophenol in rat-liver homogenates was essentially independent of the succinate concentration, between 3.75 and 30 mm, except that during the first 15 min. the higher concentrations of succinate delayed the onset of the inhibition. Dinitrophenol (1 mm) caused an inhibition of about 50 % after 15 min. and this remained at the same level during the subsequent 30 min.

DISCUSSION

Role of a reversal of oxidative phosphorylation in the reduction of acetoacetate. The fact that 88 % inhibition of oxidative phosphorylation did not affect the rate of reduction of acetoacetate indicates that oxidative phosphorylation was not of importance in the mechanism of reduction under the test conditions. This does not mean that oxidative phosphorylation could not play a part in the reduction under other conditions. The main difference between the conditions of the present experiments and those used by other authors concerns the period of incubation which was generally much longer in the present experiments (30–60 min. against 1–2 min.). A reversal of oxidative phosphorylation leading to reductions is theoretically conceivable (see Davies & Krebs, 1952; Krebs & Kornberg, 1957), and may especially be expected to operate in short-term experiments. Whilst several of the present observations could be explained either by a reversal of oxidative phosphorylation or by the kinetic concept, the fact that the reduction of acetoacetate was not necessarily diminished when oxidative phosphorylation was strongly inhibited cannot be reconciled with the assumption that oxidative phosphorylation was of major importance.

Inhibitory effects of dinitrophenol on the oxidation of succinate. Higher concentrations of dinitrophenol inhibit the oxygen consumption under certain conditions depending especially on the nature of the available substrate (see Kulka *et al.* 1961). These inhibitions by dinitrophenol are difficult to explain on the assumption that uncoupling of oxidative phosphorylation is the only action of dinitrophenol. They have been attributed

to the loss of adenosine phosphates (Emmelot & Bos, 1959), although it is by no means clear why a fall of adenosine phosphates should cause a loss of enzymic activity. The data on the fate of added adenosine triphosphate (Table 2) confirm that adenosine phosphates are degraded beyond the stage of adenosine monophosphate in the presence of high concentrations of dinitrophenol.

Among the observations which cannot yet be satisfactorily explained are the following. Dinitrophenol inhibits the oxidation of succinate more strongly than that of other substrates. This inhibition is reversed by Amytal and this reversal is abolished by adenosine triphosphate. All these effects of dinitrophenol and Amytal are found only in cell material which has been treated with special care. They disappear on washing mitochondria with saline media or on freezing, and purified preparations of succinic dehydrogenase are not at all affected by dinitrophenol. These facts indicate that the activity of the succinic dehydrogenase system *in situ* depends on agents other than the enzyme itself. It may be that in the intact mitochondrial structure access of the substrate to the enzyme is limited and controlled. Mitochondria as a whole are known to shrink and to swell, but measurements of the water content of the mitochondria in the presence of dinitrophenol and Amytal showed no parallelism between swelling and the rate of oxidation of succinate. This does not exclude the possibility, however, that regional swelling within the mitochondria, not detectable by measuring the total water content, may influence enzyme activity.

Observations of Schollmeyer & Klingenberg (1961) suggest that oxaloacetate formed from succinate may be the immediate cause of an inhibition of succinate oxidation but this was not borne out by the experimental test. The concentration of oxaloacetate which accumulated under test conditions was less than 0.05 mM and addition even of 10 mM-oxaloacetate did not inhibit the disappearance of succinate in the homogenates.

Some of the present observations differ from those reported by Azzone & Ernster (1961c) working with a somewhat different system. Thus under their conditions preincubation of rat-liver mitochondria with 2 mM-arsenate and 0.1 mM-dinitrophenol reduced the capacity for oxidizing succinate; this was restored by the addition of adenosine triphosphate.

SUMMARY

1. Dinitrophenol at high concentrations (0.5 mM) inhibits the oxygen uptake of liver homogenates in the presence of succinate. It also inhibits the removal of succinate and this inhibition was

greater than that of the oxygen uptake. At a concentration of 0.05 mM dinitrophenol increased the rate of the oxygen uptake without increasing the rate of succinate removal. Thus dinitrophenol shifts the substrate of oxidation from succinate towards other substances.

2. Amytal when added together with dinitrophenol abolished the following effects of dinitrophenol: (a) the stimulation of the oxygen consumption at 0.05 mM-dinitrophenol; (b) the inhibition of the oxygen uptake at 0.5 mM-dinitrophenol; (c) the inhibition of the reduction of acetoacetate at both dinitrophenol concentrations; (d) the inhibition of succinate oxidation at 0.5 mM-dinitrophenol. It did not remove the inhibition of oxidative phosphorylation by dinitrophenol.

3. The stimulation of respiration caused by Amytal in the presence of 0.5 mM-dinitrophenol was prevented by adenosine triphosphate. Thus adenosine triphosphate inhibited the oxygen uptake in the presence of dinitrophenol plus Amytal but not in the absence of these substances.

4. Liver mitochondria prepared in iso-osmotic sucrose solution reacted in the same way as homogenates towards dinitrophenol, Amytal and adenosine triphosphate. Mitochondria prepared in saline media behaved differently.

5. Homogenates of pigeon-breast muscle and sheep heart showed the same response as rat liver to dinitrophenol and Amytal.

6. The fact that a reduction of acetoacetate to β -hydroxybutyrate can readily occur in the presence of dinitrophenol and Amytal supports the view put forward previously that under the test conditions a reversal of oxidative phosphorylation does not play a major part in the reduction of acetoacetate.

This work was aided by grants from the Rockefeller Foundation and the National Institutes of Health, United States Public Health Service.

REFERENCES

- Aldridge, W. N. (1957). *Biochem. J.* **67**, 423.
- Azzone, G. F. & Ernster, L. (1961a). *J. biol. Chem.* **236**, 1501.
- Azzone, G. F. & Ernster, L. (1961b). *J. biol. Chem.* **236**, 1510.
- Azzone, G. F. & Ernster, L. (1961c). *J. biol. Chem.* **236**, 1518.
- Azzone, G. F., Ernster, L. & Klingenberg, M. (1960). *Nature, Lond.*, **188**, 552.
- Bartley, W. (1953). *Biochem. J.* **54**, 677.
- Berenblum, I. & Chain, E. (1938). *Biochem. J.* **32**, 295.
- Birt, L. M. & Bartley, W. (1960). *Biochem. J.* **76**, 427.
- Chance, B. (1961a). *Nature, Lond.*, **189**, 719.
- Chance, B. (1961b). *J. biol. Chem.* **236**, 1544, 1569.
- Chance, B. & Fugmann, U. (1961). *Biochem. biophys. Res. Commun.* **4**, 317.

- Chance, B. & Hagihara, B. (1960). *Biochem. biophys. Res. Commun.* **3**, 6.
- Chance, B. & Hollunger, G. (1957). *Fed. Proc.* **16**, 163.
- Chance, B. & Hollunger, G. (1960). *Nature, Lond.*, **185**, 666.
- Chance, B. & Hollunger, G. (1961). *J. biol. Chem.* **236**, 1534, 1555, 1562, 1577.
- Davies, R. E. & Krebs, H. A. (1952). *Symp. biochem. Soc.* **8**, 77.
- Emmelot, P. & Bos, C. J. (1959). *Brit. J. Cancer*, **13**, 520.
- Hanes, C. S. & Isherwood, F. A. (1949). *Nature, Lond.*, **164**, 1107.
- Klingenberg, M. & Schollmeyer, P. (1960). *Biochem. Z.* **333**, 335.
- Klingenberg, M. & Schollmeyer, P. (1961). *Biochem. biophys. Res. Commun.* **4**, 38, 323.
- Klingenberg, M., Slenczka, W. & Ritt, E. (1959). *Biochem. Z.* **332**, 47.
- Krebs, H. A., Eggleston, L. V. & d'Alessandro, A. (1961). *Biochem. J.* **79**, 537.
- Krebs, H. A. & Kornberg, H. L. (1957). *Ergebn. Physiol.* **49**, 212.
- Kulka, R. G., Krebs, H. A. & Eggleston, L. V. (1961). *Biochem. J.* **78**, 95.
- Schollmeyer, P. & Klingenberg, M. (1961). *Biochem. biophys. Res. Commun.* **4**, 43.
- Wade, H. E. & Morgan, D. M. (1954). *Biochem. J.* **56**, 41.
- Webster, H. L. (1953). *Nature, Lond.*, **172**, 453.

Biochem. J. (1962) **82**, 143

Mitochondrial Preparations from the Leaves of Tobacco (*Nicotiana tabacum*)

4. SEPARATION OF SOME COMPONENTS BY DENSITY-GRADIENT CENTRIFUGING*

BY W. S. PIERPOINT

Biochemistry Department, Rothamsted Experimental Station, Harpenden, Herts.

(Received 31 July 1961)

Extracts of animal tissues have frequently been fractionated by centrifuging in stabilized density gradients. Sometimes the process is used to fractionate the extracts into nuclear, mitochondrial and microsomal fractions (Anderson, 1956), but more often to examine, in greater detail, mitochondrial and microsomal preparations made by differential centrifuging (de Duve *et al.* 1960). The process has occasionally been applied to plant-tissue extracts, as when James & Das (1957) used a simplified system to separate chloroplasts from leaf extracts. However, the full resolving power of the technique, as illustrated by the experiments of de Duve *et al.* (1960), has not been exploited with plant extracts.

This paper describes a sucrose-gradient procedure, which uses only low centrifugal forces, and its application to the chloroplast and mitochondrial fractions prepared from tobacco leaves (Pierpoint, 1959). It partially separates succinoxidase activity from the chloroplast-bearing particles, and enables the distribution of fumarase, cytochrome oxidase and glycolic oxidase among the different particles to be measured.

EXPERIMENTAL

Chloroplast preparation. Tobacco leaves (*Nicotiana tabacum*, White Burley strain) (25–40 g.) were disrupted in five to six times their own weight of medium (sucrose,

0.4M; tris-HCl buffer, 0.2M; K_2HPO_4 , 0.01M; EDTA, 5 mM; trisodium citrate, 0.02M; final pH 7.7), by using the method described by Pierpoint (1959). The extract was passed through two layers of cheese-cloth, and centrifuged for 1 min. at a centrifugal force of approx. 500g to remove sand, fibre residue and starch grains. A second centrifuging for 7 min. at 1000g gave a dark-green precipitate rich in intact chloroplasts, which was usually washed by suspending it in the washing medium (Pierpoint, 1959) and resedimenting it at 1000g. It was then evenly suspended in 15 ml. of a solution containing sucrose (0.15M), tris-HCl buffer, pH 7.4 (5 mM), and EDTA (1 mM).

When mitochondrial and chloroplast preparations were made from the same batch of leaves, liquid in which the chloroplasts had been washed was added to the supernatant fluid from which the mitochondria were to be sedimented.

Mitochondrial preparations. Mitochondrial preparations were made from tobacco leaves (25–40 g.) by the method described (Pierpoint, 1959, 1960a). They were not washed, but suspended in 10–15 ml. of a solution containing sucrose (0.15M), tris-HCl buffer, pH 7.4 (5 mM), and EDTA (1 mM).

Centrifuging in sucrose gradients. Sucrose gradients were made by layering sucrose solutions of decreasing densities in the 100 ml. glass tubes of an MSE Major centrifuge. The solutions of sucrose generally used were 1.76M (20 ml.), 1.4M (6 ml.), 1.1M (10 ml.), 0.8M (6 ml.), 0.5M (6 ml.) and 0.3M (10 ml.); all except the most concentrated one was buffered at pH 7.4 with tris buffer (0.01M). The chloroplast or mitochondrial preparations (10 ml.) were pipetted on the top after the interfaces of the layers had been allowed to smooth by standing for at least 5 hr. in the cold (0–5°). The tubes were then fitted with plastic caps and centrifuged in

* Part 3: Pierpoint (1960b).