- Niu, C-I & Fraenkel-Conrat, H. (1955). *J. Amer. chem. Soc.* **77**, 5882.
- Phillips, D. M. P. (1958). *Biochem. J.* **68**, 35.
- Phillips, D. M. P. & Johns, E. W. (1959). *Biochem. J.* **72**, 538.
- Phillips, D. M. P. & Simson, P. (1961). *Biochem. J.* **78**, 32F.
- Sanger, F. (1945). *Biochem. J.* **39**, 507.
- Satake, K. & Luck, J. M. (1959). *Fed. Proc.* **18**, 316.
- Satake, K., Rasmussen, P. S. & Luck, J. M. (1960). *J. biol. Chem.* **235**, 2801.
- Vendrey, R., Knobloch-Mazen, A. & Vendrey, C. (1960). *Biochem. Pharmacol.* **4**, 19.
- Wilkins, M. H. F., Zubay, G. L. & Wilson, H. R. (1959a). *Trans. Faraday Soc.* **55**, 497.
- Wilkins, M. H. F., Zubay, G. L. & Wilson, H. R. (1959b). *J. molec. Biol.* **1**, 179.

Biochem. J. (1962) **82**, 241

Purification of Human Plasminogen

By M. DERECHIN*

Lister Institute of Preventive Medicine, Chelsea Bridge Road, London, S.W. 1

(Received 11 July 1961)

Plasminogen is the inactive precursor of plasmin, a proteolytic enzyme of plasma. The euglobulin fraction of serum was first used as a source of plasminogen after Milstone's (1941) description of the preparation of the 'lytic factor'. Remmert & Cohen (1949) purified plasminogen by two precipitations at pH 5.3 at different ionic strengths, followed by adsorption on kaolin and elution with phosphate buffer. Christensen & Smith (1950) extracted plasminogen from Cohn's fraction III by means of 0.2N-sulphuric acid. Kline (1953) purified plasminogen by extracting Cohn's fraction III with 0.05N-sulphuric acid, subsequently adjusting the extract to pH 11 with NaOH and then precipitating plasminogen with a small amount of phosphate buffer, pH 6. Finally, the precipitated protein was redissolved and crystallized at pH 8.2.

Wallen & Bergström (1960) precipitated plasminogen from Cohn's fraction II + III at low ionic strength at pH 5.3. The precipitate was dissolved in ammonium acetate buffer, adsorbed on a column of diethylaminoethylcellulose and eluted by a stepwise procedure with ammonium acetate buffers, pH 9 and pH 4, containing 0.01M-lysine.

Hagan, Ablondi & De Renzo (1960) used carb-oxymethylcellulose to purify plasminogen prepared by the Kline procedure. The preparation was adsorbed from solution in formate buffer, pH 3, and gradient elution yielded plasminogen at either pH 2.4-1.8 or 5.0-5.9.

The greatest increase in activity relative to plasma or serum has been claimed for preparations obtained by methods involving acid extraction (Kline, 1953; Hagan *et al.* 1960). However, such preparations are insoluble at neutral pH (Alk-

jaersig, Fletcher & Sherry, 1958), although the original material is soluble in plasma and the reported isoelectric point is about pH 5.6 (Shulman, Alkjaersig & Sherry, 1958).

A procedure for purifying plasminogen described here consists of steps which are milder than those entailing the treatment of the proteins with sulphuric acid or sodium hydroxide at extreme pH values. A final adsorption with diethylaminoethyl-cellulose, based on the method of Wallen & Bergström (1960), provides material soluble at neutral pH with high specific activities.

MATERIALS AND METHODS

Reagents. These were: L-lysine monohydrochloride (Hopkin and Williams Ltd.); L-arginine hydrochloride (Hopkin and Williams Ltd.); ϵ -amino-n-hexanoic acid (L. Light and Co. Ltd.); diethylaminoethylcellulose, Whatman (DE 50).

Buffers. The following were used: acetate-phosphate buffers (Kekwick & Mackay, 1954): pH 4: Na_2HPO_4 (7 mm), acetic acid (43 mm); pH 5-15: Na_2HPO_4 (11 mm), acetic acid (23 mm). Citrate buffer (Kekwick, Mackay, Nance & Record, 1955), 1 1.0: trisodium citrate (0.164M), citric acid (6 mm). Phosphate buffers (Kolthoff, 1937): pH 6.5-8.0, 1 0.1-0.3; NaCl-phosphate buffers prepared by mixing 9 vol. of 0.2M-NaCl with 1 vol. of phosphate buffer, 1 0.2, covering the range 6.5-8.0. Sodium chloride-veronal-buffers, prepared by mixing 8 vol. of 0.2M-NaCl with 2 vol. of veronal buffers (Kolthoff, 1937) covering the range 6.5-9.0. Ammonium acetate buffer: 57.6 ml. of acetic acid was added to 1800 ml. of water and then conc. NH_3 solution [Hopkin and Williams Ltd.; Analytical Reagent, 35% (w/v) NH_3] was added until the pH of the mixture was 9.0; this solution was designated 0.5M-ammonium acetate buffer and was used as a stock solution from which diluted buffers were made. Glycine-NaOH-NaCl buffer (Miller & Golder, 1950), pH 9.2, 1 0.1: glycine (17 mm), NaOH (3 mm), NaCl (97 mm).

* Present address: Department of Colloid Science, Free School Lane, Cambridge, England.

Diethyl ether. The ether used for the precipitation was anaesthetic grade but free from antiperoxidants. Peroxides were not detected with KI (British Pharmacopoeia).

Determination of pH. pH values were determined electrometrically with a MacInnes-type glass electrode, the outer chamber of which contained 0.05M-potassium hydrogen phthalate; 3.5M-KCl-calomel cells were used as reference electrodes and a buffer containing 0.09M-KCl and 0.01M-HCl assumed to have a pH 2.10 was used as a standard. Measurements were made at room temperature with a valve potentiometer, having an accuracy of ± 0.01 pH unit. In systems containing ether the pH values given are apparent and no attempt was made to correct for the effect of the ether on the measurements.

Estimation of nitrogen. Samples of protein solutions were digested in 2 ml. of H_2SO_4 (microanalytical reagent) with the catalyst mixture recommended by Chibnall, Rees & Williams (1943) and estimations of N were made according to Ma & Zuazaga (1942).

Plasminogen assay. Plasminogen was estimated after converting it into plasmin with streptokinase, as described by Derechin (1961). The amount of enzyme in the reaction mixture was adjusted to ensure that three samples taken at intervals of 3–5 min. from one another would each give estimates falling on the linear portion of the reaction curve. These samples (4 ml.) were pipetted into 2 ml. of 1.7M- $HClO_4$ at room temperature and shaken at intervals for 20–30 min. The suspension was filtered and the extinction of the filtrate was read at 276 m μ in 1 cm. cells with a Unicam model SP. 600 spectrophotometer. In all, four samples were taken, the first after proteolysis for 10 min. being used as a blank. The plasmin 'unit' here is that amount of enzyme which gives rise to an increase of 0.001 unit of extinction/min., of the filtrate, during the linear part of the reaction curve with casein (2 g./100 ml.) in phosphate buffer, pH 7.6 (I 0.2), at 40°. The specific activity of a preparation is defined by the equation: specific activity = $10^3 SF/N$, where S is the slope of the linear portion of the reaction curve in units of extinction/min., F the dilution factor, and N mg. of nitrogen/ml. of the plasminogen solution assayed.

Plasma. Out-dated citric acid-sodium citrate-dextrose plasma was used throughout.

Electrophoresis studies. The electrophoresis patterns were obtained in the Tiselius (1937) apparatus with the cylindrical-lens schlieren system described by Philpot (1938). Protein solutions were run at 0°, for the first hour at 10 mA followed by 3 hr. at 15 mA. The patterns were photographed at intervals with Ilford half-tone panchromatic plates, with a high-pressure mercury arc as a source of light, and a suitable filter to isolate light of a wavelength of 546 m μ . One run was carried out in the Perkin-Elmer version of the Tiselius (1937) apparatus for 1 hr. at 15 mA.

EXPERIMENTAL AND RESULTS

Precipitation of fibrinogen from plasma

The fibrinogen was precipitated from plasma at pH 7.7 with 11% (v/v) ether at 0° as described by Kekwick *et al.* (1955). The mixture was centrifuged for 45 min. at 0°, 30 min. after adjusting the precipitation conditions, and the supernatant solution (S1) was siphoned off.

Precipitation of crude plasminogen (fraction P1) from S1

The solution S1 was adjusted to pH 5.3–5.4 with 0.5M-acetic acid and diluted with an equal volume of water at 0°. The precipitate that forms contains plasminogen and prothrombin and by diluting the S1 with an equal volume of water I is lowered sufficiently to produce a maximum precipitation of plasminogen. The mixture was centrifuged 30 min. after the precipitation conditions had been adjusted. The recovery of plasminogen was the same after 30 min. equilibration as after 16 hr., although more protein precipitated after the longer period.

Differences in the yield and specific activity were not great when the pH of precipitation was varied from 5.05 to 5.4 but the recovery was a little higher at pH 5.15 than at other pH values. However, at the next stage of the fractionation the yield of plasminogen obtained was much lower from material precipitated at pH 5.15 than from that precipitated at pH 5.3–5.4; the latter was consequently preferred.

Table 1. *Effect of pH and temperature on the precipitation of plasminogen from citrate buffers*

Column (6) gives the calculated amount of N in fraction derived from 1 l. of original plasma. Column (8) gives the calculated total activity derived from 1 l. of original plasma. Fraction P1 was the starting material. The precipitates P2 were collected after 16 hr. equilibration (Expts. 30 and 32) or 30 min. equilibration (Expt. 34).

Expt. no.	Fraction	pH	Temp.	Ether (%)	Nitrogen (mg./l. of plasma)	Specific activity	Activity (units/l. of plasma)	Recovery (%)
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)
30	P2	5.3	0°	5	647	114	95 000	98
	P2	5.8	0	5	386	144	72 000	75
	P2	6.5	0	5	203	198	55 000	57
32	P2	6.8	0	10	158	264	41 800	39
	P2	7.4	0	10	59	330	19 500	18
	P2	7.7	0	10	41	378	15 000	14
34	P2	6.5	–3	10	410	246	110 000	96
	P2	6.5	0	10	315	252	79 000	68
30	P1	—	—	—	722	135	97 000	100
32	P1	—	—	—	630	170	107 000	100
34	P1	—	—	—	625	185	115 000	100

Table 2. *Effect of protein concentration, ionic strength and pH on the precipitation of plasminogen from fraction P2 dissolved in acetate-phosphate buffer*

Column (2) gives the total protein concentration used in the precipitation mixture. Column (3) gives values measured after the conditions had been adjusted. The values in columns (6) and (7) are relative to those of the starting fraction P2 (specific activity was 354 in Expts. 43 and 58, and 336 in Expt. 44).

Expt. no. (1)	Protein (g./100 ml.) (2)	pH (3)	I of pptn. mixture (4)	Specific activity (5)	Yield of N (%) (6)	Activity (%) (7)
43	0.40	5.05	0.01	420	49	59
	0.22	5.08	0.01	474	39	52
	0.11	5.09	0.01	498	34	48
	0.07	5.05	0.01	534	27	41
58	0.30	5.12	0.03	432	53	72
	0.15	5.15	0.03	492	47	72
	0.09	5.10	0.03	492	45	68
44	0.045	5.02	0.02	450	31	42
	0.045	5.11	0.02	510	38	57
	0.045	5.23	0.02	492	40	59
	0.045	5.28	0.02	444	42	56

Table 3. *Effect of the final temperature on the plasminogen remaining in the bottom layer (fraction BL)*

The last two columns give the percentages of nitrogen and activity yielded relative to those of the starting fraction P3 (specific activity 480). The solution was in phosphate-NaCl (pH 7.5), protein concentration 0.07 g./100 ml.

Final temp. of mixture	Specific activity	Yield of N (%)	Activity (%)
-3°	1 500	11	34
0	1 380	15	44
2	1 260	16	43
4	940	26	51
9	850	28	50
17	690	68	98

Fractionation of fraction P1 in citrate buffers (fraction P2)

It was shown in a procedure devised for preparing plasminogen-free fibrinogen (Kekwick *et al.* 1955) that, by extracting crude fibrinogen with citrate buffers of suitable ionic strength between pH 6.1 and 6.5 at 0°, more plasminogen remained in the undissolved residue than was extracted. The possibility of using similar citrate buffers to purify plasminogen was explored in the range 5.3-7.7.

Fraction P1 was dissolved in citrate buffers over a pH range of 5.3-6.5 and plasminogen was precipitated from these solutions with 5% (v/v) ether at 0°. Precipitations at pH values higher than 6.5 were carried out by dissolving the fraction P1 in 9 mM-trisodium citrate and adjusting the pH of the protein solution electrometrically to the required value with 0.02N-NaOH and adding 10% (v/v) ether. The precipitate obtained in this way was designated fraction P2.

The specific activity of the fraction P2 increases on increasing the pH of the precipitation (Table 1); however, at pH values higher than 7, recoveries of the total activity were low.

The recovery of activity was influenced by the temperature of precipitation, the highest yields

being obtained at -3° and at pH values not higher than 6.5 (Table 1). Neither varying the protein concentration between 0.5 and 1.8 g./100 ml. nor *I* between 0.06 and 0.11 had any striking influence on the results.

The standard conditions chosen for the precipitation of fraction P2 were, accordingly, pH 6.3-6.5, *I* 0.06, protein concentration 1-1.5 g./100 ml., ether 10% (v/v) at -3°. The precipitation mixture was centrifuged for 120 min. at 1500 rev./min. at -3°, 30 min. after the conditions were adjusted. Yields were between 90 and 100%, the specific activity being increased by a factor of about 1.5.

Fractionation of fraction P2 in acetate phosphate buffers (fraction P3)

Fraction P2 was suspended in a volume of acetate-phosphate buffer, pH 4, about 20-30% of that of the original plasma, the resulting mixture having a pH of 4.2-4.3. Some 10% of the material did not dissolve and was removed by centrifuging. No plasminogen could be detected in the insoluble material.

The clarified solution was then titrated electrometrically to the desired pH with acetate-phosphate buffer, pH 6, diluted with water to a selected ionic strength and equilibrated at 0° for 16 hr. The supernatant solution was siphoned off and discarded and the precipitate separated by centrifuging for 40 min. at 1500 rev./min. at 0°. The fraction prepared in this way was designated P3.

From pH 5.1 to 5.4 the recovery of activity varied very little, but at pH 5.0 the recovery was 30% below that at pH 5.1; the highest specific activity was obtained at pH 5.1-5.2 (Table 2, Expt. 44).

Precipitates of higher specific activity were obtained by decreasing the total protein concentration of the mixtures but great losses of plasminogen occurred at total protein concentrations lower than 0.05 g./100 ml. The optimum conditions for the

precipitation of fraction P3 were pH 5.1–5.2, *I* 0.02–0.04, and protein 0.3–0.5 g./100 ml. Over a number of preparations the activity recovered in this fraction varied between 70 and 95 % of that of fraction P2 and the specific activity varied between 450 and 550.

Purification of the plasminogen of the fraction P3 by phase separation

Initial observations. The fractions obtained in the previous steps contain a large amount of lipoprotein, and an attempt was made to remove some of the lipids by freezing with ether (30 %, v/v) at -25° and thawing in cold water (McFarlane, 1942). Fraction P3 was treated in this manner but the mixtures were centrifuged at 0° immediately after thawing, instead of allowing them to stand for 6–8 hr. at room temperature as originally described.

When solutions of fraction P3 in phosphate–NaCl were treated in this way most of the mixture had a gel-like appearance, only a small amount of clear solution separating at the bottom of the bottle. When the mixture was reshaken and warmed to room temperature, a clear solution was obtained after centrifuging, with little gel-like material remaining. By simply shaking solutions with excess of ether several times at $0-4^{\circ}$ without freezing, a layer of material separated between the undissolved ether at the top and the bottom layer of protein solution, and after centrifuging at 0° this intermediate layer packed to give a gel-like appearance. When a bottom layer prepared at -3° to 0° and separated from the mixture immediately after centrifuging was placed in a closed container and warmed to room temperature, further material separated and rose to the surface.

The influence of these procedures on the plasminogen activity of the bottom layer was examined under various conditions.

Procedure for preparing the fraction BL. Ether (200 ml.) was added to 100 ml. of protein solution at 4° and the mixture was cooled to 0° in a bath at -3.5° . The mixture was then shaken at intervals whilst cooling to the final temperature selected, during 90–120 min. After centrifuging, the bottom layer was siphoned off and the plasminogen (fraction BL) was precipitated from it by dialysing overnight against distilled water and then for 24 hr. against two changes of acetate–phosphate buffer, pH 5.15, *I* 0.03. When the protein concentration was above 0.2 g./100 ml. the plasminogen was precipitated quantitatively from the bottom layer by adjusting the pH to 5.15 with 0.05M-acetic acid and diluting with 3 vol. of water, omitting the dialysis step.

Factors influencing the yield and specific activity of the plasminogen of the fraction BL. If after being treated at -3° the mixture was warmed before centrifuging, the amount of protein in fraction BL increased and the specific activity of the plasminogen decreased (Table 3).

With phosphate–NaCl buffers the yield of protein increased on increasing the pH throughout the ranges studied and the specific activity of the preparations remained fairly constant up to pH 7.6–8.0, but decreased at pH values higher than 8. At pH values below 6.4 the yield of protein and

activity was very low and there was no increase in specific activity. With phosphate buffers not containing NaCl, the protein yield and the specific activity of preparations increased with pH but no preparation having a specific activity higher than the starting material was obtained.

On decreasing *I* from 0.3 to 0.1 the recovery of total protein and activity decreased and the specific activity increased. On increasing the protein concentration used for preparing the bottom layer the specific activity of the resulting fraction BL tended to decrease and the recovery of protein fell. The protein concentration giving the highest specific activity at the final temperature of 0° appeared to be about 0.1–0.26 g./100 ml.

When a bottom layer separated at -3° was placed in a closed container and warmed to room temperature for 4 hr. a second bottom layer again separated and the specific activity of the precipitate from this was 26 % higher than that from the first bottom layer obtained without warming to room temperature. The total activity in the second bottom layer was 58 % of that in the first.

For routine purification plasminogen solutions (fraction P3) containing 0.5–1 g. of protein/100 ml. were prepared in phosphate–NaCl buffer, pH 8, *I* 0.2, separated as described at -3° and then warmed to 15° before centrifuging. The bottom layer obtained was adjusted to pH 5.15 with 0.05M-acetic acid, and diluted with 3 vol. of water at 0° .

Purification of plasminogen with cellulose resin

Plasminogen was adsorbed on to diethylaminoethyl-cellulose and eluted with ammonium acetate buffer containing certain amino acids, by a batch procedure. The protein, fraction P3 or BL dissolved in 0.02M-ammonium acetate buffer, was mixed with resin and stirred at intervals during 60 min., and the resin filtered off on a Buchner funnel and washed several times with buffer. The washed resin was suspended in an eluent solution for 30 min. and then filtered as before. It was necessary to saturate the resin with protein to obtain reproducible results. The sample of resin used was saturated by 130–150 mg. of protein/g. of resin, and eluted with 25 ml. of eluent/g.

Lysine and arginine (0.2M) were both very effective eluents (Table 4), yielding more than 90 % of the total activity applied, whereas ϵ -aminohexanoic acid (0.5M) eluted only 50 %. Here the increase in the specific activity was about 2.5 times that of the starting material in all instances. With 0.1M-lysine in the eluent, the specific activity of the eluate was usually five to seven times that of the fraction applied.

The eluates (fraction R) obtained from fractions BL and P3 with 0.1M-lysine in the eluent showed the same order of increase in the specific activity compared with the starting materials; 80–100 % of the total activity was recovered from fraction BL and 70 % or more from fraction P3.

The yields and specific activities of the different fractions are shown in Table 5.

Table 4. *Effect of ϵ -aminohexanoic acid, lysine and arginine on the elution of plasminogen from diethylaminoethylcellulose resin*

Specific activities of the starting material were 600, 480 and 1320 for P3, R5; P3, R6 and BL, R6 respectively. Yield of N and activity are relative to the starting material.

Expt. no.	Starting material	Eluate	Specific activity	Yield of N (%)	Activity (%)
R5	P3	ϵ -Aminohexanoic acid (0.5M)	1410	21	50
		Lys (0.2M)	1400	41	95
		Arg (0.2M)	1280	42	91
R6	P3	Lys (0.1M)	3600	10	71
	BL	Lys (0.1M)	6000	17	80

Table 5. *Yields and specific activities of the plasminogen fractions*

Fraction	N (mg./l. of plasma)	Specific activity	Activity		Purification factor*
			(units/l. of plasma)	Yield* (%)	
Plasma	7500	18†	140 000†	100	1
P1	620	185	113 000	80†	10
P2	410	250	110 000	78	15
P3	150	450-550	65 000-85 000	46-61	30
BL†	80-100	690-900	63 000-84 000	45-60	45
R	10-20	3600-6000	60 000-83 000	36-59	190-315

* Relative to the plasma used.

† BL fraction was prepared at the final temperature of 15°.

† Assumed values.

Electrophoresis of fractions

The results are shown in Fig. 1. It is difficult at present to correlate the difference in activity of the preparations with any particular changes in their electrophoretic components. Fraction P1 is a complex mixture which probably contains almost all the main electrophoretic components of plasma. When fraction P1 is compared with fractions P2 and P3, respectively, an increase in the proportion of the main peak of the electrophoretic pattern is apparent. The increase in the proportion of this peak was accompanied by an increase in the specific activity from 180 in fraction P1 to 270 and 480 respectively in fractions P2 and P3. However, the pattern of fraction BL, which had a specific activity of 1500, did not show any striking difference from that of fraction P3; both fractions containing at least three components.

The solution of fraction R in phosphate buffer became increasingly opalescent in the course of the run and this made the observation of the electrophoretic pattern very difficult. However, when this fraction was run in glycine-NaOH-NaCl buffer no opalescence occurred, the preparation having at least two electrophoretic components. The reason for the development of the opalescence in this fraction is obscure. It might result from an interaction between the phosphate ion and a protein during the electrophoresis. Kline

Fig. 1. Electrophoresis of plasminogen fractions. The experiments were run in phosphate buffer, pH 8, I 0.2 (a-d), or in glycine-NaOH-NaCl buffer, pH 9.2, I 0.1 (e). (a) Fraction P1, protein 1.75 g./100 ml.; (b) fraction P2, protein 1.65 g./100 ml.; (c) fraction P3, protein 1.16 g./100 ml.; (d) fraction BL, precipitated from a bottom layer prepared at -3° and -3° (separation and final temperature respectively), protein 1.36 g./100 ml.; (e) fraction R, protein 1.1 g./100 ml.

(1953) observed that a small amount of phosphate caused precipitation in his preparations of purified plasminogen.

DISCUSSION

One of the main difficulties encountered (Remmert & Cohen, 1949; Kline, 1953) in attempting to purify plasminogen is its tendency to co-precipitate with other proteins. In the present investigations the tendency to co-precipitate has been used to prepare fraction P2. The activity recovered in this fraction was greater from crude plasminogen (fraction P1) precipitated at pH 5.3–5.4 than at pH 5.15, although these contained similar amounts of plasminogen. Also, when fraction P1 was first fractionated in acetate-phosphate buffer at pH 5.0 and subsequently in citrate buffer at pH 6.5, no precipitate formed on adjusting the conditions to those normally producing a separation of fraction P2. These observations suggest that protein necessary for co-precipitating with plasminogen at pH 6.5 remains in solution at pH 5.0–5.15.

In the separation of fractions P1 and P2, the amount of ether present was less than that required to saturate the aqueous phase, and the precipitated protein always sedimented. In preparing the 'bottom layer', a considerable excess of ether was present and separated protein rose to the surface to form a new gel-like phase, with physical properties very different from those of the precipitates obtained previously. This suggests that, in the presence of excess of ether under these conditions, the ether and some of the proteins may interact with consequent separation of material with a density lower than that of the aqueous solution. Whatever the mechanism and interpretation of these phenomena, this procedure yielded plasminogen preparations of increased specific activity which dissolved to give very clear solutions.

Most previous workers have described their preparations as 'so many times purified relative to serum' with no indications of how these calculations were made.

Remmert & Cohen (1949) described their plasminogen as purified 136–165-fold relative to serum but do not state the yield. Christensen & Smith (1950) estimated a purification of 250-fold for their plasminogen with yields varying from 10 to 100 % of the total activity, the average being 30 %; Kline (1953) a 425-fold purification with a yield of 33 %; Hagan *et al.* (1960) a 300–400-fold purification without stating yield; and Wallen & Bergström (1960) a 20–30-fold purification compared with the starting material, Cohn's fraction II + III. Plasminogen is concentrated in this fraction, which contains 25 % of the total protein of plasma (Oncley, Melin, Richert, Cameron & Gross, 1949). If it is

assumed that fraction II + III contains all the plasminogen of plasma, Wallen & Bergström achieved 80–120-fold purification relative to the plasma.

It is difficult to relate the activity of a plasminogen preparation to that of serum directly, since serum contains potent plasmin inhibitors whose effect will vary with the dilution of serum used in any assay. It is possible that a similar limitation applies to the first plasminogen fraction obtained during purification but in these circumstances the amount of inhibitor, if present, is much lower.

No attempt was made directly to estimate the amount of plasminogen in plasma. An approximate estimate of the degree of purification achieved was obtained on the assumption that about 20 % of the plasminogen precipitates with the fibrinogen fraction, (M. E. Mackay, unpublished work) and that all the plasminogen in the solution S1 is precipitated in fraction P1. Since the protein content of the plasma used was about 46 g./l. and the protein obtained from it in fraction P1, with a specific activity of 180–190, corresponds to 3.8 g./l. of plasma, the factor for P1 is 10, indicating a specific activity of 18 in whole plasma. With 0.1M-lysine in the eluent the potency of batch eluates lay between 3600 and 6000, corresponding to purification factors of 190–315 relative to plasma; the recovery of total activity was usually not less than 70 % that of fraction P1.

The preparations obtained compare favourably in several respects with others described in the literature, not only because of the high yield but also because they are soluble at neutral pH. However, no comparison between the specific activity of the preparation described here and the results of other workers can be made because of the different assay procedures that have been used.

The properties of the proteins obtained here may be nearer to those of the native proteins than is the case with acid-alkali-treated plasminogen preparations.

SUMMARY

1. Plasminogen was purified from out-dated human plasma. A crude plasminogen (fraction P1) was first precipitated after the removal of fibrinogen from plasma at pH 5.3, and a fraction P2 obtained from this by treatment with citrate buffer.

2. Fraction P2 was dissolved in acetate-phosphate buffer, pH 4.0, and the plasminogen precipitated from the solution by adjusting it to pH 5.15, to give fraction P3.

3. By dissolving the fraction P3 in phosphate-sodium chloride buffer, pH 8.0, and shaking the solution with excess of ether at -3° , lipid material and some inactive protein were removed, and plasminogen precipitated from the aqueous layer at pH 5.15 to give fraction BL.

4. From both fractions P3 and BL purified plasminogen fractions were obtained by batch adsorption on to diethylaminoethylcellulose resin followed by elution with lysine-containing buffers. The eluates obtained by the batch adsorption of fraction BL (fraction R) had a potency 190–315 times that of plasma, the preparation being soluble at neutral pH.

This work was done during the tenure of a British Council Scholarship (1957–59) and a grant from the Consejo Nacional de Investigaciones Científicas y Técnicas, (Buenos Aires, Argentina) (1959–60). The author is indebted to Dr R. A. Kekwick for his helpful advice and criticism and for his invaluable assistance in the preparation of the manuscript and to Mr Brian Jordan for much technical assistance. Thanks are due to Dr A. E. Pearce for the facilities made available for an electrophoresis run to be made in the Perkin-Elmer apparatus.

REFERENCES

- Alkjaersig, N., Fletcher, P. & Sherry, S. (1958). *J. biol. Chem.* **233**, 81.
- Chibnall, A., Rees, M. W. & Williams, E. F. (1943). *Biochem. J.* **37**, 354.
- Christensen, L. R. & Smith, D. R. (1950). *Proc. Soc. exp. Biol., N.Y.*, **74**, 840.
- Derechin, M. (1961). *Biochem. J.* **78**, 443.
- Hagan, J. J., Ablondi, F. B. & De Renzo, E. C. (1960). *J. biol. Chem.* **235**, 1005.
- Kekwick, R. A. & Mackay, M. E. (1954). *Spec. Rep. Ser. med. Res. Coun., Lond.* no. 286.
- Kekwick, R. A., Mackay, M. E., Nance, M. H. & Record, B. R. (1955). *Biochem. J.* **60**, 671.
- Kline, D. L. (1953). *J. biol. Chem.* **204**, 949.
- Kolthoff, I. M. (1937). *Acid-Base Indicators*, pp. 249, 258. New York: The Macmillan Co.
- Ma, T. & Zuazaga, G. (1942). *Industr. Engng Chem. (Anal.)*, **14**, 280.
- McFarlane, A. S. (1942). *Nature, Lond.*, **149**, 439.
- Miller, G. L. & Golder, R. H. (1950). *Arch. Biochem.* **29**, 420.
- Milestone, H. (1941). *J. Immunol.* **42**, 109.
- Onceley, J. L., Melin, L., Richert, D. A., Cameron, J. W. & Gross, P. M., jun. (1949). *J. Amer. chem. Soc.* **71**, 541.
- Philpot, J. St L. (1938). *Nature, Lond.*, **141**, 283.
- Remmert, L. F. & Cohen, P. P. (1949). *J. biol. Chem.* **181**, 431.
- Shulman, S., Alkjaersig, N. & Sherry, S. (1958). *J. biol. Chem.* **233**, 91.
- Tiselius, A. (1937). *Trans. Faraday Soc.* **33**, 524.
- Wallen, P. & Bergström, K. (1960). *Acta chem. scand.* **14**, 217.

Biochem. J. (1962) **82**, 247

The Reaction of Normal Adult Human Haemoglobin with Heavy-Metal Reagents

BY R. CECIL AND N. S. SNOW

Department of Biochemistry, University of Oxford

(Received 31 May 1961)

All the mammalian haemoglobins which have so far been examined contain SH groups that form mercaptides immediately on the addition of heavy-metal reagents in the same way as simple thiols. Most, but not all, contain further SH groups that are unreactive in the native protein, but which react normally when the protein is denatured with acid or sodium dodecyl sulphate. It is not yet known why these groups are unreactive or what is their function.

The unreactive SH groups will react slowly with some heavy-metal reagents. This paper describes the reactions of mercuric chloride, phenylmercuric hydroxide, *p*-chloromercuribenzenesulphonate and silver nitrate with the SH groups of normal adult human haemoglobin in its various forms. Allison & Cecil (1958) have shown that this protein has a total of 6 SH groups/molecule, 2·2 of which are normally reactive and the remainder unreactive until the protein is denatured.

THEORETICAL

Abbreviations. HbO₂, Oxyhaemoglobin; HbCO, carboxyhaemoglobin; Met-Hb, methaemoglobin; Hb, reduced haemoglobin.

The slow reaction of haemoglobin with heavy-metal reagents was followed polarographically by observing the current due to the electroreduction of the heavy-metal reagent at the dropping-mercury electrode. In practice this is done by setting the applied potential to a value at the top of the polarographic wave and observing the fall in current with time. This section deals with the various factors that have to be taken into account when relating this change in current to the number of protein SH groups that have reacted to form mercaptide.

At any time during the reaction the heavy-metal ions will be distributed as follows: (a), free in solution; (b), combined as haemoglobin mercaptide