- Brooks, C. J. W., Hopkins, R. P. & Young, L. (1959). *Biochem. J.* **71**, 28p.
- Brooks, C. J. W. & Young, L. (1956). *Biochem. J.* **63**, 264.
- Chang, L. H. (1943). *J. biol. Chem.* **151**, 93.
- Chang, L. H. & Young, L. (1943). *J. biol. Chem.* **151**, 87.
- Cook, J. W. (1950). *J. chem. Soc.* p. 1210.
- Corner, E. D. S., Billett, F. S. & Young, L. (1954). *Biochem. J.* **56**, 270.
- Crieggee, R., Kraft, L. & Rank, B. (1933). *Liebigs Ann.* **507**, 159.
- Ewan, T. & Cohen, J. B. (1889). *J. chem. Soc.* **55**, 578.
- Feigl, F. (1954). *Spot tests II. Organic Applications*, p. 125. New York: Elsevier Publishing Co.
- Graebe, C. & Gfeller, E. (1892). *Ber. dtsh. chem. Ges.* **25**, 652.
- Graebe, C. & Gfeller, E. (1893). *Liebigs Ann.* **276**, 1.
- Grenby, T. H. & Young, L. (1960). *Biochem. J.* **75**, 28.
- Grimes, A. J. & Young, L. (1956). *Biochem. J.* **62**, 11p.
- Hopkins, R. P. (1959). Ph.D. Thesis: University of London.
- Hopkins, R. P. & Young, L. (1961). *Biochem. J.* **80**, 2p.
- Jack, K. M. & Rule, H. G. (1938). *J. chem. Soc.* p. 188.
- Kerr, L. M. H. & Levvy, G. A. (1951). *Biochem. J.* **48**, 209.
- Kynaston, W. & Jones, J. I. (1949). *J. Soc. chem. Ind. Lond.*, **68**, 228.
- Lindsey, A. J. (1959). *Brit. med. J.* **i**, 506.
- Loon, C. van (1919). Doctoral Thesis: Technische Hoogeschool, Delft.
- Plattner, P. A. & Pfau, A. S. (1937). *Helv. chim. acta*, **20**, 224.
- Prévost, C. (1933a). *C.R. Acad. Sci., Paris*, **196**, 1129.
- Prévost, C. (1933b). *C.R. Acad. Sci., Paris*, **197**, 1661.
- Talalay, P., Fishman, W. H. & Huggins, C. (1946). *J. biol. Chem.* **166**, 757.
- Young, L. (1947). *Biochem. J.* **41**, 417.
- Young, L. (1950). *Symp. biochem. Soc.* **5**, 27.

Biochem. J. (1962) **82**, 466

Studies on Zinc in Blood

TRANSPORT OF ZINC AND INCORPORATION OF ZINC IN LEUCOCYTES

BY E. DENNES, R. TUPPER AND A. WORMALL

Department of Biochemistry and Chemistry, The Medical College of St Bartholomew's Hospital, London, E.C. 1

(Received 25 August 1961)

It is widely recognized that zinc is essential for the normal growth of both animals and plants, but our present knowledge of zinc metabolism and transport in the animal body is still incomplete.

It has long been known that blood contains zinc and it is reasonable to expect that, since a significant fraction of the blood zinc is present in the plasma, it is the plasma form which is the main transport medium. Varying values for the plasma zinc have been recorded; thus Delezenne (1919) reported that about one-tenth of the blood zinc is contained in the plasma and a similar value of 12% for the serum zinc as a percentage of the whole-blood zinc (of 880 $\mu\text{g.}/100\text{ ml.}$) is given by Vallee (1959). Analyses on plasma and serum give identical results, and, whichever fluid is used, there is fairly general agreement that the plasma or serum zinc may be divided into two fractions, in one of which the metal is more firmly bound than it is in the other.

There appears to be no evidence of any plasma-saturation limit for zinc (Berfenstam, 1952) similar to that for iron. Indeed, the mode of transport of zinc in plasma is still not clear; studies have been made with several species of animals and it is

possible that species differences account for some of the recorded differences. Another factor is the possible exchange of zinc between the plasma proteins and the zinc compounds in erythrocytes and leucocytes, these cells accounting respectively for about 85 and 3% of the whole-blood zinc (Vallee, 1959). Part of the plasma zinc is probably associated with zinc-containing enzymes, but this amount is probably small and has not been completely identified, though Vessel & Bearn (1957) have found, in electrophoretic separation of isotopically labelled serum proteins, that part of the zinc is associated with the α_2 -globulins, coinciding with one of the peaks of lactic dehydrogenase.

In view of this uncertainty about the distribution of the zinc in normal plasma, we have investigated this problem in the rabbit, with ^{65}Zn as tracer, and have studied the labelled blood to determine which protein fractions bind zinc injected intravenously as a zinc-glycine complex.

A study has also been made of the extent to which the injected zinc is incorporated into erythrocytes and leucocytes, and the possibility of exchange of this zinc between the plasma and the cells in the blood.

MATERIALS AND METHODS

Zinc-glycine solutions containing ^{65}Zn . In previous studies we found that the addition of glycine to solutions containing ^{65}Zn prevents the precipitation of zinc as an inorganic salt when the solution is introduced into tissues or added to tissue extracts. The zinc in the zinc-glycine complex can undergo exchange with other zinc compounds, except where the zinc is an integral component of the molecule, as, for example, in the zinc-containing enzyme carbonic anhydrase (Tupper, Watts & Wormal, 1951). For the experiments described below, several zinc-containing solutions with ^{65}Zn were prepared with varying specific activities. They all contained 2–6 molecules of glycine/atom of zinc and the mixtures were treated with sufficient dilute NaOH to bring the pH to about 7.4; solid NaCl was added to the final solutions to the extent of 9 mg./ml. Many of the ^{65}Zn -containing solutions were prepared from a stock solution (soln. A), which was prepared by adding zinc-free water (18.3 ml.) to a solution of ZnCl_2 (1.7 ml.) containing 2 mc of ^{65}Zn and 13 mg. of total zinc (obtained from The Radiochemical Centre, Amersham, Bucks.). Soln. A contained 650 μg . of zinc/ml. and had an activity of 100 μC of ^{65}Zn /ml.

Soln. B1 was prepared from soln. A (1.28 ml.) and glycine (5 mg.). This solution (10 ml.) contained 83 μg . of total zinc (12.8 μC of ^{65}Zn)/ml.

Soln. B2 was prepared from soln. A (5.1 ml.) and glycine (15 mg.) and contained 332 μg . of total zinc (51.2 μC of ^{65}Zn)/ml.

Soln. C (of lower specific activity than solns. A and B) was prepared from soln. A (5.1 ml.), ZnSO_4 (18.15 mg.), glycine (30 mg.), NaCl (90 mg.) and water (to 10 ml.). It contained 743 μg . of zinc/ml. and had an activity of 51.2 μC /ml.

Soln. D (a buffered solution used in dialysis experiments) was prepared from soln. A (3 ml.), 0.05M-veronal buffer, pH 7.4 (22 ml.), ZnSO_4 (35.4 mg.) and glycine (25 mg.). It contained 400 μg . of zinc/ml. and had an activity of 12 μC /ml.

Soln. E (for the labelling *in vivo* of leucocytes and erythrocytes). Glycine (20 mg.) and water (1 ml.) were added to a solution of ZnCl_2 containing ^{65}Zn (0.3 ml.; The Radiochemical Centre) and the solution was adjusted to pH 7.4 by addition of N-NaOH (0.2 ml.). Water (4.5 ml.) and NaCl (42 mg.) were added. The solution was filtered before use; 5 ml. contained 1.9 mg. of zinc and had an activity of 1.48 mc of ^{65}Zn .

Soln. F (used for labelling *in vivo* of rabbit plasma for injection into rabbits). Glycine (25 mg.) and water (2 ml.) were added to a solution of ZnCl_2 containing ^{65}Zn (0.67 ml.; The Radiochemical Centre) and the solution was adjusted to pH 7.4 with N-NaOH (0.1 ml.). Water (3.9 ml.) and NaCl (48 mg.) were added; 5 ml. contained 10.2 mg. of zinc and had an activity of 5.9 mc of ^{65}Zn .

Soln. G (used for labelling *in vivo* of rabbit plasma, leucocytes and erythrocytes for ^{65}Zn -exchange experiments). Glycine (125 mg.) was added to a solution of ZnCl_2 containing ^{65}Zn (1.6 ml.; The Radiochemical Centre) and the whole was adjusted to pH 7.4 with 0.1N-NaOH (5.2 ml.). Water (5.4 ml.) and NaCl (78 mg.) were added; 5 ml. contained 12 mg. of zinc and had an activity of 3.5 mc.

Buffers. Tris buffer, pH 7.4 (0.05M), and veronal buffer, pH 7.4 (0.05M), were prepared as described by Gomori

(1955), and veronal buffers, pH 8.6 (0.06 and 0.09M and I 0.05 and 0.075 respectively), as described by Varley (1958).

For the dialysis studies, EDTA or glycine was added to the veronal buffer, pH 7.4, to give a concentration of 1 mM.

Ferric chloride-sodium citrate solution. Ferric chloride soln. (containing 5000 μg . of iron/ml.; 2.5 ml.) was added to 3.8% sodium citrate soln. (2.5 ml.) and the mixture was diluted to 20 ml. with 0.9% NaCl soln. This solution contained 625 μg . of iron/ml.

Zinc-free or zinc-depleted water. This was prepared by passing water through a column of mixed ion-exchange resins (Elgastat water purifier; Elga Products Ltd.). Zinc-free materials and apparatus were treated as described by Dennes, Tupper & Wormal (1961).

Labelling of plasma proteins with ^{65}Zn . Studies on the distribution of zinc between the various protein fractions of plasma were made with plasma which had been labelled with ^{65}Zn either *in vivo* or *in vitro*. Human or rabbit plasma was labelled *in vitro* by the equilibrium-dialysis method (Hughes & Klotz, 1956), buffered ^{65}Zn -containing zinc-glycine solutions with concentrations of 'carrier' zinc ranging from 4 to 100 μg /ml. being used. A veronal buffer, pH 7.4, was used to dilute the rabbit plasma before dialysis and also in the preparation of the ^{65}Zn -containing zinc-glycine solutions used for labelling the rabbit plasma proteins. In the labelling of human plasma proteins, tris buffer, pH 7.4, was used for dilution of the plasma and in the preparation of the ^{65}Zn solutions. The change of buffer was made because of the lower nitrogen content and greater stability of this buffer compared with veronal. With the veronal buffer it was observed that some barbitone was occasionally precipitated at 4°.

The dialysis tubing was prepared as described by Hughes & Klotz (1956). Rabbit plasma (5 ml.) was diluted with veronal buffer, pH 7.4 (10 ml.), before dialysis. Human plasma (5 ml.) was diluted with tris buffer, pH 7.4 (5 ml.). The diluted plasma (10 ml.) was dialysed against the appropriate buffered ^{65}Zn -containing zinc-glycine solution (10 ml.). Dialysis was carried out at 4° for 1 week in stoppered Pyrex boiling tubes. The tubes were shaken at intervals each day. After the dialysis, determinations of ^{65}Zn were made on the solutions inside and outside the sac. Protein was precipitated by addition of 10% trichloroacetic acid (10 ml.) to the dialysed solution (10 ml.); it was then washed once with 10% trichloroacetic acid (10 ml.).

The plasma used in the paper-electrophoresis studies was obtained from rabbits which had received intravenous injections of a ^{65}Zn -containing zinc-glycine solution (soln. E). Blood was collected in polythene bottles containing a solution of heparin in 0.9% NaCl (25 mg. of heparin/ml., 0.1 ml. of solution/5 ml. of blood).

Labelling of blood cells with ^{65}Zn . Rabbits were injected intravenously with soln. E (5 ml. containing 1.9 mg. of zinc with a total activity of 1.48 mc of ^{65}Zn). The animals were bled from the ear opposite to that injected and the blood was collected in polythene bottles containing heparin. Leucocytes and erythrocytes were separated and washed as described by Dennes *et al.* (1961).

Separation of leucocytes and erythrocytes from bone marrow. The long bones of the legs were removed from the rabbit immediately after the animal had been killed by a blow on the neck. The bones were divided into smaller portions by transverse cuts with a saw and the marrow was taken from the bones with either a Pasteur pipette or a

stainless-steel spatula. The marrow from the bones of each rabbit was suspended in a 5% (w/v) solution of bovine plasma albumin in 0.9% NaCl (30 ml.). The suspension was thoroughly mixed and centrifuged, and the pellet of fat which had risen to the top was removed. When this suspension was centrifuged, small particles of bone sedimented first and formed a layer at the bottom of the centrifuge tube. After removal of the fat pellet and the supernatant solution, the cells were pipetted off and transferred to a second tube, leaving the bone in the first tube. The cells were washed three times with 0.9% NaCl (5 ml. each time) and finally resuspended in 0.9% NaCl (2 ml.).

Paper electrophoresis of plasma proteins. Perspex tanks designed for horizontal electrophoresis were used and the temperature was maintained at 4° in a refrigerated cabinet. Veronal buffers, pH 8.6, 1.0.05 or 0.075, were used (Varley, 1958). The paper was Whatman no. 1 for chromatography, in strips 5 cm. by 35 cm. Plasma (0.05 ml.) was placed on the wet strip and run for 18 hr. at 240 v with a current of 0.3 ma/cm. Samples of the same plasma were placed on several strips (usually six). At the end of the run all the strips were dried at 120° for 20 min. Some of the strips were stained with bromophenol blue as described by Durrum (1950). These stained strips were used for the measurement of protein distribution with the aid of an EEL scanner (Evans Electroselenium Ltd.). Nitrogen was determined on samples of the plasma (0.05 ml.) and, from the total amount of plasma nitrogen applied to each strip, the approximate amount of nitrogen in each protein fraction could be calculated from the measured protein distribution.

The remaining unstained strips were used in the determination of ^{65}Zn distribution. A longitudinal band 5 mm. wide was removed from the centre of the strip and stained with bromophenol blue to identify the positions of the various protein fractions on the two unstained sections of strip, which were then cut into portions corresponding to the main protein fractions. The two pieces thus obtained for each protein fraction were placed in the centre of a glass dish, one on top of the other, and the ^{65}Zn content was determined in a scintillation counter.

Separation of plasma proteins by starch-gel electrophoresis. The method described by Smithies (1955) was used for the separation on starch gel of plasma proteins labelled with ^{65}Zn .

Total nitrogen determinations. These were made either by the micro-Kjeldahl method, the ammonia being distilled in the Markham (1942) apparatus into a boric acid soln. containing methyl red-methylene blue indicator, or, where greater sensitivity was required, by the method described by Russell (1944).

Determinations of ^{65}Zn . These were made by the scintillation-counter method described by Banks, Tupper, Watts & Wormall (1955).

Total zinc determinations. The method described by Vallee & Gibson (1948) was used, with the slight modifications described by Banks, Tupper, White & Wormall (1959).

RESULTS

Rate of removal of zinc from plasma after its intravenous injection into rabbits. There is a constant exchange of zinc between plasma and body

tissues. The loss of zinc from the plasma can be conveniently followed if a zinc-glycine solution containing ^{65}Zn is injected. We wished to study the effect that varying the amount of zinc injected had upon the rate at which zinc was lost from the plasma. The rabbits were injected in pairs with ^{65}Zn -containing zinc-glycine solutions of different zinc concentrations. The first, second and third pairs received respectively 83, 332 and 743 $\mu\text{g.}$ of zinc/kg. body wt. These amounts of zinc were calculated as the amounts required to increase the plasma zinc to two, five or ten times the normal values respectively. Injections were made intravenously into the ear. Solns. B1, B2 and C (1 ml./kg. body wt.) were used for the first, second and third pair of rabbits respectively.

The animals were bled from the ear opposite to the one injected and the blood was collected in oxalated tubes at intervals of 3, 10, 30, 60, 90, 120 and 180 min., 24 and 48 hr. after the injection. The plasma was immediately separated and determinations of ^{65}Zn were made. The results are shown in Fig. 1. In the calculation of the values the activity of the plasma 3 min. after the injection has been taken as 100%. Loss of zinc from the plasma was rapid; 10 min. after the injection only 50% of the initial activity could be detected in the rabbits which received the smallest amount of zinc. The effect of increasing the weight of zinc injected was to decrease the percentage rate at which the metal was lost from the plasma. For example, rabbits injected with 743 $\mu\text{g.}$ of zinc/kg. body wt. still

Fig. 1. Rate of removal of ^{65}Zn from rabbit plasma after intravenous injection of the isotope as the zinc-glycine complex. Zinc injected/kg. body wt.: ●, 83 $\mu\text{g.}$; ▲, 332 $\mu\text{g.}$; ○, 743 $\mu\text{g.}$

retained 18 % of the initial activity in the plasma 3 hr. after the injection, compared with 5 % for the rabbits receiving 83 μ g. of zinc/kg. body wt. In all the animals studied, less than 2 % of the initial activity could be found in the plasma 24 and 48 hr. after the injection and, in the range studied, the amount of zinc injected had no effect on the amount of zinc still present in the plasma at this time.

If the same β_1 -metal-binding globulin transports both iron and zinc in the blood, it might be expected that saturation of the plasma with iron before injection of a radioactive zinc solution would affect the rate at which the radioactive zinc was removed from the plasma. To test this, two rabbits were each given an intravenous injection of a ferric chloride-sodium citrate solution at a dose of 132 μ g. of iron/kg. body wt. Five minutes later a ^{65}Zn -containing zinc-glycine solution was injected into an ear vein (dose: 332 μ g. of zinc/kg. body wt.) (solution B2; 1 ml./kg. body wt.). Blood samples were collected and oxalated at intervals and ^{65}Zn determinations were made on the separated plasma. The rate of disappearance of the ^{65}Zn from the plasma of the rabbits receiving the iron injection did not differ from that of the rabbits which had received a similar amount of zinc but no iron.

Rate of removal of ^{65}Zn from plasma after intravenous injection into rabbits of ^{65}Zn -labelled plasma obtained from a rabbit injected with the isotope. To determine whether plasma-bound zinc is easily removed *in vivo*, four rabbits were given intravenous injections of plasma (5 ml./rabbit) that had been obtained from another rabbit 3 hr. after the injection of a ^{65}Zn -containing zinc-glycine solution (sol. F, 5 ml.). The labelled plasma was sterilized by filtration through a Berkefeld filter candle, sealed in sterile glass ampoules and stored in the refrigerator until needed. The plasma gave 296 600 counts/min./ml. in the scintillation counter, and contained all its ^{65}Zn in a protein-bound non-diffusible form. The four recipient rabbits were bled 5 min. and at intervals up to 24 hr. after the injection. Fewer samples of blood were taken from rabbits 449 and 491 compared with rabbits 432 and 466 to check that the results were not being affected by haemodilution. The blood samples were collected in heparinized tubes, immediately centrifuged and ^{65}Zn in the supernatant plasma was determined.

The results (Table 1) showed a rapid fall in plasma ^{65}Zn concentration during the first hour after the injection, followed by a slower fall. Comparison of the values for rabbits 432 and 466 with those for rabbits 449 and 491 shows that the small volumes of blood lost by bleeding had no great effect on the disappearance of zinc from the plasma. The ^{65}Zn bound to the injected plasma

protein is lost from the blood stream as rapidly as ^{65}Zn injected as the zinc-glycine complex (cf. Fig. 1). The initial rapid fall in concentration followed by a slower fall is seen whether the ^{65}Zn injected is plasma-bound or not.

Binding of zinc by plasma proteins

Zinc content and specific activity of plasma and globulins from rabbits injected with zinc-glycine containing ^{65}Zn . The plasma used was obtained from two rabbits injected intravenously with soln. C (1 ml./kg. body wt.; corresponding to 743 μ g. of zinc/kg. body wt.). The blood was collected 3 and 24 hr. after the injection. Total zinc and ^{65}Zn determinations were made on the whole plasma and the globulin fraction of the plasma. Globulins were precipitated by the addition of saturated ammonium sulphate solution (5 ml.) to plasma (5 ml.). The globulin precipitates were removed by centrifuging and were washed twice with half-saturated ammonium sulphate solution (5 ml. each time).

The results (Table 2) show that 3 and 24 hr. after the injection, the globulins in the plasma contained about one-third of the total plasma zinc. The zinc content of the whole plasma at the 24 hr. period was approximately one-third of the value for the 3 hr. period, and a similar decrease occurred in the globulin fraction. The specific activities of the zinc in whole plasma and globulins were almost identical 3 hr. after the injection but 24 hr. after the injection the specific activity of the zinc in the globulins was 30 % of the 3 hr. value, compared with 46 % for the whole plasma.

Amount of plasma zinc bound to the globulin fraction. For this experiment, plasma was obtained from two rabbits 3 and 24 hr. after an

Table 1. *Disappearance of ^{65}Zn from the plasma of rabbits injected with rabbit plasma labelled in vivo with the isotope*

Time after injection (min.)	^{65}Zn content of plasma (% of total injected dose)			
	Rabbit no. 432	Rabbit no. 466	Rabbit no. 449	Rabbit no. 491
5	0.51	0.46	0.47	0.42
10	0.39	0.30	—	—
30	0.11	0.13	0.10	0.08
60	0.09	0.06	—	—
120	0.05	0.04	0.09	0.02
180	0.03	0.03	—	—
240	0.02	0.01	0.02	0.01
360	0.01	0.01	—	—
24 hr.	0.01	0.01	0.01	0.01

Each rabbit was injected intravenously with 5 ml. of labelled plasma (296 000 counts/min./ml.) obtained from a rabbit 3 hr. after the intravenous injection of zinc-glycine solution containing ^{65}Zn (soln. F, 5 ml.).

Table 2. *Zinc content and specific activities of whole plasma and the plasma-globulin fraction from rabbits injected with zinc-glycine solution containing ^{65}Zn*

Rabbits were injected intravenously with a solution of zinc-glycine containing ^{65}Zn (soln. C, 1 ml./kg. body wt.). This solution contained 743 $\mu\text{g.}$ of Zn/ml. with a specific activity of 31200 counts/min./ $\mu\text{g.}$ of Zn. Globulins were precipitated by half-saturation with ammonium sulphate. Figures for the zinc content of the globulin fraction are the amounts of zinc contained in the globulins obtained from 1 ml. of plasma.

Time after injection (hr.)	Total zinc content ($\mu\text{g./ml.}$)		Specific activity (counts/min./ $\mu\text{g.}$ of Zn)	
	Whole plasma	Globulin fraction	Whole plasma	Globulin fraction
3	6.4	2.2	1280	1189
24	2.2	0.8	587	355

Table 3. *Effect of EDTA or glycine on the dialysis of ^{65}Zn from rabbit plasma labelled in vivo with the isotope*

Plasma used was obtained from a rabbit 3 hr. after the intravenous injection of zinc-glycine solution containing ^{65}Zn . The labelled plasma was dialysed for 96 hr. against the appropriate buffer at 0–4°. The buffer was changed twice daily.

Buffer	^{65}Zn content of external fluid (% of total ^{65}Zn in plasma)
Veronal (0.05M)	4
Veronal (0.05M) containing glycine (1 mM)	7
Veronal (0.05M) containing EDTA (1 mM)	100

intravenous injection of solution B1 (1 ml./kg. body wt., corresponding to 332 $\mu\text{g.}$ of zinc) and also from the two rabbits injected with soln. C. The globulins were precipitated and washed as described above, and the ^{65}Zn content of the globulins was determined and compared with that of the whole plasma. In the rabbits injected with 332 and 743 $\mu\text{g.}$ of zinc/kg. body wt., the globulins 3 hr. after the injection contained 37 and 42 % of the total plasma ^{65}Zn respectively. The corresponding figures for the 24 hr. period were 44 and 53 %.

Amount of plasma zinc diffusible at pH 7.4. To determine what proportion of the plasma zinc is loosely bound at pH 7.4, plasma from the rabbits used in the two previous experiments was dialysed against veronal buffer, pH 7.4 (5 ml. of plasma and 20 ml. of buffer), at 4°. The buffer was changed twice daily and ^{65}Zn determinations were made on the buffer samples, and, at the end of the experiment, the contents of the dialysis sacs. By the second or third change of buffers no ^{65}Zn could be detected outside the sac and the amount of diffusible ^{65}Zn was found to be 3 % or less.

Effect of glycine or EDTA on the amount of

plasma-bound zinc diffusible at pH 7.4. In an attempt to determine how firmly zinc is bound to rabbit plasma, ^{65}Zn -labelled rabbit plasma (3 hr. plasma from rabbit 464) was dialysed against veronal buffer, pH 7.4 (0.05M) and the same buffer containing either glycine or EDTA (1 mM).

Samples of the plasma (2 ml.) in Visking cellulose tubing were dialysed against the appropriate buffer (50 ml.) at 0–4° for 96 hr., with the external solutions changed twice daily; samples of each solution were taken for determinations of ^{65}Zn . At the end of the experiment the contents of the sac were pipetted out and the empty sac was then washed thoroughly with 0.9 % sodium chloride. The sac contents plus washings were then made up to a final volume of 100 ml. with 0.9 % sodium chloride and samples (5 ml.) were taken for determinations of ^{65}Zn . As a check on the efficiency of the washing, the ^{65}Zn content of the empty sacs was also measured. Recoveries were good (96–104 %).

The results (Table 3) showed that glycine removed 7 % of the ^{65}Zn in the sacs, only 3 % more than the buffer without glycine. All the ^{65}Zn was removed by EDTA, almost 85 % dialysing out in the first 6 hr.

Zinc-binding capacity of rabbit plasma

The equilibrium-dialysis method was used to study the amount of zinc which could be bound to rabbit plasma. The method of labelling the plasma has been described in the Materials and Methods section. Twelve ^{65}Zn -containing zinc-glycine solutions were used, containing amounts of zinc from 40 to 600 $\mu\text{g.}$ (cf. Table 4). The total plasma nitrogen in the sac was 29 mg.

The results (Table 4) showed that the zinc bound to the plasma protein increased in proportion to the amount of zinc added to the system. To avoid precipitation of protein amounts of zinc greater than 600 $\mu\text{g.}$ were not used.

In an attempt to measure the amount of zinc bound to the globulin fraction of rabbit plasma, we have labelled the plasma *in vitro* by the equilibrium-dialysis method, using solutions containing 160–600 $\mu\text{g.}$ of zinc. The globulins were precipitated from the labelled plasma in two different ways, by half-saturation with ammonium sulphate either at pH 4.2 or at pH 7.0. Precipitation and washing of the globulin fraction was carried out as described above.

The results (Table 5) showed that the pH at which the globulins were precipitated had a definite influence on the amount of zinc bound to the globulin. From the results obtained at pH 4.2, the concentration of zinc in the globulins appears to be lower than that in the whole plasma, but at pH 7.0 (with one exception) the concentration of zinc in the globulins was higher than that for the whole plasma. The pH of the ammonium sulphate

Table 4. *Zinc-binding capacity of rabbit plasma in vitro studied by equilibrium dialysis*

The dialysis sacs contained rabbit plasma diluted with veronal buffer, pH 7.4, 0.05 M (3.3 ml. of plasma: 6.7 ml. of buffer). The diluted plasma was dialysed at 4° for 1 week against the appropriate solution (10 ml.) of zinc-glycine containing ^{65}Zn in 0.05 M-veronal buffer. Total plasma nitrogen present in each sac was 29 mg.

Zinc ($\mu\text{g.}$) added to system as zinc- glycine complex	Zinc bound to plasma ($\mu\text{g./mg. of N}$)
40	1.3
80	2.4
120	3.7
160	5.0
180	5.5
280	8.0
320	9.1
380	9.8
440	12.8
520	14.1
600	15.7

Table 5. *Amount of zinc associated with the globulin fraction of rabbit plasma labelled in vitro with varying amounts of radioactive zinc by equilibrium dialysis*

The plasma was labelled with varying amounts of ^{65}Zn by the equilibrium-dialysis method (as in Table 4). Globulins were separated from the plasma by the addition of an equal volume of saturated ammonium sulphate soln. at pH 7.0 or pH 4.2.

Zinc added to system ($\mu\text{g.}$)	Zinc content ($\mu\text{g./mg. of N}$)		
	Whole plasma	Globulin fraction	
		pH 7.0	pH 4.2
160	5.0	7.1	4.3
320	9.1	11.9	5.7
440	12.8	10.2	9.3
520	14.1	16.4	8.5
600	15.7	15.8	11.1

solutions used for the precipitations must therefore not be so low that metal is removed from the protein.

Distribution of zinc between some protein fractions of human plasma

Human plasma was labelled *in vitro* with ^{65}Zn by equilibrium-dialysis, as described in the Materials and Methods section. Six solutions were used, containing 100, 200, 300, 500, 750 and 1000 $\mu\text{g.}$ of total zinc. The total amount of plasma nitrogen present in the system was 40 mg. After the plasma had been labelled, the Cohn low-temperature-fractionation technique (Hughes, 1954) was used to separate the proteins into the following fractions: fibrinogen, β - and γ -globulins (plus prothrombin and plasminogen), α - and β -globulins plus some

albumin, and albumin. The ^{65}Zn and nitrogen content of each fraction was measured. The results have been expressed as the ratio: (^{65}Zn content of fraction/mg. of nitrogen)/(^{65}Zn content of plasma/mg. of nitrogen). The amount of zinc added to the system did not seem to affect the ratios. Only one fraction, that containing the α - and β -globulins, had a ratio higher than unity (1.53–2.74). The lowest ratios (0.33–0.65) were found in the albumin fraction, and this finding was supported by the results of later paper-electrophoresis studies described below. Fibrinogen bound more zinc than the fraction containing β - and γ -globulins. The rather variable ratios found in the α - and β -globulin fraction may be due to the presence of different amounts of contaminating albumin.

Distribution of zinc between some protein fractions of human serum and rabbit plasma determined by paper electrophoresis

Paper electrophoresis provides a rapid method for the study of the distribution of a metal between the various plasma proteins. If the metal can be obtained in the radioactive state, the use of plasma labelled *in vivo* or *in vitro* with the isotope enables the distribution of the metal between the various plasma proteins to be determined with comparative ease. The method also has the advantage that only very small samples of plasma are needed. This was important in our experiments as most of the blood from the rabbits injected with ^{65}Zn was required for studies on the distribution of the isotope between erythrocytes and leucocytes.

It was first necessary to examine the behaviour of zinc chloride and the zinc-glycine complex in electrophoresis. Samples of zinc chloride and zinc-glycine containing ^{65}Zn were placed separately on Whatman no. 1 paper strips and run as described above. After the strips had been dried they were marked off into numbered transverse sections, 0.5 mm. wide, and the zinc-glycine strips were in addition marked longitudinally down the centre. The zinc-glycine strips were then cut down the centre and one-half of the strip was treated with ninhydrin to find the position of the glycine. The other half-strip was cut into 18 0.5 in. wide sections and the ^{65}Zn content of each section was measured. Similarly, ^{65}Zn determinations were made on the transverse sections of whole strips labelled with zinc chloride containing ^{65}Zn .

As might be expected, the ^{65}Zn on the strip marked with $^{65}\text{ZnCl}_2$ migrated towards the cathode and the major part of the activity was located about 3 cm. from the origin. The ^{65}Zn on the zinc-glycine-marked strip was found almost the same distance from the origin. The glycine also moved towards the cathode but did not migrate as rapidly as the zinc; the ninhydrin-stained band was found

almost equidistant between the origin and the position of major ^{65}Zn activity. From these experiments it may be concluded that, in the absence of protein, zinc, whether applied as zinc chloride or as the zinc-glycine complex, will migrate towards the cathode; under these conditions of paper electrophoresis most of the zinc in the zinc-glycine complex is dissociated and travels as the free ion.

Our first studies were made on human serum which had been labelled *in vitro* with a ^{65}Zn -containing zinc-glycine solution (soln. E). Human serum (2 ml.) was mixed with soln. E (0.5 ml.) and samples were taken for electrophoresis 5 min. after mixing. The albumin, α_1 -globulin, α_2 -globulin, β -globulin and γ -globulin fractions contained respectively 22–38, 25–34, 23–26, 15–17 and 1% of the total ^{65}Zn in the plasma. Expressed as the

ratio (^{65}Zn content of protein fraction/mg. of nitrogen)/(^{65}Zn content of whole plasma/mg. of nitrogen) the corresponding values were 0.4–0.6, 3.6–4.0, 3.0–3.5, 1.2–1.3 and 0.1.

A similar experiment made with rabbit plasma showed that 44–54% of the total ^{65}Zn in the plasma was in the albumin fraction, 38–44% in the combined α -globulins, 8–10% in the β -globulins and 1% in the γ -globulins.

The remaining experiments in this series were made on plasma obtained from rabbits which had been injected with soln. E (5 ml. given intravenously). Samples of blood were collected in heparinized bottles 5 min. and 1 and 3 hr. after the injection. The plasma was separated immediately and used for the paper-electrophoresis studies. The results are given in Tables 6 and 7. The maximum

Table 6. *Distribution of ^{65}Zn between protein fractions of plasma of rabbits injected intravenously with zinc-glycine solution containing ^{65}Zn*

Samples of blood were taken from the ear vein of the rabbits at the times indicated and collected in polythene bottles containing heparin. The plasma proteins were separated by paper electrophoresis as described in the text. All the results represent the average values for two (and in most cases, three) separate electrophoretic separations of the plasma samples. Results for ^{65}Zn content are expressed as percentages (to the nearest whole number) of ^{65}Zn in the whole plasma.

Time after injection (min.)	Rabbit no.	^{65}Zn content of plasma fractions (separated by paper electrophoresis)			
		Albumin	α_1 - plus α_2 -Globulin	β -Globulin	γ -Globulin
5	412	29	68	3	1
	413	35	59	6	1
	403	16	80	3	<1
	464	14	83	2	<1
60	412	27	71	1	2
	413	32	62	4	2
	403	34	58	8	0
	464	46	35	18	2
180	464	41	47	13	<1

Table 7. *Binding of ^{65}Zn by plasma proteins of rabbits injected intravenously with zinc-glycine solution containing ^{65}Zn*

Samples of blood were taken from the ear vein of the rabbits at the time indicated, and were collected in heparinized vessels. Plasma proteins were separated by paper electrophoresis as described in the text. The amount of protein in each fraction was determined by staining the strip with bromophenol blue and scanning with an EEL scanner. All the results represent the average values for two (and in most cases, three) separate electrophoretic separations on the plasma samples.

Time after injection (min.)	Rabbit no.	Relative ^{65}Zn content* of protein fractions			
		Albumin	α_1 - plus α_2 -Globulin	β -Globulin	γ -Globulin
5	412	0.5	3.3	0.2	0.2
	413	0.8	4.2	0.3	0.0
	403	0.4	3.7	0.2	<0.1
	464	0.3	7.6	0.1	<0.1
60	412	0.6	6.0	<0.1	0.1
	413	0.7	4.7	0.2	0.1
	403	0.8	2.8	0.6	0.0
	464	1.0	2.7	1.0	<0.1
180	464	0.9	4.1	0.7	<0.1

* Expressed as ratio: (^{65}Zn /mg. of protein in fraction)/(^{65}Zn /mg. of protein in whole plasma).

amount of plasma which could be applied to the paper (0.05 ml.) did not contain sufficient ^{65}Zn to enable accurate counts to be made on the 3 hr. samples, except with the plasma from rabbit 464. The results for the α_1 - and α_2 -globulins have been combined, the α -globulins being treated as one fraction. Generally, the ^{65}Zn contents of the α_1 - and α_2 -globulin fractions were about equal. The values given for each rabbit are the results of experiments on the same plasma, triplicate or duplicate samples being run at the same time in the electrophoresis tank. Results obtained with samples of the same plasma are rather variable, particularly in the β -globulin fraction, but certain conclusions may be drawn. The γ -globulins are apparently not concerned in the binding of plasma zinc. Very little activity could be found in this fraction in either the serum or plasma labelled *in vitro* or the plasma labelled *in vivo*. A major portion (40–78 %) of the plasma ^{65}Zn 5 min. or 1 hr. after the injection of the isotope may be found in the α -globulins. The albumin fraction contains 20–56 % of the ^{65}Zn .

If the ratios given in Table 7 are an index of the extent to which zinc is bound to the various protein fractions, it seems that the α -globulins have the highest concentration of ^{65}Zn . There appears to be very little change in the distribution of ^{65}Zn between the proteins during the period 5 min.–1 hr. after the injection.

Distribution of zinc between some protein fractions of rabbit plasma determined by starch-gel electrophoresis

To confirm that the distribution of zinc as determined by paper electrophoresis was not affected by the experimental conditions a few experiments with ^{65}Zn -labelled plasma were made with starch-gel electrophoresis (Smithies, 1955).

Samples of plasma on Whatman no. 3MM paper strips (3 cm. $\times$ 0.5 cm.) were inserted into gels made in borate buffer, pH 8.5. Samples were run for 18 hr. at 500 v and 5 mA/strip. The gels were then horizontally sliced into two and the top strip was stained with bromophenol blue. The lower strip was sliced transversely into sections corresponding to the stained bands and the slices were placed in glass dishes for determinations of ^{65}Zn .

A sample of plasma from rabbit 403 taken 5 min. after injection of ^{65}Zn gave the following results (expressed as percentages of the total ^{65}Zn on the strip): α_2 -globulin 48; albumin 36; β -globulin 14. A sample of plasma from the same rabbit taken 1 hr. after the injection gave: α_2 -globulin 57; albumin 12; β -globulin 26. Separation of α_1 -globulin from albumin was not achieved and the figure for the ^{65}Zn content of the albumin fraction probably includes a contribution from the α_1 -globulin fraction. The results confirm that a major

part of the ^{65}Zn in the plasma is carried by the α_2 -globulin fraction. The γ -globulin bands contained very little ^{65}Zn (1 % or less).

^{65}Zn content of leucocytes from blood and bone marrow of rabbits injected intravenously with the isotope as zinc-glycine complex

The object of this experiment was to enable a comparison to be made between the ^{65}Zn content of leucocytes and erythrocytes in blood and bone marrow at various times after an intravenous injection of ^{65}Zn as the zinc-glycine complex. Experimental details have been given in the Materials and Methods section. The results (Table 8) showed that the ratio leucocyte ^{65}Zn /erythrocyte ^{65}Zn was highest (290 and 252) in the blood 1 hr. after the injection and fell to about a quarter of the 1 hr. value 24 hr. after the injection. The values for the bone-marrow ratios were much lower than those for the blood and, with the exception of rabbits 480 and 456, were almost constant at about 30. Entry of ^{65}Zn into the circulating leucocytes was rapid and this would indicate that a free exchange of zinc occurs between plasma zinc and leucocyte zinc. The lower values for the bone-marrow leucocytes may reflect the great preponderance of immature leucocytes in the marrow.

Exchange of zinc between erythrocytes or leucocytes and the plasma

Isotope-labelling experiments have shown that zinc readily penetrates into erythrocytes and leucocytes (Tupper, Watts & Wormald, 1950), and studies on man have shown that intravenously

Table 8. ^{65}Zn content of leucocytes from blood and bone marrow of rabbits injected intravenously with the isotope

Time after injection (hr.)	Rabbit no.	^{65}Zn content* of leucocytes	
		Blood	Bone marrow
1	480	290	51
	456	252	44
	468	—	30
6	409	102	—
	413	110	—
	431	120	30
24	374	62	32
	412	51	28
	409	39	31
	413	75	30

* Expressed as ratio: (counts/min./ 10^9 leucocytes)/(counts/min./ 10^9 erythrocytes).

injected ^{65}Zn is rapidly incorporated into erythrocytes and leucocytes (Ross, Ebaugh & Talbot, 1958). It would be interesting to know how far this incorporation of zinc in these blood cells merely involves diffusion of Zn^{2+} ions through the cell membranes, and how far biological synthetic mechanisms are concerned, as, for example, the synthesis of enzymes containing zinc.

We made preliminary experiments to determine how rapid and complete is the passage of Zn^{2+} ions from the blood cells to the plasma, and in the opposite direction. Separated normal rabbit erythrocytes and leucocytes were immersed in rabbit plasma biologically labelled with ^{65}Zn , and in other experiments rabbit erythrocytes and leucocytes biologically labelled with ^{65}Zn were immersed in normal rabbit plasma. In both cases the cells were separated from the mixtures, usually after standing at room temperature for 1 hr. The cells were washed well with 0.9% sodium chloride and their ^{65}Zn contents were determined.

Two rabbits were given intravenous injections of a solution of zinc-glycine containing ^{65}Zn (soln. G; 5 ml. to each rabbit) and the animals were bled after 1 hr. The blood of these rabbits was collected in polythene bottles containing heparin, and part of the pooled blood was centrifuged for the separation of the plasma. The remainder of the blood was mixed with Dextraven and the blood cells were separated and washed as previously described (Dennes *et al.* 1961). Suspensions of the leucocytes or erythrocytes in 0.9% sodium chloride were made and well mixed with normal rabbit plasma. After 1 hr. at room temperature these mixtures were centrifuged and the cells washed well with 0.9% sodium chloride. Cell counts and determinations of ^{65}Zn were made on the suspensions, and radioactivity measurements on the washings (to determine whether the washing had been adequate to remove all loosely adhering ^{65}Zn). The biologically labelled blood used in these experiments yielded leucocytes and erythrocytes that had respectively an activity of 108×10^3 and 3.7×10^3 counts/min./ 10^6 cells; the plasma gave 3×10^6 counts/min./ml.

In the experiments where the biologically labelled blood cells were suspended in normal rabbit plasma for 1 hr., the labelled erythrocytes lost about half the ^{65}Zn they originally contained, whereas under similar conditions there was little or no loss of zinc by the biologically labelled leucocytes.

Exposure for 1 hr. at room temperature led to rapid uptake of ^{65}Zn by rabbit leucocytes and erythrocytes immersed in the biologically labelled plasma. The amount taken up in 1 hr. by the leucocytes was approximately the same as that present in the leucocytes of the blood which yielded the labelled plasma, i.e. the labelling *in vitro* seemed to be quite parallel to that *in vivo*. The

erythrocytes, however, took up from the plasma only about one-tenth of the amount that was incorporated into the erythrocytes in the biologically labelled blood.

DISCUSSION

Unlike iron and copper, which are carried in the serum firmly bound to transferrin and caeruloplasmin respectively, zinc is transported in two main fractions, one loosely bound, the other firmly bound. In the loosely bound fraction zinc is probably associated with albumin, and, in the firmly bound fraction, with globulins (Vikbladh, 1951). It has been shown that zinc will combine with the imidazole groups (approximately 16) of serum albumin *in vitro*, but this binding is not very firm and, in the presence of an excess of glycine, zinc is removed from combination with the albumin (Gurd & Goodman, 1952). Zinc will also form, *in vitro*, water-soluble complexes with α -lipoproteins, α -glycoprotein and the β_1 -metal-combining globulin (transferrin).

There is no evidence that in serum the β_1 -globulin carries firmly bound zinc. Our results with the rabbits whose plasma proteins had been saturated with iron before injection of ^{65}Zn indicate that zinc is probably not transported by this protein. However, zinc may be bound to a site on the β_1 -globulin different from that which binds the iron. The iron-transferrin complex is a particularly stable one, and zinc is unlikely to displace iron from combination with the protein.

Wolff (1956) has shown that in dogs 6 hr. after oral administration of ^{65}Zn 50% of the total serum ^{65}Zn is found in the α -globulin fraction and the major part of this ^{65}Zn is firmly bound. The β - and γ -globulin fractions contained only 12 and 9% of the total serum ^{65}Zn respectively. Most of the ^{65}Zn associated with the β -globulin was loosely bound. Our results obtained after intravenous injection of ^{65}Zn into rabbits are in agreement with those of Wolff for dogs. We found that the amount of zinc bound by albumin and β -globulins varies considerably and that this variation may be attributed to the loose binding of the metal to these proteins. The results obtained with human plasma labelled *in vitro* by using the Cohn low-temperature fractionation also support the view that the α -globulin fraction is the major carrier of firmly bound zinc.

The specific activity of the globulin fraction from plasma labelled *in vivo* with ^{65}Zn is lower than that of the whole plasma. This agrees with the view that the zinc is more firmly bound to the globulins than to the other plasma proteins. A low rate of exchange between injected ^{65}Zn and globulin-bound zinc would result in a specific activity lower than that of the whole plasma.

That the affinity of α -globulins for zinc is very great is indicated by the ratios given in Table 7, which show that the amount of zinc bound per milligram of protein is very much greater in the α -globulins than in the other plasma proteins studied. From our evidence it would appear that the γ -globulins transport a negligible fraction of the serum zinc.

Although, *in vitro*, glycine will remove zinc in combination with albumin, it was not possible to remove more than 7% of ^{65}Zn bound *in vivo* from rabbit plasma dialysed against a veronal buffer containing glycine. The 30% or so of the plasma zinc bound to albumin in the presence of the other plasma proteins may not be so easily removed by glycine as in an isolated system (see Gurd & Goodman, 1952). None of the ^{65}Zn present in the labelled plasma is as firmly bound as the zinc in some zinc-containing enzymes, e.g. carbonic anhydrase, yeast and liver alcohol dehydrogenases and glutamic dehydrogenase. All the ^{65}Zn could be removed from the labelled plasma by dialysis against veronal buffer, pH 7.4, containing EDTA.

Berfenstam (1952) stated that there seems to be no evidence for any plasma-saturation limit for zinc as there is for iron; the results of our equilibrium-dialysis studies are in agreement with this. Plasma would appear to be capable of binding zinc up to the point at which actual precipitation of proteins occurs.

In studies on precipitation of protein fractions by acidic reagents such as ammonium sulphate, the effect of pH on the metal-protein complex must be considered. As Table 5 shows, the globulin fraction obtained by precipitation at pH 7.0 contains more zinc than that obtained by precipitation at pH 4.2.

After injection of rabbits with ^{65}Zn the isotope content of the plasma falls rapidly in the first hour and there is then a slow decrease. The rapid fall is undoubtedly due to removal of plasma zinc by the tissues. The slow decrease may be caused either by a gradual release of ^{65}Zn from the α -globulin fraction, where the zinc is firmly bound, or by a gradual re-entry of ^{65}Zn from the tissues into the blood, resulting in an apparent decrease in the rate at which zinc is leaving the plasma. There is no information to show which of these possibilities is correct.

The erythrocyte contain 85% of the total blood zinc. It is generally accepted that most of this zinc is firmly bound in the carbonic anhydrase of the erythrocytes. The leucocytes contain 3% of the total blood zinc, but the concentration of the metal in these cells is, in human blood, about 15 times the concentration in the erythrocytes. The results given in Table 8 show that entry of ^{65}Zn into leucocytes is rapid. The exchange experiments between labelled leucocytes and unlabelled plasma

indicate that the zinc, once taken up by the leucocytes, is not easily lost. These findings agree with the results of some earlier experiments which showed that EDTA would not remove zinc from leucocytes *in vitro* (Dennes *et al.* 1961).

Labelled erythrocytes left in contact with normal plasma for 1 hr. lose an appreciable amount of ^{65}Zn . No haemolysis was observed in our experiments and rupture of the erythrocytes cannot be responsible for this finding. Erythrocytes suspended in ^{65}Zn -labelled plasma *in vitro* took up only about 8% of the amount of ^{65}Zn which they take up under comparable conditions *in vivo*, whereas leucocytes took up about the same amount of zinc *in vivo* or *in vitro*. It is improbable that, *in vivo*, sufficient erythrocytes containing ^{65}Zn are synthesized in 1 hr. to account for the great difference in ^{65}Zn content between labelled cells *in vivo* and *in vitro*.

Equine erythrocytes will *in vitro* take up ^{65}Zn which becomes bound to haemoglobin (Sivarama Sastry, Viswanathan, Ramaiah & Sarma, 1960). There appears to be a pool of labile zinc in erythrocytes, and the carbonic-anhydrase zinc may not account for the bulk of the total erythrocyte zinc. This loosely bound erythrocyte zinc may form part of the soft-tissue pool of labile zinc referred to by Gilbert & Taylor (1956). If the formed elements of the blood assist materially in zinc transport, the erythrocytes and not the leucocytes must be responsible for this carriage of zinc.

The amount of ^{65}Zn found in the leucocytes of the rabbit bone marrow was very much lower than in the blood leucocytes and also very constant. The bone marrow contains a high preponderance of immature leucocytes and these leucocytes may either contain less zinc or take up zinc more slowly than do mature leucocytes. The finding that leukaemic leucocytes contain much less zinc than do normal leucocytes (Gibson, Vallee, Fluharty & Nelson, 1950; Wolff, 1956; Dennes, Tupper & Wormall, 1960) may be pertinent. In leukaemia, large numbers of immature leucocytes may be present in the blood and this may account for the low zinc content of the leucocytes. Some results obtained by us in cases of chronic lymphatic leukaemia (Dennes *et al.* 1961) indicate, however, that the immaturity of the leucocytes may not be responsible for the low zinc content.

SUMMARY

1. The transport of zinc in the blood of the rabbit after the intravenous injection of zinc-glycine solutions, and the rate of removal of the metal from the blood, have been studied with ^{65}Zn as tracer.

2. The injected zinc was rapidly removed from blood and about 90 % left the blood stream in the course of 3 hr.

3. A similar rapid loss occurred when plasma labelled *in vivo* with ^{65}Zn was injected. Plasma protein-bound zinc is thus readily exchangeable with the pool of body zinc.

4. The blood plasma of the rabbits, 3–24 hr. after the injection of zinc–glycine, contained essentially all its ^{65}Zn in a bound form. About half this zinc was attached to the plasma globulin, and it has been shown that the α -globulin fraction is responsible for the carriage of the main part of the zinc.

5. Dialysis studies have shown that at pH 7.4 all the zinc is removed from rabbit plasma by ethylenediaminetetra-acetic acid, but only a small fraction (about 7 %) by glycine.

6. Intravenous injection of iron salts into rabbits in amounts sufficient to saturate the iron-carrying capacity of the plasma proteins has no significant effect on the binding powers of plasma for zinc *in vivo*.

7. It has been confirmed that the zinc-binding capacity of rabbit leucocytes is very much greater than that of the erythrocytes. Ratios of leucocyte ^{65}Zn to erythrocyte ^{65}Zn in blood and bone marrow 24 hr. after the injection of zinc–glycine containing ^{65}Zn were 39–75/1 and 30/1 respectively.

8. Studies have shown that *in vitro* there is a rapid exchange of zinc between erythrocytes and plasma, with no detectable transference of zinc from leucocytes to plasma. Uptake of zinc by leucocytes from plasma is rapid.

9. The role of the blood cells in the transport and storage of zinc is briefly discussed. There is good evidence that zinc in the blood cells, mainly that in the erythrocytes, should be regarded as a significant part of the mobilizable or 'available' zinc in the animal body.

This work was carried out with the aid of grants from the British Empire Cancer Campaign, and we are also indebted to Dr T. E. Banks, of the Physics Department of this College, for continued valuable help with the maintenance of equipment used for radioactivity measurements. We also wish to thank Dr D. J. Lawford for advice about the use of starch gel for electrophoresis of plasma proteins.

REFERENCES

- Banks, T. E., Tupper, R., Watts, R. W. E. & Wormal, A. (1955). *Biochem. J.* **59**, 149.
- Banks, T. E., Tupper, R., White, E. M. A. & Wormal, A. (1959). *Int. J. Appl. Rad. Isotopes*, **4**, 221.
- Berfenstam, R. (1952). *Acta Paediat., Suppl.*, **87**.
- Delezenne, C. (1919). *Ann. Inst. Pasteur*, **33**, 68.
- Dennes, E., Tupper, R. & Wormal, A. (1960). *Nature, Lond.*, **187**, 302.
- Dennes, E., Tupper, R. & Wormal, A. (1961). *Biochem. J.* **78**, 578.
- Durum, E. L. (1950). *J. Amer. chem. Soc.* **72**, 2943.
- Gibson, J. G., Vallee, B. L., Fluharty, R. G. & Nelson, G. E. (1950). *Unio intern. clin. contra cancerum. acta*, **6**, 1102.
- Gilbert, I. G. F. & Taylor, D. M. (1956). *Biochim. biophys. Acta*, **21**, 544.
- Gomori, G. (1955). In *Methods of Enzymology*, vol. 1, p. 138. Ed. by Colowick, S. P. & Kaplan, N. O. New York: Academic Press.
- Gurd, F. R. N. & Goodman, D. S. (1952). *J. Amer. chem. Soc.*, **74**, 670.
- Hughes, W. L. (1954). In *The Proteins*, vol. 2, Part B, p. 745. Ed. by Neurath, H. & Bailey, K. New York: Academic Press.
- Hughes, T. R. & Klotz, I. M. (1956). *Meth. Biochem. Anal.* **3**, 278.
- Markham, R. (1942). *Biochem. J.* **36**, 790.
- Ross, J. F., Ebaugh, F. G. & Talbot, T. R. (1958). *Trans. Ass. Amer. Physicians*, **71**, 322.
- Russell, J. (1944). *J. biol. Chem.* **156**, 457.
- Sivarama Sastry, K., Viswanathan, L., Ramaiah, A. & Sarma, P. S. (1960). *Biochem. J.* **74**, 561.
- Smithies, O. (1955). *Biochem. J.* **61**, 629.
- Tupper, R., Watts, R. W. E. & Wormal, A. (1950). *Biochem. J.* **48**, xxxvii.
- Tupper, R., Watts, R. W. E. & Wormal, A. (1951). *Biochem. J.* **50**, 429.
- Vallee, B. L. (1959). *Physiol. Rev.* **30**, 443.
- Vallee, B. L. & Gibson, J. G. (1948). *J. biol. Chem.* **176**, 435.
- Varley, H. (1958). *Practical Clinical Chemistry*, p. 198. London: Heinemann.
- Vessel, E. S. & Bearn, A. G. (1957). *Proc. Soc. exp. Biol., N.Y.*, **94**, 96.
- Vikbladh, I. (1951). *Scand. J. clin. lab. Invest. Suppl.* **2**, 9.
- Wolff, H. P. (1956). *Klin. Wschr.* **34**, 409.