action of penicillinase from *B. cereus* or *Staph. aureus* on penicillins. Different derivatives show a similar ability to inhibit the enzyme from *B. cereus* but differ greatly in their ability to inhibit the staphylococcal enzyme.

3. Culture fluids of strains of *B. cereus* contain a cephalosporinase which may be a β -lactamase.

4. Several derivatives of 7-aminocephalosporanic acid induce the formation of penicillinase by a strain of *Staph. aureus* and a strain of *B. cereus*. Under the conditions used the relative efficiencies of the inducers vary greatly from one organism to the other.

REFERENCES

- Abraham, E. P. & Newton, G. G. F. (1956). *Biochem. J.* **63**, 628.
 Abraham, E. P. & Newton, G. G. F. (1961a). *Endeavour*, **20**, 92.
 Abraham, E. P. & Newton, G. G. F. (1961b). *Biochem. J.* **79**, 377.
 Batchelor, F. R., Doyle, F. P., Nayler, J. H. C. & Rolinson, G. N. (1959). *Nature, Lond.*, **183**, 257.
 Crawford, K. & Abraham, E. P. (1957). *J. gen. Microbiol.* **16**, 604.
 Eadie, G. S. (1942). *J. biol. Chem.* **146**, 85.
 Hale, C. W., Newton, G. G. F. & Abraham, E. P. (1961). *Biochem. J.* **79**, 403.
 Hofstee, B. H. J. (1959). *Nature, Lond.*, **184**, 1296.
 Hunter, A. & Downs, C. E. (1945). *J. biol. Chem.* **157**, 427.
 Jeffery, J. d'A., Abraham, E. P. & Newton, G. G. F. (1961). *Biochem. J.* **81**, 591.
 Kay, D. & Fildes, P. (1950). *Brit. J. exp. Path.* **31**, 338.
 Kogut, M., Pollock, M. R. & Tridgell, E. J. (1956). *Biochem. J.* **62**, 391.
 Loder, B., Newton, G. G. F. & Abraham, E. P. (1961). *Biochem. J.* **79**, 408.
 Moat, A. G., Ceci, L. N. & Bondi, A. (1961). *Proc. Soc. exp. Biol., N.Y.*, **107**, 675.
 Novick, R. (1962). *Biochem. J.* (in the Press).
 Pollock, M. R. (1950). *Brit. J. exp. Path.* **31**, 739.
 Pollock, M. R. (1952). *Brit. J. exp. Path.* **33**, 587.
 Pollock, M. R. (1957). *Biochem. J.* **66**, 419.
 Pollock, M. R. (1959). In *The Enzymes*, 2nd ed., vol. 1, p. 619. Ed. by Sumner, J. B. & Myrback, K. New York: Academic Press Inc.
 Pollock, M. R. (1960). In *The Enzymes*, 2nd ed., vol. 4, p. 269. Ed. by Boyer, P. D., Lardy, H. & Myrback, K. New York: Academic Press Inc.
 Pollock, M. R. & Torriani, A. M. (1953). *C.R. Acad. Sci., Paris*, **237**, 276.
 Rogers, H. J. (1953). *J. Path. Bact.* **66**, 545.
 Rolinson, G. N., Stevens, S., Batchelor, F. R., Wood, J. C. & Chain, E. B. (1960). *Lancet*, ii, 564.
 Sheehan, J. C., Henery-Logan, K. R. & Johnson, D. A. (1953). *J. Amer. chem. Soc.* **75**, 3292.

Biochem. J. (1962) **83**, 63

Studies on the Mode of Action of Excess of Vitamin A

4. THE SPECIFICITY OF THE EFFECT ON EMBRYONIC CHICK-LIMB CARTILAGE IN CULTURE AND ON ISOLATED RAT-LIVER LYSOSOMES*

By HONOR B. FELL†, J. T. DINGLE AND M. WEBB‡
Strangeways Research Laboratory, Cambridge

(Received 11 August 1961)

Previous work has shown that when embryonic cartilage is grown in medium containing an excess of vitamin A, metachromatic material is progressively lost from the matrix, which rapidly disintegrates (Fell & Mellanby, 1952; Fell & Thomas, 1960); metachromatic material also disappears from the cartilage of young rabbits fed with excessive amounts of the vitamin (Thomas, McLuskey, Potter & Weissmann, 1960). This previous work

suggested that the degradation of the cartilage matrix under the influence of hypervitaminosis A might be due to an enhanced proteolytic activity of the chondroblasts; recent results (Dingle, Lucy & Fell, 1961; Lucy, Dingle & Fell, 1961) have confirmed this view and indicated that the degradation is associated with the release of protease from the lysosomes (Dingle, 1961).

The present experiments were undertaken to determine whether the action of vitamin A on cartilage in culture and on isolated lysosomes is specific, or whether a similar effect can be produced by other chemically-related compounds. The

* Part 3: Dingle (1961).

† Royal Society Foulerton Research Fellow.

‡ Member of the External Staff, Medical Research Council.

results indicate that the effect of the vitamin is highly specific on cartilage in culture and also at low concentrations on the isolated lysosomes.

MATERIALS AND METHODS

Compounds tested

Vitamin A (alcohol). A commercial preparation (Roche) was dissolved in ethanol (100 mg./ml.) and stored under nitrogen. Dilutions were made immediately before use. If the dilute solution showed a yellow colour indicative of oxidation a fresh preparation was used.

Vitamin A acid. This compound was a gift from Dr T. Moore of the Dunn Nutritional Laboratory, Cambridge. The sodium salt was obtained by neutralization of an ethanolic solution of the acid with the calculated amount of sodium hydroxide. The water-soluble preparation was dissolved in 0.02 M-sodium acetate in 0.25 M-sucrose and the pH adjusted to 7.

Vitamin A acetate and palmitate. These derivatives were commercial preparations (British Drug Houses Ltd.).

Vitamin A epoxide. This was prepared by oxidation of vitamin A (alcohol) (1 g.) according to the method of Karrer & Jucker (1947). The product (100 mg.) from the oxidation moved as a single band when rechromatographed on alumina with anhydrous benzene as the mobile phase, and had absorption maxima at 238 m μ ($E_{1\text{cm}}^{1\%}$ 337) and 281 m μ ($E_{1\text{cm}}^{1\%}$ 282), as found by Vinet & Meunier (1947), but not at 275 m μ as reported by Karrer & Jucker (1947).

Hydrogenated vitamin A. Vitamin A (alcohol) in ethanol (1 g./20 ml.) was hydrogenated at a pressure slightly greater than atmospheric and at room temperature (18°) in the presence of Adams catalyst (23.5 mg.). Uptake of hydrogen ceased when the theoretical amount for the saturation of four double bonds/molecule had been absorbed. The solution was filtered in nitrogen to remove the catalyst, diluted to 50 ml. with ethanol and stored under nitrogen. A portion of the solution was evaporated to dryness *in vacuo* and the residue dissolved in light petroleum (b.p. 80–100°). This solution (50 μ g./ml.) had an absorption maximum below the limit of the spectrophotometer (200 m μ), and exhibited no absorption above 300 m μ .

Anhydrovitamin A. This derivative was prepared in 70–75% yield by the method of Edisburg, Gillam, Heilbron & Morton, (1932), and isolated by chromatography in light petroleum (b.p. 60–80°) solution on alumina (Shantz, Cawley & Embval, 1943). Evaporation of the eluate in nitrogen under reduced pressure gave a yellow syrup which crystallized slowly at room temperature.

In light petroleum (b.p. 80–100°) solution the compound exhibited three sharp absorption bands in the range 300–400 m μ with maxima at 351, 369 and 392 m μ . At 369 m μ $E_{1\text{cm}}^{1\%}$ for different preparations varied from 2560 to 3010. Although the compound could be crystallized as deep-yellow prisms (m.p. 78–79°) from either light petroleum (b.p. 60–80°) (Shantz *et al.* 1943) or ethanol solutions at low temperature, its concentrated solutions were extremely susceptible to oxidation by traces of air, and it was difficult to avoid the formation of an insoluble resinous material during the separation of the crystals. Preparations for the biological experiments, therefore, were used without recrystallization. Under the conditions of storage in these experiments, the absorption of a solution of the compound

at 369 m μ decreased by 12.8% over a period of 70 days; the proportions of the $E_{1\text{cm}}^{1\%}$ values at 350, 369 and 391 m μ changed from 1:1.40:1.23 to 1:1.43:1.18.

β -Ionone. The commercial preparation ('perfumery quality', British Drug Houses Ltd.) was purified as described by Tiemann (1898). The intermediate semicarbazone was recrystallized from ethanol to constant m.p. (148°), the crude compound being dissolved initially in an excess of the solvent to remove less-soluble impurities. The β -ionone (b.p. 134°/14 mm., 130–131°/12 mm.; d , 0.946), obtained by hydrolysis of the semicarbazone, was stored in sealed ampoules under nitrogen.

Brij 35. This neutral detergent (Atlas Powder Co., Wilmington, Del., U.S.A.), an ether of polyethylene glycol, was kept as a 1% (w/v) solution in glass-distilled water.

Other chemicals. Other compounds tested were commercial products of the highest purity available.

Organ-culture experiments

The rudiments of the humerus, ulna, femur and tibia of 7-day-old chick embryos were used. At this stage the bones consist of a cartilaginous rod differentiated into the usual small-celled epiphyseal regions, proliferative zones of flattened cells and middle zone of hypertrophic cells; the hypertrophic cartilage is usually covered by a thin layer of periosteal bone. In sections, the cartilage matrix stains metachromatically with toluidine blue and appears a deep reddish-purple.

The following compounds were tested and their effects compared: vitamin A (alcohol), vitamin A acid, hydrogenated vitamin A, anhydrovitamin A, β -ionone and phytol.

The technique of organ culture was the same as that already described (Dingle *et al.* 1961). Eight pairs of bone rudiments, namely the humeri, ulnae, femora and tibiae from two embryos, were used for each test. The lengths of the explants were measured at the beginning of the experiment and at 2-day intervals during cultivation by the method described by Fell & Mellanby (1952). The results are expressed as the mean of the lengths of the eight rudiments in each group. Length was measured on camera lucida drawings in arbitrary units, 2.6 units being equal to 1 mm.

After eight days' growth, the explants were fixed in Zenker's fluid plus 3% (v/v) acetic acid for 30 min., followed by Zenker's fluid without acetic acid for 1½ hr. Serial sections were cut and stained with 0.5% toluidine blue (G. T. Gurr Ltd., London) in 5% ethanol for 30 min.

In the first series of experiments, all seven compounds, dissolved in ethanol, were tested against vitamin A (alcohol). Each compound was added to the plasma so as to give a concentration of 0.01 μ mole/ml. in the final medium (plasma-embryo-extract clot). The rudiments from one side of each chick were grown in medium containing the substance being tested, and those from the opposite side were cultivated in medium to which an equimolar concentration of vitamin A (alcohol) had been added. In the second series of experiments, the action of hydrogenated vitamin A, anhydrovitamin A, β -ionone and phytol in a tenfold concentration (0.1 μ mole/ml.) was investigated, and the treated rudiments were compared with controls grown in normal medium to which only ethanol had been added. All the media in both series contained the same amount of solvent (0.2% ethanol).

Table 1. *Effect of vitamin A and related compounds on the matrix and growth in length of chick-cartilaginous-limb-bone rudiments in culture*

The compounds were added to the culture as ethanolic solutions at a final concentration of 0.01 μ mole/ml. An equal amount of ethanol was added to the control. The bone rudiments of 7-day-old chick embryos were grown in organ culture for eight days. The mean length at the end of the culture period is given $\pm$ the standard error. The figures in parentheses are the number of bone rudiments. The effect on metachromasia is indicated by the following symbols: $\pm$, rather less matrix in hypertrophic cartilage of some explants; +, rather less matrix in hypertrophic cartilage; + + + +, metachromasia nearly gone, drastic effect.

Addition to medium	Effect on matrix	Mean length after 8 days in culture (arbitrary units)
None (control)	-	15.66 $\pm$ 0.282 (8)
Phytol	-	16.31 $\pm$ 0.405 (8)
Anhydrovitamin A	-	15.79 $\pm$ 0.443 (8)
β -Ionone	-	15.51 $\pm$ 0.423 (8)
Vitamin A epoxide	-	14.96 $\pm$ 0.252 (8)
Citral	-	14.64 $\pm$ 0.268 (8)
Hydrogenated vitamin A	-	14.10 $\pm$ 0.307 (8)
Vitamin A acid	+ + + +	9.88 $\pm$ 0.394 (8)
Vitamin A (alcohol)	+ + + +	9.48 $\pm$ 0.244 (64)

Lysosome experiments

Lysosome suspensions in 0.25 M-sucrose were prepared as previously described (Dingle, 1961). No attempt was made to separate the lysosomes from the mitochondria in this work. To test the action of the compounds upon the lysosomes, each substance, dissolved in ethanol, was added to a suspension of particles in 0.25 M-sucrose; the suspension was then incubated at 37° for 40 min., after which it was centrifuged in the cold for 20 min. at 15 000g. The free protease was measured in the supernatant fluid as described in earlier work (Dingle, 1961). Haemoglobin (2%, w/v) dissolved in 0.1 M-acetate buffer, pH 4.0, was used as enzyme substrate, and acid-soluble tyrosine was measured after incubation for 20 min. at 37°. In each experiment a standard concentration of vitamin A (alcohol) (0.7 μ mole/ml.) was tested, and the activity of the test compounds in releasing enzyme activity was expressed as a percentage of that released by the standard concentration of the vitamin.

RESULTS

Organ culture experiments

Series 1. At a concentration of 0.01 μ mole/ml. none of the agents tested had a significant effect on the explants except vitamin A acid [Table 1; Figs. 1 and 2; Plate 1, (a)-(d)]; this compound produced the typical vitamin A response, i.e. softening, shrinkage and distortion of the cartilage and, in sections, loss of metachromatic material from the matrix. Hydrogenated vitamin A may have slightly retarded growth in length (Table 1) but no histological effects were seen.

Fig. 1. Camera lucida drawings of five pairs of living femora after cultivation for 8 days. All the explants on the right were grown in medium containing 0.01 μ mole of vitamin A (alcohol)/ml.; the corresponding rudiments on the left were grown either in normal medium (top) or in the presence of 0.01 μ mole of an analogue of vitamin A/ml. Of the test compounds only vitamin A acid produced the same effect as vitamin A.

Fig. 2. Camera lucida drawings of three pairs of living femora after cultivation for 8 days. The explants on the right were grown in medium containing 0.01 μ mole of vitamin A (alcohol)/ml., those on the left in medium with the same concentration of a test compound. None of the test compounds produced an effect comparable with that of vitamin A.

The controls treated with vitamin A (alcohol) underwent all the characteristic changes previously described (Fell & Mellanby, 1952; Fell & Thomas, 1960). There was some variation in the degree of effect in different experiments; this was probably due both to variation in the degree of development of different embryos and to partial oxidation of the vitamin during storage, the rapidity of this process not being fully appreciated at first.

Table 2. *Effect of some related compounds on the growth of chick-limb-bone rudiments in culture*

Cultural conditions were as described in Table 1; the compounds were added in tenfold concentration (0.1 μ -mole/ml.). Each compound was tested against controls in normal medium by the paired limb-bone technique. The standard deviation of the difference and the *P* values for the difference being ascribed to chance are shown. The figures in parentheses are the number of bone rudiments. The symbols for effect on the matrix are defined in Table 1.

	Effect on metachromasia	Mean length (arbitrary units)	S.D. of diff.	
Control	-	16.75 (8)	0.415	<i>P</i> < 0.001
Hydrogenated vitamin A	+	15.53 (8)		
Control	-	17.50 (8)	0.578	<i>P</i> < 0.001
β -Ionone	$\pm$	15.86 (8)		
Control	-	17.73 (8)	1.256	<i>P</i> = 0.07
Anhydrovitamin A	$\pm$	16.70 (8)		
Control	-	16.37 (8)	0.548	<i>P</i> > 0.9
Phytol	-	16.22 (8)

Fig. 3. Comparison of the action of vitamin A and a non-ionic detergent (Brij 35) on isolated rat-liver lysosomes. The action of the compounds was tested by addition to a suspension of particles in 0.25 M-sucrose. The suspensions were then incubated at various temperatures for 45 min., after which they were centrifuged in the cold at 15 000*g* for 20 min. and the activity of the released protease was measured in the supernatant fluids. Vitamin A (●) was used at a final concentration of 0.15 mg./ml. and Brij 35 (○) at 1.0 mg./ml. Enzyme activity is expressed as the acid-soluble tyrosine (μ g.) liberated/ml. of supernatant fraction from a 2% haemoglobin substrate in 20 min. at 37°.

Fig. 4. Release of enzyme from rat-liver lysosomes by the action of vitamin A and of Brij 35. The action of the compounds was tested by addition to a suspension of particles in 0.25 M-sucrose. The suspensions were then incubated at 37° for various periods. After incubation the suspensions were centrifuged in the cold at 15 000*g* for 20 min. and the released enzyme was measured in the supernatant. Vitamin A (●) was used at a final concentration of 0.2 mg./ml. and Brij 35 (○) at 1.0 mg./ml. The activity of the enzyme liberated is expressed as a percentage of the total activity released by hypo-osmotic treatment (Dingle, 1961).

Series 2. Of the four compounds tested at the higher concentration (0.1 μ mole/ml.) (Table 2), phytol had no effect on either growth in length or histology, as compared with controls in normal medium. The remaining three agents all caused a slight reduction in growth, which was statistically significant only for hydrogenated vitamin A and β -ionone, but very little histological change was produced. In section the matrix of some of the treated explants was rather less intensely stained with toluidine blue, and the intercellular partitions in the hypertrophic cartilage were narrower than

in the corresponding controls in normal medium; these effects were greatest in the humeri, femora and tibiae exposed to hydrogenated vitamin A [Plate 1, (e)].

Protease release from lysosomes

Comparison of vitamin A and Brij 35. Previous work has shown that the release of protease from isolated lysosomes by vitamin A is temperature-dependent (Dingle, 1961). This observation has been confirmed and the action of the vitamin at various temperatures compared with that of the detergent (Fig. 3). The latter was effective at

All the explants are from 7-day-old chick embryos and were grown in culture for eight days. Sections were stained with toluidine blue.

(a) Control femur grown in normal medium.

(b) Opposite femur from the same embryo grown in medium containing 0.01 μ mole of vitamin A (alcohol). Note shrinkage, distortion and loss of metachromasia from all but the interior of the epiphyses.

(c) Femur grown in an equimolar concentration of vitamin A acid. The effect is the same as that of vitamin A.

(d) Femur grown in an equimolar concentration of phytol. No effect is apparent (cf. a).

(e) Femur grown in medium containing β -ionone at ten times the previous molar concentration. There is a slight reduction in the amount of matrix in the young hypertrophic cartilage, but otherwise no effect.

Table 3. *Action of derivatives of vitamin A and other compounds on isolated rat-liver lysosomes*

The particles, isolated at 15 000g from rat liver, were suspended in 0.25M-sucrose at a concentration equivalent to 1 g. of original tissue/5 ml. The compounds, dissolved in ethanol, were added to the sucrose suspensions which were then incubated for 40 min. at 37°. At the end of the incubation period the suspensions were centrifuged at 15 000g for 20 min. and the released enzyme was measured in the supernatant fractions. The results are expressed as a percentage of the activity released by 0.7 μ mole of vitamin A (alcohol)/ml. The vitamin A acid (sodium salt) was tested in 0.02M-sodium acetate-0.25M-sucrose solution at pH 7.0.

	0.7 μ mole/ml.	3.5 μ moles/ml.	7.0 μ moles/ml.
Vitamin A (alcohol)	100	157	189
Vitamin A acid (Na salt)	7	62	—
Vitamin A acid (free)	92	—	—
Vitamin A acetate	11	22	—
Vitamin A palmitate	0	—	—
Vitamin A oxidized in air	25	—	—
Vitamin A epoxide	0	115	165
Hydrogenated vitamin A	0	17	76
Anhydrovitamin A	0	0	0
Isoprene	7	12	28
Citral	0	85	164
Geraniol	27	98	92
α -Ionone	21	92	110
β -Ionone	25	—	—
<i>p</i> -Cymene	20	—	98
Pinene	24	80	170
Phytol	7	3	3
Dodecanol	0	0	0
Octanol	0	—	—
Glycerol trioleate	—	—	0
Glycerol	—	—	3
Isobutanol	—	—	15
2-Methylbutan-2-ol	—	—	0
Ethanol	0	0	0
Toluene	—	—	9
Acetone	—	—	7
Cysteine	—	—	22
Calciferol	—	—	17

temperatures at which no action of the vitamin could be detected, but to release about the same amount of enzyme at 40° it was necessary to add approximately seven times as much detergent (by weight) as vitamin A. The action of the detergent was also extremely rapid, most of the proteolytic activity being released within a few seconds of its addition, unlike the action of the vitamin, which required incubation (Fig. 4).

Action of derivatives of vitamin A. The effect of vitamin A derivatives on isolated lysosomes is shown in Table 3. The concentration of vitamin used as the standard (0.7 μ mole/ml.) was sufficient to liberate approximately two-thirds of the bound enzyme under the conditions of assay in these experiments. At this concentration, only vitamin A acid had an effect comparable with that of vitamin A (alcohol); the water-soluble sodium salt of vitamin A acid in the buffered medium was not active.

Oxidation of the vitamin by bubbling a stream of oxygen through an alcoholic solution reduced the activity by approximately 75%. This reduction in activity was related to changes in the ultraviolet absorption which indicated a 70% loss of the

vitamin. Inactivation of the vitamin by oxidation was confirmed by studies with the epoxide, which was ineffective at the standard concentration; at five times this concentration the activity was still less than that of the vitamin.

Hydrogenated vitamin A had very little effect on the isolated lysosomes. Anhydrovitamin A showed no activity at any of the concentrations tested, but this may have been partly due to the rapidity with which this compound polymerized when exposed to air.

Action of terpenes and related compounds. At the standard concentration most of the terpenes tested had about one-fifth of the activity of vitamin A, but isoprene and citral were ineffective. At the higher concentrations, however, all of these compounds, except isoprene, were active, though in varying degrees.

Action of long-chain alcohols and other compounds. The C₂₀-alcohol phytol resembles the basic skeleton of the vitamin A molecule, but lacks the conjugated double bonds that are characteristic of vitamin A and the terpenes. It is possible, at least in theory, to derive vitamin A from phytol by dehydrogenation and ring closure. In spite of this similarity in

structure, however, phytol, when tested in seven experiments on various occasions, exhibited only minimal activity against the lysosomes, even at the highest concentration. All the other alcohols tested, from dodecanol down to ethanol, had little effect.

DISCUSSION

Vitamin A appears to act as specifically on cartilage in organ culture as it does in reversing hypovitaminosis-A changes in animals [see Moore (1957) for review]. The concentration of vitamin A (alcohol) (10 i.u./ml.) used as the standard dose in the culture experiments is comparable with that found in the plasma of hypervitaminotic patients; in such cases Gribetz, Silverman & Sobel (1951) found levels of 20–30 i.u./ml., of which as much as 70% was present as the free alcohol. Our results have shown that at this concentration relatively small modifications of the vitamin A molecule can completely abolish its action on chick-limb-bone rudiments in culture. Increasing the dose tenfold, however, gives a statistically significant inhibition of growth by both the hydrogenated vitamin and β -ionone; at this dose these compounds may also have a very slight histological effect on the matrix of the rudiments.

The use of these high doses in the organ-culture experiments was suggested by the results obtained with the isolated lysosomes. In the latter system the specificity of the vitamin was high at a concentration that released approximately two-thirds of the enzymic activity; at this concentration anhydrovitamin A, hydrogenated vitamin A and vitamin A epoxide were ineffective, as also was isoprene, and the terpenes gave only about 25% of the activity of the vitamin. A five- or ten-fold increase in the dose, however, diminished the specificity (see Table 3). It would appear that, both in culture and on the isolated lysosomes, vitamin A (alcohol) is active at concentrations at which none of the other compounds tested produce much effect.

Vitamin A acid has recently been shown to cause marked resorption in the skeleton of young rats (Thompson & Pitt, 1961), and we have found it to be at least as potent as the free alcohol on cartilage in culture. In the lysosome system, too, it is about as active as the free alcohol, provided that it is added to the sucrose suspension as an ethanolic solution. If, however, the water-soluble sodium salt is used at neutral pH, very little activity is found. Thus this derivative acts only when it is in a lipid-soluble form, presumably owing to the 'lipoprotein' structure of the lysosomal membrane. The activity of the vitamin A derivatives cannot be explained solely in terms of lipid-solubility, however, since both vitamin A

acetate and palmitate had a negligible effect (see Table 3).

The non-ionic detergent Brij 35 probably acts in a different manner from the vitamin, since its effect is far more rapid and is not altered greatly by temperature (Fig. 4). It is also significant that, weight for weight, only about one-seventh of the amount of vitamin is required to give approximately the same amount of release as that given by the detergent; in fact the amount of vitamin is probably much smaller than this, since a considerable proportion is lost as an insoluble precipitate when the ethanolic solution is added to the sucrose suspension. It is thus unlikely that the action of the vitamin resembles that of this surface-active agent.

The degree of specificity reported in this work both in organ culture and on the isolated lysosomes lends further support to the hypothesis (Dingle, 1961; Lucy *et al.* 1961) that the effect of excess of vitamin A is due at least in part to an action on the intracellular particles.

SUMMARY

1. A number of compounds, including derivatives of vitamin A, terpenes and alcohols, have been examined for ability to reproduce the effects of vitamin A on embryonic chick-limb cartilage in culture and on the release of cathepsin from isolated lysosomes.

2. In the organ cultures at a concentration of 0.01 μ mole/ml. none of the compounds examined, except vitamin A acid, had a significant effect on the explants; the effects of this derivative were indistinguishable from those produced by vitamin A. At a concentration of 0.1 μ mole/ml. hydrogenated vitamin A and β -ionone slightly reduced the growth in length of the explants, but caused little histological change.

3. Approximately two-thirds of the bound protease was released from isolated rat-liver lysosomes by 0.7 μ mole of vitamin A/ml. Of the compounds studied at this concentration and under the same conditions, only vitamin A acid had an effect comparable with that of the vitamin; other derivatives of vitamin A were inactive, and most of the terpenes had about 25% of the activity of the vitamin. At higher concentrations the specificity of the vitamin was reduced.

4. The results indicate that the action of vitamin A is highly specific on cartilage in culture and also at low concentrations on the isolated lysosomes.

The authors are indebted to Professor R. A. Morton, F.R.S., for helpful information; they express their grateful thanks to Mr L. J. King and Miss P. A. Carter for their skilful technical assistance and to Mr M. Applin for preparing the photographs. The work was supported by the Medical Research Council and the British Empire Cancer Campaign.

REFERENCES

- Dingle, J. T. (1961). *Biochem. J.* **79**, 509.
 Dingle, J. T., Lucy, J. A. & Fell, H. B. (1961). *Biochem. J.* **79**, 497.
 Edisburg, J. R., Gillam, A. E., Heilbron, I. M. & Morton, R. A. (1932). *Biochem. J.* **26**, 1164.
 Fell, H. B. & Mellanby, (1952). *J. Physiol.* **116**, 320.
 Fell, H. B. & Thomas, L. (1960). *J. exp. Med.* **111**, 719.
 Gribetz, D., Silverman, S. H. & Sobel, A. E. (1951). *Pediatrics*, **7**, 372.
 Karrer, P. & Jucker, E. (1947). *Helv. chim. acta*, **30**, 559.
 Lucy, J. A., Dingle, J. T. & Fell, H. B. (1961). *Biochem. J.* **79**, 500.
 Moore, T. (1957). *Vitamin A*. Amsterdam: Elsevier Publishing Co.
 Shantz, E. M., Cawley, J. D. & Embval, N. D. (1943). *J. Amer. chem. Soc.* **65**, 901.
 Thomas, L., McLusky, R. T., Potter, J. L. & Weissmann, G. (1960). *J. exp. Med.* **111**, 705.
 Thompson, J. N. & Pitt, G. A. J. (1961). *Biochem. J.* **79**, 33p.
 Tiemann, F. (1898). *Ber. deutsch. chem. Ges.* **31**, 867.
 Vinet, A. & Meunier, P. (1947). *Bull. Soc. Chim. biol.* **29**, 25.

Biochem. J. (1962) **83**, 69

Incorporation of Radioactive Amino Acids into Serum Albumin by Isolated Rat-Liver Ribosomes

By A. KORNER

Department of Biochemistry, University of Cambridge

(Received 15 September 1961)

It is generally accepted that protein synthesis occurs in the ribosome part of the microsomes of cells and, indeed, that much of the intracellular synthesis of protein takes place in these ribonucleo-protein particles (see Hoagland, 1960, for review). Ribosomes can now be isolated from rat liver (as well as from other tissues) in a state in which they will incorporate amino acids into protein (Korner, 1959*a*, 1960*a*, 1961*a*; Takamami, 1960; Rendi & Hultin, 1960). Amino acid incorporation into protein is not, however, synonymous with synthesis of protein and, in the hope of providing some support for the idea that isolated ribosomes can synthesize protein, it was decided to attempt to demonstrate the radioactive labelling of a specific protein by isolated rat-liver ribosomes.

Serum albumin is synthesized in the liver in considerable quantity (Miller, Bly, Watson & Bale, 1951; Jensen & Tarver, 1956; Gordon & Humphrey, 1960). The net synthesis of albumin has been studied with perfused rat liver (Miller *et al.* 1951), with slices of chick liver (Peters & Anfinsen, 1950) and with slices of rat liver (Campbell & Stone, 1957). Ogata, Hirokawa & Omori (1960) and Campbell, Greengard & Kernot (1960) showed that serum albumin becomes radioactively labelled when rat-liver microsomes are incubated *in vitro* with radioactive amino acids, although in these experiments net synthesis of albumin could not be demonstrated. Further, serum albumin can be isolated by chemical and immunological means, and therefore the ability of isolated ribosomes to incorporate radioactive

amino acids into this protein was studied. The successful result has been reported briefly (Korner, 1960*b*), and this paper gives fuller details of the methods used and the results achieved. Lingrel & Webster (1961) have reported the labelling of albumin by isolated microsomes and ribosomes, and Decken & Campbell (1962) have obtained similar results with ribosomes prepared by the method of Rendi & Hultin (1960).

METHODS

Radioactive amino acids. Generally labelled L-leucine (7.1 mc/m-mole) and DL-[1-¹⁴C]leucine (5.4 mc/m-mole) were obtained from The Radiochemical Centre, Amersham, Bucks.

Chemicals. The sodium salts of ATP and GTP were obtained from Sigma Chemical Co. The sodium salt of creatine phosphate was made by the method of Ennor & Stocken (1948) and creatine phosphokinase was prepared by the method of Kuby, Noda & Lardy (1954).

Preparation of albumin. Rat-serum albumin was separated from rat serum by the method of Korner & Debro (1956) and Debro, Tarver & Korner (1957). Rat-serum proteins were precipitated by adding to the serum an equal volume of cold 10% (w/v) trichloroacetic acid. The proteins were spun down and the albumin was removed from the precipitate by three extractions each with a volume of cold 96% (v/v) ethanol-1% (w/v) trichloroacetic acid equal to the original volume of serum. Material that did not dissolve was removed by centrifuging, the alcohol-acid extract was dialysed at 4° against several changes of water to remove alcohol and acid, and the albumin was freeze-dried. Chicken-serum albumin was prepared similarly.