SUMMARY

1. Compounds containing *N*-terminal residues have been shown to be present in preparations of soluble collagen which account for most if not all the reaction of fluorodinitrobenzene with α -amino groups.

2. The significance of the residues has been discussed in relation to the known molecular weight of collagen as determined by physical means. The terminal residues are thought to be due to non-protein nitrogen firmly attached to collagen and not to belong to collagen as true *N*-terminal residues in the usually accepted meaning of the term '*N*-terminal residue'.

3. The thermal conversion of collagen into gelatin has been shown to take place as a physical dissociation with the minimal rupture of peptide bonds.

We wish to thank Imperial Chemical Industries Ltd. for a Research Fellowship (to F.S.S.) and the Department of Scientific and Industrial Research for a Senior O.E.E.C. Visiting Fellowship which enabled one of us (F.S.S.) to visit the Carlsberg Laboratory. We are very grateful to

Professor Ottesen and his colleagues for their hospitality at the Carlsberg Institute.

Our thanks are also due to the British Gelatine and Glue Research Association for financial support, and to Dr A. Courts for the gift of dinitrophenyl-amino acids.

REFERENCES

- Bowes, J. H., Elliott, R. G. & Moss, J. A. (1957). In *Connective Tissue Symposium*, p. 264. Ed. by Tunbridge, R. E. *et al.* Oxford: Blackwell Press.
- Courts, A. (1954). *Biochem. J.* **58**, 70, 74.
- Courts, A. (1959). *Biochem. J.* **73**, 596.
- Fraenkel-Conrat, H. & Singer, B. (1954). *J. Amer. chem. Soc.* **76**, 180.
- Heyns, K. & Legler, G. (1957). *Hoppe-Seyl. Z.* **306**, 165.
- Heyns, K. & Legler, G. (1958). In *Recent Advances in Gelatine and Glue Research*, p. 186. Ed. by Stainsby, G. London: Pergamon Press.
- Orekhovitch, V. N. & Shpikiter, V. O. (1957). In *Connective Tissue Symposium*, p. 281. Ed. by Tunbridge, R. E. *et al.* Oxford: Blackwell Press.
- Phillips, D. M. P. (1955). *Biochem. J.* **60**, 403.
- Sanger, F. (1945). *Biochem. J.* **39**, 507.
- Steven, F. S. & Tristram, G. R. (1962). *Biochem. J.* **83**, 240.
- Ward, A. G. (1960). *J. Soc. Leath. Tr. Chem.* **44**, 505.

Biochem. J. (1962) **83**, 248

The Degradative Metabolism of L-Cysteine and L-Cystine *in vitro* by Liver in Cystinosis

By A. D. PATRICK

Institute of Child Health, Hospital for Sick Children, Great Ormond Street, London, W.C. 1

(Received 13 November 1961)

Since Abderhalden (1903) first detected widespread storage of crystalline cystine in the tissues of an infant at autopsy, and noted the familial incidence of the disorder, an increasing number of cases have been reported in which cystinosis has been diagnosed by post- and ante-mortem examination (Lignac, 1924; McCune, Mason & Clarke, 1943; Worthen & Good, 1958). The associated clinical features of this disease include vitamin D-resistant rickets, glucosuria, generalized hyper-amino aciduria, and acidosis, symptoms which characterize a group of diseases based on the failure of reabsorptive and base-conserving function of the renal tubules, and often referred to as the Fanconi syndrome (Fanconi, 1936). Several hypotheses seeking to explain cystine storage and to relate this to the Fanconi syndrome have been put forward. Bickel *et al.* (1953) reported increased concentrations of a number of plasma amino acids in cystinosis, and suggested that these resulted

from a generalized failure of their normal incorporation into proteins. The intracellular deposition of the poorly soluble cystine and the renal anomalies were considered to be manifestations of this primary defect. However, essentially normal plasma amino acid concentrations have been found by several authors using chromatographic methods (Dent, 1947; Evered, 1956; Brigham, Stein & Moore, 1960), and these are now generally accepted as proof that the amino aciduria is of renal origin.

A second suggestion proposes a specific enzymic defect in cysteine-cystine metabolism, leading to storage of the resultant excess of cystine in the tissues and consequent nephrotoxic effects. The reduction of cystine to cysteine by cystine reductase has been considered the most probable site of such a defect (see Worthen & Good, 1958). Since no investigation of these latter proposals had been reported, a comparison of cystinotic- and normal-

liver preparations, with respect to the degradation of cysteine and cystine, was undertaken. Particular attention has been given to the enzymic reduction of cystine.

Recently, deficiencies in the reduction of cystine by two different enzyme systems of blood have been reported in cases of cystinosis. Worthen & Good (1961) concluded that cystine reductase (NADH_2 -L-cystine oxidoreductase, EC 1.6.4.1) was deficient in two cases examined, whereas Seegmiller & Howell (1961) reported cases of low activity of a system catalysing the transfer of hydrogen from reduced glutathione to cystine. Neither of these discordant results is supported by the present study.

MATERIALS

Autopsy specimens. The following cases of cystinosis were studied: Case 1, female, aged 11 yr.; Case 2, male, aged 8 yr.; Case 3, female, aged 7 yr. Tissues for comparison were obtained from two cases of congenital heart disease in which no primary metabolic abnormality had been apparent. These are referred to as Normal 1 (female, aged 8 yr.) and Normal 2 (male, aged 6 yr.). Autopsies were performed within 2 hr. after death. Portions of liver and kidney were placed in polyethylene bags and frozen immediately in crushed solid CO_2 . Thereafter, specimens were stored at -25° . Only small samples of cystinotic kidney were obtained, their use being limited to tests for cystine reduction.

Chemicals. The following chemicals were obtained from the sources indicated: NADH_2 (disodium salt), NADP (monosodium salt), pyridoxal phosphate and α -oxoglutaric acid, from the Sigma Chemical Co.; glucose 6-phosphate (disodium salt), NADPH_2 (monosodium salt) and L-glutamic acid, from the British Drug Houses Ltd.; L-cystine, L-cysteine, L-cysteic acid, reduced glutathione (GSH) and NAD from L. Light and Co. Ltd.; L-cysteine-sulphinic acid, from the California Corp. for Biochemical Research, Los Angeles. All other reagents were of AnalaR grade where obtainable.

METHODS

Preparation of acetone-dried liver powder

Chopped frozen liver (10 g.) was homogenized by grinding with sand and water (20 ml.) at 0° . After standing in the cold for 1 hr. with occasional stirring, the suspension was filtered through surgical gauze, and acetone (2 vol.) at -5° was added to the filtrate with continuous stirring. The precipitate was collected by rapid filtration in the cold, and washed with cold acetone and finally with ether. The powders were stored *in vacuo* over magnesium perchlorate and liquid paraffin.

Tissue homogenates and slices

Homogenates (usually 20%) were prepared in the appropriate ice-cold media with a Potter-type glass homogenizer for 1 min. at about 700 rev./min. The layer of fat obtained from certain specimens after centrifuging at 800g for 15 min. at 2° could be removed almost completely by screening through several layers of surgical gauze.

Slices were cut free-hand from frozen blocks, and suspended in the cold buffer to be used in the particular assay. Suitable slices, selected for translucency, were collected into bundles and blotted before transfer to incubation flasks. In slice experiments, tissue-dry weights from individual flasks were determined at the end of the incubation period by removal of the total contents (washing with water if necessary to remove small fragments), to which was then added an equal volume of 10% (w/v) trichloroacetic acid. After centrifuging, and washing with water and with acetone, the residues were dried to constant weight at 100° . Homogenate (0.5 ml.) was precipitated with trichloroacetic acid, and washed and dried similarly. The protein content of other enzyme solutions was measured spectrophotometrically (Warburg & Christian, 1941).

Enzyme preparations

Glucose 6-phosphate dehydrogenase (D-glucose 6-phosphate-NADP oxidoreductase, EC 1.1.1.49) was prepared from yeast by the method of Kornberg (1950), and assayed according to Kornberg & Horecker (1955). The final solution had an activity of 30 units/ml. Yeast glutathione reductase [NAD(P)H_2 -glutathione oxidoreductase, EC 1.6.4.2] was prepared and assayed according to Racker (1955). The preparation used was that obtained at the first ethanol-precipitation stage, and possessed an activity of approximately 20 000 units/ml. Oxidized glutathione (GSSG) used in the reductase assay was prepared by the method of Rall & Lehninger (1952).

Enzyme assays

Cystine reductase. The activity of cystine reductase in acetone-dried liver powders was estimated by the initial decrease in NADH_2 extinction at 340 m μ in the presence of cystine added as a saturated solution in either 0.02M-phosphate buffer, pH 6.5, or 0.1M-tris buffer, pH 8.2. Under these conditions the solubility of cystine at 22° was 11.6 and 43.6 mg./100 ml. in the phosphate and tris buffers respectively, as determined by the method of Sullivan, Hess & Howard (1942). An extract of acetone-dried powder (0.3 g.) was prepared in 3 ml. of the appropriate buffer, and dialysed overnight in the cold against the same buffer. Clear extracts were obtained by centrifuging for 30 min. at 6000g. Kidney extracts were assayed similarly. Cystinotic kidney, owing to its fibrous nature, could not be homogenized satisfactorily in the Potter homogenizer. Extracts were prepared by grinding 1 g. with cold 0.02M-phosphate buffer (pH 6.5, 3 ml.) and sand. After dialysis for 40 hr. against this buffer, clear extracts were obtained by centrifuging at 3000g for 20 min.

Alternative methods of estimation were used for liver homogenates, which were dialysed in cellophan tubing (5 mm. diam.) for 40 hr. in the cold against the buffer to be used in the assay. In the first method, homogenate in phosphate buffer, pH 6.5, was shaken anaerobically in Warburg flasks for 2 hr. with NADH_2 and solid cystine, and the increase of thiol groups in samples deproteinized with 10% trichloroacetic acid (0.5 vol.) was determined by titration with 5 mN-iodine. The starch-iodine end point was detected on an EEL Micro-titrator (Evans Electro-selenium Ltd.). In the second method, homogenate was incubated in tris buffer, pH 8.5, containing NADH_2 and cystine, and NADH_2 extinctions at 340 m μ were measured

at zero time and after 10 min. with samples which had been deproteinized by the addition of 95% ethanol (1 vol.) followed by 10% (w/v) sodium sulphate (0.1 vol.). A 5 mm-cystine soln. used in these tests was prepared by dissolving L-cystine (30 mg. of free base) in N-sodium hydroxide (1 ml.) and adding 0.05M-tris, pH 8.5 (5 ml.); the solution was readjusted to pH 8.5 with N-hydrochloric acid, and diluted to 25 ml. with water. Solutions prepared in this way were stable for at least 4 hr.

System reducing cystine in the presence of reduced glutathione. Liver homogenate (20%) in 1.15% (w/v) potassium chloride was dialysed for 24 hr. in the cold against 0.1M-tris buffer, pH 7.6. The incubation medium, added to Warburg flasks, contained in a final volume of 5 ml.: 0.1M-tris, pH 7.6 (1 ml.); 0.5M-glucose 6-phosphate (0.2 ml.); 0.1M-magnesium chloride (0.2 ml.); glucose 6-phosphate dehydrogenase (10 units); 5 mM-NADP (0.1 ml.); 10 mM-GSH (0.2 ml.); mM-pyridoxal phosphate (0.1 ml.); homogenate (0.5 ml.); solid cystine (5 mg.). Side arms contained 0.2 ml. of a zinc acetate reagent (see below) to trap any hydrogen sulphide liberated. At zero time, a sample (2 ml.) of medium was deproteinized with 10% trichloroacetic acid (1 ml.), and the clear supernatant (2 ml.) obtained after centrifuging was titrated with 5 mM-iodine. Incubation flasks were flushed with nitrogen, and, after shaking for 2 hr. at 37°, the iodine titration for thiol groups was repeated. Trapped hydrogen sulphide was estimated as in the assay of cysteine desulphydrase.

Cysteine desulphydrase. The assay medium, consisting of 0.067M-phosphate buffer, pH 7.4 (1 ml.), liver homogenate, 20% in 0.9% sodium chloride, (1 ml.) and mM-pyridoxal phosphate (0.1 ml.), was added to the main compartments of Warburg flasks. Side arms contained 0.1M-cysteine, pH 7.4 (0.2 ml.), and centre wells a reagent (0.2 ml.) consisting of hydrated zinc acetate (6.0 g.), hydrated sodium acetate (1.7 g.) and sodium chloride (0.005 g.) dissolved in water (100 ml.). Flasks were flushed with nitrogen, and shaken for 90 min. at 37°. *p*-Amino-*NN*-dimethylaniline [0.3 ml. of a 0.05% soln. in 20% (w/v) sulphuric acid] followed by 1 drop of 10% (w/v) ferric chloride in 0.1N-hydrochloric acid was added to the centre well, and the resulting methylene blue solution, after colour development for 1 hr., was washed out and diluted to 25 ml. with water. Extinctions at 630 m μ were measured in 1 cm. cuvettes. An aq. hydrogen sulphide soln., standardized with sodium arsenite, was used to prepare a standard curve for the range 10–100 μ g. of hydrogen sulphide.

Desulphurization of cystine. This system refers to the NADH₂-dependent reduction of cystine to cysteine, followed by the liberation of hydrogen sulphide by cysteine desulphydrase [L-cysteine hydrogen sulphide-lyase (deaminating), EC 4.4.1.1]. The hydrogen sulphide, estimated as described above, gave a measure of the overall reaction. Alcohol dehydrogenase (alcohol-NAD oxido-reductase, EC 1.1.1.1), contained in the liver preparations, was used to regenerate NADH₂. A 10% liver homogenate was prepared in 0.9% sodium chloride and dialysed overnight in the cold against 0.067M-phosphate buffer, pH 7.4.

Decarboxylation of L-cysteate and L-cysteinesulphinate. This was determined for liver extracts according to the method of Hope (1955).

Oxidation of L-cysteine and L-cystine to inorganic sulphate. Liver slices were suspended in 5 ml. of bicarbonate-saline (Krebs & Henseleit, 1932), in which magnesium sulphate

was replaced by magnesium chloride (Pirie, 1934). The Warburg flasks were flushed continuously with O₂ + CO₂ (19:1). At zero time 0.4M-cysteine (0.2 ml.), pH 7.4, was added from the side arms, and the flasks were shaken at 37°. When cystine was used as substrate this was added as the solid (5 mg.) directly to the main compartments at the beginning of the experiment. After 2 hr., inorganic sulphate in the deproteinized medium (i.e. after addition of 5 ml. of 10% trichloroacetic acid) was determined by a modification of the method of Cuthbertson and Tompsett (Maw, 1954). In later experiments the method of Dodgson (1961) was found to give more reproducible results.

Oxidation of L-cysteinesulphinate. The metabolism of L-cysteinesulphinate by liver slices in the presence of α -oxoglutarate was measured in the following medium: 0.1M-phosphate, pH 7.4 (2 ml.); slices (approx. 100 mg. fresh wt.); 0.376M-L-cysteinesulphinate, pH 7.4 (0.2 ml.); water (2 ml.); added to Warburg flasks. The gas phase was air, and centre wells contained 10% (w/v) potassium hydroxide (0.2 ml.). After equilibration for 10 min. at 37°, 0.25M- α -oxoglutarate, pH 7.4 (0.2 ml.), was added from the side arm, and oxygen uptake followed for 2 hr. The flask contents were treated with cold 10% trichloroacetic acid (5 ml.), and, after centrifuging, pyruvate was estimated by the method of Friedemann & Haugen (1943). Inorganic sulphate was estimated according to Dodgson (1961). Glutamate was estimated chromatographically. Supernatant (2 ml.) was evaporated to dryness, and the residue was extracted with water (0.5 ml., or 0.1 ml. in experiments in which either α -oxoglutarate or cysteinesulphinate was omitted from the medium). Samples (30 μ l.) were chromatographed in butan-1-ol-acetic acid-water (12:3:5, by vol.) on Whatman no. 2 paper. Marker solutions containing 10, 20 and 30 μ g. of glutamate were run simultaneously. After spraying with ninhydrin, the approximate concentrations of glutamate in test solutions were estimated by visual comparison. This was found to give estimates in fair agreement with those obtained by spot elution and colour measurement.

Glutathione reductase. This was assayed essentially according to the method of Racker (1955). Acetone-dried liver powder (50 mg.) was extracted with cold water (1 ml.) for 1 hr. After centrifuging, the solution was dialysed overnight in the cold against 0.2M-phosphate, pH 7.6, and recentrifuged.

All spectrophotometric measurements were made in 1 cm. silica cells with a Unicam SP. 500 spectrophotometer (Unicam Instruments Ltd.).

RESULTS

The reduction of cystine by cystine reductase. A comparison of the activities of cystine reductase in liver homogenates and kidney extracts, and in acetone-dried liver powders of the cystinotic and normal cases, is shown in Table 1. Tests were made after the tissue samples had been stored for 2 months at -25°, except for cystinosis Case 1 (Expt. D), which had been stored for 3 months. The activities of the cystinotic- and normal-tissue preparations were in good general agreement in each experimental series, with different conditions

Table 1. *Cystine-reductase activity of liver and kidney preparations*

Silica cuvettes of 1 cm. light path contained 2.5 ml. of a saturated soln. of cystine in 0.02M-phosphate buffer, pH 6.5 (Expt. A), or in 0.1M-tris buffer, pH 8.2 (Expt. B); acetone-dried liver-powder extract (0.5 ml.) (see Methods section); and 1.4 mM-NADH₂ (0.1 ml.). Extinctions at 340 m μ were measured at 1 min. intervals for 10 min. at room temperature (22°). Blank assays contained buffer instead of buffer-cystine soln. Kidney extract (0.5 ml.) (see Methods section) was assayed similarly in 0.02M-phosphate, pH 6.5 (Expt. C). In Expt. D, incubation media contained liver homogenate (1 ml.); 0.05M-tris buffer, pH 8.5 (1 ml.); 5 mM-cystine, pH 8.5 (2 ml.), or water; 3 mM-NADH₂ (0.2 ml.); in 5 ml. final volume. Incubation temperature was 30°. Extinctions at 340 m μ of samples deproteinized at zero time and after 10 min. (see Methods section) were measured. In Expt. E, media contained liver homogenate (2 ml.); 0.02M-phosphate, pH 6.5 (2 ml.); 14 mM-NADH₂ (0.1 ml.); water (1 ml.); solid cystine (5 mg.); added to Warburg flasks. Flasks were shaken anaerobically at 37° for 2 hr., and the increase of thiol groups was determined by titration with 5 mM-iodine.

Case	Activity ($-\Delta E_{340}$ /mg. of protein/hr.): Acetone-dried-liver- powders preparation		Activity ($-\Delta E_{340}$ /mg. dry wt./hr.)		Thiol groups formed (μ moles/g. dry wt./hr.). Liver-homogenate preparation Expt. E
	Expt. A	Expt. B	Kidney- extract preparation Expt. C	Liver- homogenate preparation Expt. D	
Cystinosis 1	0.052	0.055	0.019	0.024	25.5
Cystinosis 2	0.039	0.031	0.021	0.029	33.0
Cystinosis 3	0.038	—	—	0.024	29.0
Normal 1	0.047	0.050	0.024	0.026	31.5
Normal 2	0.044	—	—	0.021	25.5

and methods of assay. It was found that the cystinotic homogenates released thiol groups in the absence of added cystine when such preparations were dialysed for only 24 hr. This activity, which was completely dependent on the addition of NADH₂, in some homogenate preparations amounted to 60 % of that obtained by full stimulation with added cystine. Dialysis for 40 hr. in cellophan tubing (5 mm. diam.) reduced this endogenous activity almost to zero without causing undue loss (not more than 10 %) of overall cystine-stimulated activity. It appears certain from these experiments that the high endogenous activity was due to stored cystine, low residual concentrations of which would be sufficient to saturate the enzyme in the most active of the preparations tested. Fig. 1 shows the effect of varying concentrations of such a preparation in assay media containing the lowest concentrations of cystine, i.e. when cystine was added as a saturated solution, pH 6.5. The specific activities given in Table 1 were confined to determinations carried out in the linear range of the activity curves. Fig. 1 also illustrates the specificity of cystine reductase for NADH₂, an equivalent amount of NADPH₂ failing to act as replacement, and shows the complete loss of activity of a boiled preparation.

Evidence that cystine-reductase activity is normal in cystinotic liver (stored frozen for 14 weeks) was also obtained from determinations of the desulphurization of cystine by homogenates. The liberation of hydrogen sulphide from cystine was completely dependent on additions of NAD and, with one exception (cystinosis Case 3, in

Fig. 1. Cystine-reductase activity of acetone-dried liver powder (cystinosis Case 3). Varying amounts of enzyme were assayed under the experimental conditions of Expt. A, Table 1. ○, Complete system; ●, cystine omitted; △, enzyme soln. boiled for 2 min.; □, NADH₂ replaced by NADPH₂; ▲, NADH₂ omitted.

which a low endogenous activity was observed), a NAD-linked oxidizable substrate. In the experiments listed in Table 2, ethanol was used as oxidizable substrate for the regeneration of NADH₂ by alcohol dehydrogenase of the liver homogenates. The results are interpreted as showing a normal reduction of cystine to cysteine by cystine reductase of cystinotic liver, followed by the liberation of hydrogen sulphide from cysteine by cysteine desulphhydrase. The slight activity of

the cystinotic samples observed in the tests from which cystine was omitted was attributed to residual stored tissue cystine.

The reduction of cystine by reduced glutathione. In the presence of reduced glutathione, cystine is converted into cysteine. Liver homogenates stimulate this reaction, and further metabolize part of the cysteine formed, mainly through the desulphhydrase reaction under anaerobic conditions. With low concentrations of glutathione the overall system requires glutathione reductase, coupled with a reaction which regenerates NADPH₂. In the experiments on liver homogenates, this latter requirement was met by the addition of the glucose 6-phosphate-dehydrogenase system. Homogenates possessed sufficient glutathione-reductase activity to maintain maximum concentrations of reduced glutathione throughout the incubation period. A comparison of the activities of glutathione reductase in cystinotic and normal liver shows good agreement of values independent of the period of storage of the frozen tissues (Table 3).

Table 4 shows the requirements of the cystine-reduced glutathione system for cystine, reduced glutathione, a NADP-linked oxidizable substrate (glucose 6-phosphate) and glutathione reductase. To account for the extent of spontaneous non-enzymic reduction of cystine by reduced glutathione in the overall reaction, it was necessary to employ a composite system in which a glutathione-reductase preparation was substituted for liver homogenate in these experiments. A comparable concentration of reduced glutathione was thereby

Table 2. *Desulphurization of L-cystine by liver homogenates*

Media contained 0.067M-phosphate, pH 7.4 (1.5 ml.); liver homogenate (0.5 ml.) (see Methods section); 3 mM-NAD (0.1 ml.); 2 mM-pyridoxal phosphate (0.1 ml.); ethanol (0.1 ml.); solid cystine (5 mg.); added to Warburg flasks. The trapping and estimation of H₂S were carried out as for the assay of cysteine desulphhydrase (see Methods section). Anaerobic incubation was at 37° for 4 hr., with constant shaking.

Case	Activity (μmoles of H ₂ S/g. dry wt. of liver homogenate/hr.)		
	Complete system	Ethanol omitted	Cystine omitted
Cystinosis 2	7.9	0	0.1
Cystinosis 3	11.8	2.2	0.3
Normal 1	7.4	0	0
Normal 2	8.2	0	0

Table 3. *Glutathione-reductase activity of liver*

Microcuvettes of 1 cm. light path contained: M-phosphate buffer, pH 7.6 (0.05 ml.); mM-NADPH₂ (0.05 ml.); 1% (w/v) serum albumin (0.05 ml.); extract of acetone-dried liver powder (0.05 ml.) (see Methods section); water (0.2 ml.); 30 mM-GSSG (0.05 ml.) or water. Extinctions at 340 mμ were measured at 30 sec. intervals for 5 min. at room temperature (22°).

Case	Period of tissue storage	Activity (-ΔE ₃₄₀ /mg. of protein/min.)
Cystinosis 1	14 months	0.106
Cystinosis 2	9 months	0.121
Cystinosis 3	3 weeks	0.110
Normal 1	11 months	0.105
Normal 2	6 months	0.088

Table 4. *Reduction of cystine in the presence of reduced glutathione*

A. For experimental details of the overall reaction (enzymic + spontaneous), see Methods section. B. Spontaneous reaction; the medium contained: 0.1M-tris buffer, pH 7.6 (1 ml.); 0.5M-glucose 6-phosphate (0.2 ml.); 0.1M-MgCl₂ (0.2 ml.); glucose 6-phosphate dehydrogenase (0.2 ml. ≡ 10 units); 5 mM-NADP (0.1 ml.); 10 mM-GSH (0.2 ml.); glutathione reductase (0.05 ml. ≡ 1000 units); solid cystine (5 mg.); in a final volume of 5 ml., contained in Warburg flasks. The experiment was otherwise performed as for (A). Results are expressed as changes occurring in 2 hr. at 37°.

A Case	Period of storage of liver	Difference in 5 mM-I ₂ required (ml./ml. of reaction medium)					Complete system		
		Complete system	Cystine omitted	Glucose 6-phosphate omitted	GSH omitted	Glutathione reductase omitted	H ₂ S* liberated (μmoles/ml.)	Thiol groups formed (μmoles/ml.)	Corrected† (μmoles/mg. dry wt. of liver homogenate/ml.)
Cystinosis 2	6 months	+1.01	0	+0.02	+0.03	—	0.18	5.20	0.66
Cystinosis 3	5 weeks	+0.82	+0.03	-0.08	+0.03	—	0.22	4.37	0.63
Normal 1	11 months	+0.31	-0.03	+0.04	+0.01	—	0.27	1.85	0.03
Normal 2	3 months	+0.86	+0.02	+0.03	+0.02	—	0.31	4.60	0.61
B Spontaneous reaction	—	+0.34	+0.01	+0.02	0	0	0	1.70‡	—

* No H₂S was liberated from incomplete media.

† Calculated after subtraction of mean value for thiol groups formed in the spontaneous reaction.

‡ Range 1.43-1.95 (mean 1.71, from four estimations).

maintained in assay media that were otherwise identical, so that the difference in cystine reduction in the two systems gave a measure of the stimulating effect of liver homogenate. Results for the spontaneous reaction (which coincided approximately with the mean values for four separate determinations) are given in Table 4B. Subtraction of this value for thiol groups formed from those obtained for the overall reaction in homogenates (Table 4A) gave good agreement for the enzymic component of cystinotic- and normal-liver specimens which had been stored frozen for up to 6 months. With further storage considerable loss of activity may occur, a 95% loss being observed for the normal sample stored for 11 months.

In view of the normal capacity of cystinotic liver for glutathione-cysteine transhydrogenation and glutathione reduction *in vitro*, it was of some interest to determine the concentrations of reduced glutathione in liver specimens, since a failure to maintain this (caused indirectly, e.g. by a deficiency of NADPH₂) could impair the subsequent reduction of cystine. Estimations were made on sulphosalicylic acid extracts by the yeast-glyoxalase method of Woodward (1935). The mean values obtained were 1.5 mg. and 1.3 mg. of reduced glutathione/g. wet wt. of liver from the cystinosis and normal cases respectively.

The oxidation of cysteine and cystine to inorganic sulphate. Slices of cystinotic liver were found to possess a normal capacity for the complete oxidation of cysteine, and probably of cystine, to inorganic sulphate (Table 5). In all cases the liver specimens had been stored frozen for 2 months before testing. The amount of endogenous sulphate derived from normal specimens was in the range reported by Maw (1954) and Medes (1939) for rat-liver slices, and is ascribed to diffusion of existing intracellular sulphate rather than to a metabolic

production such as is seen with the stimulation effected by the addition of cysteine or cystine. The high concentrations of endogenous sulphate produced by cystinotic specimens, and the apparent failure of cystine to stimulate this production, may be explained by a maximally stimulated formation of sulphate already occurring in the presence of stored cystine. The ready availability of intracellular cystine in these specimens could also account for the endogenous production of sulphate in greater amount than that derived from cystine additions to normal liver.

Further evidence of the normal oxidation of cystine in cystinosis was derived from the determination of urinary inorganic sulphate in two cases. The values obtained were 13.0 and 11.3 mg. of inorganic sulphate S/kg. body wt./24 hr., compared with 9.4 and 14.6 mg. of sulphate S/kg./24 hr. for two normal children of about the same body weight.

The metabolism of L-cysteinesulphinate. A balance study of cysteinesulphinate metabolism in liver slices is summarized in Table 6. Liver specimens

Table 5. *Oxidation of L-cysteine and L-cystine to inorganic sulphate by liver slices*

Experimental details are given in the Methods section. Inorganic sulphate production after 2 hr. at 37° was measured.

Substrate ... Case	Activity (μg. of sulphate S/100 mg. dry wt. of liver slices)		
	Cysteine	Cystine	None
Cystinosis 1	190	72	60
Cystinosis 2	151	52	54
Cystinosis 3	217	81	76
Normal 1	129	43	21
Normal 2	107	35	16

Table 6. *Metabolism of L-cysteinesulphinate in liver slices*

Experimental details are given in the Methods section. Values refer to changes occurring in 2 hr. at 37°.

Case	Substance omitted	O ₂ uptake in excess of endogenous level (μg.-atoms of O ₂ /g. dry wt. of liver slices)	Substances formed (μmoles/g. dry wt. of liver slices)		
			Pyruvate*	Sulphate	Glutamate
Cystinosis 2	None	117.0	74.5	120.0	61.0
	α-Oxoglutarate	14.7	1.7	39.5	<3.0
	Cysteinesulphinate	0	39.1	13.0	<3.0
Normal 1	None	102.5	76.5	129.5	54.0
	α-Oxoglutarate	25.5	8.8	44.8	<3.0
	Cysteinesulphinate	9.5	33.5	9.3	<3.0

* Values for pyruvate were not expected to be accurate in the presence of α-oxoglutarate in the concentrations used. Values recorded for experiments in which cysteinesulphinate was omitted were most probably due entirely to α-oxoglutarate rather than to pyruvate. However, the experiments serve to show close agreement in the net formation of pyruvate in the complete systems.

had been stored for 18 weeks. Oxidation of cysteinesulphinate to sulphate was stimulated by α -oxoglutarate, and the appearance of glutamate required the presence of cysteinesulphinate, in keeping with a reaction mechanism requiring an initial transamination with the formation of β -sulphinylpyruvate. Stoichiometric balances were not observed, but the overall similarity of results obtained with cystinotic- and normal-liver specimens was striking in view of interfering reactions, e.g. transaminations involving pyruvate and glutamate, that may occur in liver slices.

Decarboxylation of L-cysteate and L-cysteinesulphinate. Since it has been reported that taurine may be absent from urine in cases of cystinosis

(Worthen & Good, 1958), it was of interest to determine the activities of L-cysteate decarboxylase and L-cysteinesulphinate decarboxylase (L-cysteinesulphinate carboxy-lyase, EC 4.1.1.29) in the liver specimens. Results of these experiments (Table 7) indicate the normal ability of cystinotic liver to decarboxylate these substrates, of which cysteinesulphinate shows the higher decarboxylation rate.

Anaerobic reactions of cysteine. Indirect evidence of the normal activity of cysteine desulphydrase in cystinotic liver has been mentioned in the above experiments dealing with the desulphurization of cystine and with the glutathione-cysteine-transhydrogenase system. These results were confirmed by direct measurement of the cysteine desulphydrase activity of liver homogenates (Table 8).

A further anaerobic reaction of cysteine, for which no abnormality of cystinotic liver was apparent, was its transamination with α -oxoglutarate. The appearance of glutamate (followed by paper chromatography) in liver homogenates containing α -oxoglutarate was markedly stimulated by the addition of cysteine, or even more effectively by cysteic acid or cysteinesulphinic acid.

Table 7. *Decarboxylation of L-cysteate and L-cysteinesulphinate by liver extracts*

Liver extract was prepared by grinding frozen tissue with sand and 0.1 M-sodium phosphate buffer, pH 6.5, and centrifuging at 3000g for 30 min. at 2°. Reaction media contained: liver extract (1 ml. $\equiv$ approx. 0.5 g. of fresh tissue); mm-pyridoxal phosphate (0.1 ml.); water (0.5 ml.); added to main compartments of Warburg flasks. Side arms contained 10 mm-substrate (0.5 ml.) adjusted to pH 6.5 with NaOH (water for blank determinations). Centre wells contained 3 N-H₂SO₄ (0.3 ml.). The gas phase was N₂. After equilibration for 20 min. at 37°, substrate was added, and CO₂ liberation followed for 20 min. The H₂SO₄ was then added, and final CO₂ readings were taken after a further 10 min. Enzyme activities are expressed as q_{CO_2} , i.e. as μ l. of CO₂ formed/mg. fresh wt. of liver/hr.

Substrate ...	q_{CO_2}	
	L-Cysteate	L-Cysteinesulphinate
Case*		
Cystinosis 1	0.12	0.36
Cystinosis 2	0.11	—
Cystinosis 3	—	0.31
Normal 1	0.10	0.29
Normal 2	0.11	0.26

* All liver specimens had been stored frozen for 4 months.

Table 8. *Cysteine-desulphydrase activity of liver homogenates*

Experimental details are given in the Methods section. Liver specimens had been stored frozen for 3 months (except Normal 1, which was stored for 4 months) before assay.

Case	H ₂ S produced (μ moles/g. dry wt. of liver homogenate/hr.)
Cystinosis 1	32.0
Cystinosis 2	31.6
Normal 1	33.3
Normal 2	27.2

DISCUSSION

A comparative investigation of the kind described suffers the limitation that experimental tissues, particularly in cases of a rare disease such as cystinosis, may have to be accumulated over a considerable period of time, with consequent possible changes in enzymic activity due to variable periods of frozen storage. As far as possible in this work such uncertainties have been controlled by determining the activities of individual enzymes at identical ages of the cystinotic- and normal-tissue specimens.

Accepting this as a valid basis for comparison it may be concluded that no differences attributable to a simple enzyme deficiency could be observed, under conditions of assay *in vitro*, between cystinotic- and normal-liver specimens in their capacity to degrade cystine, cysteine, and certain of their metabolic intermediates.

Theories proposing a specific defect in cystine metabolism to explain the aetiology of cystinosis have understandably selected the reduction of cystine to cysteine as the most probable site of such an abnormality, and recent preliminary reports have claimed the defective reduction of cystine by two different enzyme systems. For the NADH₂-dependent cystine-reductase activity of whole blood, in two cases Worthen & Good (1961) reported values that were about 60 % of the lower limit observed for normal controls, but it was not stated whether excess cystine, which occurs in

cystinotic plasma (Brigham, Stein & Moore, 1960), had been removed before assay. In the present study, endogenous activity of cystine reductase due to excess of cystine in cystinotic-liver homogenates, which had been dialysed for only 24 hr., was such that the apparent effect of further cystine addition was only about 40 % of that for the fully stimulated reaction of normal liver. Prolonged dialysis of the homogenates was necessary to compare the true effects of standard additions of cystine. Seegmiller & Howell (1961) reported glutathione-reductase activity of peripheral blood to be in the normal range in cystinosis, but concluded that the enzyme catalysing the reduction of cystine by reduced glutathione, and called 'cystine transhydrogenase' by them, may have reduced activity in the presence of plasma in some patients with cystinosis. These authors suggested either an enzyme deficiency or a plasma inhibitor to account for the low activity observed.

The present study supports neither of these suggestions of a failure of cystine reduction resulting from a simple enzyme deficiency in cystinosis, and the occurrence of two distinct primary enzyme defects is highly improbable. Results have indicated the normal reduction of cystine *in vitro* by cystine reductase of cystinotic liver and kidney, and by the glutathione-cysteine-transhydrogenase system of liver. Moreover, the possibility of indirect impairment of the latter system *in vivo*, due to a failure of NADPH₂ supply to the glutathione-reductase reaction, is unlikely in view of the demonstration of normal concentrations of reduced glutathione in cystinotic liver. A similar indirect limitation of cystine-reductase activity, due to a defective link in the supply of NADH₂, cannot be ruled out, although it might be argued that tissues already possessing a completely operable alternative means for the reduction of cystine would not suffer such a severe effect as cystinosis if cystine-reductase activity were impaired. However, it cannot be assumed from the above results that cystine reductase and glutathione-cysteine transhydrogenase coexist uniformly throughout all liver cells. It is possible that the reticulo-endothelial cells, to which cystine storage is specifically confined, depend entirely on cystine reductase for the reduction of cystine.

It is conceivable that conditions leading to the failure of cystine reductase could result from a deficiency in a system with which the reductase is coupled preferentially, or from a rate-limiting effect or deficiency at an early point in a sequence of reactions supplying NADH₂. These possibilities have also been considered by Worthen & Good (1958). The citric acid cycle is a prominent sequence of this kind, and this is reported to contribute only

little activity to the overall metabolism of cornea, a further specific site of cystine deposition in cystinosis. An abnormality of respiratory metabolism might further reduce this activity to negligible proportions, thereby creating a situation of specific or generalized failure of NADH₂ supply. It is not known whether the citric acid cycle also operates generally at low levels in reticulo-endothelial tissues, certain aspects of which, relating to the possible role of NADH₂ depletion in cystinotic tissues, are now being investigated. Whether or not such a relationship could constitute the primary defect in cystinosis, excess tissue cystine, formed by whatever means, might be expected to induce some degree of progressive depletion of NADH₂ by inhibiting thiol-group-dependent dehydrogenase systems. Such inhibition is suggested by the hyperpyruvaemia which occurs in cystinosis, and by the lowering of blood pyruvate concentration to normal values in patients treated with the thiol compounds Dimer-caprol (2,3-dimercaptopropan-1-ol) and penicillamine ($\beta\beta$ -dimethyleysteine) (Clayton & Patrick, 1961).

Though cystine reduction has received most attention in the present study, it is possible that a defect at some other point in the intermediary metabolism of cystine and cysteine might result in cystinosis. Such a relationship would be more difficult to conceive, and no evidence of a direct abnormality of this kind has emerged from the above investigations on the degradative reactions of cystine and cysteine via both oxidative and non-oxidative pathways. Confirmation of these results, and their extension to include synthetic and other pathways of cystine utilization, would perhaps best be met by the administration of S-labelled cystine to cystinosis patients. In agreement with the present results, Seegmiller & Howell (1961) have demonstrated the complete oxidation of cystine in cystinosis patients by the appearance of labelled urinary inorganic sulphate after [³⁵S]cystine infusions. These authors have also commented on the absence of urinary taurine in some of their cases. There is no obvious explanation of this finding, particularly in view of the probable role of cysteic acid and cysteinesulphinic acid as intermediates in the oxidation of cysteine, and the demonstration of normal ability of cystinotic liver to decarboxylate these substrates. Also, no abnormality of plasma taurine concentration in cystinosis has been reported. However, the possible existence of an alternative pathway of taurine formation may have some bearing on this question. For example, there is some evidence for a direct oxidative route from cystine to taurine via the disulphoxides of cystine and cystamine, as originally postulated by Medes & Floyd (1942).

SUMMARY

1. Autopsy samples of liver from three cases of cystinosis have been compared with normal liver with respect to the degradative metabolism of cystine and cysteine, with particular emphasis on the enzymic reduction of cystine.

2. Under the conditions of these experiments the reduction of cystine to cysteine by cystine reductase and by glutathione-cysteine transhydrogenase was normal in cystinotic liver. Also, no abnormality was found in the cystine-reductase activity of kidney.

3. Glutathione-reductase activity and reduced glutathione content of cystinotic liver were normal.

4. Cysteine, cystine and cysteinesulphinic acid were completely oxidized by liver of the cystinosis cases, and these findings were substantiated by the normal urinary excretion of inorganic sulphate by cystinosis patients.

5. The liberation of hydrogen sulphide from cysteine and cystine by cystinotic liver was normal, the reactivity of cystine depending on its prior reduction to cysteine.

6. No abnormality of cystinotic liver was observed for the transamination of cysteine, cysteic acid and cysteinesulphinic acid with α -oxoglutarate.

7. Normal decarboxylation of cysteic acid and cysteinesulphinic acid was observed for cystinotic liver.

8. These findings are discussed with respect to the aetiology of cystinosis.

I wish to thank Dr Barbara Clayton for her interest in this work, and Mr A. E. Whitfield for technical assistance. I am also indebted to Dr Barbara Ockenden and Dr M. Bodian, who performed the autopsies, and to Professor A. A. Moncrieff, Dr P. R. Evans and Dr R. Lightwood for permission to study their cases.

REFERENCES

- Abderhalden, F. (1903). *Hoppe-Seyl. Z.* **38**, 557.
 Bickel, H., Baar, H. S., Astley, R., Douglas, A. A., Finch, E., Harris, H., Harvey, C. C., Hickmans, E. M., Philpott, M. G., Smallwood, W. C., Smellie, J. M. & Teall, C. G. (1953). *Acta paediatr., Stockh.*, **42**, suppl. 90.
 Brigham, M. P., Stein, W. H. & Moore, S. (1960). *J. clin. Invest.* **39**, 1633.
 Clayton, B. E. & Patrick, A. D. (1961). *Lancet*, ii, 909.
 Dent, C. E. (1947). *Biochem. J.* **41**, 240.
 Dodgson, K. S. (1961). *Biochem. J.* **78**, 312.
 Evered, D. F. (1956). *Biochem. J.* **62**, 416.
 Fanconi, G. (1936). *Jb. Kinderheilk.* **147**, 299.
 Friedemann, T. E. & Haugen, G. E. (1943). *J. biol. Chem.* **147**, 415.
 Hope, D. B. (1955). *Biochem. J.* **59**, 497.
 Kornberg, A. (1950). *J. biol. Chem.* **182**, 805.
 Kornberg, A. & Horecker, B. L. (1955). In *Methods in Enzymology*, vol. 1, p. 323. Ed. by Colowick, S. P. & Kaplan, N. O. New York: Academic Press Inc.
 Krebs, H. A. & Henseleit, K. (1932). *Hoppe-Seyl. Z.* **210**, 33.
 Lignac, G. O. E. (1924). *Dtsch. Arch. klin. Med.* **145**, 139.
 McCune, D. J., Mason, H. H. & Clarke, H. T. (1943). *Amer. J. Dis. Child.* **65**, 81.
 Maw, G. A. (1954). *Biochem. J.* **58**, 665.
 Medes, G. (1939). *Biochem. J.* **33**, 1559.
 Medes, G. & Floyd, N. (1942). *Biochem. J.* **36**, 259.
 Pirie, N. W. (1934). *Biochem. J.* **28**, 305.
 Racker, E. (1955). *J. biol. Chem.* **217**, 855.
 Rall, T. W. & Lehninger, A. L. (1952). *J. biol. Chem.* **194**, 119.
 Seegmiller, J. E. & Howell, R. R. (1961). *Clin. Res.* **9**, 189.
 Sullivan, M. X., Hess, W. C. & Howard, H. W. (1942). *J. biol. Chem.* **145**, 621.
 Warburg, O. & Christian, W. (1941). *Biochem. Z.* **310**, 384.
 Woodward, G. E. (1935). *J. biol. Chem.* **109**, 1.
 Worthen, H. G. & Good, R. A. (1958). *Amer. J. Dis. Child.* **95**, 653.
 Worthen, H. G. & Good, R. A. (1961). *Amer. J. Dis. Child.* **102**, 494.

Biochem. J. (1962) **83**, 256

Studies on α -Amylase Formation by *Bacillus subtilis*

By G. COLEMAN*† AND W. H. ELLIOTT

Department of Biochemistry, John Curtin School of Medical Research,
 Australian National University, Canberra, A.C.T., Australia

(Received 26 September 1961)

The synthesis of α -amylase (α -1,4-glucan 4-glucanhydrolase, EC 3.2.1.1) by cultures of *Bacillus subtilis* has been studied by several groups of

workers, who have defined the basic conditions required for the formation of high yields of the enzyme (Hagihara, 1951, 1954; Fukumoto, Yamamoto, Tsuru & Ichikawa, 1957*b*; Fukumoto, Yamamoto & Tsuru, 1957*a*). For α -amylase production a carbohydrate source is required and this

* Australian National University Research Scholar.

† Present address: Microbiology Unit, Department of Biochemistry, University of Oxford.