- Brent, L., Medawar, P. B. & Ruskiewicz, M. (1961). *Brit. J. exp. Path.* **42**, 464.
- Castermans, A. (1961a). *Nature, Lond.*, **189**, 504.
- Castermans, A. (1961b). *Transplant. Bull.* **27**, 98.
- Converse, J. M. & Rapaport, F. T. (1960). *Ann. N.Y. Acad. Sci.* **87**, 6.
- Crompton, M. J. & Davies, D. A. L. (1956). *Proc. Roy. Soc. B*, **145**, 109.
- Davies, D. A. L. (1957). *Biochem. J.* **67**, 253.
- Davies, D. A. L. (1960). *Advanc. Carbohydr. Chem.* **15**, 271.
- Davies, D. A. L. (1961). *Abstr. 5th int. Congr. Biochem., Moscow*, **20**, p. 415.
- Davies, D. A. L. (1962a). *Nature, Lond.*, **193**, 34.
- Davies, D. A. L. (1962b). In *Ciba Foundation Symp., Transplantation*. Ed. by Wolstenholme, G. E. W. & Cameron, M. P. London: J. and A. Churchill Ltd.
- Davies, D. A. L. (1962c). *Ann. N.Y. Acad. Sci.* (in the Press).
- Davies, D. A. L. & Hutchison, A. M. (1961). *Brit. J. exp. Path.* **42**, 587.
- Dische, Z. & Borenfreund, E. (1951). *J. biol. Chem.* **192**, 583.
- Dische, Z. & Shettles, L. B. (1948). *J. biol. Chem.* **175**, 595.
- Earle, W. R. (1943). *J. nat. Cancer Inst.* **4**, 165.
- Gorer, P. A. (1956). *Advanc. Cancer Res.* **4**, 149.
- Gorer, P. A. & Mikulska, Z. B. (1954). *Cancer Res.* **14**, 651.
- Gorer, P. A. & Mikulska, Z. B. (1959). *Proc. Roy. Soc. B*, **151**, 57.
- Hašková, V. & Hilgert, I. (1961). *Folia biol., Prague*, **7**, 81.
- Hašková, V. & Hrušeová, M. (1958). *Nature, Lond.*, **182**, 61.
- Haughton, G. (1962). *Ann. N.Y. Acad. Sci.* (in the Press).
- Haughton, G. & Davies, D. A. L. (1962). *Brit. J. exp. Path.* (in the Press).
- Herzenberg, L. A. & Herzenberg, L. A. (1961). *Proc. nat. Acad. Sci., Wash.*, **47**, 762.
- Kaliss, N. (1958). *Cancer Res.* **18**, 992.
- Kandutsch, A. A. (1960). *Cancer Res.* **20**, 264.
- Kandutsch, A. A. (1961). *Transplant. Bull.* **27**, 135.
- Kandutsch, A. A. & Reinert-Wenck, V. (1957). *J. exp. Med.* **105**, 125.
- Ma, T. S. & Zuazaga, G. (1942). *Industr. Engn. Chem. (Anal.)*, **14**, 280.
- Markham, R. (1942). *Biochem. J.* **36**, 790.
- Martland, M. & Robison, R. (1926). *Biochem. J.* **20**, 847.
- Medawar, P. B. (1958a). *Proc. Roy. Soc. B*, **149**, 145.
- Medawar, P. B. (1958b). *Nature, Lond.*, **182**, 62.
- Möller, G. (1961). *J. exp. Med.* **114**, 415.
- Moore, S. & Stein, W. H. (1948). *J. biol. Chem.* **176**, 367.
- Morgan, W. T. J. (1960). *Bull. Soc. Chim. biol., Paris*, **42**, 1591.
- Rondle, C. J. M. & Morgan, W. T. J. (1955). *Biochem. J.* **61**, 586.
- Snell, G. D. (1957). *Annu. Rev. Microbiol.* **11**, 439.
- Snell, G. D. (1958). *J. nat. Cancer Inst.* **20**, 787.
- Snell, G. D., Winn, H. J., Stimpfling, J. H. & Parker, S. J. (1960). *J. exp. Med.* **112**, 293.
- Synge, R. L. M. & Wood, J. C. (1958). *Biochem. J.* **70**, 321.
- Warren, L. (1959). *J. biol. Chem.* **234**, 1971.

Biochem. J. (1962) **84**, 317

Production of Aminoacetone by *Rhodopseudomonas spheroides**

By A. NEUBERGER AND G. H. TAIT

Department of Chemical Pathology and M.R.C. Research Group in Enzymology, St Mary's Hospital Medical School, London, W. 2

(Received 13 February 1962)

Shemin (1955) and Nemeth, Russell & Shemin (1957) were the first to suggest a possible metabolic role for aminoacetone. On administration of amino-[1-¹⁴C]acetone to rats the label appeared in respiratory carbon dioxide, in urinary formate and in the ureido groups of uric acid. These findings were analogous to those obtained with δ -aminolaevulinic acid, on the basis of which the succinate-glycine cycle was postulated (Shemin & Russell, 1953). Mauzerall & Granick (1956) detected a small amount of an aminoketone, with properties similar to those of aminoacetone, in normal human urine. Gibson, Laver & Neuberger (1958) and Kikuchi, Kumar & Shemin (1959) showed that δ -amino-

laevulate synthetase was not specific for succinyl-coenzyme A, but would catalyse the condensation of glycine with various acyl-coenzyme A compounds, including acetyl-coenzyme A. Elliott (1958, 1959, 1960a) reported that an aminoketone is produced when *Staphylococcus aureus* is incubated aerobically in a medium containing glycine and glucose or in one containing threonine. He characterized the aminoketone produced from threonine as aminoacetone by the preparation of derivatives, but the one from glycine and glucose is produced in very small quantities and has only been identified as aminoacetone on the basis of its chromatographic properties. Elliott (1959) postulated that aminoacetone could be metabolized by a cyclic pathway similar to the succinate-glycine cycle for δ -aminolaevulate, and supported this

* Some of this work forms part of a Ph.D. Thesis submitted to the University of London in July 1961 by G. H. Tait.

hypothesis by finding an enzyme in ox plasma which oxidatively deaminated aminoacetone to methylglyoxal (Elliott, 1960b). Urata & Granick (1961) described the formation of aminoacetone from pyruvate and glycine and from threonine in liver mitochondria and also detected the breakdown of aminoacetone in liver. In addition, Mijovic & Walker (1960) identified aminoacetone as a constituent of the antibiotic micrococcin P. These reports indicate that aminoacetone may be an important natural metabolite.

The work described in this paper deals with the finding that an aminoketone accumulates when *Rhodopseudomonas spheroides* is incubated under suitable conditions. This aminoketone has been identified as aminoacetone. In addition, a NAD-dependent enzyme which converts L-threonine into aminoacetone (L-threonine dehydrogenase) has been detected in extracts of these organisms. This enzyme has been partially purified and some of its properties have been studied. Part of this work has already been briefly reported (Neuberger & Tait, 1960).

MATERIALS AND METHODS

Chemicals

NAD⁺, NADH, NADP, α -oxoglutarate and reduced glutathione were obtained from C. F. Boehringer und Soehne, Mannheim, Germany; oxidized nicotinamide-hypoxanthine dinucleotide [described as 'deamino-DPN⁺' by Pullman, Colowick & Kaplan (1952)] and oxidized 3-acetylpyridine-adenine dinucleotide [described as '3-acetylpyridine-DPN⁺' by Kaplan & Ciotti (1956)] were from Mann Research Laboratories Inc., New York, U.S.A. D-Threonine, L-allothreonine and D-allothreonine were kindly given by Dr D. F. Elliott, National Institute for Medical Research, Mill Hill, London, N.W. 7. Aminoacetone hydrochloride was prepared by catalytic hydrogenation (Pd-charcoal at atmospheric pressure) of oximinacetone (A. Neuberger & J. J. Scott, unpublished work); it was recrystallized from dry ethanol-ether. γ -Amino- β -hydroxybutyric acid was prepared as described by Neilands, Neuberger & Scott (1960). Phosphate buffers were made from K₂HPO₄ and KH₂PO₄, and tris-HCl buffers according to Gomori (1955).

Organisms

The strain of *Rps. spheroides* was obtained from Dr J. Lascelles (National Collection of Industrial Bacteria no. 8253) and was maintained in stab culture as described by Lascelles (1956).

In all experiments the organisms used were grown anaerobically in the light at 32–34° for 44 hr. in 'medium S' of Lascelles (1956) in 500 ml. or 1000 ml. conical flasks filled almost to the top. No special precautions were taken to make the conditions strictly anaerobic. The illumination was supplied and the temperature maintained by 60W bulbs placed at an appropriate distance from the flasks.

After growth the organisms were harvested by centrifuging at 0°, washed with a volume of cold 0.05M-phos-

phate buffer, pH 7.4, equal to half the original culture volume, and suspended at a concentration of 80 mg. dry wt./ml. This suspension was used immediately or kept frozen until required. The dry weight of organisms in suspensions was determined by the method of Lascelles (1956).

Procedure for the study of aminoketone synthesis by suspensions of organisms

Aerobic conditions. Organisms, grown as described above, were suspended at 4 mg. dry wt./ml. in 'mixture I' of Lascelles (1956) or in 'mixture I' modified in various ways. The composition of 'mixture I' is: α -oxoglutaric acid, 0.01M; glycine, 0.01M; fumaric acid, 0.02M; (NH₄)₂HPO₄, 3 mM; MgSO₄, 0.8 mM; CaCl₂, 0.4 mM; MnCl₂, 5 μ M; phosphate buffer, pH 6.8, 6 mM. The pH was adjusted to 6.8 with NaOH. The suspensions, 5–10 ml. in volume, were contained in 2.5 cm. $\times$ 15 cm. tubes. These tubes were connected in series and the suspensions vigorously aerated by bubbling air through from a cylinder. Incubations were done at 35°. Samples were removed at various times, 0.06 vol. of 100% (w/v) trichloroacetic acid was added and aminoketone estimated colorimetrically in the supernatant, after removal of the precipitate by centrifuging, by the modification of the method of Mauzerall & Granick (1956) described by Gibson *et al.* (1958). Quantities of aminoketone formed are expressed by using aminoacetone as a standard.

Anaerobic conditions. Organisms were illuminated at a concentration of 1 mg. dry wt./ml. in 'mixture I' of Lascelles (1956) for 18–24 hr. at 32–34°. The heat and light were supplied by 60W bulbs placed at an appropriate distance from the tubes.

Purification of the aminoketones

The aminoketone which accumulates on aerobic incubation in the dark ('aerobic' aminoketone) and the one which accumulates anaerobically in the light ('anaerobic' aminoketone) were produced in large amounts, by using the optimum conditions worked out (see Results section), and were purified as follows.

At the end of the incubation the pH was adjusted to 5 with HCl, the organisms were removed by centrifuging, and the supernatants used for purification.

Step 1. The supernatants, after dilution (1:2) with water, were passed through 2 cm. $\times$ 20 cm. columns of Zeo-Karb 225 (H⁺ form; 40–80 mesh). Solution was passed until the columns were almost saturated, as judged by the colour. The columns were washed with water until the effluents were neutral. The aminoketone was eluted with N-HCl followed by 1.5N-HCl, and 10 ml. fractions were collected. All fractions containing aminoketone, as determined colorimetrically, were combined and dried.

Step 2. The aminoketone and amino acids were extracted from the large residue, consisting mostly of KCl, obtained from step 1 with acetone containing 5% of 6N-HCl (Smith, 1958). The extract was dried and dissolved in water.

Step 3. The solution from step 2 was applied to a 1 cm. $\times$ 24 cm. column of Zeo-Karb 225 (H⁺ form; 8% cross-linked; 50 μ particle size). The column was washed with water until the effluent was neutral. Elution was performed with 0.5N-HCl at a flow rate of 5 ml./20 min., and 5 ml. fractions were collected. Both 'aerobic' and 'anaero-

bic' aminoketones were eluted with approx. 200 ml. of 0.5N-HCl. Analysis of the fractions showed a symmetrical peak with very little tailing. The fractions containing aminoketone were pooled, dried and the residue was dissolved in water.

Step 4. On paper chromatography in butanol-acetic acid-water (12:3:5, by vol.) the fraction from step 3 was found to contain small traces of an amino acid. The final purification was therefore achieved by paper chromatography in butanol-acetic acid-water and the aminoketone eluted off the paper with dilute HCl.

Gas-partition chromatography

Gas-partition chromatography was carried out according to James & Martin (1952). The apparatus was built by Dr R. D. Marshall, and we are grateful to him for instruction in its use. The column was made from kieselguhr (Celite 545, Johns Manville Co. Ltd.) which was mixed with half its weight of silicone (MS 550, Hopkin and Williams Ltd.) containing 10% (w/w) of stearic acid. The column (internal diameter 0.5 cm. $\times$ 130 cm. long) was contained in a steam jacket. Partition was accomplished by passing nitrogen at a pressure of 45 cm. Hg. The acids coming off the column were titrated with 0.02N-NaOH with 0.001% phenol red as indicator.

With the column used and running it at the pressure stated and at 100°, the elution time of each acid was as follows: formic acid, 7 min.; acetic acid, 20 min.; propionic acid, 35 min.; butyric acid, 65 min.

Preparation of enzyme extracts

Suspensions of organisms, 80 mg. dry wt./ml., were disrupted in the Hughes (1951) press and then diluted with 2 vol. of 0.05M-phosphate buffer, pH 7.4. The suspensions were centrifuged for 90 min. at 105 000g in the Spinco model L ultracentrifuge. The clear yellow supernatants were removed and kept at -20° until required for assay. This extract is referred to below as 'crude enzyme extract'. Protein in the extracts was measured by the biuret method (Layne, 1957).

RESULTS

Lascelles (1956) noted that *Rps. spheroides* incubated aerobically in the dark in 'mixture I' did not form porphyrins. On repeating this experiment this observation was confirmed, but an aminoketone, which was at first assumed to be δ -aminolaevulinic acid, was found. This aminoketone behaved on colorimetric estimation like aminoacetone, and unlike δ -aminolaevulinic acid, in giving a colour immediately the Ehrlich reagent was added. The pyrroles formed on condensation with acetylacetone from authentic aminoacetone and the unknown aminoketone could both be extracted from aqueous solution at pH 8 into ether. The pyrrole formed from δ -aminolaevulinic acid and acetylacetone, having a carboxyl group, could not be extracted from aqueous solution under these conditions. This indicated that the unknown aminoketone lacked an acidic function. Aerobic incuba-

tion of organisms with threonine also gave an aminoketone with similar properties to the one just described.

Accumulation of aminoketone

Aerobic conditions. In 'mixture I', the buffer concentration is low (6 mM), and during aerobic incubation the pH rose from 6.8 to 9 or higher. At a phosphate buffer concentration of 0.2M the pH did not rise markedly, and the amount of aminoketone which accumulated in 16 hr. (63 μ m-moles/ml.) was six times the amount which was found after similar incubation in 'mixture I', but without added buffer (10 μ m-moles/ml.). All further aerobic incubations were therefore done in the presence of 0.2M-phosphate buffer. Omission of diammonium hydrogen phosphate from the medium did not affect the yield of aminoketone.

The accumulation of aminoketone was linear for 4 hr. (Fig. 1), then it stopped completely, and the amount of aminoketone detected remained constant over the next 18 hr. α -Oxoglutarate and fumarate, if added after 4 hr., gave a further increase in aminoketone, but the addition of glycine had no effect.

When the incubation was done in a medium lacking metal ions, the accumulation of aminoketone was low and variable, and it was concluded

Fig. 1. Effect of metal ions on the accumulation of aminoketone. Organisms (4 mg. dry wt./ml.) were incubated aerobically in a medium containing sodium α -oxoglutarate (0.02M), sodium fumarate (0.04M), glycine (0.02M) and phosphate buffer, pH 6.6 (0.2M). Samples were removed at various times and estimated for aminoketone. Additions: ○, Mg²⁺ ions (1.6 mM); Δ, Ca²⁺ ions (0.8 mM); ●, Mn²⁺ ions (10 μ M); □, no additions.

that one or more of the metals in 'mixture I'— Mg^{2+} , Ca^{2+} , Mn^{2+} —were required. Fig. 1 shows that Mn^{2+} ions greatly enhance the yield of aminoketone, whereas Ca^{2+} and Mg^{2+} ions have little or no effect. An increase in the concentration of Mn^{2+} ions in the medium from $10\ \mu M$ to $0.5\ mM$ increased the yield of aminoketone approximately twofold, but higher concentrations had no further effect.

With $0.2\ M$ -phosphate buffer the pH optimum for accumulation of aminoketone under aerobic conditions was 6.2. At pH 7 the amount accumulating was only about half that found at pH 6.2. No aminoketone was detected if α -oxoglutarate and fumarate were replaced by acetate.

Anaerobic conditions. When organisms were illuminated in 'mixture I' under anaerobic conditions a small amount of aminoketone accumulated ($12\ \mu M$ -moles/ml. in 20 hr.). The yield was increased to $80\ \mu M$ -moles/ml. in 20 hr. by raising the concentration of potassium phosphate buffer in the medium to $0.15\ M$. This effect is not dependent on the cation present in the buffer; a similar yield of aminoketone was obtained with $0.15\ M$ -sodium phosphate or tris buffer. Under these conditions the production of aminoketone was linear for 16 hr. and after this the rate decreased and the production appeared to stop at about 20 hr.

Identification of the aminoketone

The aminoketones produced under the two different incubation conditions were purified and identified separately. Large amounts were obtained by employing the optimum incubation conditions described above. The yields obtained during the purification were about 80–90 % for each step, resulting in a 60 % overall yield up to the end of step 3. However, the final purification by paper chromatography gave rather low yields.

The following properties of both 'aerobic' and 'anaerobic' aminoketones are identical with those of aminoacetone: (a) paper chromatography in butanol–acetic acid–water (12:3:5, by vol.) gave a single spot with R_F 0.39 (aminoacetone, R_F 0.39;

δ -aminolaevulinic acid, R_F 0.29); (b) on paper they give yellow spots with ninhydrin, which turn brown and then purple on heating as do aminoacetone and δ -aminolaevulinic acid; (c) the pyrroles formed from the aminoketones on condensation with acetylacetone have the same R_F as the pyrrole from aminoacetone and acetylacetone on paper chromatography in butanol–acetic acid–water (63:10:27, by vol.) (R_F 0.87) and in butanol–1.5*N*-ammonia (1:1, v/v) (R_F 0.85).

Final proof of the identity of the aminoketones was obtained by periodate oxidation and identification of the fatty acid formed by gas-partition chromatography. Shemin, Russell & Abramsky (1955) reported that periodate oxidation of δ -aminolaevulinic acid gives a good yield of succinic acid. On similar treatment aminoacetone ought to give acetic acid. Samples of authentic aminoacetone and of the purified aminoketones were treated with periodate at neutral pH and the reaction was allowed to proceed for 16 hr. in the dark at room temperature. The periodate and iodate were precipitated with excess of barium chloride. After centrifuging, the pH of the supernatant was adjusted to 3 with sulphuric acid, and the solution was steam-distilled and the distillate collected until no more acid came over. The distillates were kept alkaline and dried *in vacuo*. The residues were dissolved in a small volume of water, and 1 drop of 100 % orthophosphoric acid was added to each. The fatty acids were extracted into ether, the ether was dried over anhydrous magnesium sulphate, and the solution was evaporated to a small volume with a stream of nitrogen and applied to the end of the gas-partition chromatography column. The chromatography was performed as described in the Materials and Methods section and the results are shown in Table 1. The columns were run for a total of from 60 to 80 min. but no more acid was eluted.

These results show that both the 'aerobic' aminoketone and the 'anaerobic' aminoketone are aminoacetone.

Table 1. *Identification of aminoketones*

The conditions of periodate oxidation of the aminoketones and gas chromatography of the resulting volatile fatty acids are described in the text. Sodium acetate was acidified with H_3PO_4 and treated in the same way as the fatty acids produced by oxidation of the aminoketones. The aminoketones were estimated colorimetrically as described in the Materials and Methods section.

Sample	Amount of sample (μ moles)	Acid recovered on gas chromatography		Elution time (min.)
		(μ moles)	(%)	
Reagent blank	—	0.0	—	—
Aminoacetone	10	4.2	42	20
Aminoacetone	20	11.4	57	20
Aminoacetone	50	26.6	53	20
'Aerobic' aminoketone	13.7	9.8	59	20
'Anaerobic' aminoketone	16.8	11.2	67	20
Sodium acetate	50	25.0	50	20

L-Threonine-dehydrogenase activity in extracts

An enzyme has been found in extracts of *Rps. spheroides* which produces aminoacetone from *L*-threonine in the presence of NAD^+ , indicating that the enzyme responsible is a threonine dehydrogenase (Table 2).

NADP^+ cannot replace NAD^+ as hydrogen acceptor, and there is no reaction in the absence of the latter. Pyridoxal phosphate does not enhance the activity.

The aminoketone formed in the above reaction has not been rigorously identified, but its properties are identical with those of aminoacetone in that the pyrrole formed by condensation with acetylacetone can be extracted into ether at pH 8, and the spectrum of the pyrrole-*p*-dimethylaminobenzaldehyde adduct formed in the modified Mauzerall & Granick (1956) colour reaction is identical with that formed from aminoacetone.

The production of aminoacetone is linear with increasing amounts of enzyme up to at least 4.0 mg. of protein/0.75 ml. of assay system, and linear for at least 2 hr.

The pH optimum is between 8.8 and 9.1 in tris (Fig. 2). This is more clearly shown below with partially purified enzyme. In carbonate-bicarbonate buffer the activity is reduced when compared with that in tris. The addition of sodium hydrogen carbonate to the reaction mixture with 0.1 M-tris buffer also produces a strong inhibition; thus a 50% inhibition is produced with 0.13 M-bicarbonate.

Partial purification of L-threonine dehydrogenase

Protein estimations were performed by the biuret method (Layne, 1957) or by measuring the absorption of solutions at 260 and 280 $m\mu$ (Warburg & Christian, 1942). All purification steps were carried out at 0–4°.

Step 1: treatment with manganous chloride. To crude enzyme extract, prepared as described above, was added 0.05 vol. of M- MnCl_2 . The mixture was left for 15 min. and then centrifuged. The supernatant was dialysed overnight against 0.05 M-phosphate, pH 7.4. A precipitate which formed was removed by centrifuging.

Step 2: fractionation with ammonium sulphate. The supernatant was fractionated with solid $(\text{NH}_4)_2\text{SO}_4$ successively at 35, 45 and 60% saturation. Each precipitate was collected by centrifuging, dissolved in buffer and dialysed.

Step 3: treatment with calcium phosphate gel. The enzyme was found mainly in the solution containing the 45–60% precipitate. This solution was adjusted to pH 6 and treated successively with 0.1, 0.1 and 0.2 vol. of calcium phosphate gel (13.5 mg./ml.), the supernatant being recovered from one gel treatment before addition of the next batch of gel. To the last supernatant was added 0.6 vol. of gel. The precipitate obtained by centrifuging was eluted once with water and twice with 0.05 M-phosphate buffer, pH 7.4.

The yields and purification are described in Table 3. The first buffer eluate, referred to below as 'purified enzyme', shows a purification of 4.8-fold with respect to protein and contains very little nucleic acid. This purified enzyme has been used in a number of experiments which are described below.

With the purified enzyme a linear production of aminoacetone was obtained with increasing amount of enzyme, and the pH optimum curve in tris is similar to that with crude enzyme (Fig. 2), giving an optimum activity at pH 8.8–8.9.

Specificity of L-threonine dehydrogenase

The specificity of both crude and purified enzyme preparations towards the four isomers of α -amino- β -hydroxybutyric acid is shown in Table 4.

Table 2. Requirements for *L*-threonine-dehydrogenase activity

L-Threonine (25 μmoles), pyridoxal phosphate (0.05 μmole) where added, NAD^+ or NADP^+ (as indicated), crude enzyme extract (1.0 mg. of protein) and phosphate buffer, pH 7.4 (75 μmoles), in a total volume of 0.75 ml., were incubated at 37° for 1 hr. The reaction was stopped with trichloroacetic acid. The supernatant, after removal of protein by centrifuging, was analysed for aminoacetone.

Coenzyme	Amount of coenzyme (μmole)	Pyridoxal phosphate	Aminoacetone ($\mu\text{m-moles}$)
NAD^+	0.5	+	32
NAD^+	1.0	+	42
NADP^+	0.5	+	4
NADP^+	1.0	+	5
NAD^+	1.0	–	38
NADP^+	1.0	–	5
No coenzyme	.	+	2

Fig. 2. pH-activity curve for *L*-threonine dehydrogenase. *L*-Threonine (25 μmoles), NAD^+ (0.3 μmole), buffer (75 μmoles) and crude enzyme extract (1.0 mg. of protein), in a total volume of 0.75 ml., were incubated at 37° for 1 hr. The reaction was stopped with trichloroacetic acid and aminoacetone was estimated. Buffers: ●, tris; ○, phosphate; △, carbonate-bicarbonate.

Table 3. *Partial purification of L-threonine dehydrogenase*

For the assay of activity, DL-threonine (50 μ moles), NAD⁺ (0.2 μ mole), tris, pH 8.8 (75 μ moles), and enzyme, in a total volume of 0.75 ml., were incubated at 37° for 1 hr. The reaction was stopped with trichloroacetic acid and aminoacetone was estimated. The purification steps are described in the text. The fractions designated '(NH₄)₂SO₄ precipitates' were obtained by successive precipitation of the MnCl₂ supernatant with (NH₄)₂SO₄ added to give the percentage saturation shown.

Fraction	Specific activity (μ -moles of aminoacetone/mg. of protein/hr.)	Purification (based on protein)	Recovery (%)
Crude extract	113	1.0	100
MnCl ₂ supernatant	126	1.1	65
35% (NH ₄) ₂ SO ₄ precipitate	—	—	0
45% (NH ₄) ₂ SO ₄ precipitate	155	1.4	15
60% (NH ₄) ₂ SO ₄ precipitate	224	2.0	40
80% (NH ₄) ₂ SO ₄ precipitate	132	1.2	12
First buffer eluate	545	4.8	24
Second buffer eluate	465	4.1	8

Table 4. *Specificity of L-threonine dehydrogenase*

α -Amino- β -hydroxybutyric acid isomers (as stated), NAD⁺ (0.5 μ mole), tris, pH 8.8 (75 μ moles), and crude enzyme (0.96 mg. of protein) or purified enzyme (0.11 mg. of protein), in a total volume of 0.75 ml., were incubated at 37° for 1 hr. The reaction was stopped with trichloroacetic acid and aminoacetone was estimated.

Substrate	Amount of substrate (μ moles)	Amino- acetone (μ -moles)
With crude enzyme		
L-Threonine	5.0	33
D-Allothreonine	5.0	11
D-Threonine	5.0	0
L-Allothreonine	5.0	0
With purified enzyme		
L-Threonine	5.0	17
L-Threonine	10.0	32
D-Allothreonine	5.0	9
D-Allothreonine	10.0	15
D-Threonine	10.0	0
L-Allothreonine	10.0	0
L-Threonine	5.0	17
+		
D-Threonine	5.0	17
L-Threonine	5.0	
+		23
L-Allothreonine	5.0	
L-Threonine	5.0	8
+		
D-Allothreonine	5.0	8
D-Allothreonine	5.0	
+		8
D-Threonine	5.0	
D-Allothreonine	5.0	8
+		
L-Allothreonine	5.0	

Only L-threonine and D-allothreonine are acted on. D-Threonine and L-allothreonine are inactive both as substrates and inhibitors.

Serine does not inhibit the activity, and γ -amino- β -hydroxybutyric acid is not oxidized to form an aminoketone, nor does it interfere with L-threonine-dehydrogenase activity.

Oxidized nicotinamide-hypoxanthine dinucleotide cannot replace NAD⁺ for L-threonine-dehydrogenase activity and oxidized 3-acetylpyridine-adenine dinucleotide has only 40% of the activity of NAD⁺. Neither analogue inhibits the activity with NAD⁺.

The ratio of aminoacetone formed to NADH formed is approximately unity. Thus, in an experiment where DL-threonine (200 μ moles), NAD⁺ (20 μ moles), tris, pH 8.8 (300 μ moles), and purified enzyme (0.21 mg. of protein) were incubated in a total volume of 3.0 ml. at 37° in the thermostatically controlled cell-holder of the Unicam SP. 500, 70 μ -moles of NADH and 84 μ -moles of aminoacetone were found after 30 min.

Of the two mechanisms for the enzymic conversion of L-threonine into aminoacetone, discussed below, one involved D-1-aminopropan-2-ol as an intermediate. DL-1-aminopropan-2-ol was therefore tested to see whether it could be dehydrogenated to aminoacetone: with crude enzyme extracts no aminoacetone was detected in the presence of either NAD⁺ or NADP⁺ as hydrogen acceptors; in addition DL-1-aminopropan-2-ol did not inhibit the conversion of L-threonine into aminoacetone.

Effect of inhibitors

A range of known enzyme inhibitors has been tried for an effect on L-threonine-dehydrogenase activity (Table 5). The controls were performed in each case to allow: (a) for interference of the added compound with the colour reaction, and (b) for non-enzymic destruction of aminoacetone. None of the compounds tested had any appreciable effect on the colorimetric estimation of aminoacetone, but aminoacetone disappears when incubated at 37° in the presence of mercuric sulphate, copper sulphate or manganous sulphate. *p*-Chloromercuribenzoate, Ag⁺ ions and EDTA strongly inhibit, and Zn²⁺ ions, Pb²⁺ ions, sodium cyanide,

Table 5. *Effect of enzyme inhibitors on L-threonine dehydrogenase*

DL-Threonine (50 μ moles), NAD⁺ (0.3 μ mole), tris, pH 8.8 (75 μ moles), enzyme (MnCl₂ supernatant containing 0.26 mg. of protein and with specific activity as stated in Table 3) and inhibitors (as stated), in a total volume of 0.75 ml., were incubated at 37° for 1 hr. The reaction was stopped with trichloroacetic acid and aminoacetone was estimated. For the controls, aminoacetone (52 μ m-moles), tris, pH 8.8 (75 μ moles), and inhibitor, in a total volume of 0.75 ml., were mixed. Trichloroacetic acid was added immediately to the non-incubated control, and after incubation at 37° for 30 min. to the incubated control. Results are expressed as percentages of aminoacetone in the tubes without inhibitor. The non-incubated control and the incubated control gave the same amount of aminoacetone in the absence of inhibitor.

Added compound	Concn. of added compound (mm)	Enzyme activity		
		Experimental	Controls	
			Non-incubated	Incubated
None	—	100	100	100
HgSO ₄	1.0	10	95	10
CuSO ₄	1.0	6	100	24
AgNO ₃	2.0	12	107	99
ZnSO ₄	1.0	77	100	98
Lead acetate	1.0	54	105	103
FeSO ₄	1.0	125	105	106
FeCl ₃	1.0	125	111	112
MgSO ₄	1.0	119	103	101
MnSO ₄	1.0	62	81	61
CaCl ₂	1.0	142	105	103
Thiosemicarbazide	4.0	116	111	105
Sodium <i>p</i> -chloromercuribenzoate	0.5	16	104	100
NaCN	0.2	50	99	95
L-Penicillamine	2.0	112	114	105
D-Penicillamine	2.0	119	115	107
L-Cysteine	2.0	125	115	107
Iodoacetamide	5.4	74	101	120
NaN ₃	0.2	112	114	105
$\alpha\alpha'$ -Bipyridyl	0.2	97	101	98
8-Hydroxyquinoline	5.4	65	103	100
EDTA	1.0	17	106	100

iodoacetamide, and 8-hydroxyquinoline also inhibit but less markedly.

The inhibition by EDTA has been investigated further. Under assay conditions identical to those described in Table 5 a 50% inhibition was obtained at a concentration of 56 μ M.

Michaelis constants of L-threonine dehydrogenase

The K_m values for L-threonine and NAD⁺ for L-threonine dehydrogenase have been determined with crude enzyme extract. The assays were done in tris buffer, pH 8.8. For L-threonine the assays were done in the presence of enough NAD⁺ to saturate the enzyme, and for NAD⁺ in the presence of enough L-threonine to saturate the enzyme. The results, obtained by plotting the reciprocal of the velocity of aminoacetone production against the reciprocal of the substrate concentration (Lineweaver & Burk, 1934) are shown in Fig. 3 and give K_m values (at 37°) of 74 mM for L-threonine and 0.17 mM for NAD⁺.

D- β -Hydroxybutyrate-dehydrogenase activity in extracts

L-Threonine is very similar structurally to D- β -hydroxybutyric acid, and DL- β -hydroxybutyric

acid strongly inhibited aminoacetone production. Investigation showed that the apparent inhibition was due to the extremely rapid reduction of NAD⁺ by the crude enzyme extract and by the partially purified L-threonine dehydrogenase in the presence of β -hydroxybutyric acid.

Some of the properties of this soluble β -hydroxybutyrate dehydrogenase have been studied, since it was thought that the same enzyme might be responsible for the dehydrogenation of both β -hydroxybutyric acid and threonine.

With crude enzyme extract the production of NADH was linear for 3 min., and the amount of NADH formed was proportional to the enzyme added up to at least 0.25 mg. of protein/3.0 ml. of assay mixture.

Partial purification of the enzyme was carried out exactly as described for L-threonine dehydrogenase. A comparison of purifications and yields of both activities showed that no substantial separation was obtained. In general the activities towards the two substrates were parallel in the various fractions. However, there were sufficient differences to indicate the possibility that the oxidation of L-threonine and D- β -hydroxybutyric acid might be catalysed by two separate enzymes.

With the assay conditions described in Table 6 the pH optimum in 0.1 M-tris was found to be 8.4–8.6. At a given pH the activity is less in carbonate-hydrogen carbonate buffer than it is in tris. This apparent inhibition is also shown when 0.2 M-

Fig. 3. Michaelis constants for L-threonine dehydrogenase. In the upper diagram the amount of NAD^+ was kept constant ($0.3 \mu\text{mole/assay}$) and the concentration of L-threonine was varied. In the lower diagram the amount of L-threonine was kept constant ($150 \mu\text{moles/assay}$) and the concentration of NAD^+ was varied. In both sets of experiments a total volume of 0.75 ml. also containing tris, pH 8.8 ($75 \mu\text{moles}$), and crude enzyme extract ($0.27 \text{ mg. of protein}$) was incubated at 37° for 30 min. The reaction was stopped with trichloroacetic acid and aminoacetone was estimated. The initial reaction velocity, v , is expressed as $\mu\text{moles of aminoacetone formed in 30 min.}$

sodium hydrogen carbonate is added to an assay system containing 0.1 M-tris, when the activity is reduced to 39 % of the control. As described above, a similar concentration of sodium hydrogen carbonate lowers the L-threonine-dehydrogenase activity to 37 % of the control. NADP^+ and oxidized nicotinamide-hypoxanthine dinucleotide are not reduced in the presence of DL- β -hydroxybutyric acid, and oxidized 3-acetylpyridine-adenine dinucleotide is reduced at 8 % of the rate of NAD^+ . None of these nucleotides inhibits NAD^+ reduction.

The effect of a number of known enzyme inhibitors on β -hydroxybutyrate-dehydrogenase activity is shown in Table 6.

The effect of EDTA has been investigated further with the same assay system as that described in Table 6. When, instead of starting the reaction by addition of enzyme, as was done in the experiments described in Table 6, the enzyme was preincubated with EDTA in the presence of either substrate or coenzyme for 5 min. at 37° and then the reaction started by addition of coenzyme or substrate, only 5 % of the activity of the control was obtained with concentrations of EDTA between 0.1 and 1.0 mM. On further addition of enzyme to the mixture inhibited by mM-EDTA, the same percentage of the control activity as in Table 6 was obtained, showing that EDTA takes a few minutes to react with the enzyme to produce full inhibition of activity. Under these conditions, where EDTA and enzyme were preincubated for 5 min. before the activity was tested, 50 % inhibition was obtained with about $10 \mu\text{M}$ -EDTA.

The Michaelis constants for β -hydroxybutyric acid and NAD^+ were determined with crude enzyme extract. Each estimation was made at a saturating concentration of the other component. From the results of Carr & Lascelles (1961) and of Doudoroff, Merrick & Contopoulou (1961) only the D-isomer is acted on. The results gave K_m values (at 37°) of

Table 6. *Effect of enzyme inhibitors on β -hydroxybutyrate dehydrogenase*

Sodium DL- β -hydroxybutyrate ($10 \mu\text{moles}$), NAD^+ ($1.0 \mu\text{mole}$), tris, pH 8.8 ($300 \mu\text{moles}$), crude enzyme extract ($0.14 \text{ mg. of protein}$) and inhibitor (as stated), in a total volume of 3.0 ml., were incubated at 37° in the thermostatically controlled cell-holder of the Unicam SP. 500. The reaction was started by adding the enzyme and the production of NADH was followed at $340 \text{ m}\mu$. Readings were taken every 30 sec. for 3 min.

Inhibitor	Concn. of inhibitor (mm)	Activity ($\mu\text{m-moles of NADH/min.}$)	Percentage of control activity
None	—	34	100
Sodium <i>p</i> -chloromercuribenzoate	0.5	0	0
NaCN	0.2	29	85
Iodoacetamide	4.0	33	97
EDTA	1.0	19	56
HgSO_4	1.0	0	0
MgSO_4	1.0	31	90
CaCl_2	1.0	39	115

0.63 mM for D- β -hydroxybutyric acid and 0.04 mM for NAD⁺.

The reversibility of the β -hydroxybutyrate-dehydrogenase reaction was shown by a rapid oxidation of NADH when sodium acetoacetate and NAD⁺ were incubated with crude enzyme extract.

Role of aminoacetone in controlling porphyrin and bacteriochlorophyll synthesis

Since aminoacetone is an analogue of δ -aminolaevulinic acid it was thought that it might exert some control on porphyrin and bacteriochlorophyll synthesis. However, aminoacetone was found to be completely inactive in this respect. Aminoacetone did not inhibit δ -aminolaevulate-synthetase activity, nor was it a substrate or an inhibitor of the bacterial δ -aminolaevulate dehydratase or of the δ -aminolaevulate-pyruvate aminotransferase found in extracts of organisms (Gibson, Neuberger & Tait, 1962a). Its addition to organisms illuminated in 'mixture I' did not affect porphyrin or bacteriochlorophyll synthesis and increasing the buffer concentration in 'mixture I', which leads to accumulation of aminoacetone by illuminated organisms, also had no effect on pigment synthesis.

DISCUSSION

Aminoacetone production by suspensions of organisms. The aminoketone which accumulates when organisms are incubated aerobically in the dark or anaerobically in the light in a medium containing α -oxoglutarate, fumarate and glycine was shown to be aminoacetone. However, the results do not rule out the possibility that other aminoketones may be formed in small amounts. The conditions which give the highest yield of aminoacetone were arrived at empirically and it is not known why they should be so suitable. It may be that aminoacetone synthesis is stimulated by the high buffer concentrations and the relatively high concentrations of Mn²⁺ ions or alternatively that the degradation of aminoacetone is inhibited by these conditions. The possible importance of aminoacetone as a biological intermediate has been indicated by the results mentioned in the introduction but no attempt has been made in this study to extend these findings, nor have any further results been reported in the recent literature. Aminoacetone can be synthesized by two different mechanisms. First, the enzyme δ -aminolaevulate synthetase whose normal function appears to be the formation of δ -aminolaevulate from succinyl-CoA and glycine will also catalyse the formation of aminoacetone from acetyl-CoA and glycine (Gibson *et al.* 1958; Kikuchi *et al.* 1959). Secondly, as indicated by the results of Elliott (1958, 1959, 1960a), there is a direct conversion of threonine

into aminoacetone. The nature of this second reaction has been elucidated by our discovery of a L-threonine dehydrogenase in extracts of *Rps. spheroides*. Although the mechanism by which aminoacetone is produced by whole organisms under the conditions used has not been directly tested, it would appear to be by the first pathway. The yields of aminoacetone are highest at pH values below 7, whereas the L-threonine dehydrogenase has only a very low activity at acid and neutral pH, its optimum being at pH 8.8–8.9. There are certain objections to the proposition that aminoacetone is synthesized by δ -aminolaevulate synthetase since when acetate replaces α -oxoglutarate and fumarate in the aerobic incubation medium no aminoacetone is produced. However, acetate is relatively poorly utilized by these organisms for growth (A. Neuberger & G. H. Tait, unpublished work).

Aminoacetone is produced aerobically in the dark, where no δ -aminolaevulate, porphyrin or bacteriochlorophyll are synthesized, and also anaerobically in the light where porphyrin and bacteriochlorophyll are formed. It had previously been proposed that the inability of organisms grown aerobically in the dark to form porphyrins and bacteriochlorophyll could be explained by the low activity of δ -aminolaevulate synthetase (Lascelles, 1959) or by the presence of an inhibitor (Kikuchi, Kumar, Talmage & Shemin, 1958). Since aminoacetone is formed under these conditions where no δ -aminolaevulate or porphyrins are produced, and since this aminoketone is at least likely to be formed by δ -aminolaevulate synthetase, the low activity of this enzyme or its inhibition is probably not the only explanation for the absence of δ -aminolaevulate, porphyrin and bacteriochlorophyll under aerobic conditions in the dark. It is proposed that at least an additional factor governing the production of δ -aminolaevulate is the stationary concentration of succinyl-CoA in the cell. Thus it is possible that under aerobic conditions succinyl-CoA is not formed at a sufficient rate or that it is utilized too quickly in other reactions for it to be converted into δ -aminolaevulate. At low oxygen pressures in the dark or anaerobically in the light the organisms can form porphyrins and bacteriochlorophyll and this might be due, in addition to the increase in δ -aminolaevulate-synthetase activity shown by Lascelles (1959), to changes in the rates of reactions involving succinyl-CoA, making some of this compound available to condense with glycine to form δ -aminolaevulate. The answer to the control of pigment synthesis is probably not as simple as this, since Gibson, Neuberger & Tait (1962a, b) have considered evidence for other control mechanisms. In favour of the above hypothesis being at least on the

correct lines is the observation of Urata & Granick (1961) that when glycine plus any one of a number of the intermediates of the tricarboxylic acid cycle are incubated aerobically with liver homogenates only aminoacetone, but no δ -aminolaevulate, can be detected. It seems likely that aminoacetone is synthesized by δ -aminolaevulate synthetase from acetyl-CoA and glycine and that its synthesis is dependent mainly on the availability of acetyl-CoA. The pathway from threonine may also be operative under certain conditions. The results also show that aminoacetone does not interfere with any of the enzymic reactions of δ -aminolaevulate nor has it any effect in whole organisms on porphyrin or bacteriochlorophyll production.

L-Threonine dehydrogenase. Although the occurrence of an enzyme capable of dehydrogenating threonine to aminoacetone was inferred by Elliott (1960a) on the basis of his experiments with whole organisms, this is the first time that the enzyme has been demonstrated in cell-free systems and some of its properties studied. There are two possible ways, also considered by Elliott (1960a), in which L-threonine can be converted into aminoacetone (Scheme 1). The present work has shown that 1-aminopropan-2-ol is not converted into aminoacetone nor does it inhibit the production of aminoacetone from L-threonine. For these reasons it may be assumed that pathway (I) represents the mechanism of action of L-threonine dehydrogenase. α -Amino- β -oxobutyric acid has a half-life of less than 1 min. at pH 7 (Laver, Neuberger & Scott, 1959) and therefore no enzyme need be present for its decarboxylation to aminoacetone. The possibility remains that the decarboxylation occurs simultaneously with the dehydrogenation on the enzyme surface by a concerted reaction similar to that put forward by Ingraham (1959).

Serine, γ -amino- β -hydroxybutyric acid and 1-aminopropan-2-ol are inactive as substrates or inhibitors, but hydrogen carbonate inhibits the reaction markedly. From these results it would appear that, apart from the β -hydroxyl group, both the carboxyl and the methyl groups of

threonine are essential for the compound to be a substrate. The enzyme is completely specific for the isomers of α -amino- β -hydroxybutyric acid having the D-configuration on the β -carbon atom, but the configuration on the α -carbon atom does not affect activity to such a marked extent. Thus D-allothreonine, where the amino group is in a cisoid position with regard to the hydroxyl group, is acted on more slowly than L-threonine where the amino group and the hydroxyl group are in a transoid relationship to one another.

The inhibitor studies suggest that both thiol groups and a metal ion are essential for enzymic activity, but that pyridoxal phosphate is not involved. Complete description of the properties of the enzyme will have to await more extensive purification.

Since the metabolic importance of aminoacetone is still unknown the significance of the L-threonine-dehydrogenase reaction is likewise unknown. It may simply be a mechanism for the degradation of threonine, the aminoacetone formed being excreted as a waste product. Since the K_m for L-threonine is high—74 mM—it may be that the enzyme is only operative to remove excess of L-threonine. This may be of some importance in *Rps. spheroides* at least under some conditions since addition of L-threonine interferes with growth and bacteriochlorophyll synthesis by competitively inhibiting homoserine dehydrogenase, one of the enzymes in the pathway of biosynthesis of both threonine and methionine (Gibson, Matthew, Neuberger & Tait, 1961). A threonine dehydrogenase has been detected in extracts of *Staph. aureus* (W. H. Elliott & M. Green, personal communication) and its presence in mammalian liver is indicated by the results of Urata & Granick (1961), so that the enzyme may be of widespread occurrence.

D- β -Hydroxybutyrate dehydrogenase. In the present work a soluble D- β -hydroxybutyrate dehydrogenase utilizing NAD⁺ has been found in cell-free extracts. Crude enzyme extracts were shown to reduce NAD⁺ about 100 times as fast in the presence of β -hydroxybutyric acid as in the presence of threonine, and it was thought that both activities might be due to a single enzyme acting on β -hydroxybutyric acid and its α -substituted derivatives. The difference in the rates of reaction could be caused by steric hindrance by the α -amino group of threonine. If this were so the enzyme would have to be specific for the D-isomer of β -hydroxybutyric acid as shown by the results above on the substrate specificity of the L-threonine dehydrogenase. Recently, Carr & Lascelles (1961) have shown that extracts of *Rps. spheroides* dehydrogenate the D- but not the L-isomer of β -hydroxybutyric acid, and Doudoroff *et al.* (1961)

α -amino- β -oxobutyric acid

1-aminopropan-2-ol

Scheme 1

have described a D- β -hydroxybutyrate dehydrogenase with properties very similar to the one described here in the related micro-organism *Rhodospirillum rubrum*. Some of our results are compatible with the assumption that both reactions are catalysed by the same enzyme, but other results suggest that two different enzymes are involved. Further studies are required before this problem can be resolved.

In the report by Williamson, Mellanby & Krebs (1962) on the partial purification of the D- β -hydroxybutyrate dehydrogenase from *Rps. spheroides* it is stated that the enzyme is not inhibited by EDTA. In our study a marked inhibition was found, and this was greater when EDTA and enzyme were preincubated before addition of substrate or coenzyme. Enzymes with similar activities in *Rsp. rubrum* (Doudoroff *et al.* 1961) and in *Bacillus megaterium* (Gavard, Combre & Tuffet, 1960) are both inhibited by EDTA.

We cannot explain this discrepancy, but it may be relevant that the conditions of growth of the organisms and the extent of purification of the enzyme were different in the two sets of experiments. Williamson *et al.* (1962) grew their organisms aerobically in the dark in the presence of acetate whereas our enzyme was obtained from organisms grown anaerobically in the light on malate. In addition, the Oxford workers purified their enzyme tenfold, whereas in the present study crude enzyme was used. The inhibition found with EDTA in the present work does not necessarily show requirement for metal ions by the enzyme, as other enzymes, e.g. aminolaevulinate dehydratase (Gibson, Neuberger & Scott, 1955), for which no metal ion has been indicated, are markedly inhibited by EDTA.

SUMMARY

1. A non-acidic aminoketone accumulates when *Rhodopseudomonas spheroides* is incubated aerobically in the dark or anaerobically in the light in a medium containing α -oxoglutarate, fumarate and glycine. Aerobically in the dark the accumulation is increased by the presence of Mn^{2+} ions in the medium. The aminoketone was purified and identified as aminoacetone.

2. A L-threonine dehydrogenase, which converts threonine into aminoacetone in the presence of NAD^+ , has been detected in extracts of this organism. The enzyme has been partially purified and its properties have been studied.

3. The enzyme will not utilize $NADP^+$ in place of NAD^+ , and it is specific for the isomers of α -amino- β -hydroxybutyric acid having the D-configuration on the β -carbon atom. 1-Aminopropan-2-ol is not an intermediate. The mechanism of the reaction is discussed.

4. A soluble β -hydroxybutyrate dehydrogenase, obtained from the same organism, and having properties similar to that of the L-threonine dehydrogenase, has also been studied. The evidence obtained is not considered sufficient to decide whether or not both activities investigated are catalysed by the same enzyme.

We are grateful to Dr D. F. Elliott of the National Institute for Medical Research, Mill Hill, London, N.W. 7, for the gift of the isomers of α -amino- β -hydroxybutyric acid. One of us (G.H.T.) wishes to thank the Carnegie Trust for the Universities of Scotland for a Scholarship.

REFERENCES

- Carr, N. G. & Lascelles, J. (1961). *Biochem. J.* **80**, 70.
 Doudoroff, M., Merrick, J. M. & Contopoulou, R. (1961). *Fed. Proc.* **20**, 272.
 Elliott, W. H. (1958). *Biochim. biophys. Acta*, **29**, 446.
 Elliott, W. H. (1959). *Nature, Lond.*, **183**, 1051.
 Elliott, W. H. (1960a). *Biochem. J.* **74**, 478.
 Elliott, W. H. (1960b). *Nature, Lond.*, **185**, 467.
 Gavard, R., Combre, C. & Tuffet, A. (1960). *C.R. Acad. Sci., Paris*, **251**, 1931.
 Gibson, K. D., Laver, W. G. & Neuberger, A. (1958). *Biochem. J.* **70**, 71.
 Gibson, K. D., Matthew, M., Neuberger, A. & Tait, G. H. (1961). *Nature, Lond.*, **192**, 204.
 Gibson, K. D., Neuberger, A. & Scott, J. J. (1955). *Biochem. J.* **61**, 618.
 Gibson, K. D., Neuberger, A. & Tait, G. H. (1962a). *Biochem. J.* **83**, 539.
 Gibson, K. D., Neuberger, A. & Tait, G. H. (1962b). *Biochem. J.* **83**, 550.
 Gomori, G. (1955). In *Methods in Enzymology*, vol. 1, p. 138. Ed. by Colowick, S. P. & Kaplan, N. O. New York: Academic Press Inc.
 Hughes, D. E. (1951). *Brit. J. exp. Path.* **32**, 97.
 Ingraham, L. L. (1959). In *Enzymes in Health and Disease*, p. 37. Ed. by Greenberg, D. M. & Harper, H. A. Springfield, Ill.: Charles C. Thomas.
 James, A. T. & Martin, A. J. P. (1952). *Biochem. J.* **50**, 679.
 Kaplan, N. O. & Ciotti, M. M. (1956). *J. biol. Chem.* **221**, 823.
 Kikuchi, G., Kumar, A. & Shemin, D. (1959). *Fed. Proc.* **18**, 259.
 Kikuchi, G., Kumar, A., Talmage, P. & Shemin, D. (1958). *J. biol. Chem.* **233**, 1214.
 Lascelles, J. (1956). *Biochem. J.* **62**, 78.
 Lascelles, J. (1959). *Biochem. J.* **72**, 508.
 Laver, W. G., Neuberger, A. & Scott, J. J. (1959). *J. chem. Soc.* p. 1483.
 Layne, E. (1957). In *Methods in Enzymology*, vol. 3, p. 447. Ed. by Colowick, S. P. & Kaplan, N. O. New York: Academic Press Inc.
 Lineweaver, H. & Burk, D. (1934). *J. Amer. chem. Soc.* **56**, 658.
 Mauzerall, D. & Granick, S. (1956). *J. biol. Chem.* **219**, 435.
 Mijovic, M. P. V. & Walker, J. (1960). *J. chem. Soc.* p. 909.
 Neilands, J. B., Neuberger, A. & Scott, J. J. (1960). *J. Amer. chem. Soc.* **82**, 214.

- Nemeth, A. M., Russell, C. S. & Shemin, D. (1957). *J. biol. Chem.* **229**, 415.
- Neuberger, A. & Tait, G. H. (1960). *Biochim. biophys. Acta*, **41**, 164.
- Pullman, M. E., Colowick, S. P. & Kaplan, N. O. (1952). *J. biol. Chem.* **194**, 593.
- Shemin, D. (1955). In *Ciba Foundation Symp. Porphyrin Biosynthesis and Metabolism*, p. 4. London: J. and A. Churchill Ltd.
- Shemin, D. & Russell, C. S. (1953). *J. Amer. chem. Soc.* **75**, 4873.
- Shemin, D., Russell, C. S. & Abramsky, T. (1955). *J. biol. Chem.* **215**, 613.
- Smith, I. (1958). In *Chromatographic Techniques*, p. 36. Ed. by Smith, I. London: William Heinemann Medical Books Ltd.
- Urata, G. & Granick, S. (1961). *Biochem. biophys. Res. Commun.* **4**, 96.
- Warburg, O. & Christian, W. (1942). *Biochem. Z.* **310**, 384.
- Williamson, D. H., Mellanby, J. & Krebs, H. A. (1962). *Biochem. J.* **82**, 90.

Biochem. J. (1962) **84**, 328

A Magnesium Yeast and its Properties

By E. J. CONWAY AND MARY E. BEARY

Department of Biochemistry, University College, Dublin

(Received 4 January 1962)

This is one of three papers describing some of the chief properties of baker's yeast in which almost all the K^+ ion of the normal yeast cells has been replaced by the NH_4^+ ion (Conway & Breen, 1945), the Na^+ ion (Conway & Moore, 1954), or, as here, by the Mg^{2+} ion. The ammonium yeast was produced by fermentation in the presence of ammonium chloride and the sodium yeast by a series of fermentations in sodium citrate. In these ammonium and sodium yeasts 98% of the normal potassium content was replaced by NH_4^+ or Na^+ ions. The yeast cells were undamaged by these procedures, being readily restored to the functional condition of normal potassium yeast by fermentation for an hour or so in the presence of potassium chloride solution.

In the process of forming the magnesium yeast, by fermentation in magnesium acetate, large amounts of magnesium were accumulated through the activity of the cation-carrier (or physiological potassium-carrier) in the yeast-cell wall (Conway & Beary, 1958). Almost no concomitant loss of potassium occurred from the yeast cells as it did during the formation of sodium yeast, so that cells very rich in magnesium but with an almost normal potassium content resulted. It appeared, therefore, that this procedure would not produce magnesium yeast similar to the ammonium yeast and the sodium yeast. The difficulty in producing such a magnesium yeast was overcome by passing first from a normal potassium yeast to a sodium yeast and then fermenting this repeatedly in the presence of magnesium acetate. In this way the high sodium content of the sodium yeast was replaced by magnesium.

Some of the principal properties of this magnesium yeast have been studied, as well as those of the magnesium-rich yeast with normal potassium content, obtained by repeated fermentations in the presence of magnesium acetate.

METHODS

Preparation of sodium yeast. This was carried out by the method of Conway & Moore (1954) from baker's yeast (supplied by the Irish Yeast Co.) which was used throughout after being washed twice.

Preparation of magnesium yeast with minimal potassium content. The sodium yeast was suspended (1:20, w/v) in a solution containing (final concentrations) 0.2M-magnesium acetate, 5% (w/v) glucose and 25 mM-tris-HCl buffer, pH 7.2, at room temperature (18°). The suspending fluid was changed after centrifuging, every 2 hr. throughout the day. This procedure was repeated the next day. The magnesium, potassium and sodium contents of this yeast were determined at regular intervals, and the procedure was continued until no further appreciable change occurred.

Preparation of a magnesium-rich yeast with normal potassium content. This was prepared as described by Conway & Beary (1958), at room temperature, and the samples were bubbled with oxygen during the preparation. The pH of the suspension mixture was maintained at 7.2 by the addition of tris buffer (final concentration 25 mM), the pH being adjusted with 0.1 N-HCl.

Sodium and potassium determinations. Both were determined with the Beckman flame photometer. A sample of twice-washed yeast was boiled in 10–20 vol. of water, centrifuged, and the supernatant diluted as required to obtain a convenient reading on the photometer. The standards used for comparison were in the range 1.25–5.0 mg./l. for both sodium and potassium.