nucleotide was increased, and, assuming that the product was derived from oligonucleotides and deoxyribonucleoside triphosphates, the synthesis amounted to 74% of the deoxyribonucleic acid primer.

4. The nucleotidyltransferase was inhibited and the lag phase extended by 3'-phosphoryl-terminal oligonucleotides from a splenic-deoxyribonuclease digest of deoxyribonucleic acid.

5. The compounds obtained by removal of terminal phosphoryl groups from the 3'- and 5'-phosphoryl-terminal oligonucleotides by bacterial alkaline phosphatase abolished the lag phase, and they supported synthesis at a level similar to that observed under standard assay conditions.

6. It was concluded that 5'-phosphoryl-terminal oligonucleotides associate by specific hydrogen-bonding with single-stranded regions of the deoxyribonucleic acid primer and stimulate synthesis of polydeoxyribonucleotide by providing 3'-hydroxyl groups to initiate polymerization along the adjacent single-stranded template. The 5'-phosphoryl-terminal group on these oligonucleotides is partly responsible for the stimulation observed. Inhibition by 3'-phosphoryl-terminal oligonucleotides is attributable to competition with 3'-hydroxyl groups formed on the deoxyribonucleic acid primer by deoxyribonuclease during the nucleotidyltransferase assay, and the lag phase observed under standard conditions is a period of production of stimulatory end groups in the deoxyribonucleic acid primer.

I wish to thank Professor J. N. Davidson, F.R.S., for his interest and support. The investigation was aided by grants from the British Empire Cancer Campaign, the Jane Coffin Childs Memorial Fund for Medical Research and the Rankin Fund of the University of Glasgow, which are gratefully acknowledged. I also wish to thank Dr R. M. S. Smellie for many helpful discussions, and Mr C. Macleod and Miss Helen Moss for expert technical assistance.

REFERENCES

- Allen, R. J. L. (1940). *Biochem. J.* **34**, 858.
 Bollum, F. J. (1960a). *J. biol. Chem.* **235**, 2399.
 Bollum, F. J. (1960b). *J. biol. Chem.* **235**, PC18.
 Burton, K. (1959). In *Data for Biochemical Research*, p. 74. Ed. by Dawson, R. M. C., Elliott, D. C., Elliott, W. H. & Jones, K. M. London: Oxford University Press.
 Davidson, J. N. (1961). *Symp. biochem. Soc.* **21**, 29.
 Dixon, M. & Webb, E. C. (1958). *Enzymes*, pp. 71-73. London: Longmans, Green and Co.
 Doty, P., Boedtker, H., Fresco, J. R., Haselkorn, R. & Litt, M. (1959). *Proc. nat. Acad. Sci., Wash.*, **45**, 482.
 Garen, A. & Levinthal, C. (1960). *Biochim. biophys. Acta*, **38**, 470.
 Gray, E. D., Weissman, S. M., Richards, J., Bell, D., Keir, H. M., Smellie, R. M. S. & Davidson, J. N. (1960). *Biochim. biophys. Acta*, **45**, 111.
 Griswold, B. L., Humoller, F. L. & McIntyre, A. R. (1951). *Analyt. Chem.* **23**, 192.
 Heppel, L. A., Harkness, D. R. & Hilmo, R. J. (1962). *J. biol. Chem.* **237**, 841.
 Jordan, D. O. (1960). *The Chemistry of the Nucleic Acids*, pp. 221-223, 249-256. London: Butterworths.
 Keir, H. M. & Aird, G. L. (1962). *Biochem. J.* **84**, 44P.
 Keir, H. M., Binnie, B. & Smellie, R. M. S. (1962). *Biochem. J.* **82**, 493.
 Koerner, J. F. & Sinsheimer, R. L. (1957a). *J. biol. Chem.* **228**, 1039.
 Koerner, J. F. & Sinsheimer, R. L. (1957b). *J. biol. Chem.* **228**, 1049.
 Laskowski, M. (1961). In *The Enzymes*, 2nd ed., vol. 5, p. 123. Ed. by Boyer, P. D., Lardy, H. & Myrback, K. New York: Academic Press Inc.
 Mantsavinos, R. & Canellakis, E. S. (1959). *Cancer Res.* **19**, 1239.
 Rosenkranz, H. S. & Bendich, A. (1961). *Biochim. biophys. Acta*, **51**, 186.
 Sarkar, N. K. (1961). *Arch. Biochem. Biophys.* **93**, 328.
 Sinsheimer, R. L. (1959). *J. molec. Biol.* **1**, 37, 43.
 Smellie, R. M. S. (1961). *J. Chim. phys.* **58**, 965.
 Smellie, R. M. S. & Eason, R. (1961). *Biochem. J.* **80**, 39P.
 Staehelin, M. (1961). *Biochim. biophys. Acta*, **49**, 20.
 Vanecko, S. & Laskowski, M. (1961). *J. biol. Chem.* **236**, 3312.

Biochem. J. (1962) **85**, 276

The Degradation of Acid-Soluble Collagen by Rat-Liver Preparations

By D. MARY FRANKLAND AND C. H. WYNN

Rheumatism Research Centre, Clinical Sciences Building, University of Manchester, Manchester 13

(Received 9 April 1962)

In the development of normal connective tissue, the mature collagen fibres are stable and, metabolically, relatively inert (Neuberger, Perrone & Slack, 1951). In certain physiological and pathological conditions, however, a rapid turnover of collagen has been demonstrated. Thus Harkness &

Harkness (1954) showed that the collagen of the involuting uterus was rapidly resorbed, and Jackson (1957) showed that, although connective tissue growth was stimulated by carrageenin, the newly formed granulation tissue was metabolically unstable and was rapidly degraded, being replaced

by adipose tissue. It seems probable therefore that the mammal possesses a mechanism, which in normal conditions is latent, for the rapid turnover of collagen.

The existence of a bacterial collagenase is well known and contemporary knowledge of this enzyme has been reviewed by Mandl (1961). It has been established that many mammalian proteolytic enzymes are capable of degrading gelatin, but most evidence suggests that they are inactive against native undenatured collagen. Conflicting reports of the nature of the action of trypsin on collagen have been presented. Thus Hodge, Highberger, Deffner & Schmitt (1960) have postulated that trypsin has a depolymerizing action on linear polymers of tropocollagen, but this 'proteolytic activity is confined to very limited susceptible regions extending beyond the triple-helix body of the native tropocollagen macromolecule'. However, Kühn, Kühn & Hannig (1961) have attributed the action of trypsin on collagen to impurities in the collagen used.

Lucy, Dingle & Fell (1961) and Dingle (1961) have postulated that intracellular proteases may have a role in the degradation of cartilage matrix. They demonstrated that an enzyme, localized in rat-liver lysosomes, can split the non-collagenous protein-polysaccharide complex of cartilage. Woessner & Boucek (1961*a, b*) have also implicated lysosome particles in the degradation of connective tissue. However, intracellular proteases are reported to be inactive against native collagen (Orekhovitch, Shpikiter, Mazurov & Kounina, 1960).

The present investigation was an attempt to characterize the mammalian system by which collagen is degraded and to determine the intracellular localization of this system. Preliminary communications of part of this work have already been made (Frankland & Wynn, 1961, 1962).

MATERIALS AND METHODS

Acid-soluble-collagen solutions. A portion (1 g.) of calf-skin dermis, which had previously been exhaustively extracted with neutral salt solutions and freeze-dried (see Wood & Keech, 1960), was cut into small pieces (approx. 2 mm. square) and extracted for 18 hr. with 60 ml. of 0.5*N*-acetic acid at 4°. The suspension was filtered on a coarse sintered-glass funnel and the extraction was repeated. Combination of the two filtrates gives an approximately 0.16% solution of acid-soluble collagen which can be stored at 4° for at least 2 weeks without appreciable change in physical and chemical properties. For some experiments described below the more concentrated acid-soluble collagen obtained by a single extraction of dermis was used.

¹⁴C-Labelled collagen solutions. The ability of injected carrageenin solutions to stimulate the rapid formation of

easily separated connective tissue in comparatively large amounts (Robertson & Schwartz, 1953) provides a useful method for the synthesis of labelled collagens.

Male and female guinea pigs were used, their weights at the time of injection varying between 350 and 450 g. The animals were fed on Diet no. SG1 (Oxoid Division, Oxo Ltd.), hay and greens. Ascorbic acid was supplemented by giving orally, every other day, 50 mg. dissolved in 10% (w/v) sucrose.

A portion (5 ml.) of a 1% (w/v) sterile solution of carrageenin (Seaplant Corp., New Bedford, Mass., U.S.A.) in 0.9% NaCl was injected subcutaneously into the lower abdominal region, under ether anaesthesia. Five days later daily intraperitoneal injection of 1 ml. of generally labelled L-¹⁴C-proline (15 μ C/ml.; specific activity 5 mc/m-mole) in 0.9% (w/v) NaCl was commenced. On the tenth day after the injection of carrageenin, the animals were killed and the new tissue growth between the dermis and the abdominal wall was dissected out. Acid-soluble collagen was isolated from this tissue by the method of Jackson (1957), the collagen being purified by precipitation and re-solution three times. The specific activity of the purified acid-soluble collagen (0.2% solution) was 236 counts/min./mg. of protein.

Viscometric determination of collagen degradation. Previous workers have shown that the effect of proteolytic enzymes on collagen may be conveniently measured viscometrically (Gallop, Seifter & Meilman, 1957; Hodge *et al.* 1960). Accordingly, this method was used in the earlier experiments to investigate the effect of various purified extracts and subcellular-particle preparations of rat liver on collagen. A solution (1.5 ml.) of acid-soluble collagen and 1.5 ml. of rat-liver preparation buffered with 0.1*M*-sodium acetate buffer, pH 5.0, previously warmed for 5 min. at 35°, were mixed, and the viscosity of a 2 ml. sample of the mixture was measured by flow-time determinations at intervals for up to 1 hr. The pH of the final incubation mixture was 4.0 $\pm$ 0.1. With subcellular-particle preparations sucrose was also present in the incubation mixture at a final concentration of 0.125*M*.

All viscosity measurements were made with a miniature U-tube viscometer (BS/U/M; size M2 or M3) in a constant-temperature water bath at 35°.

Study of degradation of ¹⁴C-labelled collagen. In the study of the degradation of collagen the following method was used: A solution (0.5 ml.) of ¹⁴C-labelled acid-soluble collagen was preincubated at 35° and mixed with 0.5 ml. of the rat-liver preparation under study, adjusted so that it contained 0.1*M*-sodium acetate buffer, pH 5.0. After incubation for 18 hr. at 35°, 1 ml. of 99–100% (v/v) ethanol was added, the tubes were kept at room temperature for 30 min. and then centrifuged at 59 000*g* for 20 min. at 0° in a Spinco ultracentrifuge model L (rotor no. 40). Preliminary experiments revealed that the collagen was completely precipitated under these conditions. A sample (0.2 ml.) of the supernatant was pipetted on aluminium planchets, spread by applying a piece of lens tissue of the same internal diameter as the planchets, dried with an infrared lamp and the radioactivity of the sample counted by an end-window counter until the s.e. of the determination when corrected for background radioactivity was less than 1%. The release of radioactivity into the supernatant is approximately proportional to the enzyme concentration and this procedure provides a semiquantitative method

of following the conversion of collagen into a form not precipitable by 50% ethanol under the conditions specified.

Release of hydroxyproline-containing material from acid-soluble collagen. Wood & Keech (1960) have shown that acid-soluble collagen is precipitated from neutral solutions of suitable ionic strength at 30°.

Acid-soluble collagen solutions were incubated with rat-liver preparations under the same conditions as specified for the viscometric assay for periods up to 24 hr. The solutions were then cooled to 30° and NaOH-KH₂PO₄ buffer was added so that the final phosphate concentration was 0.04 M and the pH was 7.0. After being incubated for 2 hr. at 30° the solutions were centrifuged at 20 000g for 30 min. at 15° and a sample of the supernatant was hydrolysed in 6N-HCl in a sealed tube for 18 hr. at 100°. The hydroxyproline content of the hydrolysate was determined by the method of Stegemann (1958).

Preparation of subcellular fractions. Rat-liver homogenates were divided into nuclear, heavy-mitochondrial, light-mitochondrial, microsomal and supernatant fractions by ultracentrifuging by the method of de Duve, Pressman, Gianetto, Wattiaux & Appelmans (1955). Before assay each fraction was subjected to ultrasonic disintegration.

Determination of acid-phosphatase activity. The acid-phosphatase activity of the subcellular fractions was determined by the method of Gianetto & de Duve (1955).

Determination of cathepsin activity. The cathepsin activity of the various rat-liver extracts was determined by the method of Anson (1937), with denatured haemoglobin as substrate. The results are expressed as m-equivalents of tyrosine released from the haemoglobin. For present purposes 'cathepsins' is thus taken to mean enzymes in rat liver which hydrolyse haemoglobin in acid solution.

Protein determination. Protein was determined by the method of Lowry, Rosebrough, Farr & Randall (1951), with normal human serum as a standard; it was assumed that the serum contains 70 mg. of protein/ml. Nitrogen was determined by a micro-Kjeldhal method.

EXPERIMENTAL AND RESULTS

Action of rat-liver extract on acid-soluble collagen

Recent evidence (Lucy *et al.* 1961; Woessner & Boucek, 1961a, b) has suggested that the lysosomal particle contains enzymes involved in the degradation of connective tissue and it has been reported that the soluble hydrolytic enzymes in this particle have an acid pH optimum (de Duve, 1959). Therefore in preliminary attempts to locate a collagenolytic enzyme in rat liver the final pH of the reaction mixture was adjusted to 4.0.

Stage 1: preparation of liver extract. Livers from six adult albino rats were pooled and chilled, pressed through a fine sieve and 20 g. of the resultant pulp was homogenized in 100 ml. of 0.1 M-sodium acetate adjusted to pH 5.0, with 0.1 M-acetic acid, in a glass-Teflon tissue homogenizer; the homogenate was kept at 0° throughout. Disruption of any intact membranes was achieved by alternately freezing (-70°) and thawing (37°) the homogenate ten times. The resulting suspension was centrifuged at 30 000 rev./min.

(80 000g) for 1 hr. at 1° in a Spinco model L ultracentrifuge (rotor no. 30) and the clear supernatant (designated S₁) was decanted and stored at 0°.

Fig. 1 shows the effect of the supernatant, S₁, on the viscosity of acid-soluble collagen at 35°. There was a marked decrease in the viscosity over the first 20 min. of incubation, this rate of decrease falling off with longer incubation periods. The viscosities of both the collagen solution and the supernatant S₁, when incubated separately under the appropriate conditions, were constant.

Gallop *et al.* (1957) have shown that, when bacterial collagenase is assayed viscometrically with ichthyocol as substrate, the results are best calculated by making a semilogarithmic plot of specific viscosity against time, when a linear graph is obtained. Fig. 1 also shows the action of S₁ on collagen plotted in this manner, when a linear relationship is obtained initially. To compare the activities of different preparations and fractions we have defined one activity unit as that amount of enzyme which under the specified conditions of assay (see the Materials and Methods section) gives a decrease in the log(specific viscosity) of 0.01 in 10 min.

Stage 2: acetone fractionation of supernatant S₁. The supernatant (100 ml.) was cooled to -5° and ice-cold acetone was added dropwise, with constant stirring, until the acetone concentration was 30% (v/v). The mixture was allowed to stand for 30 min. at -5° before the precipitated protein was centrifuged off at -5° at 1610g for 20 min. The supernatant was decanted and the fractionation continued in the same manner, the acetone concentration being made up to 50% and 70% successively.

The three precipitates obtained were each resuspended in 20 ml. portions of 0.1 M-sodium acetate, adjusted to pH 5.0 with 0.1 M-acetic acid, and dialysed at 0° for 18 hr. against 2 l. of the same buffer to remove traces of acetone. The fractions were clarified by centrifuging at 25 000g for

Fig. 1. Change of viscosity with time during the action of rat-liver extract, S₁, on acid-soluble collagen under incubation conditions specified in the text. ●, Specific viscosity; ▲, log(specific viscosity).

20 min. at 0° and stored at 4° before use. These 0–30%, 30–50% and 50–70% acetone fractions were designated P₁, P₂ and P₃ respectively.

Both P₁ and P₂, when incubated with collagen, effected a marked decrease in the viscosity of the incubation mixture with time, and when the results were plotted on a semilogarithmic scale the graph was linear over the first 15 min. of incubation. P₃ produced an increase in viscosity over this period. The viscosities of collagen and the fractions when incubated separately were constant over this period.

Although both P₁ and P₂ show activity, Table 1 shows that P₁ is more active than P₂, and when the results are expressed in terms of the protein content of these fractions this difference in activity is very marked. P₂ was therefore discarded and P₁ was purified further by ammonium sulphate fractionation.

Stage 3: ammonium sulphate fractionation of P₁. The 0–30% acetone fraction was cooled to 0° and solid (NH₄)₂SO₄ was added to give a final concentration equivalent to 25% saturation [calculated from the nomogram of Dixon (1953) without correcting for temperature]. The solution was allowed to stand for 3 hr. at 0° and centrifuged at 2000 g for 30 min. at 0°. The supernatant was decanted off and further portions of solid (NH₄)₂SO₄ were added to give final concentrations equivalent to 40% and 60% saturation. All the precipitates were redissolved in 5 ml. portions of 0.1 M-sodium acetate adjusted to pH 5.0 with 0.1 M-acetic acid and dialysed at 4° against two changes of the same buffer (total volume 4 l.). These fractions were designated P₁A, P₁B and P₁C respectively.

Both fractions P₁A and P₁B possess collagenolytic activity but fraction P₁C was inactive against collagen.

Comparison of the collagenolytic and catheptic activity of the various enzyme fractions

Table 1 summarizes the protein content, collagenolytic activity and catheptic activity of the original extract, acetone fractions and ammonium sulphate fractions.

The 0–25% ammonium sulphate fraction, P₁A, is the most active preparation when expressed in terms of protein content, and is 15 times as active as the original supernatant. There is a considerable loss of activity during the purification procedure and the recovery of the total activity in this fraction is only 11%. The 25–40% ammonium sulphate fraction, P₁B, contains 10% of the total activity, and as its specific activity is only slightly less than that of the 0–25% fraction these fractions may be reasonably combined for further study.

The catheptic activity of the original supernatant was extremely small and not capable of accurate measurement, but the concentration in the acetone fractions was measured. The distribution of catheptic activity of these fractions is different from that of the collagenolytic activity. Thus the 0–30% fraction which is the most active towards collagen is the least active to haemoglobin. No detectable catheptic activity was found in the ammonium sulphate fractions.

Intracellular distribution of collagenolytic activity

Table 2 shows the intracellular distribution of collagenolytic activity as determined viscometrically and also by using ¹⁴C-labelled collagen. The intracellular distribution of acid-phosphatase activity is included for comparison.

The distribution of the collagenolytic activity of the fractions, as determined by either method, follows closely the acid-phosphatase activity, the highest relative specific activity being found in the light-mitochondrial or lysosomal fraction. The main difference is in the supernatant fraction, which contains considerably more collagenolytic activity than acid-phosphatase activity.

Collagenolytic activity of a concentrated lysosomal extract and its inhibition by thermal denaturation

Lysosomes were prepared from 50 g. of fresh rat liver by the method of de Duve *et al.* (1955) and the final pellet was

Table 1. *Protein content, collagenolytic activity and catheptic activity of rat-liver fractions*

Rat liver was fractionated as described in the text, and collagenolytic and catheptic activities were determined as indicated in the Materials and Methods section.

Fraction	Total protein content (g.)	Collagenolytic activity		Total cathepsin activity (m-equiv. of tyrosine)
		Total activity	Activity/mg. of protein	
Supernatant S ₁	1.68	890	0.56	0
Acetone fractions				
P ₁	0.26	400	1.6	2.2 × 10 ⁻³
P ₂	0.83	220	0.24	1.2 × 10 ⁻³
P ₃	0.33	0	0	2.5 × 10 ⁻³
(NH ₄) ₂ SO ₄ fractions				
P ₁ A	0.012	96	8	0
P ₁ B	0.018	88	4.8	0
P ₁ C	0.024	13	0.56	0

Table 2. Intracellular distribution of acid-phosphatase and collagenolytic activities

The results were obtained by subcellular fractionation of 20 g. of rat liver. The assay methods were as described in the Materials and Methods section. Relative specific activity is defined as the percentage of the total activity found in a fraction divided by the percentage of the total nitrogen in that fraction. Where 'total' activity is specified, this refers to the activity of the whole fraction isolated from 20 g. of rat liver, calculated from an assay of a suitable portion. The collagenolytic activities measured by the ¹⁴ C assay method are given ± s.e., calculated from the difference in s.e. of test and background counts (both determinations being continued for 10 000 counts, i.e. until the s.e. of each is less than 1% of the count).									
Fraction	N content (mg.)	Percentage of total N	Total acid phosphatase (μmoles of phosphate released)	Collagenolytic activity			Relative specific activity		
				¹⁴ C assay method		Viscometric method		Acid phosphatase	¹⁴ C
				'Counts/min. released in 0.2 ml. of supernatant'	Total 'counts/min. released'				
Nuclei	187	34.8	108.7	0.20 ± 0.20	54.4	54	0.65	0.97	0.40
Heavy mitochondria	50.5	9.4	63.4	0.21 ± 0.20	18.5	33	0.82	2.10	0.91
Light mitochondria	14.7	2.7	81.5	0.48 ± 0.20	42.2	35	6.54	9.37	3.34
Microsomes	21.7	4.0	44.5	0.22 ± 0.20	18.5	8	1.94	3.45	0.54
Supernatant	263	49.0	23.8	0.36 ± 0.20	105.4	260	0.90	0.15	1.35

again homogenized in 10 ml. of 0.25 M-sucrose. The homogenate was treated in a MSE Ultrasonic Disintegrator at 20 kcyc./sec. with an output of 60 W for 2 min. at 0° and subsequently clarified by centrifuging at 80 000g for 10 min. at 0°. The clear supernatant was dialysed for 18 hr. at 0° against 2 l. of water. The non-diffusible material was centrifuged at 80 000g for 10 min. at 0° and the collagenolytic activity of the final supernatant assayed viscometrically with the concentrated solution of acid-soluble collagen obtained by a single extraction of dermis (see the Material and Methods section).

Fig. 2. Change of specific viscosity with time during the action of rat-liver lysosomal extract under incubation conditions specified in the text. ●, Collagen control; ○, lysosome control; ■, lysosome extract acting on collagen; ▲, heated lysosome extract (see text) acting on collagen.

Fig. 3. Release of hydroxyproline during the action of rat-liver lysosomal extract under incubation conditions specified in the text. ●, Lysosome extract acting on collagen; ▲, collagen control.

To investigate the effect of heat on the collagenolytic activity of the lysosomal extract, a sample of this extract was heated at 65° in a water bath for 40 min., cooled rapidly to 0° in ice and then centrifuged at 16 000*g* for 15 min. The collagenolytic activity of the supernatant was then assayed.

Fig. 2 shows the results of these viscometric collagenolytic-activity assays. The lysosomal extract possesses considerable activity, but this is abolished by heating to 65°. Both collagen and lysosomal extract when incubated separately under the appropriate conditions show a slight viscosity decrease over 1 hr.

Release of hydroxyproline-containing material. Fig. 3 shows the steady increase in hydroxyproline-containing material released with time on incubation of collagen with lysosomal extract and precipitation of unchanged collagen by warming at neutral pH. After 24 hr. 85 % of the hydroxyproline-containing material was no longer precipitable.

When collagen is incubated in the absence of lysosomes under the appropriate conditions there is little release of hydroxyproline-containing material.

DISCUSSION

The results described in this paper show that there is present in rat liver an enzyme which can hydrolyse acid-soluble collagen at pH 4. The nature and specificity of the enzyme is as yet uncertain but some indirect information is given by the observation that the collagenolytic activity is destroyed by heating to 65° for 40 min. Tallan, Jones & Fruton (1952) have shown that such treatment results in only a slight loss of cathepsin C activity and thus the collagenolytic activity is unlikely to be due to this enzyme. The finding that during fractionation of the rat-liver extract collagenolytic enzymes were partially separated from enzymes which hydrolyse haemoglobin in acid solution provides further evidence against their identity.

The localization of the collagenolytic activity in the lysosome particles of the rat-liver cell is in agreement with recent hypotheses of the mechanism of the degradation of connective tissue. Woessner & Boucek (1961*a, b*) showed that during development of connective tissue in poly(vinyl alcohol)-sponge implants, the activities of enzymes believed to be of lysosomal origin, e.g. acid phosphatase, continue to increase after the DNA and collagen content of the connective tissue have reached their maximum levels. The activities of other enzymes thought to be associated with cell proliferation, e.g. alkaline phosphatase, pass through a maximum and decrease over this period. The finding that lysosomes contain a protease active against the non-collagenous protein-poly-

saccharide complex of cartilage (Lucy *et al.* 1961) provides further evidence that these particles may be implicated in the degradation of connective tissue.

The finding that, after treatment with a lysosomal extract, the collagen is no longer precipitable in the usual manner and that large amounts of hydroxyproline-containing material are present in the supernatant contrasts with the results of Kühn *et al.* (1961), who showed that after trypsin treatment collagen was still precipitable. It is obvious therefore that this lysosomal enzyme has a far more drastic action than trypsin even at far lower concentrations.

Although the enzyme described in the present paper is capable of hydrolysing collagen there is no direct evidence that this is a physiological mechanism for collagen turnover and resorption. The main difficulty in extrapolating these results *in vitro* to situations *in vivo* is that the enzyme acts at pH 4. However, all the hydrolytic enzymes associated with the lysosome have an acid pH optimum and it seems likely that local changes in pH may occur.

SUMMARY

1. A collagenolytic activity which is active at acid pH has been demonstrated in rat-liver extracts.
2. The preliminary purification of this activity by acetone and ammonium sulphate fractionation is reported.
3. The collagenolytic activity has been shown to be lysosomal in origin.
4. Unlike cathepsin C activity, the collagenolytic activity is readily lost on moderate heating. A partial separation of collagenolytic and catheptic activity by acetone fractionation has also been demonstrated.

We thank Professor J. H. Kellgren and Dr G. C. Wood for their generous interest in this work and for their helpful criticism, Miss J. Smith and Miss S. Hatchett for technical assistance, and the Medical Research Council for financial assistance to one of us (D.M.F.).

REFERENCES

- Anson, M. L. (1937). *J. gen. Physiol.* **20**, 565.
de Duve, C. (1959). In *Subcellular Particles*, p. 128. Ed. by Hayashi, T. New York: The Ronald Press Co.
de Duve, C., Pressman, B. C., Gianetto, R., Wattiaux, R. & Appelmans, F. (1955). *Biochem. J.* **60**, 605.
Dingle, J. T. (1961). *Biochem. J.* **79**, 509.
Dixon, M. (1953). *Biochem. J.* **54**, 457.
Frankland, D. M. & Wynn, C. H. (1961). *Biochem. J.* **81**, 25*F*.
Frankland, D. M. & Wynn, C. H. (1962). *Biochem. J.* **84**, 20*F*.
Gallop, P. M., Seifter, S. & Meilman, E. (1957). *J. biol. Chem.* **227**, 891.

- Gianetto, R. & de Duve, C. (1955). *Biochem. J.* **59**, 433.
- Harkness, M. L. R. & Harkness, R. D. (1954). *J. Physiol.* **123**, 492.
- Hodge, A. J., Highberger, J. H., Deffner, G. G. J. & Schmitt, F. O. (1960). *Proc. nat. Acad. Sci., Wash.*, **46**, 197.
- Jackson, D. S. (1957). *Biochem. J.* **65**, 277.
- Kühn, K., Kühn, J. & Hannig, K. (1961). *Hoppe-Seyl. Z.* **326**, 50.
- Lowry, O. H., Rosebrough, N. J., Farr, A. L. & Randall, R. J. (1951). *J. biol. Chem.* **193**, 265.
- Lucey, J. A., Dingle, J. T. & Fell, H. B. (1961). *Biochem. J.* **79**, 500.
- Mandl, I. (1961). *Advanc. Enzymol.* **23**, 163.
- Neuberger, A., Perrone, J. C. & Slack, H. G. B. (1951). *Biochem. J.* **49**, 199.
- Orekhovitch, V. N., Shpikiter, V. O., Mazurov, V. I. & Kounina, O. V. (1960). *Bull. Soc. Chim. biol., Paris*, **42**, 505.
- Robertson, van B. & Schwartz, B. (1953). *J. biol. Chem.* **201**, 689.
- Stegemann, H. (1958). *Hoppe-Seyl. Z.* **311**, 41.
- Tallan, H. H., Jones, M. E. & Fruton, J. S. (1952). *J. biol. Chem.* **194**, 793.
- Woessner, J. F. & Boucek, R. J. (1961a). *Arch. Biochem. Biophys.* **93**, 85.
- Woessner, J. F. & Boucek, R. J. (1961b). *Arch. Biochem. Biophys.* **93**, 95.
- Wood, G. C. & Keech, M. K. (1960). *Biochem. J.* **75**, 588.

Biochem. J. (1962) **85**, 282

Immunochemical Significance of L- and D-Fucose Derivatives

BY G. F. SPRINGER* AND P. WILLIAMSON

Immunochemistry Section, William Pepper Laboratory and Departments of Microbiology and Pathology, Medical School and Graduate School of Medicine, University of Pennsylvania, Philadelphia, U.S.A.

(Received 26 January 1962)

The elucidation of the chemical structures responsible for the serological independence of the various blood-group characters and the specific combining sites of their complementary antibodies constitutes a fundamental challenge of immunochemistry. Knowledge of the structures determining specificity may yield information on the forces decisive in antigen-antibody interactions, as well as on genetical mechanisms controlling blood-group characters.

It was first observed by Watkins & Morgan (1952) and by Morgan & Watkins (1953) that L-fucose inhibited the agglutination of human blood-group O erythrocytes by anti-H(O) agglutinins in eel serum and in seed extracts of *Lotus tetragonolobus*. This inhibition was shown to possess considerable specificity in that L-fucose was the only active sugar of the four constituent monosaccharides of blood-group mucoids. Of numerous other sugars only 2-deoxy-L-fucose, 6-deoxy-L-talose, D-arabinose and D-fructose showed some, though less, activity; D-fucose was inactive. Watkins & Morgan noted that methyl α -L-fucopyranoside was 4-5 times as active as L-fucose; the β -glycoside was less active than the parent compound. Kuhn & Osman (1956) confirmed and extended the earlier findings of Watkins & Morgan and postulated the following necessary conditions

for the serological activity of fucosides: (1) The fucose residue must belong to the L-series; (2) it must be in the pyranose form; (3) if linked, it must be present in α -glycosidic linkage.

Springer, Ansell & Ruelius (1956) reported that the polysaccharide with high blood-group H(O) activity isolated from *Taxus cuspidata* contained no L-fucose, but instead serologically active 2-O-methyl-L-fucose, not then known to occur in Nature. It is now recognized that this sugar is widely distributed in plants (Andrews & Hough, 1958): it has also been found in mycobacteria (MacLennan, Randall & Smith, 1961). It was also found that di-O-methylation of L-fucose at C-2 and C-3 was compatible with high serological activity; introduction of a third methyl group at C-4 led to complete inactivation of the compound (Springer, Williamson & Ruelius, 1959).

In view of the apparent influence of methyl groups on the serological activity of the fucose molecule, we decided to investigate the serological properties of various fucose ethers, glycosides and other derivatives. The effect of substituents in the various positions of the fucose molecule might yield information on the nature of the forces acting between haptens and antibodies or antibody-like substances, not only for the blood-group H(O) system but for other antigen-antibody systems as well.

During the extensive study on L-fucose derivatives reported here, the surprising observation was

* Established Investigator of the American Heart Association.