place at interfaces, but not in homogeneous solutions: this contrasts with the present study, where reaction appears to take place in either homogeneous or heterogeneous systems. With kidney dialkyl fluorophosphatase there is no increase in the rate of hydrolysis of sparingly soluble substrates when the solubility limit is reached (Mounter & Dien, 1956).

It would appear from the data of Table 5 that there are a number of bases on which classification could be made but no single property permits a general distinction between esterases and lipases. It seems that by any criterion the wheat-germ enzyme should be considered an esterase rather than a lipase.

SUMMARY

1. The specificity of wheat-germ esterase was studied in homogeneous and in heterogeneous systems.

2. The enzyme will hydrolyse glyceryl, *n*-alkyl, aromatic and choline esters.

3. Wheat-germ esterase is inhibited by organophosphorus compounds.

4. Maximal rates of hydrolysis of sparingly soluble esters are frequently only attained at substrate concentrations in excess of the solubility limit of the substrate.

5. The specificity and kinetics of esterases, including cholinesterase and lipase, are discussed.

This investigation was supported by a research grant [no. CY-3652 (C3)] from the National Cancer Institute, National Institutes of Health, United States Public Health Service.

REFERENCES

- Adams, D. H. (1949). *Biochim. biophys. Acta*, **3**, 1.
 Aldridge, W. N. (1953*a*). *Biochem. J.* **53**, 110.
 Aldridge, W. N. (1953*b*). *Biochem. J.* **53**, 117.
 Aldridge, W. N. (1954). *Biochem. J.* **57**, 692.
 Bier, M. (1955). In *Methods in Enzymology*, vol. 1, p. 627. Ed. by Colowick, S. P. & Kaplan, N. O. New York: Academic Press Inc.
 Desnuelle, P. (1953). *Bull. Soc. Chim. biol., Paris*, **33**, 909.
 Desnuelle, P. (1961). *Advanc. Enzymol.* **23**, 129.
 Eadie, G. S. (1952). *Science*, **116**, 688.
 Gowron, O., Grelecki, C. J. & Duggan, M. (1953). *Arch. Biochem. Biophys.* **44**, 455.
 Jansen, E. F., Jang, R. & MacDonnel, L. R. (1947). *Arch. Biochem.* **15**, 413.
 Jansen, E. F., Nutting, M. D. F. & Balls, A. K. (1948). *J. biol. Chem.* **175**, 975.
 Mounter, L. A. (1951). *Biochem. J.* **50**, 122.
 Mounter, L. A. & Dien, L. T. H. (1956). *J. biol. Chem.* **219**, 685.
 Mounter, L. A., Tuck, K. D., Alexander, H. C. & Dien, L. T. H. (1957). *J. biol. Chem.* **226**, 873.
 Mounter, L. A. & Whittaker, V. P. (1950). *Biochem. J.* **47**, 525.
 Murray, D. R. P. (1929). *Biochem. J.* **23**, 292.
 Sarda, L. & Desnuelle, P. (1958). *Biochim. biophys. Acta*, **30**, 513.
 Schenheyder, F. & Volqvartz, K. (1945). *Acta physiol. scand.* **9**, 57.
 Singer, T. P. & Hofstee, R. H. J. (1948*a*). *Arch. Biochem.* **18**, 229.
 Singer, T. P. & Hofstee, R. H. J. (1948*b*). *Arch. Biochem.* **18**, 245.
 Sturge, L. M. & Whittaker, V. P. (1950). *Biochem. J.* **47**, 518.
 Whittaker, V. P. (1951). *Physiol. Rev.* **31**, 312.

Biochem. J. (1962) **85**, 580

The Biosynthesis of Muramic Acid by Intact Staphylococci

BY M. H. RICHMOND AND H. R. PERKINS

Bacterial Physiology Division, National Institute for Medical Research, Mill Hill, London, N.W. 7

(Received 29 May 1962)

The amino sugar muramic acid was first clearly characterized by Strange & Dark (1956) and was shown by Cummins & Harris (1956) to be present in the cell walls of all the Gram-positive bacteria that they examined. It has since come to be recognized as the key compound in the structural mucopeptide polymer of all bacterial cell walls, in which it is probably present as the *N*-acetyl derivative. Its structure has been shown to be 3-*O*- α -carboxyethylglucosamine (Strange & Kent, 1959) (Fig. 1). Muramic acid has been synthesized

chemically by various methods (Strange & Kent, 1959; Zilliken, 1959; Gigg & Carroll, 1961), but so far there has been little decisive evidence on the mechanism of its biosynthesis.

An approach to this problem was made by Strominger (1958), who incubated extracts of *Staphylococcus aureus*, *Escherichia coli* and *Aerobacter aerogenes* with various radioactively labelled C_3 compounds in the presence of uridine diphosphate *N*-acetylglucosamine. He found that under his conditions only phosphoenolpyruvate would

condense with the glucosamine nucleotide to yield a compound substituted in the C-3 position; 2-phospho-D- or 2-phospho-L-glycerate, 3-phospho-D-glycerate and pyruvate or lactate with ATP were all inactive. No muramic acid was formed, but the substituted glucosamine produced was thought to be 3-O-enolpyruvyl-N-acetylglucosamine, although it was not conclusively identified. In view of these findings, it was decided to investigate the biosynthetic pathway of muramic acid in intact organisms, since this might yield evidence on whether a reaction of the type envisaged by Strominger could be involved in the synthesis of muramic acid.

It had been shown by Mandelstam & Rogers (1958, 1959) and Hancock & Park (1958) that staphylococci can synthesize the mucopeptide component of their cell walls (containing virtually all the muramic acid of the cell) without synthesizing significant quantities of protein. This system was therefore convenient for studying the biosynthesis of muramic acid in intact organisms.

It seems likely that muramic acid would be synthesized in the cell by the condensation of some precursor with glucosamine (probably in the form of uridine diphosphate N-acetylglucosamine) to yield the 3-O-carboxyethyl side chain. The main point examined in the present experiments was the nature of the biosynthetic precursor of the side chain of muramic acid. To achieve this object it was necessary to be able to isolate from muramic acid the carbon atoms of its side chain and if possible to characterize them. For this purpose the reaction described by Strange & Kent (1959) was used, in which heating with concentrated sulphuric acid converts part of the 3-O-carboxyethyl substituent into acetaldehyde and carbon monoxide. The acetaldehyde was converted into its 2,4-dinitrophenylhydrazone and the carbon monoxide, by treatment with iodine pentoxide, into carbon

dioxide and thence into barium carbonate, and the specific radioactivity of each was measured.

In a preliminary account of part of this work (Richmond & Perkins, 1960) we stated that alanine depressed by 75% the labelling of the side chain of muramic acid by [^{14}C]glucose. It has not been possible to repeat this result in several subsequent experiments; the regularly observed depression was only of the order of 5%. At this stage we cannot offer any explanation for this difference.

The various carbon atoms in muramic acid are referred to below by the numbers shown in Fig. 1.

METHODS

Organisms. *Staphylococcus aureus* 524 SC (Rogers, 1953) was grown and prepared for the experiments as described by Mandelstam & Rogers (1959). For experiments with radioactive lactic acid the growth medium contained 0.2% of sodium lactate in place of glucose.

Mucopeptide synthesis. The conditions for mucopeptide synthesis were those reported by Mandelstam & Rogers (1959); glucose was always present, but its concentration in the incubation medium was lowered from 1 to 0.02%. In certain experiments alanine was omitted and on other occasions DL-aspartic acid or pyruvic acid was added. The washed organisms were incubated for 90 min. in the presence of chloramphenicol (60 $\mu\text{g./ml.}$) at an initial density of about 0.5 mg./ml., and the total increase in mucopeptide was followed turbidimetrically by reference to the curves published by Mandelstam & Rogers (1959). Increases in individual components of the mucopeptide fraction were measured by taking samples at the beginning and end of incubation.

Preparation of mucopeptide. Samples were precipitated with an equal volume of 10% (w/v) trichloroacetic acid and then fractionated to yield the 'protein + cell wall' fraction (Richmond, 1959). The mucopeptide was prepared from the 'protein + cell wall' fraction by successive treatments with pepsin (1 mg./ml. in 0.05N-HCl at 35° for 16 hr.) and trypsin (1 mg./ml. in 0.05M- Na_2HPO_4 - KH_2PO_4 buffer, pH 7.6, at 35° for 4–8 hr.). The insoluble material obtained after washing closely resembled the mucopeptide fraction prepared by Mandelstam & Rogers (1959) and contained only trace amounts of amino acids characteristic of protein (cf. Hancock & Park, 1958).

Hydrolysis and separation of components of mucopeptide. Treatment with 4N-HCl at 105° for 4 hr. hydrolysed the mucopeptide fractions completely to give free amino acids and amino sugars. The HCl was removed *in vacuo*. Muramic acid was separated from the other components of the mucopeptide hydrolysates (Perkins & Rogers, 1959). The remaining amino acids and amino sugars were separated by quantitative chromatography on Whatman paper no. 3 developed in butan-1-ol-acetic acid-water (63:10:27, by vol.; upper phase) or butan-1-ol-pyridine-water (6:4:3) (Mandelstam & Rogers, 1959; Perkins & Rogers, 1959).

Estimation of specific activities. Quantitative estimations of glucosamine and muramic acid were obtained by the method of Randle & Morgan (1955), as described by Perkins & Rogers (1959). Amino acids were estimated after elution from chromatograms by a ninhydrin method (Yemm

Fig. 1. Muramic acid.

& Cocking, 1955). The radioactivity of these substances and of the acetaldehyde 2,4-dinitrophenylhydrazone (see below) was determined by plating representative samples (Richmond, 1959). The samples were assumed to be infinitely thin if they weighed less than 0.1 mg./cm.².

Degradation of muramic acid. Treatment of muramic acid with 86% H₂SO₄ for 17 min. at 100° liberates C-8 and C-9 atoms as acetaldehyde in about 35% yield (Strange & Kent, 1959). The carboxyl group of normal and substituted α -hydroxy acids is liberated as CO on treatment with conc. H₂SO₄ (Fearon, 1918), and this is the fate of some, at least, of C-7 atoms of muramic acid under these conditions. The acetaldehyde was trapped as the 2,4-dinitrophenylhydrazone by the following procedure.

A known quantity of muramic acid was placed in a glass ampoule, evaporated to dryness and cooled to -79° in a solid CO₂-methanol mixture. Sulphuric acid (86%; 0.2 ml.) was added down the side of the ampoule, which was maintained at -79° during the process. Usually the H₂SO₄ froze before reaching the muramic acid, but, even if the reactants mixed, little or no reaction occurred.

The ampoule was sealed and heated at 100° for 17 min. It was then cooled to -79° to solidify the acetaldehyde formed in the previous step, opened and 2.0 ml. of 2,4-dinitrophenylhydrazine solution (2% in *N*-HCl) added rapidly. The contents of the ampoule were warmed to room temperature and allowed to react for 30 min.

The acetaldehyde 2,4-dinitrophenylhydrazone and unchanged reagent were separated from the acid aqueous solution by repeated extraction with benzene. The benzene extracts were evaporated to dryness *in vacuo*, and the acetaldehyde 2,4-dinitrophenylhydrazone was purified by paper chromatography in the light petroleum-*NN*-dimethylformamide solvent (see below). The method allowed specific activity determinations on as little as 2 μ g. of acetaldehyde with an error of about $\pm 10\%$.

Attempts were also made to measure the specific activity of the CO liberated from C-7 atoms of muramic acid under these conditions by conversion into CO₂ on passing over I₂O₅ at 150° (Gautier, 1898; Adams & Simmons, 1951), in the apparatus shown in Fig. 2. Muramic acid was placed in the special ampoule and degraded with 86% H₂SO₄ as described above. After insertion of the ampoule, the apparatus was evacuated and the CO formed in the reaction was converted into CO₂ by heating I₂O₅ in the other limb to 150°. The CO₂ was frozen by immersing the tube containing the ampoule in liquid air, and was then allowed to distil into 0.1 *N*-Ba(OH)₂, which replaced the I₂O₅ in the other limb.

Degradation of alanine. Alanine was degraded to acetaldehyde + CO₂ by treatment with ninhydrin at 100°. The acetaldehyde was trapped as the 2,4-dinitrophenylhydrazone, and the CO₂ as BaCO₃. The experiment was also carried out in the apparatus shown in Fig. 2. The procedure was as follows.

A known amount of alanine was dried in the bottom of the special ampoule (Fig. 2) and cooled to -79°. The ampoule was kept at -79°, and 0.2 ml. of ninhydrin solution [2% (w/v) in aq. 4% (v/v) pyridine] was run into the bleb on the ampoule and frozen solid.

The ampoule was sealed, the reactants being kept solid, heated for 30 min. at 105°, and then cooled to room temperature. It was inserted in the apparatus in position A (Fig. 2) and a tube containing 2 ml. of 2,4-dinitrophenyl-

hydrazine (2%, v/v, in *N*-HCl) placed on the other side. The apparatus was evacuated and the tube containing the dinitrophenylhydrazine cooled to -190° in liquid air. The tube containing the ampoule was twisted to position B (Fig. 2), thus fracturing the ampoule. The acetaldehyde and CO₂ liberated immediately distilled into the cooled side of the apparatus and solidified.

With the dinitrophenylhydrazine tube still in liquid air, the vacuum was released through a soda-lime trap and the ampoule tube replaced with a tube containing 0.1 *N*-Ba(OH)₂. The apparatus was evacuated and allowed to warm to room temperature. The acetaldehyde was then caused to react with the 2,4-dinitrophenylhydrazine; the CO₂ distilled over to react with the baryta to form BaCO₃.

Separation and estimation of degradation products of alanine and muramic acid. Acetaldehyde 2,4-dinitrophenylhydrazone arising from either alanine or muramic acid was separated from unchanged reagent by ascending chromatography on Whatman no. 1 paper developed overnight in the upper phase of light petroleum (b.p. 100-120°)-*NN*-dimethylformamide (20:7, v/v) mixture. A period of equilibration of several hours in the presence of the lower phase was essential to obtain well-defined spots. The *R_F* values of the 2,4-dinitrophenylhydrazones of aliphatic aldehydes were: formaldehyde, 0.13; acetaldehyde, 0.20; propionaldehyde, 0.57; *n*-butyraldehyde, 0.79. This solvent was found to give more reproducible chromatography for our purpose than the systems described by Meigh (1952) and Huelin (1952). Acetaldehyde dinitrophenylhydrazone was eluted from the chromatograms with benzene containing 5% (v/v) of methanol and its specific absorption at 353 μ compared with that of known quantities of a reference sample of acetaldehyde dinitrophenylhydrazone that had been run on the same chromatogram. The 2,4-dinitrophenylhydrazine used in these studies contained material with *R_F* similar to that of acetaldehyde 2,4-dinitro-

Fig. 2. Apparatus for opening a sealed ampoule in an evacuated system. The ampoule is inserted with the tube in position A; turning the tube to position B opens it.

phenylhydrazone and appreciable absorption at 353 m μ . Because of this, experimental results were corrected by separating a control quantity of 2,4-dinitrophenylhydrazine on the same chromatogram as the experimental and standard samples, eluting the relevant area under standard conditions and measuring the quantity of interfering material absorbing at 353 m μ . This value was then subtracted from E_{353} of the experimental samples. The amount of interfering material was usually 10–20% of the experimental values.

Barium carbonate samples were washed three times with CO₂-free water, once with acetone, once with ether, dried in a stream of air and the specific radioactivity was determined on weighed samples at infinite thickness.

Materials. Generally labelled [¹⁴C]glucose, L-[1,2,3-¹⁴C]-alanine, DL-[1-¹⁴C]alanine, [2-¹⁴C]pyruvate and generally labelled L-[¹⁴C]aspartic acid were obtained from The Radiochemical Centre, Amersham, Bucks. They were used in these experiments at the following final average specific activities/C atom: [¹⁴C]glucose, 312 counts/min./ μ g. of C; L-[1,2,3-¹⁴C]₃alanine, 1380 counts/min./ μ g. of C; DL-[1-¹⁴C]alanine, 1864 counts/min./ μ g. of C; [¹⁴C]₄aspartic acid, 1400 counts/min./ μ g. of C; [2-¹⁴C]pyruvate, 1836 counts/min./ μ g. of C; sodium [2-¹⁴C]lactate, 7000 counts/min./ μ g. of C-2.

Other compounds. Iodine pentoxide and 2,4-dinitrophenylhydrazine were AnalaR standard reagents. The I₂O₅ was heated for 10–15 min. at 160–180° immediately before use. Acetaldehyde 2,4-dinitrophenylhydrazone was synthesized by distilling acetaldehyde into a solution of 2,4-dinitrophenylhydrazine (2% in *n*-HCl). The product was recrystallized from aq. ethanol.

RESULTS

[¹⁴C]Glucose (generally labelled) as a precursor of muramic acid

Since Strominger (1958) has suggested that muramic acid may be synthesized by a reaction between uridine diphosphate *N*-acetylglucosamine and phosphoenolpyruvate, and Roseman, Moses, Ludowig & Dorfman (1953, 1954) have shown that glucose is an effective precursor for the synthesis of glucosamine, a preliminary experiment was carried out to measure the extent to which radioactive glucose would label the components of the mucopeptide, particularly muramic acid. Exponentially growing organisms were harvested by centrifuging, washed once in 0.05M-phosphate buffer, pH 7.0, containing 0.02% of glucose and suspended in two flasks at an initial density of 0.43 mg. dry wt. of organisms/ml. in a medium containing [¹⁴C]glucose (200 μ g./ml.; 312 counts/min./ μ g. of C) and non-radioactive lysine, glycine and glutamic acid. Non-radioactive alanine (400 μ g./ml.) was added to one culture to discover whether this amino acid was an effective competitor for the flow of carbon atoms to the components of the mucopeptide. After 90 min., when considerable mucopeptide synthesis had taken place (Mandelstam & Rogers, 1959), the bacteria were precipi-

tated with an equal volume of 10% (w/v) trichloroacetic acid. The mucopeptide was prepared as described above and hydrolysed, the muramic acid being further degraded to liberate C-8 and C-9 atoms as acetaldehyde. Table 1 shows the specific radioactivity of the various components of the mucopeptide. In the absence of exogenous alanine [¹⁴C]glucose labels the glucosamine, muramic acid and alanine heavily and to approximately the same extent, whereas the lysine, glycine and glutamic acid had average specific activities about 1/20 to 1/100 of those of glucosamine. Since the carbon atoms of the acetaldehyde liberated from the muramic acid have the same specific activity as those of both the glucosamine and the complete molecule of muramic acid, it is likely that each carbon atom of the muramic acid had the same specific activity. Addition of non-radioactive alanine as competitor had no significant effect on the specific activity of any of the carbon atoms in the glucosamine, muramic acid, lysine, glycine and glutamic acid but lowered the specific activity of the alanine in the mucopeptide. Since the non-radioactive DL-alanine does not lower the specific activity of C-8 and C-9 atoms of muramic acid, it follows that both optical isomers of this amino acid must be poor precursors of the C-7, C-8 and C-9 positions of muramic acid.

Possible precursors of the side chain of muramic acid

The results obtained with radioactive glucose show the close biosynthetic relationship between the amino sugars and this compound but do not

Table 1. Average specific activities of carbon atoms of the components of the cell-wall mucopeptide formed in the presence of [¹⁴C]glucose

The mucopeptide-synthesizing system contained the following constituents (μ g./ml. of the 0.05M-phosphate buffer, pH 7.0): [¹⁴C]glucose (generally labelled, specific activity: 312 counts/min./ μ g. of C), 100; sodium DL-glutamate, 400; L-lysine hydrochloride, 400; glycine, 400; chloramphenicol, 60; DL-alanine, where required, 400.

Substance	C atoms	Specific activity (counts/min./ μ g. of C)	
		Without alanine	With alanine
Glucose*	1–6	312	312
Glucosamine	1–6	272	293
Muramic acid	1–9	254	268
	8, 9	281	268
	Calc. 1–6	240	268
Alanine	1–3	212	40
Glycine	1–2	2.7	4.2
Lysine	1–6	12.5	14.4
Glutamic acid	1–5	5.1	7.6

* Average specific activity of glucose added to incubation system.

throw any light on the immediate source of the side chain of muramic acid. Further experiments were conducted with various compounds that might be expected to be related to a possible precursor. Since alanine and the muramic acid side chain are obviously related structurally, yet alanine is not a direct precursor, it seemed likely that the side-chain precursor should be another C₃ compound such as pyruvate, phosphoenolpyruvate or lactate. Unfortunately it was not possible to test phosphoenolpyruvate directly in this system because of the impermeability of bacterial cells to phosphate esters. Alanine, however, can freely enter the cells and radioactive alanine was used in an attempt to find out more about biosynthesis of the side chain, since the use of specifically labelled alanine should give some indication of the chemical changes undergone by the molecule before becoming part of muramic acid. Thus if C-1, C-2 and C-3 atoms of alanine become C-7, C-8 and C-9 atoms of muramic acid without randomization, then [1-¹⁴C]alanine should label C-7 atoms of muramic acid only, and acetaldehyde prepared from C-8 and C-9 atoms should not be radioactive. If, however, [1-¹⁴C]-alanine labels the acetaldehyde as well as C-7 atoms, the alanine must be converted into some symmetrical compound before becoming the side chain of muramic acid. A number of organisms can produce phosphoenolpyruvate by transdeamination of aspartate followed by decarboxylation of the resulting oxaloacetic acid (Utter & Wood, 1951;

Meister, 1955), and generally labelled [¹⁴C]aspartic acid was therefore tested as well as pyruvate and lactate as a source of C-7, C-8 and C-9 atoms of muramic acid.

Alanine. Mucopolypeptide was prepared as in the previous experiment from organisms that had been incubated in a solution containing non-radioactive glucose, glutamic acid, glycine and lysine, and either DL-[1,2,3-¹⁴C₃]alanine (1380 counts/min./μg. of C) or DL-[1-¹⁴C]alanine (1864 counts/min./μg. of C). The amino acid and amino sugar components were liberated from the mucopolypeptide by hydrolysis and the labelling of the whole muramic acid molecule and the acetaldehyde derived from C-8 and C-9 atoms was determined. Table 2 shows the results of this experiment. [1,2,3-¹⁴C₃]Alanine labels muramic acid with an average specific radioactivity of 4.0 counts/min./μg. of C, and the acetaldehyde derived from C-8 and C-9 atoms was found to have a specific activity of 12.3 counts/min./μg. of C. If C-7 atom has the same specific activity as C-8 and C-9 atoms, then C-1 to C-6 atoms of muramic acid are not radioactive. As might have been expected from the experiments with radioactive glucose, the glucosamine too was not radioactive. When [1-¹⁴C]alanine was added to the mucopolypeptide-synthesizing system, the specific activity of the muramic acid averaged over the entire molecule was 2.2 counts/min./μg. of C. If alanine is a precursor of C-7, C-8 and C-9 atoms without randomization, then the C-7 atom should

Table 2. *Average specific activity of carbon atoms of the amino sugars from mucopolypeptide formed in the presence of DL-[1,2,3-¹⁴C₃]alanine or DL-[1-¹⁴C]alanine*

The mucopolypeptide-synthesizing medium contained (μg./ml. of 0.05M-phosphate buffer, pH 7.0): glucose, 200; sodium DL-glutamate, 400; L-lysine hydrochloride, 400; glycine, 400; chloramphenicol, 60; either DL-[1,2,3-¹⁴C₃]alanine (specific activity: 1380 counts/min./μg. of C) or [1-¹⁴C]alanine (specific activity: 1864 counts/min./μg. of C-1), 400.

	Amino sugar C atoms		Specific activity of newly formed substance (counts/min./ μg. of C)
	Muramic acid	Glucosamine	
DL-[1,2,3- ¹⁴ C ₃]Alanine	Total, C-1-C-9		4.0
	Calc., C-1-C-6		Nil
	Calc., C-7, C-8, C-9*		12.0
	Found, C-8, C-9		12.3
		C-1-C-6	Nil
DL-[1- ¹⁴ C]Alanine	Total, C-1-C-9		2.2
	Calc., C-1-C-6		Nil
	Calc., C-7*		19.8
	Calc., C-8, C-9*		Nil
	Found, C-7		Positive†
	Found, C-8, C-9		1
		C-1-C-6	Nil

* Calculated assuming that [¹⁴C]alanine does not label the ring of muramic acid and that the order C-1, C-2 and C-3 atoms of alanine is completely conserved as C-7, C-8 and C-9 atoms of muramic acid.

† In the experiment with [1-¹⁴C]alanine, the CO liberated from the C-7 atom of muramic acid on digestion with 86% H₂SO₄ was converted into CO₂ by passage over I₂O₅. The BaCO₃ derived from the CO₂ thus formed was radioactive but the recovery of the CO was too poor for a satisfactory specific radioactivity to be measured.

contain all the activity of the molecule, namely a specific activity of 19.8 counts/min./ μ g. of C. In the experiment it was found that the acetaldehyde derived from C-8 and C-9 atoms had an average specific activity of 1 count/min./ μ g. of C, and, as the glucosamine from the same experiment was not radioactive, this suggests that virtually all the radioactivity derived from the [14 C]alanine has gone to the C-7 position of muramate. This is the carbon atom that is evolved as carbon monoxide on digestion with 86% sulphuric acid. Conversion of the carbon monoxide into carbon dioxide by passing over iodine pentoxide showed that this carbon atom was indeed labelled, but the overall reaction was not quantitative. No radioactive carbon dioxide was produced by the direct reaction between this muramic acid and 86% sulphuric acid. The results show that the precursor of the side chain of muramic acid is a compound in which the identities of C-1, C-2 and C-3 atoms of alanine are maintained to give C-7, C-8 and C-9 atoms of muramic acid.

Aspartic acid. The medium contained non-radioactive glucose, lysine, glycine and glutamic acid together with generally labelled L-[14 C]aspartic acid (50 μ g./ml.; specific activity, 1400 counts/min./ μ g. of C). A further experiment was carried out with non-radioactive alanine added. The specific activities of the carbon atoms of the newly formed alanine, glucosamine and muramic acid obtained in these experiments together with the specific activity of the carbon in the acetaldehyde derived from C-8 and C-9 atoms of muramic acid are shown in Table 3. [14 C]Aspartic acid labels alanine and muramic acid. If the radioactivity observed in the muramic acid were averaged over all the carbon atoms the specific activity of each would be 23 counts/min./ μ g. of C. The glucosamine in this experiment is almost non-radioactive (less than 1 count/min./ μ g. of C), and it is therefore unlikely that C-1 to C-6 atoms of muramic acid are labelled. The acetaldehyde has a specific activity of 70 counts/min./ μ g. of C, sufficient to

account for the whole radioactivity of the muramic acid if aspartic acid labels C-7, C-8 and C-9 atoms equally without labelling C-1 to C-6 atoms. The alanine in the mucopeptide had an average specific activity of 68 counts/min./ μ g. of C, and an experiment in which this alanine was degraded with ninhydrin to acetaldehyde and carbon dioxide showed that both products of the alanine were radioactive. Addition of non-radioactive alanine with [14 C]aspartic acid resulted in a fall of 74% in the specific activity of the alanine from the mucopeptide but a slight increase of the average activity of C-7, C-8 and C-9 atoms of muramic acid. This result shows that aspartic acid is a much more efficient source of carbon atoms for the side chain of muramic acid than alanine, and is consistent with the idea that phosphoenolpyruvate may be the direct precursor.

Pyruvate. The mucopeptide-synthesizing system contained non-radioactive glucose, lysine, glycine and glutamic acid with [14 C]pyruvate (specific activity: 1836 counts/min./ μ g. of C). As before, the alanine, glucosamine and muramic acid were isolated from the mucopeptide together with the acetaldehyde derived from muramic acid.

Since pyruvate is in ready equilibrium with alanine in most organisms it is likely that it will be converted into C-7, C-8 and C-9 atoms of muramic acid in the same way as alanine. In this experiment therefore [14 C]pyruvate would be expected to label the C-8 atom of muramic acid; that is, the acetaldehyde should be labelled only in the carbonyl carbon atom. Likewise, alanine in the mucopeptide should be labelled at the C-2 position. The results of this experiment are shown in Table 4. In the absence of non-radioactive alanine, both muramic acid and the alanine from the mucopeptide are radioactive, whereas glucosamine is not. Isolation of the acetaldehyde showed that C-8 atoms contained all the radioactivity present in the muramic acid molecule (assuming that the C-9 atom was not labelled). Addition of non-radioactive alanine reduced the specific activity of the

Table 3. *Radioactivity incorporated from generally labelled [14 C]aspartic acid into the newly formed muramic acid, glucosamine and alanine of the mucopeptide*

The mucopeptide-synthesizing system contained (μ g./ml. of 0.05M-phosphate buffer, pH 7.0): glucose, 200; sodium DL-glutamate, 400; glycine, 400; L-lysine hydrochloride, 400; chloramphenicol, 60; generally labelled L-[14 C]aspartic acid (average specific activity: 1400 counts/min./ μ g. of C), 50; DL-alanine, where required, 400.

	Average specific activity (counts/min./ μ g. of C of newly formed substance)				
	Muramic acid			Glucosamine C-1-C-6	Alanine C-1-C-3
	C-1-C-9	C-8, C-9	C-1-C-6 (calc.)		
Non-radioactive alanine					
Absent	23	70	Nil	0.92	68
Present	31	94	Nil	1.47	17

Table 4. *Radioactivity incorporated from [2-¹⁴C]pyruvate into the muramic acid, glucosamine and alanine isolated from mucopeptide*

The mucopeptide-synthesizing system contained ($\mu\text{g./ml.}$ of 0.05M-phosphate buffer, pH 7.0): glucose, 200; L-lysine hydrochloride, 400; sodium DL-glutamate, 400; glycine, 400; chloramphenicol, 60; [2-¹⁴C]pyruvate (specific activity: 1836 counts/min./ $\mu\text{g.}$ of C-2), 500; DL-alanine, where required, 400.

Non-radioactive alanine	Average specific activity (counts/min./ $\mu\text{g.}$ of C)				
	Muramic acid			Glucosamine	Alanine
	C-1-C-9	C-8	C-1-C-6 (calc.)	C-1-C-6	C-2
Absent	26	230	Nil	0.94	900
Present	22	200	Nil	0.62	175

Table 5. *Radioactivity incorporated from [2-¹⁴C]lactate into newly formed components of mucopeptide by organisms previously adapted by growth in the presence of sodium lactate*

The mucopeptide-synthesizing system contained ($\mu\text{g./ml.}$ of 0.05M-phosphate buffer, pH 7.0): glucose, 200; sodium DL-glutamate, 400; glycine, 400; L-lysine hydrochloride, 400; chloramphenicol, 60; sodium [2-¹⁴C]lactate (specific activity: 7000 counts/min./ $\mu\text{g.}$ of C-2), 100; sodium pyruvate, where required, 400.

Non-radioactive pyruvate	Average specific activity (counts/min./ $\mu\text{g.}$ of C of newly formed substance)						
	Muramic acid			Glucosamine	Alanine	Lysine	Glutamic acid
	C-1-C-9	C-8	C-1-C-6 (calc.)	C-1-C-6	C-2	C-1-C-6	C-1-C-5
Absent	101	503	68	75	1890	54	95
Present	53	261	36	36	1010	30	43

alanine in the mucopeptide by 81%, whereas that of the side chain of muramic acid was reduced by only about 20%. In the absence of added alanine the specific activity of the C-2 atom of the alanine isolated from the mucopeptide was much higher than that of the C-8 atom of the muramic acid, whereas if pyruvate were the immediate precursor for both these products their specific activities would be expected to be the same. These results strongly suggest that pyruvate is not the direct precursor of C-7, C-8 and C-9 atoms of muramic acid, but that some other compound (such as phosphoenolpyruvate), which lies closer to glucose in the glycolytic pathway, could well act in this way.

Lactate. Sodium lactate is poorly used by staphylococci that have been grown in the presence of glucose, but a period of adaptation in casein-hydrolysate medium containing sodium lactate (200 $\mu\text{g./ml.}$) produces organisms capable of using [2-¹⁴C]lactate as a precursor of mucopeptide. An experiment was therefore carried out with adapted cells in a solution containing non-radioactive glucose, glycine, glutamic acid and lysine and [2-¹⁴C]lactate (specific activity: 7000 counts/min./ $\mu\text{g.}$ of C-2). An additional flask contained non-radioactive pyruvate, to see whether it affected the flow of radioactive carbon from the lactate to the

alanine and the muramic acid of the mucopeptide. The results of this experiment (Table 5) show that with lactate-adapted organisms, [2-¹⁴C]lactate labelled all components of the mucopeptide with the exception of glycine, but the specific activity of the C-2 atom of the newly formed alanine was about four times that of the C-8 atom of muramic acid. Addition of non-radioactive pyruvate depressed the labelling of the alanine and C-8 atoms of muramic acid to approximately the same extent. This experiment shows that lactate is not a direct precursor of the side chain of muramic acid (even in lactate-adapted cells) but that the carbon atoms of lactate probably reach the muramate side chain by way of pyruvate.

DISCUSSION

The results obtained with radioactive alanine suggest that the carbon atoms from alanine flow to the side chain of muramate without loss of identity, i.e. C-1 $\rightarrow$ C-7, C-2 $\rightarrow$ C-8, C-3 $\rightarrow$ C-9, and do not label C-1 to C-6 atoms of muramic acid or glucosamine. When the specific activity of the most heavily labelled carbon atoms of muramic acid formed in the experiment involving [1,2,3-¹⁴C₃]-alanine is compared with that of the muramic acid synthesized from [¹⁴C]glucose (see Table 1), it can be calculated that, in a medium containing 200 $\mu\text{g.}$

of glucose and 400 μg . of DL-alanine/ml., more than 80% of the carbon atoms for C-1 to C-6 positions of muramic acid come from exogenous glucose, whereas about 2% of the carbon atoms for C-7, C-8 and C-9 positions come from exogenous alanine, and more than 80% from glucose. This result suggests that under the experimental conditions used the side chain of muramic acid is more closely related metabolically to the breakdown products of glucose than it is to alanine.

The experiments described above involving labelled alanine, pyruvate and aspartic acid are all consistent with the idea that some C_3 molecule of related structure is the immediate precursor of the side chain of muramic acid. The results with radioactive lactate make it clear that this precursor molecule cannot be lactic acid. In all cases the addition of non-radioactive alanine depressed markedly the flow of carbon atoms from the radioactive molecule to the alanine in the mucopeptide but not to the side chain of muramic acid (see Table 6). The possible reason for this is that the immediate precursor of the side chain of muramic acid is phosphoenolpyruvate, as suggested by Strominger (1958), and that the equilibrium in the reaction

lies well to the right (see Krebs & Kornberg, 1957), thus reducing the effect of the addition of alanine on the flow of carbon atoms from glucose to phosphoenolpyruvate. A scheme of this type is favoured by the high efficiency with which aspartic acid labels the side chain. This amino acid is known to yield phosphoenolpyruvate in certain biological systems (Utter & Wood, 1951; Meister, 1955), and this could well account for the finding that exo-

genous non-radioactive alanine has no inhibitory effect on the flow of carbon atoms from aspartate to C-1, C-8 and C-9 positions of muramic acid.

Although reasoning of this kind makes it probable that phosphoenolpyruvate is the immediate precursor of the side chain of muramic acid, it should be pointed out that none of the metabolic intermediates between phosphoenolpyruvate and dihydroxyacetone phosphate—glyceraldehyde phosphate in the Embden-Meyerhof glycolytic pathway can be excluded by our experiments since the pathway is freely reversible (see Krebs & Kornberg, 1957).

At first sight it might seem that phosphoenolpyruvate is activated on the wrong carbon atom to lead to the formation of an ether bond at C-2. However, it has recently been shown that shikimic acid reacts with phosphoenolpyruvate to produce such a compound (Levin & Sprinson, 1960). This compound has a C_3 side chain linked to the ring with an ether bond at the C-2 atom of the side chain, and its formation is therefore analogous to the step in the synthesis of muramic acid.

The finding that aspartic acid is such a good precursor of the side chain of muramic acid is interesting in connexion with the experiments reported by Gale & Folkes (1953) on the uptake of amino acids by bacteria. They found that aspartic acid labelled *Staphylococcus aureus* under conditions that are now recognized as being characteristic of cell-wall synthesis (Mandelstam & Rogers, 1958, 1959; Hancock & Park, 1958; Gale, Shepherd & Folkes, 1958). Since aspartic acid does not occur in the cell wall of *S. aureus* to any great extent (Mandelstam & Rogers, 1959), it seems probable that the uptake of radioactivity from $[^{14}\text{C}]$ aspartic acid into the cell wall was predominantly due to incorporation of radioactive carbon atoms from the amino acid into the side chain of muramic acid. During the course of this work O'Brien, Glick & Zilliken (1960) showed that methyl *N*-acetylglucosaminide was used in the synthesis of muramic acid by *Lactobacillus bifidus* var. *pennsylvanicus*. Since the specific activity per mole was the same in the isolated muramic acid as in the glucosaminide used, it seems probable that only the ring and not the side chain was labelled in these experiments.

Table 6. Effect of non-radioactive alanine in depressing the average specific activity of carbon atoms of alanine and the muramic acid side chain in compounds isolated from mucopeptide synthesized in the presence of various radioactive precursors

Results are taken from Tables 1, 3 and 4.

Radioactive precursor	Depression in average specific activity* (%)	
	Muramic acid (C-7, C-8, C-9)	Alanine (C-1, C-2, C-3)
$[^{14}\text{C}]$ Glucose	4.6	81
$[^{14}\text{C}]$ Aspartic acid	-15.4†	75
$[2\text{-}^{14}\text{C}]$ Pyruvate	15.2	80.5

* Calculated as

$$\frac{(\text{av. specific activity without added alanine}) - (\text{av. specific activity with added alanine})}{(\text{av. specific activity without added alanine})} \times 100$$

† A negative percentage indicates a stimulation.

SUMMARY

1. The biosynthesis of muramic acid by intact staphylococci has been studied under cultural conditions in which the synthesis of cell walls but not of protein occurs.

2. Glucose was found to be a precursor of all carbon atoms in muramic acid.

3. Radioactive DL-alanine, pyruvate, lactate or aspartate was found to label the side chain of

muramic acid (C-7, C-8 and C-9) in preference to the pyranose ring (C-1 to C-6) when the incubation medium contained non-radioactive glucose. With the C_3 compounds the identity of the carbon atoms was maintained, i.e. C-1 $\rightarrow$ C-7; C-2 $\rightarrow$ C-8; C-3 $\rightarrow$ C-9.

4. Experiments with the above-named tracers and non-radioactive competitors suggest that phosphoenolpyruvate is the most likely immediate precursor of the side chain of muramic acid. It is not possible, however, completely to exclude any of the compounds involved in glucose degradation by the glycolytic pathway between triose phosphate and phosphoenolpyruvate.

REFERENCES

- Adams, E. G. & Simmons, N. T. (1951). *J. appl. Chem.* **1**, suppl. 1, S20.
- Cummins, C. S. & Harris, H. (1956). *J. gen. Microbiol.* **14**, 583.
- Fearon, W. R. (1918). *Biochem. J.* **12**, 179.
- Gale, E. F. & Folkes, J. P. (1953). *Biochem. J.* **55**, 721.
- Gale, E. F., Shepherd, G. & Folkes, J. P. (1958). *Nature, Lond.*, **182**, 592.
- Gautier, A. (1898). *C.R. Acad. Sci., Paris*, **126**, 931.
- Gigg, R. H. & Carroll, P. M. (1961). *Nature, Lond.*, **191**, 495.
- Hancock, R. & Park, J. T. (1958). *Nature, Lond.*, **181**, 1050.
- Huelin, F. E. (1952). *Aust. J. sci. Res. B*, **5**, 328.
- Krebs, H. A. & Kornberg, H. L. (1957). *Ergebn. Physiol.* **49**, 212.
- Levin, J. G. & Sprinson, D. B. (1960). *Biochem. biophys. Res. Commun.* **3**, 157.
- Mandelstam, J. & Rogers, H. J. (1958). *Nature, Lond.*, **181**, 956.
- Mandelstam, J. & Rogers, H. J. (1959). *Biochem. J.* **72**, 654.
- Meigh, D. F. (1952). *Nature, Lond.*, **170**, 579.
- Meister, A. (1955). *Advanc. Enzymol.* **16**, 185.
- O'Brien, P. J., Glick, M. C. & Zilliken, F. (1960). *Biochim. biophys. Acta*, **37**, 357.
- Perkins, H. R. & Rogers, H. J. (1959). *Biochem. J.* **72**, 647.
- Richmond, M. H. (1959). *Biochem. J.* **73**, 155.
- Richmond, M. H. & Perkins, H. R. (1960). *Biochem. J.* **76**, 1 p.
- Rogers, H. J. (1953). *J. Path. Bact.* **66**, 545.
- Rondle, C. J. M. & Morgan, W. T. J. (1955). *Biochem. J.* **61**, 586.
- Roseman, S., Moses, F. E., Ludowieg, J. & Dorfman, A. (1953). *J. biol. Chem.* **203**, 213.
- Roseman, S., Moses, F. E., Ludowieg, J. & Dorfman, A. (1954). *J. biol. Chem.* **206**, 665.
- Strange, R. E. & Dark, F. A. (1956). *Nature, Lond.*, **177**, 186.
- Strange, R. E. & Kent, L. H. (1959). *Biochem. J.* **71**, 333.
- Strominger, J. L. (1958). *Biochim. biophys. Acta*, **30**, 645.
- Utter, H. F. & Wood, H. G. (1951). *Advanc. Enzymol.* **12**, 43.
- Yemm, E. W. & Cocking, E. C. (1955). *Analyst*, **80**, 209.
- Zilliken, F. (1959). *Fed. Proc.* **18**, 966.

Biochem. J. (1962) **85**, 588

Control of the Citric Acid Cycle by Glyoxylate

1. A NEW INHIBITOR OF ACONITASE FORMED BY THE CONDENSATION OF GLYOXYLATE WITH OXALOACETATE

BY A. RUFFO, E. TESTA, ANNA ADINOLFI AND GIUSEPPINA PELIZZA

Department of Biological Chemistry, University of Perugia, and Department of Organic Synthesis, Lepetit, S.p.A., Milan, Italy

(Received 4 June 1962)

Ruffo, Romano & Adinolfi (1959) showed that aconitase was inhibited when incubated in the presence of small amounts of glyoxylate and oxaloacetate. Since chromatographic and radioautographic evidence was given (Ruffo, Romano, Adinolfi & Verga, 1960) of the formation during the incubation of a new compound, probably a C_6 tricarboxylic acid, the condensation between sodium glyoxylate and sodium oxaloacetate was investigated under physiological conditions of pH and temperature, and is reported in the present paper.

The product of the reaction was isolated, its structure determined, and its mode of action on aconitase was studied.

EXPERIMENTAL

Condensation between sodium glyoxylate and sodium oxaloacetate. Oxaloacetic acid (1.32 g.) and sodium glyoxylate (1.14 g.) were dissolved in 50 ml. of water, and the pH was adjusted to 7.4 with about 20 ml. of N -NaOH. The solution was incubated for 3 hr. at 40° and then freeze-dried; the disappearance of the aldehyde group was determined