Hydrolysis with silver oxide. The amino acid hydrochloride (7.7 mg.) was refluxed with silver oxide (23.8 mg.) in 5 ml. of water for 3 hr. The hydrolysate was filtered, evaporated to dryness at room temperature under reduced pressure and the residue redissolved in 1 ml. of 10% (v/v) propan-2-ol. Chromatography and ionophoresis as above gave a strong spot of unchanged homoarginine and two weaker spots corresponding to lysine and homocitrulline.

Identification of homocitrulline on paper. Ureides normally give a yellow colour on paper with Ehrlich's reagent, whether applied directly or after development with ninhydrin or isatin in a multiple dipping sequence (Jepson & Smith, 1953). Homocitrulline gives the expected yellow when treated directly, but when the Ehrlich's reagent is applied after ninhydrin or isatin the colours given with these reagents rapidly fade and are replaced by an intense rose red. This reaction is more sensitive than the simple Ehrlich reaction. Hydroxyproline (Jepson & Smith, 1953) and 4-methylenepyrroline (Gray & Fowden, 1962) are reported to show colour changes with these reagents, but the development of a stable rose red appears to be specific for homocitrulline.

DISCUSSION

L-Homoarginine was detected in the seeds of those 17 species of *Lathyrus* known to contain lathyrine and also in those of 19 other species of this genus (Bell, 1962a). In the 17 species containing lathyrine the concentrations of L-homoarginine and lathyrine appear complementary, a high concentration of one being associated with a low concentration of the other. This relationship and the structural similarity of the two compounds suggest that lathyrine may be formed by the cyclization and dehydrogenation of L-homoarginine and that the fundamental difference between the species containing lathyrine and those containing

only homoarginine is the presence or absence of an enzyme system capable of bringing about this transformation.

The high nitrogen content and extremely low solubility of lathyrine at normal biological pH values agree with its supposed role in nitrogen storage. L-Homoarginine, with a similar nitrogen content, is more soluble.

SUMMARY

1. The natural occurrence of L-homoarginine has been confirmed by isolation from seeds of *Lathyrus cicera*.

2. The possible role of L-homoarginine as the precursor of lathyrine is discussed.

3. A colour reaction for the detection of homocitrulline is described.

The author thanks Professor H. Harris for his interest in this work, Dr Yu. Ya. Gofman for a gift of seed and Mr C. E. Hurst for determining the infrared absorption spectra.

REFERENCES

- Bell, E. A. (1961). *Biochim. biophys. Acta*, **47**, 602.
Bell, E. A. (1962a). *Biochem. J.* **83**, 225.
Bell, E. A. (1962b). *Nature, Lond.*, **193**, 1078.
Bell, E. A. & Foster, R. G. (1962). *Nature, Lond.*, **194**, 91.
Cox, G. J. (1928). *J. biol. Chem.* **78**, 475.
Gray, D. O. & Fowden, L. (1962). *Nature, Lond.*, **193**, 1285.
Gross, D. (1961). *J. Chromat.* **5**, 194.
Jepson, J. B. & Smith, I. (1953). *Nature, Lond.*, **172**, 1100.
Smith, I. (1953). *Nature, Lond.*, **171**, 43.
Stevens, C. M. & Bush, J. A. (1950). *J. biol. Chem.* **183**, 139.

Biochem. J. (1962) **85**, 93

Univalent Cations and the Respiration of Brain-Cortex Slices

BY L. HERTZ AND M. SCHOU

Institute of Neurochemistry, State Mental Hospital, Risskov, Denmark

(Received 5 March 1962)

The peculiar ability of brain slices to respond metabolically to certain stimuli was first observed with potassium as the stimulating agent (Ashford & Dixon, 1935; Dickens & Greville, 1935). Studies on the metabolic response to electrical pulses have been numerous (McIlwain, 1951, 1956), and the potassium effect, although frequently confirmed in a qualitative way, has not been examined quantitatively to a comparable extent.

The metabolic response to stimulation comprises increased respiratory rate and aerobic glycolysis, breakdown of creatine phosphate and ATP, and an increase in inorganic phosphate; it is found in brain tissue, but is not shown by slices from kidney, liver, testis, and tumour tissue. L. Hertz & T. Clausen (unpublished work) showed that even within the central nervous system there is considerable variation from one area to another

in the metabolic response to potassium. This may indicate a connexion with more highly developed functions of the central nervous system.

Also, ions other than potassium, e.g. caesium and rubidium, can, when added in sufficient amount to the medium, stimulate the rates of oxygen uptake and glycolysis, and ammonium ions seem to have a similar effect (Weil-Malherbe, 1938; Gore & McIlwain, 1952; Takagaki & Tsukada, 1957). There have been few experiments with lithium, and their results are inconclusive (Dickens & Greville, 1935; Canzanelli, Rogers & Rapport, 1941-42; Rybová, 1959). The effects of this ion on brain tissue may be related to its therapeutic efficacy against manic phases of the manic-depressive psychosis (Cade, 1949; Schou, 1959; Gershon & Yuwiler, 1960).

Neither electrical pulses nor potassium produce a metabolic response from the brain slices if sodium is absent from the medium. These ionic effects may gain new significance from the discovery of enzyme systems requiring the presence of both sodium and potassium for full activity (Skou, 1957; Hess & Pope, 1957).

The present work is a reinvestigation of the influence exerted by the ionic composition of the medium on the oxygen uptake of isolated rat-brain slices. The respiratory rate of the slices was not constant during the incubation, but showed an exponential fall with time. This fall was moderate in a control medium with a composition resembling that of the extracellular fluid, but it became more pronounced when sodium was replaced by various other ions or when a stimulating agent such as potassium was added to the system. The observation of this gradual decrease of the oxygen uptake led to a more detailed study of the rate of decline of respiration and of the factors that influence it under various conditions. Further, by extrapolation of the rate of decline of respiration to a point immediately after application of the stimulating agent, information about transient metabolic events that would otherwise have escaped observation was obtained. The work includes studies on the effects of sodium, potassium, lithium, rubidium, caesium, choline, ammonium, magnesium, and calcium, singly and in combination, and at a wide range of concentrations. The course as well as the level of the oxygen uptake of the slices was studied.

METHODS

Adult rats of the Wistar strain were decapitated while under light ether anaesthesia; the anaesthesia did not affect the oxygen uptake of the brain slices. The brains were removed immediately after decapitation and transferred to the cold room (2-4°) within 90 sec. after the death of the animals (Wallgren & Kulonen, 1960).

Brain-cortex slices were cut without moistening by means of a Stadie-Riggs microtome designed to cut slices about 0.5 mm. thick (Stadie & Riggs, 1944). The slices were cut from the lateral surface of each hemisphere, slice 1 being cut from the outer convexity of the hemisphere and slices 2 and 3 successively from the remaining tissue block. No more than three slices were cut from each hemisphere.

Each slice was immediately placed in a Warburg vessel containing 3.0 ml. of the ice-cold medium of incubation. The fresh weights of the slices were determined as the differences between the weights of the vessels before and after introduction of the slices; they usually amounted to 50-100 mg.

Within 30 min. after the decapitation of the rats, the vessels were placed in a water bath at 37°. They were oxygenated for 5 min. and then equilibrated for 5-10 min. Readings were started at 40 min. after the death of the first rat.

The oxygen-uptake rate of the slices was determined with conventional Warburg constant-volume respirometers. The manometers were read every 10 min. The oxygen-uptake rates were expressed as $\mu\text{moles/g. fresh wt./hr.}$, usually as the average of three 10 min. periods. Additions from the side arm took place after incubation for 30 min.; the first 10 min. period after the addition was disregarded. Frequently the incubations lasted several hours, and the manometers had to be readjusted to their starting-points; the first 10 min. period after the adjustment was disregarded.

Media were prepared from solutions of purest-grade chemicals. Usually the chlorides were used; magnesium was used as its sulphate. Owing to the hygroscopicity of lithium chloride and calcium chloride, the exact concentration of stock solutions of these salts was determined by a chloride titration. The media usually contained sodium chloride (125 mM); but the ionic composition was changed by substitution or addition of other salts or of sucrose as described in the Results section. Media in which all sodium chloride was replaced by iso-osmolar concentrations of sucrose, choline chloride, lithium chloride etc. are referred to as 'sucrose medium', 'choline medium', 'lithium medium' etc. The media were, unless otherwise indicated, prepared without potassium, but rapidly acquired a potassium concentration of about 2 mM owing to leakage of potassium from the cells.

The media all contained glucose (6 mM) as substrate and, unless otherwise stated, magnesium sulphate (1.0 mM) and calcium chloride (1.5 mM). They were buffered with 20 mM-tris [tris(hydroxymethyl)aminomethane plus tris(hydroxymethyl)aminomethane hydrochloride] to pH 7.4. The pH of the media was determined before and after the incubation; if the pH after the incubation had risen above 7.5 or fallen below 6.8, the experiment was discarded.

RESULTS

When brain-cortex slices are incubated in a medium with an ionic composition close to that of extracellular fluid, i.e. with a high sodium and a low potassium concentration, they consume oxygen at a rate of 70-90 $\mu\text{moles/g. fresh wt./hr.}$, and they maintain this respiratory rate relatively well. Nevertheless, on prolonged incubation of the

slices, the oxygen uptake shows a gradual fall and at the end of 2–2½ hr. is 10–15% lower than that at the start of the incubation (Fig. 1, upper half).

Such a decrease of the respiratory rate is observed with slices originating from the surface of the cerebral hemisphere (slices 1) as well as with slices cut from deeper layers of the organ (slices 2 and 3). But the absolute levels of the oxygen uptakes vary considerably among the slices. It is about 10% lower in systems containing slices 2 than in systems with slices 1, and still lower with slices 3.

Differences between the oxygen uptakes of slices cut from different layers of the brain have been a consistent finding in our experiments, and we have further observed that the oxygen uptakes of the slices may vary in relation to one another as the experimental conditions are changed. We conclude that comparisons between different experimental situations are valid only when slices cut from the same layer of the brain, i.e. with the same cutting number, are compared. The results presented below are, unless otherwise indicated, all derived from experiments with slices 1; but qualitatively the same phenomena can be observed with slices 2 and 3.

The rather good maintenance of the oxygen uptake depends on the presence of sodium in the medium. In the lower half of Fig. 1 are shown oxygen uptakes of slices 1, 2 and 3 incubated in media in which all the sodium chloride has been replaced by an equiosmolar concentration of sucrose. Under these circumstances the respiration of all three slices shows a rapid fall. This cannot be due to a toxic action of the sucrose, since respiration is maintained in media containing both sodium and sucrose; it must therefore be caused by the lack of sodium in the medium. Slices 1 are affected more by the lack of sodium than are slices 2 and 3.

Fig. 2 shows the respiration of slices 1 in media with sodium concentrations varying between 125 and 0 mM. The oxygen uptakes are in Fig. 2 plotted semilogarithmically, and decreases of the respiratory rates, except at very low values, follow rather closely exponential courses. When the sodium concentration is lowered from 125 to 20 mM, the oxygen uptake decreases somewhat, but it is equally well maintained, as shown by the identical slopes of the two lines. With a sodium concentration of 5 mM in the medium there is a decline in respiration which is intermediate between those obtained with concentrations of 20 and 0 mM.

When brain slices are exposed to media with a very low sodium concentration, they rapidly suffer irreversible damage. When sodium chloride is added to slices that have been incubated for a time in a sucrose medium, the fall of the respiration

becomes less steep, but the oxygen uptake does not rise and remains at a lower level than that observed in slices kept in a control medium from the start of the incubation.

With a sodium concentration of 250 mM the respiration is somewhat lower and less well maintained than in the control medium (125 mM); this is presumably mainly a result of the considerable hyperosmolarity, since a similar or somewhat steeper fall is seen when hyperosmolarity is produced by adding 250 mM-sucrose to the control medium. A moderate hyperosmolarity, such as that produced by increasing the sodium chloride in the control medium by about 50 mM or by about 100 mM, does not alter the oxygen uptake of the slices significantly.

Fig. 3 shows the effect of replacing all the sodium chloride in the medium by iso-osmolar concentrations of choline chloride, caesium chloride, rubidium

Fig. 1. Oxygen-uptake rates of rat-brain slices incubated in media with 125 mM-sodium chloride (open symbols) or with 250 mM-sucrose (closed symbols). Slices 1: ○ and ●; 2: △ and ▲; 3: □ and ■. Results are the means of two to six slices.

Fig. 2. Oxygen-uptake rates of rat-brain slices in media with 125 mM- (○), 20 mM- (△), 5 mM- (□) and 0 mM- (●) sodium chloride, iso-osmolarity being maintained with sucrose. Results are the means of two to six slices 1.

Fig. 3. Oxygen-uptake rates of rat-brain slices in media with 125 mM-sodium chloride (O) and in media in which the sodium chloride has been replaced by 125 mM-choline chloride (Δ), 125 mM-caesium chloride (□), 125 mM-rubidium chloride (▲), 125 mM-potassium chloride (■), 125 mM-lithium chloride (▽) or 250 mM-sucrose (●). Results are the means of two to six slices l.

Fig. 4. Oxygen-uptake rates of a single rat-brain slice before and after the addition (arrow) of potassium chloride (final concn. 48 mM) to a medium containing 125 mM-sodium chloride.

Fig. 5. Oxygen-uptake rates of rat-brain slices before and after the addition (arrow) of 16 mM- (□), 32 mM- (Δ), 48 mM- (O) and 384 mM- (●) potassium chloride to a medium containing 125 mM-sodium chloride. Extrapolated oxygen-uptake rates are indicated by broken lines. Results are the means of two or three slices l.

chloride, potassium chloride, lithium chloride or sucrose. In these experiments also the oxygen uptakes fall exponentially, and an extrapolation to the start of the incubation (indicated in Fig. 3 by broken lines) seems therefore permissible. In most of the media the initial rate of oxygen uptake is almost the same as in the sodium medium; only the systems containing potassium and lithium have higher initial rates of oxygen uptake. The systems differ considerably as regards the rapidity with which respiration falls during the incubation, the decline being slowest in the sodium and choline media and highest in the lithium and sucrose media. Rates of oxygen consumption have also been studied in media with ammonium chloride (125 mM) instead of sodium chloride (results not presented); in this system the respiration falls even more rapidly than in a sucrose medium. Experiments carried out with lower concentrations of the ions (and iso-osmolarity maintained with sucrose) showed that, although choline at a concentration of 125 mM maintains respiration as well as does sodium, a lowering of the choline concentration to 40 mM leads to a fall of the oxygen uptake which is as steep as that observed in a medium with a sodium concentration of only 5 mM. The cations may consequently be arranged according to their ability to maintain respiration under these conditions in the following decreasing order: Na^+ , choline, Cs^+ , Rb^+ , K^+ , Li^+ , NH_4^+ .

The media employed were prepared without potassium. Brain slices *in vitro* lose some of their potassium, and flame-photometric analyses of filtered media at the end of the incubation show that they contain potassium at a concentration of about 2 mM, which may affect the respiration.

The addition of potassium chloride to slices which are incubated in a sodium medium leads to a prompt rise of the oxygen consumption. This is followed by a relatively rapid decrease, and the respiratory rate eventually falls to values lower than those that would have been observed if no potassium chloride had been added (Fig. 4). The oxygen uptake rises from an unstimulated value of 90 μmoles/g. fresh wt./hr. to a value which in the first 10 min. period recorded after the addition is 150 μmoles/g. fresh wt./hr. (increase of 65%).

Since the incubation period started 5–10 min. after the addition of potassium chloride, and since the decline in the respiration was rapid, the initial value of the stimulated respiration might be more correctly obtained by extrapolating backwards to a point immediately after the addition of potassium chloride. This has been done in Fig. 5 where each point is the average of three 10 min. periods and the average of experiments with two or three slices. Since the stimulated respiration has an exponential decline, extrapolation is justifiable.

Fig. 5 also shows the results of experiments with the addition of potassium chloride to give lower and higher final concentrations. When the potassium concentration is lower than 20 mM, it does not affect the oxygen uptake of the slices measurably. With a potassium concentration of 32 mM, the respiration shows a moderate increase, though its rate of decline remains unchanged. At a concentration of 48 mM potassium gives a stimulation of respiration of about 90% followed by an accelerated decline. When the potassium concentration is raised to still higher levels, the initial stimulation is less and the fall of the respiration becomes gradually steeper.

The results presented in Fig. 5 are derived from experiments with slices 1 only. Similar experiments performed with slices 2 and 3 show that the initial stimulation of the respiration after the addition of potassium chloride is somewhat smaller than that observed with slices 1; but, when it is expressed as a percentage of the unstimulated respiration, it is of the same magnitude. The fall of the unstimulated oxygen uptake is somewhat less steep in experiments with slices 2 and 3 than in those with slices 1.

The effect produced by potassium varies with the concentration of sodium in the medium. When potassium (final concentration 48 mM) is present in a system which contains a sodium concentration of 250 mM instead of 125 mM (Fig. 6), there is only a moderate stimulation of the oxygen uptake, and the stimulated respiration falls very slowly; the effect resembles closely that which is observed with a potassium concentration of 32 mM in a medium whose sodium concentration is 125 mM. If the sodium concentration in the medium is lowered to 20 mM, the respiration falls rapidly; since under these conditions the decrease in respiratory rate is not exponential, the initial stimulation of the respiration, although present, cannot be evaluated quantitatively.

When a much higher potassium concentration is present in media with sodium concentrations of 20, 125 and 250 mM, results like those shown in Fig. 7 are obtained. Under these conditions the stimulated respiration falls in all three media with slopes that are identical and rather steep. The initial value of the stimulated respiration is highest in the system with a sodium concentration of 125 mM. It is somewhat lower, both absolutely and when expressed as a percentage of the unstimulated respiration, with a sodium concentration of 250 mM, and still lower with one of only 20 mM.

Fig. 8 shows the initial value of the stimulated respiration (obtained by extrapolation) as a function of the final potassium concentration; experiments were carried out in media with sodium concentrations of 125 and 250 mM. The unstimu-

lated respiration (no potassium added) is somewhat lower in the hyperosmolar medium than in the control medium. In neither of the two media is any stimulation produced by potassium concentrations that are lower than 10–15 mM, but as the potassium concentration is raised to 48 mM, a stimulation becomes apparent, and it increases more rapidly

Fig. 6. Oxygen-uptake rates of rat-brain slices before and after the addition (arrow) of potassium chloride (final concn. 48 mM) to media containing 250 mM- (Δ), 125 mM- (○) and 20 mM- (□) sodium chloride. Results are the means of two to six slices 1.

Fig. 7. Oxygen-uptake rates of rat-brain slices before and after the addition (arrow) of potassium chloride (final concn. 384 mM) to media containing 250 mM- (Δ), 125 mM- (○) and 20 mM- (□) sodium chloride. Results are the means of two or three slices 1.

Fig. 8. Oxygen-uptake rates of rat-brain slices immediately after the addition of potassium chloride to media containing 250 mM- (Δ) and 125 mM- (○) sodium chloride. The values are obtained by extrapolation. Results are the means of two or three slices 1.

and reaches a higher level with 125 mM than with 250 mM as the sodium concentration. In the latter medium the initial stimulated respiration shows a further rise with the potassium concentration greater than 48 mM and within the entire range of the potassium concentrations studied (upper level 192 mM). With a sodium concentration of 125 mM it decreases gradually when the potassium concentration is raised from 48 to 192 mM; it remains at a higher level with 125 mM than with 250 mM as the sodium concentration.

Ions other than K^+ produce, when added to brain-cortex slices incubated in a medium with a sodium concentration of 125 mM, an initial rise of the oxygen uptake and a subsequent exponential fall which is steeper than in the unstimulated system. The addition of a wide range of concentrations of the following ions, ammonium, caesium, rubidium, lithium, and choline, produces effects similar to those of potassium, but the ions differ markedly in the concentrations necessary to produce these effects.

When the ammonium concentration is as little as 3 mM with slices in a control medium, there is a noticeable rise of the respiratory rate, and a final ammonium concentration of 12 mM produces a stimulated oxygen uptake which initially is about 50 % higher than the unstimulated value (Fig. 9). At all ammonium concentrations greater than 3 mM the respiratory rate falls rapidly, and with more than about 25 mM an extrapolation of the oxygen uptake to the initial value is no longer possible.

Caesium is almost as active as ammonium, but apparently less toxic. The initial stimulated respira-

tion is maximal, 50–60 % above the unstimulated oxygen uptake, when caesium is added to give a final concentration of about 20 mM (Fig. 9). The stimulated oxygen uptake at this caesium concentration falls with about the same rate as that observed after addition of potassium (final concentration 125 mM) to the control medium. Little change is produced in the decline of respiration when the caesium concentration is raised to 192 mM, whereas the initial value of the stimulated respiration becomes slightly lower.

Rubidium resembles potassium closely in its influence on both the initial stimulated respiration and the rate of the subsequent fall of the respiration.

When lithium is used as a stimulating ion in a system containing a sodium concentration of 125 mM, it produces qualitatively the same effects as do potassium, ammonium, caesium, and rubidium, but quantitative differences are apparent. Fig. 9 shows that lithium under these conditions is somewhat less active than the others, since a concentration of about 100 mM is needed to evoke a full respiratory response. At this lithium concentration the stimulated oxygen uptake falls rapidly (Fig. 10).

The choline ion has been found to possess some stimulating activity. The respiration of the slices is not influenced by a choline concentration of 125 mM in the control medium; but a concentration of 250 mM produces a rise of the oxygen uptake which initially amounts to about 25 % of the unstimulated value. The initial rise is followed by an increased rate of decline in respiration. This does not, in the concentrations used by us, affect respiration.

Fig. 9. Oxygen-uptake rates of rat-brain slices immediately after the addition of ammonium chloride (●), caesium chloride (◻), rubidium chloride (▲), potassium chloride (○), lithium chloride (▽) and choline chloride (△) to media containing 125 mM-sodium chloride. The values are obtained by extrapolation. Results are the means of two to four slices l.

As with potassium, the effects of the other stimulating ions are influenced by the sodium concentration in the medium. In a medium with a sodium concentration of 20 mM, a stimulation can be seen with lower concentrations of the test ions than when it is 125 mM, whereas with a sodium concentration of 250 mM more test ion must be added to produce the same stimulation. The stimulated oxygen uptake shows a less rapid fall in the presence of a high sodium concentration. Figs. 10 and 11 illustrate these phenomena with lithium.

A certain amount of sodium must be present in the medium for the stimulating ions to exert their activity; if sodium is removed from the medium, no stimulation can be produced. Fig. 3 showed the oxygen uptakes of brain slices incubated in media in which all sodium had been replaced by equimolar (125 mM) concentrations of potassium and other stimulating ions. In media with choline, caesium, and rubidium instead of sodium the initial respiratory rate is the same as in the sodium medium. But even if in the potassium medium none of the oxygen uptake values actually recorded exceeds the unstimulated respiration, an extrapolation to the start of the incubation strongly indicates an initial respiration which is at a slightly, but distinctly, higher level. The brain slices incubated in a lithium medium show an initial oxygen uptake which is almost twice as high as the unstimulated value.

These observations on lithium and potassium might mean either that these can stimulate the respiration without the assistance of sodium as a cofactor, or that their requirement for sodium is (at least partially) fulfilled by the small amounts of sodium still present in or attached to the slices under the experimental conditions employed. To clarify this point experiments were carried out in which slices were 'leached' in the cold for 30 min. in media with sodium concentrations varying from 0 to 250 mM and then incubated at 37°. Such experiments were performed both with potassium (125 mM) and with lithium (125 mM) in the medium and Fig. 12 shows the oxygen uptakes immediately after the start of the incubation; the values were obtained in the usual way by extrapolation.

A difference between potassium and lithium is brought to light by the procedure described. For potassium, the 'leaching' in a sodium-free medium with its removal of traces of sodium from the slices has abolished any potassium stimulation. With a sodium concentration of 5 mM, potassium produces a slight, but significant, stimulation, and with a sodium concentration of 10 mM the potassium-stimulated respiration reaches a high value. When the sodium concentration is raised from 10 to 125 mM, the absolute values of the stimulated

respiration show a moderate increase; but when the stimulated respiration is expressed as a percentage of the corresponding unstimulated oxygen uptake, it is essentially constant between the sodium concentrations 10 and 125 mM. With a sodium concentration of 250 mM the stimulated respiration is somewhat lower.

Fig. 10. Oxygen-uptake rates of rat-brain slices before and after the addition (arrow) of 192 mM- (●), 96 mM- (○), 48 mM- (△) and 12 mM- (□) lithium chloride to media containing 125 mM-sodium chloride. Results are the means of two to four slices 1.

Fig. 11. Oxygen-uptake rates of rat-brain slices before and after the addition (arrow) of 192 mM- (●), 96 mM- (○) and 48 mM- (△) lithium chloride to media containing 250 mM-sodium chloride. Results are the means of two slices 1.

Fig. 12. Oxygen-uptake rates of rat-brain slices immediately after the start of incubation in media containing 125 mM-potassium chloride (■) or 125 mM-lithium chloride (▽) and with sodium chloride concentrations varying from 0 to 250 mM. Results are the means of two to five slices 1.

Brain slices leached and thereafter incubated in a sodium-free medium containing lithium (125 mM) consume oxygen at a very high rate; their respiration is almost twice that observed in the corresponding potassium medium. Addition of sodium to the medium leads to a decrease of the lithium-stimulated respiration, and when the sodium concentration is 50–125 mM, the initial respiration in the lithium media is equal to or somewhat lower than that obtained in the corresponding potassium media. We conclude that lithium can produce a 'stimulated' respiration also in the absence of sodium in the medium; this stimulation is counteracted by the addition of sodium to the system.

The concentration of sodium necessary for a maximum potassium stimulation, i.e. a rise of the oxygen uptake to 85–90 % above the unstimulated value, varies somewhat with the amount of potassium added. It appears from Fig. 12 that at a final potassium concentration of 125 mM the 'critical' sodium concentration is about 10 mM. It is almost the same at a potassium concentration of 50 mM. But if considerably more potassium is added, e.g. to give a concentration of 384 mM (cf. Fig. 7), not even a sodium concentration of 20 mM suffices to give a full stimulation. The critical sodium concentration has under these circumstances risen to above 50 mM.

To investigate whether ions other than sodium might be able to serve as cofactors for the stimulating ions, experiments have been carried out with combinations, two by two, of potassium, caesium, rubidium, and choline at a variety of concentrations. No evidence of a sodium-like effect was obtained, and sodium seems to be unique in its ability to support the action of these stimulating ions. The addition of potassium to a choline medium leads to a decrease of the oxygen uptake to a slightly, but significantly, lower level; the maintenance of the respiration at this new level is as good as in a choline medium to which no potassium had been added. This observation may indicate that a certain amount of choline has been displaced by potassium.

The bivalent cations of magnesium and calcium also exert an influence on the respiration of brain-cortex slices. In the experiments described so far the media contained magnesium (1.0 mM) and calcium (1.5 mM). Omission of magnesium from the medium does not affect either the unstimulated or the stimulated oxygen uptake significantly; but an increase of the magnesium concentration to 10 mM counteracts the initial stimulation as well as the subsequent fall of the respiration which is produced by a potassium concentration of 50 mM. The effect of a high magnesium concentration is still more pronounced if calcium is omitted from the medium; the stimulating action of potassium is

almost completely abolished, and neither an increase of the potassium concentration to 200 mM nor of the sodium concentration to 250 mM is able to restore it. Also the lithium effect is diminished in a medium with magnesium (10 mM) and no calcium, although to a smaller extent.

The omission of calcium from a medium with unaltered magnesium concentration increases the oxygen uptake. But under the experimental conditions employed by us this rise in respiratory rate was negligible. An increase of the calcium concentration to 10 mM, on the other hand, led to a significant lowering of the unstimulated oxygen uptake, though it did not interfere with the potassium stimulation of the respiration. An increase of the sodium concentration to 250 mM did not counteract the calcium inhibition of the unstimulated oxygen uptake.

DISCUSSION

When brain slices are incubated *in vitro* in media of different compositions they do not form a completely stable system. The metabolism of the slices undergoes a gradual change as the incubation proceeds, and since this can be observed even when the system has been fortified with sufficient substrate to cover the nutritional requirements of the slices, we must presumably be dealing with other mechanisms, such as an exhaustion of endogenous cofactors, an accumulation within the cells of metabolic inhibitors, or a gradual disintegration of those tissue structures which are responsible for the metabolic survival.

The change in metabolic activity is strongly influenced by the conditions of incubation, and among these by the ionic milieu to which the slices are exposed. In a 'balanced' medium with a composition resembling that of extracellular fluid, the respiratory rate of the slices changes only slowly, though under other circumstances, e.g. in a sodium-free medium or in a medium to which a high concentration of potassium has been added, the change is much more rapid. Since usually the decline in respiratory rate is exponential, the initial respiration may be estimated by extrapolation to the start of the incubation.

The requirement for sodium in the maintenance of the unstimulated respiration is in agreement with reports in the literature concerning the effect of sodium-free media (Elliott & Libet, 1942; Gore & McIlwain, 1952). But according to Takagaki & Tsukada (1957) the unstimulated oxygen uptake is not influenced by a removal of sodium from the medium. In their experiments the sodium chloride in the medium was, however, replaced by choline chloride, and our experiments show that the choline ion exerts a maintenance activity on the

unstimulated respiration which is equal to that of the sodium ion. The similarity of the two ions is limited to systems in which glucose is the substrate; Takagaki & Tsukada (1957) showed that a substitution of choline for sodium leads to a considerable increase of the respiration in media with pyruvate as substrate and a decrease with such substrates as fructose, lactose, citrate, and glutamate.

These observations emphasize that choline does not fulfil only an osmotic function. Such an application of choline must be based on experiments which directly exclude any specific effects of this ion in the system under investigation.

Also univalent cations other than sodium and choline exhibit maintenance activities, although to very different extents. When the ions are substituted, one by one, for sodium in the medium, their ability to maintain the unstimulated oxygen uptake decreases in the order: Na^+ , choline, Cs^+ , Rb^+ , K^+ , Li^+ , NH_4^+ . Lithium presents the double feature of an initial stimulation and a rapid decline of the oxygen uptake, and this presumably explains the apparent disagreement between our results and those of Pappius, Rosenfeld, Johnson & Elliott (1958), who with incubations for 1 hr. found identical oxygen uptakes of brain slices in sodium and in lithium media. Further, our media were prepared without potassium, whereas Pappius *et al.* (1958) used media with a potassium concentration of 5 mM; the lithium effects are less pronounced under the latter than under the former circumstances.

Brain and muscle seem to be peculiar in their requirement for sodium to maintain respiration during incubation (L. Hertz & T. Clausen, unpublished work). Slices of kidney, liver and peripheral nerve maintain their oxygen consumption equally well in a sodium-free sucrose medium and in a medium with a sodium concentration of 125 mM, even if in some cases the oxygen uptake in a sodium-free medium is at a lower level.

When potassium is added to brain slices incubated in a sodium-containing medium, a number of metabolic changes take place (McIlwain, 1959; Heald, 1960); in the present work only the oxygen uptake was followed. It appears from our experiments that the respiratory rate after the addition of potassium passes through two phases: first, a prompt rise to a value which, under optimum conditions, is about 180–190 % of the unstimulated oxygen uptake; secondly, a relatively rapid exponential fall. This double effect has not been noted regularly in previous studies of the effect of potassium. Some authors (e.g. Dickens & Greville, 1935) mention that there is a fall of the respiratory rate; but most reports do not allude specifically to the course of the unstimulated oxygen uptake.

However, in some reports the stimulated oxygen uptake does show a fall and this fall is more rapid than in an unstimulated system (Tsukada, Takagaki & Hirano, 1958; Beer & Quastel, 1958*a, b*). A fall of the stimulated respiratory rate may also be observed when the stimulation has been produced with electrical pulses (Cummins & McIlwain, 1961; Wallgren & Kulonen, 1960).

Potassium is not the only cation that is able to stimulate the oxygen uptake of brain slices. Dickens & Greville (1935) noted that when caesium and rubidium were added, to give concentrations of 0.1 M, to brain slices, the respiratory rate showed an increase similar to that obtained after the addition of potassium. Lithium did not influence respiration noticeably in their experiments; but Canzanelli *et al.* (1941–42) and Rybová (1959) reported a stimulating action also of this ion, even if a higher concentration of lithium than of potassium was needed for a maximum metabolic response. A stimulating action of ammonium has been observed by Weil-Malherbe (1938), Gore & McIlwain (1952) and Takagaki & Tsukada (1957).

According to our experiments also the choline ion can stimulate respiration in the presence of sodium, but the maximum value of the stimulated oxygen uptake is only half that obtained with all the other ions. The explanation is presumably that such a high choline concentration is needed for stimulation that the hyperosmolarity of the medium depresses the oxygen uptake.

The stimulating ions differ considerably as regards the concentration of each which is required to produce stimulation. Ammonium is found to be, for example, 10 times as active as rubidium and potassium and 25 times as active as lithium (cf. Fig. 9), and the ions may be arranged in the following order with respect to their stimulating activity:

It is likely that these 'stimulating activities' are an expression of the relative affinities of the ions to particular 'stimulating' sites in or on the brain cells.

From the findings shown in the Results section as well as others not presented we can arrange the ions in the following order as regards the activity with which they produce a more rapid fall of the stimulated respiration:

This series is very similar to that given above for the stimulating activity (or affinity) of the ions, and a close connexion between the two phenomena is highly probable. Only lithium occupies a different position in the two series; its somewhat stronger effect on the fall of the stimulated

respiration than on the initial rise is presumably a consequence of the particular potency of this ion in the acceleration of the respiratory decrease which was observed also in the unstimulated system.

It is characteristic of the respiratory stimulation by potassium that there is an obligatory requirement for sodium in the medium (Dickens & Greville, 1935; Canzanelli *et al.* 1941-42; Tsukada & Takagaki, 1955). Ammonium, caesium, rubidium, and choline all resemble potassium in this respect, whereas lithium is able to raise the oxygen uptake of the slices without the aid of this cofactor, presumably owing to a partial resemblance of this ion to both sodium and potassium (Schou, 1957).

Not only is a certain sodium concentration in the medium necessary for the production of a potassium stimulation, but the ion is unique in this respect and cannot be replaced by any other ion. Not even choline, which shows such a similarity to sodium as regards the maintenance of the unstimulated oxygen uptake, is able to replace it in the stimulated system.

There are accordingly different functions for the sodium ion: an ability to maintain respiration in the unstimulated system (this is shared by other ions) and a function as a cofactor for stimulating agents such as electrical pulses and potassium or other cations; this latter is specific for sodium.

Pappius *et al.* (1958) showed that the potassium content of incubated brain slices depends on the presence of sodium in the medium, and there is apparently a 'critical' sodium concentration at about 10 mM (Fig. 1 of Pappius *et al.* 1958). The resting potentials which could be recorded from brain slices by Hillman & McIlwain (1961) disappeared in sodium-free media, but had their full value in a medium with a sodium concentration of 26 mM.

A sodium concentration of 10-20 mM is required in the medium surrounding isolated muscles and nerves to preserve their excitability to electrical pulses (Overton, 1902; Höber, 1905). This effect of sodium is also specific; sodium cannot be replaced by such cations as rubidium, caesium, and ammonium. Only lithium can replace sodium as the sole alkali metal ion, but, although under these circumstances the excitability initially remains unchanged, it does eventually suffer an irreversible breakdown. This may be related to the rapid fall of the respiration of brain slices in a lithium medium.

In addition to the two sodium effects on respiration outlined above one may, at sodium concentrations exceeding about 20 mM, observe what appears to be yet a third action of this ion: an antagonism against the effect of the stimulating ions. Both the initial stimulation and the subsequent fall of the stimulated respiration are

counteracted by an increase of the sodium concentration in the medium. This effect of sodium can be opposed by addition of more stimulating ion, and we may therefore be dealing with a competition for one or more sites on an enzyme or at a cellular structure.

Such a competition may be of two different types. It may be one in which potassium and the other stimulating ions compete with sodium for the sites at which the latter exerts its two functions: a maintenance of the respiration and an action as cofactor for the stimulating ions. But the reverse is also possible. Sodium may compete with and displace the stimulating ions from the sites at which they exert their double action: the production of an initial rise and a subsequent fall of the respiration.

If the first hypothesis is correct, one might expect that the minimum concentration of sodium required for the production of a full stimulation would vary somewhat with the amount of stimulating ion. This, in fact, does occur. In a system with a potassium concentration of 125 mM a sodium concentration of about 10 mM is sufficient for the production of an initial respiration which is 180-190% of the unstimulated value (Fig. 12), whereas a sodium concentration greater than 50 mM is needed when potassium is added to give a final concentration of 384 mM (Fig. 7).

According to the same hypothesis one might consider the increased decline in respiration, not as a phenomenon directly related to the stimulation, but rather as the result of a displacement of sodium by an ion less able to maintain the respiration. The relatively rapid fall of the respiration after stimulation with lithium and ammonium would agree with this assumption.

A displacement of sodium by lithium is also consistent with the observation made by Pappius *et al.* (1958) that to maintain the potassium content of incubated brain slices a much higher sodium concentration is necessary in the medium when iso-osmolality is produced with sodium chloride plus lithium chloride than when it is produced with sodium chloride plus choline chloride (Fig. 1 of Pappius *et al.* 1958).

The phenomena observed after choline stimulation cannot, however, be explained in terms of the first hypothesis. When choline at a concentration of 250 mM is present with slices in a sodium medium, there is not only an initial stimulation, but also a subsequent fall of the respiration. This is more rapid than that observed in control media of the same osmolality and containing either sodium or choline alone. The increased decline in respiration can hardly be due to a displacement of sodium by choline, because the choline ion itself has such a high maintenance activity; it must be an integral

part of the stimulating action of the choline ion, i.e. in agreement with the second of the hypotheses mentioned. Both mechanisms are probably at work.

Tissue analyses have shown that during electrical stimulation brain-cortex slices lose potassium and gain sodium, whereas in the post-stimulation period the opposite ionic movements take place (Varon & McIlwain, 1962; Cummins & McIlwain, 1961). It is possible that this potassium loss during electrical stimulation constitutes the link between the respiratory stimulations produced by ions and by electrical pulses. A high potassium concentration immediately adjacent to the cell surface may, whether the ion emerged from the cell or was added to the medium, cause a depolarization (Hillman & McIlwain, 1961) and thereby, directly or indirectly, influence the metabolic processes in the cell; or it may stimulate an enzyme involved in the active transport of sodium and potassium across the cell wall.

There is a resemblance, at least qualitatively, between the brain slices investigated by us and the enzyme system described first by Skou (1957) from studies on crab nerve and found also in mammalian brain and kidney (Hess & Pope, 1957; Skou, 1961) and in human erythrocytes (Post, Merritt, Kinsolving & Albright, 1960). The enzyme, which is supposedly involved in the active transport of sodium and potassium across the cell wall, has adenosine triphosphate as its substrate, and it is characterized by a requirement for magnesium (in low concentration) and sodium. If sodium is replaced by lithium, the enzymic activity becomes lower; if it is replaced by ammonium, caesium, rubidium, or potassium, the enzymic activity becomes much lower. The addition of potassium to the system when both magnesium and sodium are present leads to a strong stimulation of the activity of the enzyme. A similar stimulation in the presence of magnesium and sodium can be produced by the addition of ammonium, rubidium, caesium, and lithium. There appears to be a competition between sodium and the stimulating ions for certain sites on the enzyme or enzyme-substrate complex (Skou, 1960).

SUMMARY

1. Univalent cations fall into three distinct groups as regards their influence on the respiratory rate of rat-brain-cortex slices incubated in media with glucose as substrate. The first group comprises sodium, which exerts the following functions: (i) It maintains the unstimulated oxygen uptake of the slices during prolonged incubation; this function is to some extent shared by choline, caesium, rubidium, and potassium. (ii) It serves, in

low concentration, as cofactor for stimulating agents (electrical pulses; potassium and other univalent ions); this function is unique for sodium. (iii) It antagonizes, in higher concentration, the stimulating cations.

2. In the second group of ions are found ammonium, caesium, rubidium, potassium, and choline; when added to slices in a sodium-containing medium, they produce an initial stimulation and a subsequent, relatively rapid, fall of the respiratory rate. The various ions evoke almost the same maximal stimulation, but the concentrations required for stimulation differ considerably among the ions, with ammonium as the most and choline as the least active ion.

3. Lithium constitutes the third group; even in the absence of sodium, it produces a high initial respiratory rate followed by a rapid fall. The ion presumably acts as its own cofactor.

4. The stimulating ions of the second and third group counteract the effect of high sodium concentrations. There is probably a competition between these ions and sodium for sites on an enzyme or at a cellular structure.

5. The unstimulated and the stimulated respiratory rates are affected by variations in the magnesium and calcium concentrations in the medium, but a competition between these ions and the univalent ions is unlikely.

The skilled technical assistance of Mrs I. Andresen and Mrs J. Lyberth is gratefully acknowledged.

REFERENCES

- Ashford, C. A. & Dixon, K. C. (1935). *Biochem. J.* **29**, 157.
- Beer, C. T. & Quastel, J. H. (1958*a*). *Canad. J. Biochem. Physiol.* **36**, 531.
- Beer, C. T. & Quastel, J. H. (1958*b*). *Canad. J. Biochem. Physiol.* **36**, 543.
- Cade, J. F. J. (1949). *Med. J. Aust.* **36**, 349.
- Canzanelli, A., Rogers, G. & Rapport, D. (1941-42). *Amer. J. Physiol.* **135**, 309.
- Cummins, J. T. & McIlwain, H. (1961). *Biochem. J.* **79**, 330.
- Dickens, F. & Greville, G. D. (1935). *Biochem. J.* **29**, 1468.
- Elliott, K. A. C. & Libet, B. (1942). *J. biol. Chem.* **143**, 227.
- Gershon, S. & Yuwiler, A. (1960). *J. Neuropsychiat.* **1**, 229.
- Gore, M. B. R. & McIlwain, H. (1952). *J. Physiol.* **117**, 471.
- Heald, P. J. (1960). *Phosphorus Metabolism of Brain*, p. 129. Oxford: Pergamon Press.
- Hess, H. H. & Pope, A. (1957). *Fed. Proc.* **16**, 196.
- Hillman, H. H. & McIlwain, H. (1961). *J. Physiol.* **157**, 263.
- Höber, R. (1905). *Zbl. Physiol.* **19**, 390.
- McIlwain, H. (1951). *Biochem. J.* **49**, 382.
- McIlwain, H. (1956). *Physiol. Rev.* **36**, 355.

- McIlwain, H. (1959). *Biochemistry and the Central Nervous System*. London: J. and A. Churchill.
- Overton, E. (1902). *Pflüg. Arch. ges. Physiol.* **92**, 346.
- Pappius, H. M., Rosenfeld, M., Johnson, D. M. & Elliott, K. A. C. (1958). *Canad. J. Biochem. Physiol.* **36**, 217.
- Post, R. L., Merritt, C. R., Kinsolving, C. R. & Albright, C. D. (1960). *J. biol. Chem.* **235**, 1796.
- Rybová, R. (1959). *J. Neurochem.* **4**, 304.
- Schou, M. (1957). *Pharmacol. Rev.* **9**, 17.
- Schou, M. (1959). *Psychopharmacologia*, **1**, 65.
- Skou, J. C. (1957). *Biochim. biophys. Acta*, **23**, 394.
- Skou, J. C. (1960). *Biochim. biophys. Acta*, **42**, 6.
- Skou, J. C. (1961). In *Symposium on Membrane Transport and Metabolism*, p. 228. Prague: Czechoslovak Academy of Sciences.
- Stadie, W. C. & Riggs, B. C. (1944). *J. biol. Chem.* **154**, 687.
- Takagaki, G. & Tsukada, Y. (1957). *J. Neurochem.* **2**, 21.
- Tsukada, Y. & Takagaki, G. (1955). *Nature, Lond.*, **175**, 725.
- Tsukada, Y., Takagaki, G. & Hirano, S. (1958). *J. Biochem. Tokyo*, **45**, 489.
- Varon, S. & McIlwain, H. (1962). *J. Neurochem.* **8**, 262.
- Wallgren, H. & Kulonen, E. (1960). *Biochem. J.* **75**, 150.
- Weil-Malherbe, H. (1938). *Biochem. J.* **32**, 2257.

Biochem. J. (1962) **85**, 104

Glutathione, Oxidized and Reduced, in some Dipterans Treated with 1,1,1-Trichloro-2,2-di-(*p*-chlorophenyl)ethane

BY H. LIPKE* AND JILL CHALKLEY

Ross Institute, London School of Hygiene and Tropical Medicine (University of London), London, W.C. 1

(Received 26 February 1962)

It has been shown that DDT-resistant houseflies (*Musca domestica*) are capable of dehydrochlorinating massive doses of 1,1,1-trichloro-2,2-di-(*p*-chlorophenyl)ethane (DDT) to the innocuous olefin DDE, 1,1-dichloro-2,2-di-(*p*-chlorophenyl)ethylene. In susceptible strains this detoxicating mechanism operates to a very limited extent (Lipke & Kearns, 1960). Extracts of resistant flies catalyse this conversion to a degree several thousand-fold that of preparations from susceptible strains when fortified with glutathione, indicating that DDT-dehydrochlorinase activity may play a major role in the survival of resistant populations. With the partial exception of cysteinylglycine, other cofactors or thiols do not function as activators of the dehydrochlorinase. Under the conditions of the dehydrochlorinase assay *in vitro*, oxidation of GSH cannot be detected (Lipke & Kearns, 1959*a, b*), and in the intact animal measurements of GSH oxidation during DDT treatment have not been reported. Cotty & Henry (1958) found minor differences in rates of incorporation of [³⁵S]cysteine into GSH in resistant and susceptible strains of flies; the effect of DDT treatment was not investigated.

DDT poisoning is characterized by a striking increase in the respiratory rate of susceptible strains. A variety of substrates have been reported for the DDT-dependent increment of respiration (Winteringham & Barnes, 1955; Winteringham, 1958), but oxidation of thiols has not been investigated. In vertebrates, GSH apparently does not serve as a

major substrate or carrier in electron transport although some involvement has been proposed in special cases (Waley, 1959; Mapson, 1959). GSH concentrations in DDT-poisoned insects are of added interest since dipterans are provided with high activities of quinone-forming oxidases (Ohnishi, 1959), whose action may interfere with thiol metabolism (Barron, 1951).

A NADPH-dependent mixed-function oxidase that hydroxylates DDT has been described in resistant and susceptible flies by Agosin, Michaeli, Miskus, Nagasawa & Hoskins (1961). These workers suggest that utilization of NADPH by the hydroxylase may limit the supply of GSH available to the dehydrochlorinase. This proposal assumes that GSH is limiting in the fly and is metabolized during dehydrochlorination. This situation would be reflected by appropriate changes in GSH and GSSG content or major differences in GSSG-reductase activity in resistant and susceptible flies. The relation of concentrations of GSH to the metabolic changes occurring during DDT poisoning and detoxication have consequently been studied in the housefly and a DDT-resistant malaria vector, *Anopheles albimanus*.

EXPERIMENTAL

Resistant and susceptible strains of flies and mosquitoes are referred to throughout as R and S respectively. Quantities and concentrations of oxidized glutathione are described in terms of the equivalents of glutathione which they would yield on reduction. The insecticide 1,1,1-tri-

* Present address: Physiology Division, Army Chemical Center, Maryland, U.S.A.