REFERENCES

- Baddiley, J., Buchanan, J. G., Handschumacher, R. E. & Prescott, J. F. (1956). *J. chem. Soc.* p. 2818.
 Bender, M. L. & Glasson, W. A. (1960). *J. Amer. chem. Soc.* **82**, 3336.
 Crook, E. M., Mathias, A. P. & Rabin, B. R. (1960). *Biochem. J.* **74**, 234.
 Findlay, D., Mathias, A. P. & Rabin, B. R. (1960). *Nature, Lond.*, **187**, 601.
 Heppel, L. A. & Whitfield, P. R. (1955). *Biochem. J.* **60**, 1.
 Herries, D. G., Mathias, A. P. & Rabin, B. R. (1962). *Biochem. J.* **85**, 127.
 Morales, M. L. (1955). *J. Amer. chem. Soc.* **77**, 4169.
 Verhoek, F. H. (1936). *J. Amer. chem. Soc.* **58**, 2577.
 Weissberger, A., Proskauer, E. S., Riddick, J. A. & Toops, E. E., jun. (1955). *Techniques of Organic Chemistry. VII. Organic Solvents; Physical Properties and Methods of Purification*, 2nd ed. London: Interscience Publishers Inc.

Biochem. J. (1962) **85**, 139

The Active Site and Mechanism of Action of Bovine Pancreatic Ribonuclease

5. THE CHARGE TYPES AT THE ACTIVE CENTRE

By DOROTHY FINDLAY, A. P. MATHIAS AND B. R. RABIN

Department of Biochemistry, University College London, Gower Street, London, W.C. 1

(Received 15 March 1962)

An understanding of the effects of solvents on the acid dissociation constants of buffers and ionizing groups on proteins is an essential prerequisite for interpreting solvent effects on enzyme reactions. According to Mandel (1958) the shift in the pK of an acid in passing from water to another solvent depends primarily on the nature of the solvent and the ionizing group (CO_2H , OH , NH_3^+) and rather little on the exact nature of the complete molecule. This conclusion is substantiated by the work of Mandel & Decroly (1960) on carboxylic acids in formamide. Although dioxan and formamide differ sharply in their physical properties, both cause the pK of acetic acid to increase, but to different extents.

Acids may conveniently be divided into two main groups: (a) neutral acids, whose dissociation involves charge separation, e.g. carboxylic acids:

(b) cationic acids, whose dissociation does not involve charge separation, e.g. amines:

The pK values of neutral acids have been measured by a number of workers (Bacarella, Grunwald, Marshall & Purlee, 1955, 1958; Bates & Schwarzenbach, 1955; Harned & Owen, 1950; Mandel & Decroly, 1960; Shedlovsky & Kay, 1956; Verhoek, 1936) in a variety of solvents and mixed solvent systems. In all instances the pK increased as water was replaced by an organic solvent. The magnitude of this increase depends on the solvent,

its concentration and to a minor extent on the acid itself. The pK of acetic acid increases from 4.76 in water to 6.82 in formamide (Mandel & Decroly, 1960), 9.72 in anhydrous methanol (Bacarella *et al.* 1955) and 10.14 in aq. 82% (w/w) dioxan (Harned & Owen, 1950, p. 581).

Cationic acids have been studied less thoroughly than carboxylic acids. The pK values of substituted anilines decrease on the addition of dioxan (Willi, 1957) or methanol (Bacarella *et al.* 1955), and the pK of triethanolamine decreases on the addition of ethanol (Bates & Schwarzenbach, 1955). Verhoek (1936) found that the pK values of neutral acids increased whereas those of cationic acids decreased on the addition of formamide. The dissociation constants of some amino acids and related compounds have been determined in water and in aq. 20% (w/w) dioxan by Duggan & Schmidt (1942). They found that the decrease in the pK of the amino group was much less than the increase in the pK of the carboxyl group.

The existence of a pH optimum in an enzyme reaction implies a requirement for at least two ionizing groups for activity, one in the base form and the other in the acid form. Solvents will in general affect the binding of the substrate to the enzyme in a non-predictable fashion. Accordingly it is best to conduct experiments involving solvents under conditions where the substrate concentration is not rate-limiting. The effects of solvents will then be due to: (a) changes in the conformation of the protein; (b) changes in the pK values of buffers and the groups at the catalytic site; (c) effects on

the solvation of the transition state relative to the reactants; (d) changes in the concentration of water, if this is involved as a reagent.

Variation of the nature and concentration of the solvent can be used to decide whether (a) and (c) are important factors. If conditions can be found where (b) is the dominant factor then solvents can be used to determine the charge types of groups at the catalytic site. In general effects due to (d) can be detected and allowance made for them. There are four possible combinations of ionizing pairs that could occur at the active site of an enzyme, which control the rising and falling sides of pH-activity curves, namely: (i) neutral-cation, (ii) cation-neutral, (iii) neutral-neutral, (iv) cation-cation. These can be distinguished by the use of solvents provided that all factors other than (b) are allowed for or eliminated.

To analyse the situation it is convenient to consider the effects on the buffer and protein separately. If the addition of solvents changes the pK of a buffer by an amount $+x$ then the pH of the solution will change by $+x$. If the pH-activity curves are plotted against the pH the solution would have in the absence of solvent (not the pH in the actual solvent mixture) the curve will be displaced along the pH-axis by an amount equal to $-x$, ignoring for the moment the effects on the ionizing groups of the protein. The effects due to changes in the pK values of groups at the catalytic site of the protein are in the opposite direction to those due to changes in the pK values of the buffers. Thus if the pK value changes by an amount $+x$ the appropriate side of the pH-activity curve will shift by an amount $+x$, in the absence of changes in the buffer pK . It can be assumed that changes due to the altered pK values of the buffer and protein are additive. In the following analysis we have assumed that all neutral acids, whether buffers or groups in the protein, change their pK values by $+n$ and all cationic acids by an amount $-c$ on addition of solvent. The effects to be expected are shown in Table 1 and Fig. 1. The curves of Fig. 1 are drawn for a hydrolytic reaction and the maximum activity in the presence of the solvent

has been shown as less than that in the absence of solvent to take account of the lowered molar concentration of water.

The pH (water)-activity curves in Fig. 1 for the case (i), neutral-cation in solvent-water, will tend

Fig. 1. Theoretical pH (water)-activity curves in water (full lines) and in a solvent-water mixture (broken lines) for the possible ionizing pairs at the active centre of an enzyme: (i) neutral-cation, (ii) cation-neutral, (iii) neutral-neutral and (iv) cation-cation in two series of buffers: (a) neutral-acid buffers and (b) cationic-acid buffers. The ordinate is a measure of the maximum velocity. The abscissa is the pH with water replacing the solvent which is assumed to affect the dissociations in the manner shown in Table 1.

Table 1. Predicted shifts along the pH-axis of a side of the pH-activity curve on the addition of a solvent

It is assumed that the pK values of neutral acids and cationic acids shift an amount $+n$ and $-c$ respectively. The pH-axis is that of the solutions with water replacing the solvent.

Nature of group on protein	Nature of buffer	Shift of pH-activity curves due to change of buffer pK	Shift of pH-activity curves due to change of pK of group on protein	Net shift
Neutral acid	Cationic	$+c$	$+n$	$+(c+n)$
	Neutral	$-n$	$+n$	0
Cationic acid	Cationic	$+c$	$-c$	0
	Neutral	$-n$	$-c$	$-(c+n)$

to be narrower than the corresponding curves in water. Since the solvent tends to bring the two ionizations closer together, the pH range over which the enzyme is active in solvent-water will be narrower than the pH range of activity in water. In case (ii), cation-neutral, the pH-activity curve in dioxan will be spread over a wider range of pH values than in water, since dioxan tends to separate the two ionizations involved. In this case the maximum activity in dioxan may well exceed that in water as the pH range over which the active enzyme exists is broadened and it may be possible to have a higher proportion of the enzyme in its active form in a mixed solvent system than in water.

The curves in Fig. 1 are distinctly different for each of the four pairs that could occur. The types or charges involved at the active centre of an enzyme can, theoretically, be determined by measuring the pH-activity curve in water and in solvent-water mixtures.

EXPERIMENTAL

Materials. Cytidine 2',3'-phosphate and ribonuclease were as described by Herries, Mathias & Rabin (1962). Formamide was purified by the method of Verhoek (1936). Dioxan, AnalaR grade, freed from peroxides, was stored over sodium before use. Buffer salts were purified (recrystallized or redistilled) by standard procedures. Hydrochloric acid was the redistilled constant-boiling acid. Sodium hydroxide was carbonate-free prepared by diluting a saturated solution of AnalaR NaOH in a CO₂-free atmosphere.

Buffers. The concentrations of buffer components were calculated to give a buffer of the desired pH and *I* 0.5 (Tables 2 and 3). The pH of the buffer at the dilution used for activity measurements was checked before use. The pH values in dioxan-water mixtures were measured by the procedures of Bacarella *et al.* (1958). Acetate buffers were used to standardize the pH meter (Harned & Owen, 1950, p. 581).

Methods. The substrate and enzyme were separately dissolved in the buffer. Immediately before addition of the enzyme the balance of water, dioxan or formamide

Table 2. *Neutral-acid buffers*

All buffers had <i>I</i> 0.25.			
Buffer	pK _a at 25°	pH determined in water	pH determined in 50% (v/v) dioxan
Acetate	4.756	4.00	5.80 (5.79)*
		4.52	6.30 (6.29)*
		4.99	6.85 (6.85)*
		5.39	7.20 (7.21)*
Maleate	6.23	5.56	8.00
		5.96	8.60
		6.53	9.05
Pentane-3,3-dicarboxylic acid (<i>K</i> ₂)	7.29	6.82	9.40
		7.40	10.00
Phenol- <i>p</i> -sulphonate (<i>K</i> ₂)	9.053	7.65	9.70
		7.87	9.90
		8.08	10.20
		8.59	10.80
		9.05	11.35

* Values calculated from the data of Harned & Owen (1950, p. 581).

Table 3. *Cationic-acid buffers*

All buffers had <i>I</i> 0.25.			
Buffer	pK _a at 25°	pH determined in water	pH determined in 50% (v/v) dioxan
Ethylenediamine hydrochloride (<i>K</i> ₁)	6.838	6.38	6.45
		6.75	6.65
		6.91	6.90
		7.23	7.20
		7.39	7.50
<i>N</i> -Ethylmorpholine hydrochloride	7.70	7.60	7.60
		8.10	8.10
		8.53	8.50
Ethylenediamine hydrochloride (<i>K</i> ₂)	9.960	8.92	8.90
		9.30	9.50

was added to the appropriate volume of substrate. The final ionic strength in the reaction mixture was 0.25 and the final concentrations of cytidine 2',3'-phosphate and ribonuclease were 0.05M and 0.0125 mg./ml. respectively. The reaction was followed by paper chromatography as described by Findlay, Mathias & Rabin (1962).

RESULTS AND DISCUSSION

Fig. 2 shows the effect of increasing the dioxan concentration up to 50% (v/v) for a reaction carried out in maleate buffers at pH 5.56 and 6.53 (pH measurements in water). As the dioxan concentration is raised the rate increases at the lower pH and decreases at the higher pH. Fig. 3 shows the effect of changing the buffer type at a fixed pH (8.1) over the range 0–50% (v/v) dioxan. In *N*-ethylmorpholine buffer (cationic acid) dioxan has little effect on the rate; in phenol-*p*-sulphonate buffer (neutral acid) the relative initial velocity falls to approximately 5% of its original value in 50% (v/v) dioxan.

There are many cases reported in the literature where an organic solvent either increased or decreased the activity of an enzyme; however, it is apparent that these results are very misleading unless the pH-activity curves in water and in the solvent system have been determined. Inagami & Sturtevant (1960) have carried out a study of the kinetics of trypsin-catalysed hydrolysis of benzoyl-L-arginine ethyl ester in dioxan-water mixtures, in which they attempted to control the pH of their reaction by using an isoelectric buffer which was unaffected by changes in the solvent composition. Yon & Arbel-Sadron (1960), in their study of the effect of pH and dielectric constant on the hydrolysis of β -lactoglobulin by trypsin, used a titrimetric method without buffer.

The pH-activity curves for ribonuclease have been determined in water, 50% (v/v) dioxan and 50% (v/v) formamide in a series of neutral-acid buffers (Fig. 4) and in a series of cationic-acid buffers (Fig. 5). The high ionic strength of the buffer (*I* 0.25) did not interfere with the assay procedure. Technical difficulties prevented the completion of the ascending portion of the pH-activity curve in cationic-acid buffers. The neutral-acid buffers are listed in Table 2 and the cation buffers in Table 3, together with their *pK* values in water and their pH values in water and in 50% (v/v) dioxan-water. No standard is available to measure the pH in formamide-water mixtures. The pH values of the neutral-acid buffers increase by about 2 pH units when the medium is changed from water to 50% dioxan, though the pH values of the cationic-acid buffers do not vary appreciably under the same conditions.

Both formamide and dioxan shift the entire curve to lower pH values in neutral-acid buffers

though they have little effect on the pH-activity curve in cationic-acid buffers. In Fig. 4 (neutral-acid buffers) the mid-point of the ascending portion of the pH (water)-activity curve is 1.7 pH units lower in dioxan-water than the corresponding point on the curve in water. The mid-point of the descending portion of the curve in dioxan-water is 1.2 pH units lower than the corresponding point for the curve in water (Table 4). Formamide

Fig. 2. Effect of dioxan on the initial velocity of hydrolysis of cytidine 2',3'-phosphate (relative to the velocity of hydrolysis in water) in maleate buffer, at pH 5.56 and *I* 0.25 (○), and at pH 6.53 and *I* 0.25 (●). The concn. of substrate was 0.05M, and that of ribonuclease 0.0125 mg./ml.

Fig. 3. Effect of dioxan on the initial velocity of hydrolysis of cytidine 2',3'-phosphate (relative to the velocity of hydrolysis in water) in phenol-*p*-sulphonate buffer, pH 8.1 and *I* 0.25 (○), and in *N*-ethylmorpholine hydrochloride buffer, pH 8.1 and *I* 0.25 (●). The concn. of substrate was 0.05M, and that of ribonuclease 0.0125 mg./ml.

showed the same effect with pK shifts of 1.6 and 1.0 respectively. These shifts are of the same order of magnitude as the buffer shifts shown in Table 3.

In Fig. 5 (cationic-acid buffers) the mid-points of the ascending portions of the pH-activity curve in water and in dioxan-water differ by 0.05 pH unit, whereas the mid-points of the descending portions of the curves differ by 0.2 pH unit. The pH-activity curve in water, 50% (v/v) dioxan-water and 50% (v/v) formamide-water are all essentially the same. The mid-point pH values are given in Table 4.

Comparison of Figs. 4 and 5 with Fig. 1 clearly exclude the situations (ii), neutral-cation, and (iii), neutral-neutral, for the charge types at the active centre of ribonuclease. In case (i), neutral-cation, the activity in neutral-acid buffers at low pH (the ascending portion of the pH-activity curve) should be comparable in water and in dioxan-water or, if the concentration of water is an important factor, the activity in dioxan-water should be less than in water. However, in Fig. 4 (neutral-acid buffers) the activity in dioxan-water in the ascending portion of the pH (water)-activity curve exceeds the activity in water. Further, in

Fig. 5 (cationic-acid buffers) there is no appreciable shift in either the ascending or the descending portions of the pH-activity curve on the addition of the organic solvent. These facts appear to eliminate case (i) (neutral-cationic). The results shown in Figs. 4 and 5 are as would be expected if the charge types at the active centre of ribonuclease are type (iv) (cationic-cationic).

Each rate determination shown in Figs. 4 and 5 was carried out at two concentrations of ribonuclease and in all cases the rate was proportional to the enzyme concentration. Ideally one should measure the variation of k_2 or K_m with pH but, since this is technically impractical, the variation of V (zero-order kinetics) with pH has been determined. There is some doubt, however, about measurements at the higher pH values as, at the substrate concentration used (0.053M), these may not have been high enough to give zero-order kinetics. The substrate concentration was chosen on the basis of the data of Hummel, Flores & Nelson (1958), who found that, at pH 6.7, K_m was 0.99 mM. However, our more recent results (Herries *et al.* 1962) suggest that this measurement

Fig. 4. pH-activity curves for ribonuclease in a series of neutral-acid buffers: in water (Δ), in 50% (v/v) dioxan ($\bullet$), and in 50% (v/v) formamide ($\circ$). The concn. of substrate was 0.05M, and that of ribonuclease 0.0125 mg./ml. Details of the buffers are given in Table 2.

Fig. 5. pH-activity curves for ribonuclease in a series of cationic-acid buffers: in water (Δ), in 50% (v/v) dioxan ($\bullet$), and in 50% (v/v) formamide ($\circ$). The concn. of substrate was 0.05M, and that of ribonuclease 0.0125 mg./ml. Details of the buffers are given in Table 3.

Table 4. pH values of the mid-points of the ascending and descending portions of the pH (water)-activity curves in water, dioxan-water and formamide-water

Buffer type	Limb of pH curve	Water	Dioxan-water	Formamide-water
Neutral	Ascending	6.4	4.7	4.8
	Descending	8.0	6.8	7.0
Cationic	Ascending	6.3	6.4	
	Descending	8.1	8.3	7.7

Table 5. *Variation of rate with substrate concentration*

The conditions used were: (a) *N*-Ethylmorpholine hydrochloride buffer, pH (water) 7.6, *I* 0.25, in 50% (v/v) dioxan; the concn. of ribonuclease was 0.0125 mg./ml. (b) Maleate buffer, pH (water) 6.53, *I* 0.25, in 50% (v/v) dioxan; the concn. of ribonuclease was 0.0125 mg./ml.

Concn. of cytidine 2',3'-phosphate (M)	Conditions	Rate of hydrolysis (mm/min.)
0.053	(a)	0.31
	(b)	0.21
0.079	(a)	0.29
	(b)	0.22
0.106	(a)	0.29
	(b)	0.25

was in error and that K_m varies considerably with pH. At least up to pH 8.0 (where K_m is 0.04M), the substrate concentration (0.053M) exceeded the K_m value, but the kinetics are probably mixed at the higher pH values. Even so, the ascending portions and the first part of the descending portions of both series of pH-activity curves are in the zero-order region.

This also appears to hold for 50% (v/v) dioxan-water mixtures (Table 5). In the cationic-acid buffer [pH (water) 7.6, pH (dioxan-water) 7.6] there was no variation in the overall rate in dioxan-water when the substrate concentration was doubled. The rate in a neutral buffer [pH (water) 6.53, pH (dioxan-water) 9.05] increased only slightly under the same conditions. If dioxan does have an effect on K_m , the most likely one would be the association of the enzyme and the substrate with a resultant lowering of K_m .

It is concluded that the active site of ribonuclease involves two cationic groups. The order of magnitude of the numerical values suggests that both are imidazole groups of histidine.

SUMMARY

1. The activity of ribonuclease can be enhanced or depressed by solvents depending on the pH and the nature of the buffer.

2. A general method of determining the charge types of the groups at the active site of an enzyme, based on the use of organic solvents, is presented and applied to ribonuclease.

3. The pH-activity curve of ribonuclease has been measured in water, 50% (v/v) dioxan and 50% (v/v) formamide in cationic-acid and in neutral-acid buffers.

4. The active site of ribonuclease contains two groups whose acid species are cationic.

REFERENCES

- Bacarella, A. L., Grunwald, E., Marshall, H. P. & Purlee, E. L. (1955). *J. org. Chem.* **20**, 747.
 Bacarella, A. L., Grunwald, E., Marshall, H. P. & Purlee, E. L. (1958). *J. phys. Chem.* **62**, 856.
 Bates, R. G. & Schwarzenbach, G. (1955). *Helv. chim. acta*, **38**, 699.
 Duggan, E. L. & Schmidt, C. L. A. (1942). *Arch. Biochem.* **1**, 453.
 Findlay, D., Mathias, A. P. & Rabin, B. R. (1962). *Biochem. J.* **85**, 134.
 Harned, H. S. & Owen, B. B. (1950). *The Physical Chemistry of Electrolytic Solutions*, 2nd ed. New York: Reinhold Publishing Corp.
 Herries, D. G., Mathias, A. P. & Rabin, B. R. (1962). *Biochem. J.* **85**, 127.
 Hummel, J. P., Flores, M. & Nelson, G. (1958). *J. biol. Chem.* **233**, 717.
 Inagami, T. & Sturtevant, J. M. (1960). *Biochim. biophys. Acta*, **38**, 64.
 Mandel, M. (1958). *Industr. chim. belge*, **23**, 721.
 Mandel, M. & Decroly, P. (1960). *Trans. Faraday Soc.* **56**, 29.
 Shedlovsky, T. & Kay, R. L. (1956). *J. phys. Chem.* **60**, 151.
 Verhoek, F. H. (1936). *J. Amer. chem. Soc.* **58**, 2577.
 Willi, A. V. (1957). *Helv. chim. acta*, **40**, 2019.
 Yon, J. & Arbel-Sadron, G. (1960). *Bull. Soc. chim. Biol., Paris*, **42**, 209.