On carrying out the above oxidation at room temperature for 24 hr., approx. 0.50 g. of starting product yielded (from the initial ether extract) an unknown acid (0.15 g.), which, after washing with ether, crystallized from water in colourless needles, m.p. 142–144° (decomp.). From the ether washings, pale-yellow prisms (9 mg.), m.p. 175°, were obtained, which were identified as 3:5-dimethoxyphthalic acid by comparison of colour reactions (Raistrick *et al.* 1953) and also by methylation of a sample (6 mg.) with diazomethane. Sublimation of the resulting ester in a high vacuum produced colourless needles (4.5 mg.), m.p. 93°, undepressed on admixture with authentic dimethyl 3:5-dimethoxyphthalate.

The major product of the room temperature oxidation was obtained on evaporation of the continuous ether extract, which yielded a colourless crystalline residue (0.37 g.) which crystallized from water in needles, m.p. 180–185° (decomp.). On admixture with 4:6-dimethoxyphthalonic acid, m.p. 174°, the m.p. was depressed.

SUMMARY

1. Alternariol monomethyl ether, $C_{15}H_{12}O_5$, a metabolite of *Alternaria tenuis* Auct., has been shown to be 3:4'-dihydroxy-5-methoxy-6'-methylbenzo- α -pyrone (III).

2. The position of the methoxyl group was established (a) by comparison of the infrared carbonyl frequencies of alternariol and alternariol monomethyl ether with the fully methylated product, and (b) by sequential methylation of alternariol monomethyl ether with [^{14}C]methyl iodide and dimethyl sulphate, followed by oxidative degradation of the resulting labelled trimethyl ether.

3. A tentative biosynthetic interrelationship between alternariol monomethyl ether and the other phenolic constituents of *A. tenuis* is proposed on the basis of their known properties.

Isotopic assaying equipment was generously loaned by Imperial Chemical Industries Ltd., who also provided a grant for the purchase of [^{14}C]methyl iodide. This investigation was carried out during the tenure of a Wellcome Research Fellowship, the award of which is gratefully acknowledged. The author wishes to thank Professor J. H. Birkinshaw for his interest in this work.

REFERENCES

- Allport, D. C. & Bu'Lock, J. D. (1960). *J. chem. Soc.* p. 654.
 Bassett, E. W. & Tanenbaum, S. W. (1960). *Biochim. biophys. Acta*, **40**, 535.
 Bellamy, L. J. (1958). *Infra Red Spectra of Complex Molecules*, p. 185. London: Methuen and Co. Ltd.
 Birch, A. J., Fitton, P., Pride, E., Ryan, A. J., Smith, H. & Whalley, W. B. (1958). *J. chem. Soc.* p. 4576.
 Birkinshaw, J. H. (1953). *Annu. Rev. Biochem.* **22**, 371.
 Bu'Lock, J. D. & Ryan, A. J. (1958). *Proc. chem. Soc., Lond.*, p. 222.
 Chaplen, P. & Thomas, R. (1960). *Biochem. J.* **77**, 91.
 Dean, F. M., Robertson, A., Roberts, J. C. & Raper, K. B. (1953). *Nature, Lond.*, **172**, 344.
 Duncanson, L. A., Grove, J. F. & Zealley, J. (1953). *J. chem. Soc.* p. 1207.
 Gibbs, H. D. (1927). *J. biol. Chem.* **72**, 649.
 Haworth, R. D. & Grimshaw, J. (1955). *Chem. & Ind.* p. 199.
 Mackenzie, J. B. D., Robertson, A. & Whalley, W. B. (1950). *J. chem. Soc.* p. 2965.
 Mosbach, K. (1960). *Acta chem. scand.* **14**, 457.
 Raistrick, H. (1949). *Proc. Roy. Soc. A*, **199**, 141.
 Raistrick, H. & Rudman, P. (1956). *Biochem. J.* **63**, 395.
 Raistrick, H. & Stickings, C. E. (1951). *Biochem. J.* **48**, 53.
 Raistrick, H., Stickings, C. E. & Thomas, R. (1953). *Biochem. J.* **55**, 421.
 Rosett, T., Sankhala, R. H., Stickings, C. E., Taylor, M. E. U. & Thomas, R. (1957). *Biochem. J.* **67**, 390.
 Ruelius, H. W. & Gauhe, A. (1950). *Liebigs Ann.* **569**, 38.
 Sankhala, R. H. (1956). Ph.D. Thesis: University of London.
 Schmidt, O. Th. (1956). *Angew. Chem.* **68**, 103.
 Schmidt, O. Th. & Eckert, R. (1956). *Z. Naturf.* **116**, 757.
 Schmidt, O. Th. & Mayer, W. (1951). *Liebigs Ann.* **571**, 1.
 Shaw, P. D., Beckwith, J. R. & Hager, L. P. (1959). *J. biol. Chem.* **234**, 2560.
 Stickings, C. E. (1959). *Biochem. J.* **72**, 332.
 Tanenbaum, S. W. & Bassett, E. W. (1959). *J. biol. Chem.* **234**, 1861.
 Thomas, R. (1951). Ph.D. Thesis: University of London, p. 126.
 Thomas, R. (1961a). *Biochem. J.* **78**, 748.
 Thomas, R. (1961b). *Biochem. J.* **78**, 807.

Biochem. J. (1961) **80**, 240

Estimation of Succinic Acid in Biological Materials

By KATHLEEN RODGERS

M.R.C. Cell Metabolism Research Unit, Department of Biochemistry, University of Oxford

(Received 9 December 1960)

The method described in this paper determines between 0.02 and 0.20 μ mole of succinate in biological material. It is based on the reduction of 2:6-dichlorophenol-indophenol by succinate in the presence of succinic dehydrogenase (Green, Mii &

Kohout, 1955) and the measurement of the extinction change of the dye at 600 m μ in a spectrophotometer.

It has the advantage over similar methods that the enzyme is more easily prepared than that used

by Massey (1959); that the sensitivity is greater than for the methods of Bril (1954) and Bach & Hibbitt (1959); and that the reaction is carried out in an atmosphere of air instead of argon (Waitzman & Adams, 1958).

Succinic acid is first extracted from the sample with ether (Krebs, 1937). Phosphate buffer is added to the ether extract, the ether distilled off, and the aqueous solution is incubated with semicarbazide to prevent interference by α -oxoglutarate, which reduces the dye under the assay conditions, and by oxalacetate, which inhibits the enzyme.

EXPERIMENTAL

Reagents. Standard succinic acid solution, 0.1 mM: A.R. succinic acid is dissolved in 0.06M-KH₂PO₄-Na₂HPO₄ buffer, pH 7.0. Perchloric acid (A.R.) is 3% (w/v). Diethyl ether: A.R. ether is shaken twice with acidified saturated FeSO₄ solution and redistilled immediately before use. Potassium phosphate-sodium phosphate buffer: 0.06M, pH 7.0, and 0.01M, pH 7.4, are made up with KH₂PO₄ (A.R.) and Na₂HPO₄ (A.R.). Semicarbazide solution, 50 mM: 557.5 mg. of semicarbazide hydrochloride (A.R.) is dissolved in 50 ml. of 0.06M-phosphate buffer, pH 6.0, readjusted to pH 6.0 with N-NaOH, and diluted with buffer to 100 ml. Cyanide reagent: 2.72 g. of KH₂PO₄ (A.R.), 1 g. of crystalline bovine plasma albumin (Armour and Co. Ltd.) and 60 mg. of ethylenediaminetetra-acetate are dissolved in 900 ml. of water, and the pH is adjusted to 7.0 with KOH; 780 mg. of KCN is added and the mixture diluted to 1 l. This reagent can be kept for at least 8 months if it is stored at 3°. 2,6-Dichlorophenol-indophenol (British Drug Houses Ltd.; oxidation-reduction indicator): stock solution (0.15%) is diluted 1:10 with 0.06M-phosphate buffer, pH 7.0, immediately before use.

Preparation of the enzyme. Pigeon breast muscle (50–100 g.) is minced in a Latapie mincer (Etablissement Jouan, Paris) and washed four times with 10 vol. of ice-cold 0.01M-phosphate buffer, pH 7.4. The suspension is filtered through muslin under reduced pressure after each washing. The washed tissue can be stored for several weeks at –15°. For use the preparation is resuspended in 10 vol. of 0.06M-phosphate buffer, pH 7.0, and homogenized for 2 min. in a Waring Blendor. The homogenate remains active for at least 5 days when kept at 5°. It contains approximately 1 mg. of N/ml.

Preparation of samples for analysis. In the earlier experiments samples were deproteinized with tungstic acid before extraction with ether. As phosphotungstate inhibits succinic dehydrogenase (Krebs & Eggleston, 1948), HClO₄ has been adopted for deproteinization in the later experiments. The sample is ground with acid-washed sand and 12 ml. of 3% HClO₄ in a mortar. The suspension is filtered through Whatman no. 30 filter paper. Portions of the filtrate (up to 10 ml.) are extracted with ether in Kutscher-Steucl type extractors (see Umbreit, Burris & Stauffer, 1959). The efficiency of each extractor must be determined by estimating the recovery of known amounts of succinate. An extraction time of 2 hr. has been found necessary with extractors made to hold 30 ml. up to the side arm. Phosphate buffer (2.0 ml.), 0.06M, pH 7.0, is

added to the ether extract and the ether removed by distillation. Since a small amount of HClO₄ is extracted into the ether, this must be neutralized with N-KOH, by using phenol red as indicator. The extract is transferred with a Pasteur pipette to a graduated stoppered Pyrex tube, and made up to 5.0 ml. with phosphate buffer, 0.06M, pH 7.0. It is then kept at 5° for 1 hr. when the KClO₄ separates out; the amount of perchlorate precipitated is so small that the volume measurement is not seriously affected. The supernatant is transferred to another stoppered tube. Interference in the succinate estimation by α -oxoglutarate and oxaloacetate is prevented by a preliminary incubation with semicarbazide. The pH of the sample is adjusted to about 6.0 with 0.1 ml. of N-HCl, and 0.2 ml. of 50 mM-semicarbazide in phosphate buffer, pH 6.0, is added. The mixture is incubated at 37° for 2 hr. or overnight at 18°. The pH is then adjusted to about 7.0 with 0.1 ml. of N-NaOH.

Spectrophotometric estimation of succinic acid. The reaction between the succinic acid and 2,6-dichlorophenol-indophenol is usually completed within 20 min. and it is therefore convenient to carry out the estimations in batches of 20 samples. Each batch includes a blank containing no added succinate and three standards containing 0.02, 0.10 and 0.20 μ mole of succinic acid. Each sample (total volume not exceeding 1.5 ml.) is mixed with 1 ml. of cyanide reaction mixture, 1.0 ml. of 0.015% 2,6-dichlorophenol-indophenol solution, and 0.06M-phosphate buffer, pH 7.0, to give a total volume of 4.0 ml. in a stoppered 10 ml. Pyrex tube. Enzyme (2 ml.) is added to the tubes at 1 min. intervals and the tubes are shaken.

After 20 min. some of the mixture from the first tube is transferred to a 1 cm. cuvette and the extinction measured at 600 m μ in a Unicam SP. 600 spectrophotometer or similar instrument against a mixture containing 1.0 ml. of cyanide reagent, 3.0 ml. of 0.06M-phosphate buffer, pH 7.0, and 2.0 ml. of enzyme. The remaining tubes are treated in the same way at 1 min. intervals. The extinction measurements are repeated at 20 min. intervals until the blank and the samples are reducing the dye at the same rate, when the reaction is considered complete.

The extinction of each sample on completion of the reaction is corrected for the extinction value of the blank, and the change in extinction due to the sample used to calculate the amount of succinate present from a calibration curve prepared as follows. Standard succinic acid solutions (0.1, 1.0 and 2.0 ml.) are mixed with cyanide reagent (1.0 ml.), 2,6-dichlorophenol-indophenol (1.0 ml.) and 0.06M-phosphate buffer, pH 7.0, and the spectrophotometric analysis is carried out as described above. The plot of $E_{1\text{cm.}}$ on completion of the reaction after subtraction from the blank value, against the amount of succinate added to each tube, is linear.

Enzyme preparations that take longer than 20 min. to complete the oxidation of 0.1 μ mole of succinic acid should be discarded. The mixture of cyanide reagent (1.0 ml.), 0.06M-phosphate buffer, pH 7.0, (3.0 ml.) and enzyme (2.0 ml.) has an extinction of approximately 1.4 measured against water; during the spectrophotometric estimation, the Unicam SP. 600 instrument is operated with the slit approximately half open.

The estimation can also be carried out with a Cary recording spectrophotometer, when the standard error for the range 0.02–0.2 μ mole of succinic acid is $\pm 0.004 \mu$ mole

(12 estimations). The total volume per cuvette is 3.0 ml. (0.5 ml. of cyanide reagent, 0.2 ml. of 0.075% 2,6-dichlorophenol-indophenol, sample up to 0.75 ml., 1.0 ml. of enzyme preparation, and 0.06M-phosphate buffer, pH 7.0). A blank cuvette for each sample contains all the above reagents with the exception of succinate.

RESULTS AND DISCUSSION

Recovery of succinate. A typical calibration curve showing the relation between the amount of succinate added in the spectrophotometric assay and the resulting change in extinction ($\Delta E_{1\text{ cm.}}$) at the end of the reaction is shown in Fig. 1. Although the curve is always linear, the slope varies from one enzyme preparation to another, and from day to day for the same enzyme preparation. The mean value of $\Delta E_{1\text{ cm.}}$ after the addition of 0.1 μmole of succinate is $0.22 \pm \text{s.d. } 0.04$ (14 estimations). Because of this variation a calibration curve must be prepared for each batch of samples analysed. The amount of dye reduced by the enzyme in the absence of added succinate (blank reduction) increases with the age of the enzyme, and the change in extinction in the presence of 0.1 μmole of succinate is least when the blank reduction is greatest. With a freshly prepared enzyme that has a low blank reduction, $\Delta E_{1\text{ cm.}}$ for 0.1 μmole of succinate is about 0.29; the theoretical value of $\Delta E_{1\text{ cm.}}$ is approximately 0.33. These results would be expected if the blank reduction is partially suppressed by the addition of succinate.

The samples containing added succinate con-

Fig. 1. A calibration curve for the spectrophotometric estimation of succinic acid. Amounts of succinic acid between 0.02 and 0.2 μmole were added to phosphate buffer, 2,6-dichlorophenol-indophenol and cyanide reagent, as described in the text, and the difference in $E_{1\text{ cm.}}$ between the sample and the blank was measured at 600 $\text{m}\mu$ on completion of the reaction.

tinue to reduce the dye at the same rate as the blank for periods between 20 and 40 min. after the completion of the reaction. After this time the dye is reoxidized, the reoxidation being most rapid in the samples containing the greatest amount of reduced dye. The calibration curve remains linear, although there is a decrease in the slope. The rate of reoxidation of the dye is much slower than that reported by Waitzman & Adams (1958).

Estimation of succinate in the presence of other compounds

Glucose, glucose 6-phosphate, glycerol 1-phosphate and most amino acids are insoluble in ether and therefore do not interfere.

Effect of ether-soluble metabolites. These compounds may cause interference in the spectrophotometric estimation by reacting enzymically or chemically with 2,6-dichlorophenol-indophenol, or by inhibiting succinic dehydrogenase. Washed pigeon breast muscle contains many dehydrogenases, but as most of its coenzymes are removed by the washing procedure, and as coenzymes in the sample to be analysed are not extracted by ether, the presence of the dehydrogenases is not important. To test the specificity of the method, or possible interference, 10 μmoles of various compounds were added to the dye and succinic dehydrogenase preparation, alone, and in the presence of succinate (0.1 μmole).

Acetate, acetoacetate, β -hydroxybutyrate, L-glutamate, lactate, pyruvate, citrate, isocitrate and fumarate caused no reduction of the dye nor interfered with the recovery of added succinate. When fumarate was added to succinate, the reaction time was increased although the amount of dye reduced was not altered. L-Malate decreased the recovery of added succinate when present in amounts greater than 0.2 μmole , but was not itself oxidized.

α -Oxoglutarate (prepared in this Department by Mr T. Gascoyne) reduced the dye in the presence of

Table 1. *Effect of malonate on the spectrophotometric estimation of succinic acid and α -oxoglutaric acid*

The amount of 2,6-dichlorophenol-indophenol reduced by portions (0.05–0.10 ml.) of 0.9 mM- α -oxoglutaric acid and 1 mM-succinic acid was determined spectrophotometrically as described in the text, in the presence or absence of malonate (3.4 mM). Results are expressed in terms of μmole of 'succinic acid'.

	'Succinic acid' (μmole)
α -Oxoglutarate (0.09 μmole)	0.180
α -Oxoglutarate (0.09 μmole) + malonate	0.089
Succinate (0.1 μmole) + malonate	0.000
α -Oxoglutarate (0.045 μmole) + succinate (0.05 μmole)	0.147
α -Oxoglutarate (0.045 μmole) + succinate (0.05 μmole) + malonate	0.046

the enzyme preparation, the amount of dye reduced being twice that of an equimolar amount of succinate. Malonate (3.3 mM) decreased the amount of dye reduced by α -oxoglutarate to half its original value (Table 1). Since this malonate concentration inhibits succinic dehydrogenase completely, but has no effect on α -oxoglutarate oxidase (Stumpf, Zarudnaya & Green, 1947), α -oxoglutarate must have been converted first into succinate and then into fumarate. Reduction of the dye by succinate and α -oxoglutarate was additive. The amounts of these acids in a mixture of the two can be determined by measuring $\Delta E_{1\text{ cm.}}$ in the presence or absence of 3.3 mM-malonate (see Table 1).

Ascorbic acid reduces the dye, but its solubility in ether is so slight that amounts up to 1 mg. do not interfere. When the sample contains a large amount of ascorbic acid, the amount of succinic acid present is determined by measuring the difference between the amounts of dye reduced in the presence or absence of 3.3 mM-malonate.

Oxaloacetate is a very active inhibitor of succinic dehydrogenase, and preliminary incubation with semicarbazide is necessary to prevent interference.

Effect of semicarbazide. Semicarbazide reduces the dye very slowly in the absence of enzyme. The error in the spectrophotometric estimation of 0.1 μ mole of succinic acid is negligible if the concentration of semicarbazide does not exceed 0.6 mM, i.e. under the conditions described above. If the molar ratio of keto acids to succinic acid in the biological material is expected to exceed 16:1, the amount of semicarbazide added to the ether extract should be increased to a molarity at least twice that of the total keto acids present; this ensures the complete absence of free keto acid after incubation for 2 hr. at 37°. In this case it is necessary to add a quantity of semicarbazide, equal to that in the unknown samples, to the standard succinate solutions used in preparing the calibration curve for the estimation. The concentration of semicarbazide may be increased to 8 mM without interference in the action of succinic dehydrogenase; this permits the estimation of 0.1 μ mole of succinic acid in the presence of 24 μ moles of keto acids.

Application of the method to biological materials

In order to test the method on a bacterial sample, succinate (0, 0.5 and 1.0 μ mole) was added to portions (4 ml.) of a washed suspension of *Rhodospseudomonas spheroides* N.C.I.B. no. 8254 grown aerobically in the dark (Lascelles, 1959) and containing 21 mg. dry wt./ml. The samples were diluted to 10 ml. with water, heated for 20 min. at

100° and centrifuged, and the residue was washed with water and recentrifuged. The supernatants were combined, concentrated to about 5 ml., mixed with 1.0 ml. of 18N-sulphuric acid and 1.0 ml. of 1% (w/v) sodium tungstate solution, and extracted with ether. The recoveries of the added succinate were found to be 92 and 101% respectively. The aqueous solutions which contained the extracted succinate did not reduce 2:6-dichlorophenol-indophenol in the absence of succinic dehydrogenase.

The concentration of succinate in a suspension of *R. spheroides* was 10.5 μ m-moles/mg. dry wt. This result was not affected by an eightfold variation in the amount of biological material.

The amount of succinate (μ moles/g. wet wt.) found in tissues from a freshly killed normally fed rat were: liver, 0.94; lung, 0.15; heart, 0.48; kidney cortex, 1.10; brain, 0.21; testis, 0.57. The liver of a second rat contained 0.99 μ mole of succinate/g. wet wt. Other published values for succinate in rat liver are 1.2 μ moles/g. wet wt. (see Busch & Potter, 1952), and 0.2 μ mole/g. wet wt. (Frohman & Orten, 1953).

The aqueous solutions containing succinate extracted from liver did not reduce 2:6-dichlorophenol-indophenol, and 3.3 mM-malonate completely inhibited dye reduction in the assay. This confirmed that the reduction of the dye was due solely to succinate.

SUMMARY

1. A spectrophotometric method for the estimation of from 0.02 to 0.20 μ mole of succinate is described. It is based on the reduction of 2:6-dichlorophenol-indophenol by succinate in the presence of succinic dehydrogenase.

2. Both succinate and α -oxoglutarate reduce the dye enzymically under the assay conditions whereas pyruvate does not. Interference by α -oxoglutarate may be prevented by a preliminary incubation with semicarbazide.

3. The advantages of the method are its sensitivity and the ease of preparation of the succinic dehydrogenase.

4. Application of the method to biological material showed that rat tissues contained between 0.15 and 1.10 μ moles of succinic acid/g. wet wt. Washed cells of *Rhodospseudomonas spheroides* contained 10.5 μ m-moles/mg. dry wt.

The author wishes to thank Professor Sir Hans Krebs, F.R.S., and Dr D. E. Hughes for their advice and interest in the work, Dr J. Lascelles for providing the suspensions of *R. spheroides*, and Mr T. Gascoyne for assistance with the estimations on rat tissues. The work was aided by grants from the Rockefeller Foundation, and from the U.S. Public Health Department.

REFERENCES

- Bach, S. J. & Hibbitt, K. G. (1959). *Biochem. J.* **72**, 87.
 Bril, C. (1954). *Biochim. biophys. Acta*, **15**, 258.
 Busch, H. & Potter, V. R. (1952). *J. biol. chem.* **198**, 71.
 Frohman, C. E. & Orten, J. M. (1953). *J. biol. Chem.* **205**, 717.
 Green, D. E., Mii, S. & Kohout, P. M. (1955). *J. biol. Chem.* **217**, 551.
 Krebs, H. A. (1937). *Biochem. J.* **31**, 2095.
 Krebs, H. A. & Eggleston, L. V. (1948). *Biochem. J.* **43**, 17.
 Lascelles, J. (1959). *Biochem. J.* **72**, 508.
 Massey, V. (1959). *Biochim. biophys. Acta*, **34**, 255.
 Stumpf, P. K., Zarudnaya, K. & Green, D. E. (1947). *J. biol. Chem.* **167**, 817.
 Umbreit, W. W., Burris, R. H. & Stauffer, J. F. (1959). *Manometric Techniques*, p. 203. Minneapolis: Burgess Publishing Co.
 Waitzman, M. B. & Adams, E. Q. (1958). *Nature, Lond.*, **182**, 129.

ADDENDUM

The Intracellular Concentration of Succinic Acid in Photosynthetic Bacteria

BY JUNE LASCELLES* AND KATHLEEN RODGERS

Microbiology Unit and Medical Research Council Unit for Cell Metabolism, Department of Biochemistry, University of Oxford

(Received 9 December 1960)

The method of Rodgers (1961) for the estimation of succinic acid has been applied to photosynthetic bacteria of the family Athiorhodaceae. These organisms can grow either aerobically in the dark or anaerobically in the light, but it is only under the latter conditions that they form the photosynthetic pigments bacteriochlorophyll and carotenoids (van Niel, 1944; Cohen-Bazire, Sistrom & Stanier, 1957; Lascelles, 1959). Succinyl-coenzyme A is a key intermediate in the formation of bacteriochlorophyll and it was thought that the ability to form this pigment might be correlated with the concentration of intracellular succinate, but the analyses provided little evidence for this.

MATERIALS AND METHODS

The strains of Athiorhodaceae and their maintenance have been described by Kornberg & Lascelles (1960).

Growth and harvesting of cultures. The media and conditions for aerobic and photosynthetic growth were as described previously (Lascelles, 1959; Kornberg & Lascelles, 1960).

Organisms were harvested at the end of the logarithmic phase of growth unless stated otherwise. To minimize possible changes occurring in the intracellular concentrations of succinate during harvesting, the cultures were chilled rapidly by plunging the vessels into a mixture of alcohol and solid carbon dioxide and then centrifuged at 0°. The organisms were washed with ice-cold 0.02M-potassium phosphate buffer, pH 7.0, equivalent to one-half of the original culture volume. Analysis of concentrated washings showed little or no succinate to be extracted by this procedure. The washed organisms were finally suspended in water at 0° to a cell concentration of 25–50 mg. dry wt./ml. Portions of the suspension containing 100–250 mg. dry wt.

of cells were extracted, and the succinic acid content was estimated as described above. Since paper chromatography by the method of Elliott (1954) showed that only traces of α -oxoglutarate were present, incubation with semicarbazide was omitted.

RESULTS

Succinate concentration in organisms grown in the dark or light

The intracellular concentration of succinate was slightly higher in light-brown cultures of *Rhodopseudomonas spheroides* on malate plus glutamate than in those grown aerobically in the dark on the same medium (Table 1). This difference is statistically significant (*t* test: $P = 0.02-0.01$). The concentration of succinate was similar in organisms grown on butyrate plus bicarbonate as carbon source (Table 1). No significant variation was found in cultures harvested at various stages of the growth cycle under aerobic-dark or anaerobic-light conditions.

Succinate concentrations were determined in butyrate-grown cells of *Rhodopseudomonas palustris* and *Rhodopseudomonas capsulatus* and were of the same order as that found in *R. spheroides*.

As a check that the material assayed was succinate and not α -oxoglutarate [shown by Rodgers (1961) to interfere with the method], extracts prepared from organisms grown aerobically and photosynthetically on malate plus glutamate were assayed in the presence or absence of malonate. This compound inhibited activity of the extracts by 90% or more showing that the active material was largely succinate (Rodgers, 1961). This was confirmed by paper chromatography of the extracts by the method of Elliott (1954).

* External Scientific Staff, Medical Research Council, at the time of this work.

Table 1. Succinate concentrations in cultures of *Rhodospseudomonas*

Organisms were harvested at the end of the logarithmic phase of growth and extracted as described in the Materials and Methods section.

Organism	Growth medium	Incubation conditions	Succinate (μ m-moles/ mg. dry wt.)	No. of experiments
<i>R. spheroides</i>	Malate-glutamate	Aerobic, dark, 14-18 hr.	6.4 (Range 5.2-8.0)	5
		Anaerobic, light, 14-18 hr.	8.2 (Range 7.0-10.5)	8
	Butyrate-bicarbonate	Aerobic, dark, 24 hr.	5.8	1
		Anaerobic, light, 24 hr.	6.5	1
<i>R. palustris</i>	Butyrate-bicarbonate	Aerobic, dark, 46 hr.	3.2	1
		Anaerobic, light, 38 hr.	4.3	1
<i>R. capsulatus</i>	Butyrate-bicarbonate	Aerobic, dark, 48 hr.	3.4	1
		Anaerobic, light, 24 hr.	3.4	1

DISCUSSION

It is not known if the extraction procedure involving heating at 100° in water liberates all the intracellular succinate. The same treatment has been shown to be effective in releasing the free amino acids from various bacteria (Gale, 1947; Hancock, 1958). Succinyl-coenzyme A, which hydrolyses spontaneously at pH 7 at room temperature (Simon & Shemin, 1953), would, if present, presumably contribute to the succinate estimated by the procedure used.

The amounts of succinate in the Athiorhodaceae are similar to those isolated from other bacteria by chromatography of ethereal extracts on Celite columns. Suspensions of *Micrococcus lysodeikticus* oxidizing acetate contained 5 μ m-moles/mg. dry wt. (Saz & Krampitz, 1955), whereas in *Escherichia coli* under the same conditions 1.7 μ m-moles/mg. wet wt. were found (Swim & Krampitz, 1954). Using similar techniques Ladd & Nossal (1954) found the intracellular concentration in baker's yeast to vary between 1.4 and 34 μ m-moles/mg. wet wt. in different samples.

The results do not show any striking correlation of the intracellular concentration of succinic acid with the conditions necessary for the formation of photosynthetic pigments. The initial reaction in bacteriochlorophyll synthesis is a condensation of succinyl-coenzyme A and glycine to give δ -amino-laevulinic acid and the enzyme system responsible for this reaction is considerably more active in pigmented than in unpigmented organisms (Lascelles, 1959, 1960). It is therefore possible that succinyl-coenzyme A is diverted rapidly towards bacterio-

chlorophyll synthesis under the appropriate conditions, and this might account for the inability to detect considerable increases in the intracellular concentration of succinate.

SUMMARY

1. The intracellular concentration of succinate has been estimated in *Rhodospseudomonas spheroides*, *Rhodospseudomonas palustris* and *Rhodospseudomonas capsulatus* grown anaerobically in the light or aerobically in the dark.

2. On a malate-glutamate medium the concentration of succinate in *R. spheroides* grown anaerobically in the light is slightly though significantly higher than in aerobically grown organisms (8.2 and 6.4 μ m-moles/mg. dry wt. respectively).

REFERENCES

- Cohen-Bazire, G., Sistrom, W. R. & Stanier, R. Y. (1957). *J. cell. comp. Physiol.* **49**, 25.
- Elliott, D. C. (1954). *J. exp. Bot.* **5**, 353.
- Gale, E. F. (1947). *J. gen. Microbiol.* **1**, 53.
- Hancock, R. (1958). *Biochim. biophys. Acta*, **28**, 402.
- Kornberg, H. L. & Lascelles, J. (1960). *J. gen. Microbiol.* **23**, 511.
- Ladd, J. N. & Nossal, P. M. (1954). *Aust. J. exp. Biol. med. Sci.* **32**, 523.
- Lascelles, J. (1959). *Biochem. J.* **72**, 508.
- Lascelles, J. (1960). *J. gen. Microbiol.* **23**, 487.
- Rodgers, K. (1961). *Biochem. J.* **80**, 240.
- Saz, H. J. & Krampitz, L. O. (1955). *J. Bact.* **69**, 288.
- Simon, E. J. & Shemin, D. (1953). *J. Amer. chem. Soc.* **75**, 2520.
- Swim, H. E. & Krampitz, L. O. (1954). *J. Bact.* **67**, 419.
- van Niel, C. B. (1944). *Bact. Rev.* **8**, 1.