How To Do It

Be a patient

B T MARSH

I shall be forever grateful to Stephen Potter for his four classic books on how to cope with the vagaries of life, namely, *Lifemanship*, *Gamesmanship*, *One-Upmanship*, and *Supermanship*.¹⁻⁴ *Lifemanship* was written in 1947, and in my schooldays reading the first three books was de rigueur in the sixth form. In fact you could almost do an "A" level in the subject, and I still find his advice both apposite and welcome some 20 to 30 years later. Many other larger and more erudite treatises on how people react to life have been published over the succeeding years, yet I doubt if any have bettered Potter's acute, albeit exaggerated, observations on how to make the most of one's opportunities and better the opposition, whoever or whatever that may be. I have no intention of making this article a poor man's Potter or an exposition of his various gambits, but I am aware that throughout my experiences of being a patient I have both consciously and subconsciously tried to beat the system with a form of "patientship."

As an opening gambit, "Don't" or "Be lucky" is the best advice I can give to anyone contemplating being a patient, but this fails to take into consideration that most of us have very little choice in the matter. If you are worried about your health it is sensible to consult an expert. Once this consultation begins inevitably you become a patient and are drawn, either willingly or unwillingly, into a whole new world of instability and anxiety, indeed a role reversal with a vengeance. Although I have suggested that you should not be a patient if you can sensibly avoid it, by the same token you should avoid acting as your own physician. This is not because of any doubts about diagnostic competence, as you may get it right, but you may do so for the wrong reasons, in the wrong order, with the wrong conclusion, or with the wrong sense of urgency.

Nobody knows how you feel

I think it important to state that being a patient is an intensely personal experience that we all approach for differing reasons and with differing expectations. Because of this, one thing I have learnt is that no one can truly say, "I know how you feel." Indeed as a patient I became quite angry when well meaning sympathisers made this statement. It is one that should always be associated with some indication that the speaker does not know, but is at least making some attempt to appreciate the situation. This usually prevents the upsurge of anger and frustration that may otherwise be provoked and allows the patient to explain in great detail what he does feel, which certainly helps him, or her, if nobody else.

In June 1984, after a pleasant dinner, I developed severe retrosternal chest pain that was relieved by metoclopramide and antacids. This pain, although less severe, recurred over the next two weeks and was related in my mind to food and to bending. Angina was clearly a differential diagnosis but I was quite unable to provoke the pain by exercise, and it came well below plain indigestion, reflux

oesophagitis, and hiatus hernia in my estimation of likelihood. As indigestion was an unusual symptom with me I was eventually persuaded by my wife to make an appointment to see a physician, and I was on the path to becoming a patient. The day before my appointment I had a massive myocardial infarction.

My indigestion had turned out to be oesophageal angina, which has a reputation of being both a mimic and unpredictable. After the infarct I successively underwent cardiac arrest and resuscitation, admission to an intensive care unit, intracardiac pacing, septicaemia, infective endocarditis, insertion of an aortic balloon pump, and, finally, 15 days after the initial event, cardiac transplantation. Discharge home came 12 days later with a convalescence of six months before I returned to work part time, now almost full time. Regular outpatient visits are still necessary (and will be, presumably, for ever) with, at six monthly intervals, an exercise electrocardiogram, a 24 hour electrocardiographic tape, thallium scan, echocardiography, heart muscle biopsy, and, in addition, a coronary angiogram every 12 months. I have described my experiences in greater detail elsewhere,⁵ and much of this advice will be specific to these experiences, but I hope that there will be a generic basis that will help those yet to be patients, confirm some of the impressions of those who have been patients, and, for the rest, allow some insight into the feelings of a reluctant but lucky patient.

What is going on

The first thing to realise is that although being in the profession does have many benefits, you are still a pawn to be moved around the chessboard under the aegis of—you hope—a benevolent dictator. Life as a patient is totally different from what you expected, what it was like in your day as a medical student, and what you would like it to be. Day merges into night and nights seem to go on for ever. Time has no relevance and it is quite astonishing how you can lose track of the date and overall time scale. If you are really ill, meals lose their importance at the same time as their taste. This is no reflection on the catering staff, who labour valiantly to excite your appetite. Patients' appetites are still, I think, one of the best indicators of their state of health. Night time sedatives or hypnotics become highly important. Long periods in bed mean frequent naps and consequent difficulty in getting off to sleep at night, with a very light and disturbed sleep pattern, especially in a busy ward or intensive care unit. Judicious use of sedatives helps but a vicious circle is often set up if the sedative is not a short acting one. The patient is then sedated during the morning, often confused, and then becomes even more awake in the evenings.

Even if you have your own room, there never seems to be enough easily accessible space to put things on. Grapes, other fruit, and flowers are, I think, only good for visitors, and "get well" cards are nice to receive but difficult to keep on show. Do not consider your stay in hospital as a chance to catch up on that heavy reading you have been putting off. Solzhenitsyn is definitely out and it will take you all your time to stagger through the most basic newspapers and magazines. A radio cassette player such as a Walkman and, especially, a television with a remote control are absolute musts.

The television programmes both entertain—using the word in its broadest sense—and take your mind off your current discomfort. The remote control allows you to jump from station to station as concentration rarely lasts longer than a few minutes, particularly with some of the more erudite programmes.

Information on what is happening to you is often at a premium even though you are in the trade. Do not be frightened to ask what is the real diagnosis and what is the plan of treatment. Invariably you will get a full explanation and be brought into the discussion. It is more difficult, however, to get from your advisers the results of your electrocardiogram, chest radiograph, and other investigations. If you feel aware of what is going on within your body do not hesitate to suggest a diagnosis, particularly if you think that something is going wrong or if you are conscious of a change. You may not be believed and you may be wrong, but at least you should stimulate further examination and thought. Do object if two way conversations occur across the bed between relatives and the attending doctor and you are not included. This is usually inadvertent and arises because it is thought that you are too tired, or sedated, or unwell. Always have your say, as you may want to say "I told you so" at a later date.

Tricks of the trade

There are certain things you should not believe, such as: "It will be just a small prick and it won't hurt"; "We are only going to put in one stitch and there is no need for a local anaesthetic because the injection would be just as painful"; "Everything is fine and don't worry"; "The doctor will be along in a minute"; "You aren't written up for it, so you can't have it"; "We have no idea when you are going to be discharged"; "No, you are not going to be moved from this bed or room"; "You must take this drug, it is essential" (for something that you well know is not essential, such as a vitamin C tablet).

There are other things that you should always ask for if you want them, such as sedatives or hypnotics at night and analgesics if you are in pain or discomfort. Long periods in bed inevitably lead to a tender sacrum and buttocks and an aching back and neck. Despite what some nurses declare, regular massage of the buttocks and sacrum is a tremendous relief, nay, pleasure, even if there is no sign of skin deterioration. A rubber ring and a lambskin undersheet also help to relieve the problem. If you are a male always have at least one spare urine bottle available, and if you have a bedpan ensure that the emergency call button is within easy reach, as sitting on one of these for half an hour is not to be recommended.

A sense of isolation may occur if you are in a room on your own, and the frequent popping in and out of nurses and domestic staff is a great comfort. Needless to say the visit of your spouse plus or minus other sundry relatives and friends is often the highlight of the day. This event becomes so important that at times unreasonable pettiness on the part of the patient can erupt. Comments such as, "Why are you late?" when your spouse has moved heaven and earth to get the family fed and clothed, friends and relatives informed, and other essential jobs performed, are not conducive to harmony. In the same vein, the simple "Thank you" is all that most of the nursing, paramedical, medical, and domestic staff require for the countless small favours that they perform for you during a stay in hospital—it is so easy to say, but so often forgotten.

There are certain essential items that you should take into hospital with you, and tricks that you should know. Always have an adaptor plug set with you. You can bet that the one on the ward doesn't fit, won't work, is lost, or is being used by someone else. Have a mirror with you, preferably a polished metal one. The mirror in the room is rarely in the best spot for shaving, hair drying, or whatever else you are trying to watch yourself do. Have soap, toothpaste, flannel, talcum powder, and other toiletries of your own choice with you, as the make you prefer is never on the trolley. Heels often feel sore after a long period in bed, and it becomes impossible to get your legs in a comfortable position. Ask for a frame to keep the bedclothes off your legs and suggest that someone fills up two rubber gloves with water and ties a knot in the wrist ends. These

"donkeys" provide wonderfully gentle supports for the heels, but be prepared for rubber fatigue and a wet bed from time to time. If you have difficulty in pulling yourself up, a knotted rope to the foot end of the bed makes it much easier. Depending upon your particular problem there are many other tricks of the trade that are available—if you ask the right nurse.

Getting out

As you come to discharge time never underestimate the weakness from weight loss and loss of muscle mass that will have occurred if you have been seriously ill and bedbound for some time. You will need help from the physiotherapy department, and this is always readily and efficiently given provided it is asked for or is routine for your type of case. It is remarkable how quickly you can lose strength, particularly for such everyday tasks as getting out of a chair unaided or walking upstairs. This is particularly important, as most bedrooms and bathrooms are on the first floor of a house. This is the time also to ensure that you know exactly what medicine to take, what exercise to perform, and what to avoid when you are discharged. Don't forget to ask what the drill is if an emergency occurs—do you go back to your own general practitioner, ring the consultant, or get yourself admitted to the hospital? When you are being driven home, usually at a sedate 20 to 30 mph, be prepared for this to seem as if you are taking part in a Formula I Grand Prix race. I suggest sitting in the rear seat with your eyes closed.

Finally, a word about outpatients. Find out the critical timings and days associated with your outpatient clinics. Which ones are over booked or under booked, always have plenty of nurses present or appear to be as unpopular with the nurses as they are with doctors? Is it better to go for radiograph, electrocardiogram, or laboratory tests first or after your consultation? Find out the people who determine the pace of the clinic—what is known in the Royal Navy as "making your number" and in hospital as "getting on the right side of Sister." Ensure that you know the treatment and investigation protocols and then keep a check on these and the results. Familiarity breeds contempt, and minor slips can quickly build up into major problems. When in doubt ask. Follow the instructions unless they appear to be clearly wrong. If they do appear wrong you will often be proved correct. Make a note of your symptoms, reactions, and problems, especially if they are related to something like drugs or food. Pass them on to the physician or nursing staff on your next visit. Much of the lore about a particular condition or its progress is often anecdotal from patients (although many findings get confirmed in the waiting room), and you never know if your experience will be helpful to someone else. If you need regular admissions it is worth while suggesting the most convenient dates for you—surprisingly enough it often works.

Being a patient is a major step that should not be taken lightly. It is an enormous psychological experience that will in most cases entirely change your outlook on life and certainly make you more understanding and sympathetic to other patients. Most patients undergo at least one period of self pity, sometimes many. There is no answer to the querulously put, "Why me?" Each must work out his or her own rationale for what has happened and what will happen in the future. It is also important to realise that your illness might have been equally upsetting to your spouse or relatives and that they may be asking the same questions. Luckily, most people are optimists and although such a question is not unreasonable, it is not too difficult to persuade oneself that all is going to be well. This attitude is essential if there is to be any return to normality. On the other hand, there is no doubt that being a patient can become addictive and may damage your health.

References

- 1 Potter S. *Gamesmanship*. London: Hart-Davis, 1947.
- 2 Potter S. *Lifemanship*. London: Hart-Davis, 1950.
- 3 Potter S. *One-Upmanship*. London: Hart-Davis, 1952.
- 4 Potter S. *Supermanship*. London: Hart-Davis, 1958.
- 5 Marsh BT. A second chance. *Br Med J* 1986;292:675-6.