XCVI. STRUCTURE AND ENZYME REACTIONS. PART IV.

THE SYSTEM GLYCOGEN-AMYLASE-LIPOIDS.

BY RICHARD TRUSZKOWSKI.

From the Biochemical Laboratory, Faculty of Medicine, Warsaw University.

(Received April 23rd, 1928.)

IN the preceding parts of this series [Przylecki and co-workers, 1927, 1928] it was shown that the introduction of an adsorbent into a system in which either the enzyme, the substrate or the products of reaction undergo adsorption may profoundly influence the velocity of the enzymic reaction. The results obtained were applied to certain biological problems as an explanation of the mechanism whereby the structure of a cell may influence reactions.

In Part I of this series it was established that for the system urea-urease-charcoal, in which only the enzyme underwent adsorption, no retardation of the enzymic decomposition of urea was brought about by the presence of this adsorbent. Part II dealing with the system polysaccharide-amylase-charcoal, exhibited a different behaviour. Here both the enzyme and the substrate undergo adsorption, that of the latter being reversible in the presence of an appropriate concentration of narcotics. The adsorbed enzyme retained its amylolytic activity, so that it was able to decompose glycogen in solution with its usual efficiency. Where, however, the enzyme and the substrate were simultaneously adsorbed, considerable retardation of hydrolysis took place. Finally, in Part III, it was shown that the introduction of coagulated egg-albumin instead of charcoal yields parallel results.

In this paper, the same problem will be considered, with the use of a different cell-constituent, namely the ether-soluble substances (lipoids), as the adsorbent.

The theoretical aspects of this question have been fully dealt with in the previous parts of this series, to which those interested should refer.

EXPERIMENTAL.

The experimental programme consisted in elucidating the following points: adsorption of enzyme and of substrate on lipoids, fats, etc., their elution, and the effect of the presence of the adsorbent upon the kinetics of the reaction.

The adsorbent consisted of an intimate mixture in equal proportions of cholesterol, lecithin, palmitic acid and calcium glycerophosphate, this mixture being referred to under the general term of lipoids. The amylase used was purchased from Merck in powder form and the substrate taken was glycogen.

1. *Adsorption and elution of amylase.*

A series of flasks containing 23 cc. of buffer solution, p_H 6.5, and 2 cc. of 0.5 % amylase solution containing 1 % of sodium chloride together with, in certain cases, 2 g. of the above mixture, and the same with 0.5 *M* propyl or 0.07 *M* butyl alcohol solution, were placed for 3 hours in a mechanical shaker and the contents of the flasks then filtered. To 20 cc. of each filtrate 10 cc. of 1 % glycogen was added, and the flask kept for 4 hours at 37°, after which the reducing sugar content was determined by Benedict's method. That of Bertrand was not here practicable, as the filtrate when rendered alkaline became unfilterable. The results, given in Table I, indicate that amylase is under these conditions adsorbed to an extent of 70 %, and that the addition of propyl alcohol up to 0.5 *M* or of butyl alcohol up to 0.07 *M* solution leads to an almost quantitative elution of the enzyme, *i.e.* only 5 % remains in adsorption.

Table I. *Adsorption and elution of amylase.*

Mg. glucose in 30 cc. solution, using 1 % glycogen and 2 g. of lipoids.

No. of exps.	Without adsorbent	With adsorbent					
		No alcohol	% adsorp- tion	0.07 <i>M</i> butyl alcohol	% adsorp- tion	0.5 <i>M</i> propyl alcohol	% adsorp- tion
2	53.8	16.2	67	50.7	6	52.0	4

2. *Adsorption of glycogen.*

A series of flasks containing 25 cc. of 1 or 0.5 % glycogen were shaken for 3 hours with and without the addition of lipoids and alcohols, filtered, and a measured volume of the filtrate was diluted to 30 cc. and hydrolysed by heating for 4 hours at 100° with 1.5 cc. of concentrated hydrochloric acid, after which the acid was neutralised by the addition of excess sodium carbonate and reducing sugars determined. The results (Table II) indicate that glycogen is not adsorbed upon these lipoids.

Table II. *Adsorption of glycogen.*

Mg. glucose in 5 cc. of filtrate, using 0.5 and 1 % glycogen and 2 g. of lipoids.

Glycogen %	No. of exps.	Without adsorbent	With adsorbent			
			No alcohol	0.07 <i>M</i> butyl alcohol	0.5 <i>M</i> propyl alcohol	% adsorption
1	2	19.1	20.62	19.27	20.55	0
1	2	20.62	21.22	—	—	0
0.5	2	9.46	9.79	9.14	10.13	0
0.5	2	10.25	10.34	—	—	0

3. *Elution of adsorbed ferment by glycogen or water.*

A number of experiments were necessary for the elucidation of this matter. In determining the elution of adsorbed amylase by glycogen, advantage was taken of the fact that this enzyme has practically no action at 0°. A series of flasks containing buffer solution and amylase, with or without lipoids, was shaken as before for 3 hours, after which some were filtered immediately, whilst others were cooled to 0°. An equal volume of 2 % glycogen solution at 0° was then added, and the flasks were shaken at 0° for 3 hours, after which the contents were filtered in an ice-chest, and placed in a thermostat at 37° for 4 hours, together with those flasks whose contents had been filtered immediately and to which an equal volume of 2 % glycogen solution had been added subsequently. The results, given in Table III, show that a certain amount of elution has taken place.

Table III. *Elution of amylase due to glycogen and water.*

Mg. glucose in 10 cc. of filtrate, using 2 % glycogen solution.

No. of exps.	Without adsorbent	With adsorbent					
		Glycogen added before filtering	% hydrolysis	Water added before filtering	% hydrolysis	Glycogen and water added after filtering	% hydrolysis
2	14.45	7.2	50	5.1	35	4.8	33
2	21.7	10.2	47	—	—	7.9	36
2	17.6	8.7	49	—	—	—	—

In order to verify whether this is due to the specific elutive action of glycogen, or whether it is merely due to dilution, an additional experiment was performed.

25 cc. of a 0.04 % solution of amylase, p_H 6.5, was shaken with 3 g. of lipid mixture for 3 hours, cooled to 0°, 25 cc. of 2 % glycogen solution at 0° were added, and the whole was shaken for a further 3 hours at 0°, after which the mixture was filtered at 0°, half the volume of the filtrate of water added, and the whole kept at 37° for 4 hours. A second pair of flasks was put through the same procedure, except that the order in which the glycogen and water were added was reversed. A third pair of solutions was filtered immediately after shaking and a double volume of 1 % glycogen added to the filtrate, and finally a fourth pair underwent the same manipulations as the first pair, except that lipoids were absent. The glucose content of all the solutions was then determined. The results of this experiment, given in Table III, show that where glycogen solution is added immediately, the amount of amylase in the filtrate is 50 % of that in the absence of adsorbent, where water is added first, 35 % and, where the amylase solution is filtered immediately, 33 %. It would hence appear that glycogen has a considerable elutive influence not possessed by water.

The dilution by water is here two-fold, and in order further to establish this point four-fold dilution was next employed. A pair of flasks containing

25 cc. of a 0.04 % amylase solution was as before shaken 3 hours with 3 g. of lipid mixture, filtered, 30 cc. of buffer solution were added to 10 cc. of the filtrate, and 40 cc. of 2 % glycogen solution. A second pair of flasks containing 100 cc. of buffer solution, with the same quantity of amylase as the first pair, was shaken and filtered, an equal volume of 2 % glycogen being added to the filtrate; a third pair containing the same quantity of solution but without lipid mixture was put through the same manipulations. After keeping at 37° for 4 hours, the reducing sugar content was determined. The results given in Table IV indicate that even a four-fold dilution of amylase solution provokes no perceptible elution.

Table IV. *Effect of dilution. Mg. glucose found in 20 cc. filtrate using 2 % glycogen.*

No. of exps.	Without adsorbent	With adsorbent	
		Diluted before shaking	Diluted after shaking
2	7.35	4.2	4.0

4. Kinetics.

A number of flasks containing 25 cc. of 0.04 % amylase solution were shaken 3 hours with 3 g. of lipid mixture, and 25 cc. of glycogen solution, either 1 or 2 %, added. The same was repeated with a second series of flasks, introducing butyl or propyl alcohol up to 0.07 and 0.5 *M* respectively, and to a third series the same volumes of solutions but no lipoids were added. All flasks were then left for 4 hours in a thermostat at 37°, after which they were filtered, and reducing sugars determined. The results (Table V) show that for 2 % glycogen the hydrolysis is only 40.4 to 49.8 % of that in the absence of an adsorbent, and for 1 % glycogen 61.2 to 75.6 %. The addition of butyl alcohol to the 1 and 2 % glycogen reaction mixtures raises hydrolysis to from 91.8 to 98.0 % of that in the absence of an adsorbent, whilst the action of propyl alcohol is less marked.

Table V. *Influence of adsorbent upon the kinetics of the hydrolysis of glycogen by amylase. Mg. glucose in 10 cc. filtrate.*

Glycogen added %	No. of exps.	Without adsorbent	With adsorbent					
			No alcohol	% hydrolysis	0.07 <i>M</i> butyl alcohol	% hydrolysis	0.5 <i>M</i> propyl alcohol	% hydrolysis
2	2	16.7	8.4	49.8	15.5	92.7	13.2	79.8
2	2	17.8	7.2	40.4	17.6	98.0	7.8	43.8
1	2	11.1	8.4	75.6	—	—	10.1	91.0
1	2	9.8	6.0	61.2	9.0	91.8	6.6	67.3

In view of the possibility that retardation of reaction is due to mechanical causes, *i.e.*, that the adsorbed enzyme, whilst remaining active, reacts more slowly as a result of a decline in the number of collisions between the molecules and those of the substrate, a similar experiment was carried out, with the

difference that the reaction flasks were shaken whilst in the thermostat. The results (see Table VI) indicate that under these conditions shaking has no influence upon the velocity of reaction.

Table VI. *Influence of shaking upon velocity of enzymic hydrolysis.*

Mg. glucose found in 10 cc. filtrate, using 1 % glycogen solution.

No. of exps.	No lipoids	Lipoids with shaking	Lipoids without shaking
2	10.3	6.6	6.6

DISCUSSION.

The above experiments show that amylase undergoes a considerable degree of adsorption upon the mixture of ether-soluble substances used. The adsorbed amylase can, however, be eluted almost quantitatively by the addition of an appropriate quantity of propyl or butyl alcohol; in this respect lipoids differ from protein, as also in the partial elution produced by the action of glycogen. Again, lipoids do not adsorb glycogen, in contrast to the adsorbents studied previously.

The system glycogen-amylase-lipoids would in the above respects appear to be similar to the systems urea-urease-charcoal and glycogen-amylase-protein (alcohol) described in Parts I and III, in so far as only the enzyme is in adsorption, the substrate being practically unadsorbed. In these cases, little retardation of reaction was observed, pointing to the unimpaired activity of the enzyme in spite of adsorption. Here, however, reaction is retarded to an extent commensurate with that of the quantity of enzyme in adsorption, pointing to the very strong probability of amylase losing its activity on adsorption upon lipoids.

This retardation cannot be due to purely mechanical causes, since shaking in no way affects the velocity of hydrolysis in the presence of an adsorbent.

The elution of amylase from adsorption on lipoids by glycogen is of interest, inasmuch as most elutives act as a result of their preferential adsorption. Here, however, glycogen is not adsorbed, so that the elution must be due to some different cause. This is possibly the decline in the concentration of free amylase in solution as a result of its combination with the substrate.

For the sake of clarity, the different properties of charcoal, protein and lipoids as adsorbents are tabulated in Table VII.

Table VII.

Adsorbent	Substrate	Adsorption of		Elution of			Activity of adsorbed enzyme
		Enzyme	Substrate	Enzyme by		Substrate	
				Alcohols	Substrate		
Charcoal	Urea	+	-	-	-	-	+
Charcoal	Glycogen	+	+	-	-	+	+
Protein		+	+	-	-	+	+
Lipoids	"	+	-	+	+	-	-

The behaviour of these two cell constituents, protein and lipids, is in almost every case contrasting, and points to the great complexity of the mechanism of the processes taking place within the cell, which possesses the means of modifying enzymic reactions in so manifold a manner.

Research is now in hand at this laboratory upon the effect of applying simultaneously both protein and lipids. In the absence of alcohols retardation of reaction should be observed. The effect of the addition of narcotics should, where protein is present in excess, be to accelerate hydrolysis as a result of elution of substrate; whether the amylase eluted from lipids is immediately adsorbed upon protein, or whether an enzyme-substrate compound, adsorbed on protein only to a small extent, is formed will be shown by experiment. In all probability, however, the behaviour of such systems will resemble that of protein systems alone, except where lipids are present in excess. Within the cell, locational influences may further come into play whereby an enzyme eluted from lipids situated in one part of a cell need not necessarily undergo adsorption upon protein situated in a different part of the cell. Thus the difference between the livers of summer and of winter frogs, mentioned by Lesser [1924] and by Przylecki, and consisting in the almost complete absence of glycogenolysis during the winter months, might be explained not only by the simultaneous adsorption of amylase and glycogen upon protein, but possibly also by the exclusive adsorption of the enzyme upon lipids.

It further appears probable that Lesser's [1927] observation, that alcohols cause augmented secretion of amylase from the submaxillary glands, can be explained on the assumption that it is adsorbed upon lipid constituents of these glands.

SUMMARY.

1. Amylase is adsorbed upon lipids to about 67 % of the quantity in solution.
2. Adsorbed amylase is quantitatively eluted by narcotics, partly by glycogen and hardly at all by water.
3. Glycogen does not undergo adsorption upon lipids.
4. The addition of lipids to the system glycogen-amylase causes retardation of hydrolysis.
5. Amylase adsorbed upon lipids is thereby inactivated; its activity can be restored by elution.
6. The enhanced secretion of amylase from the submaxillary glands under the influence of alcohol observed by Lesser is perhaps explicable on the basis of the above findings.

This paper was, similarly to the rest of the series, undertaken at the suggestion and under the supervision of Prof. St J. Przylecki, to whom I wish to express my sincere gratitude for his co-operation.

REFERENCES.

- Lesser (1924). *Biochem. Z.* **140**, 439.
— (1927). *Biochem. Z.* **184**, 125.
Przylecki, Niedzwiecka and Majewski (1927). *Biochem. J.* **21**, 1025.
Przylecki and Niedzwiecka (1928). *Biochem. J.* **22**, 34.