CCXI. CRYSTALLINE INSULIN.

By DAVID AYLMER SCOTT.

Connaught Laboratories, University of Toronto.

(Received April 28th, 1934.)

DURING the past few years the problem of isolating insulin in a crystalline form has been of great interest to the author for several reasons. When large quantities of crystalline material are required for chemical or clinical investigations the present methods of separation leave much to be desired. In the hands of an experienced worker a fair degree of success may be attained. Nevertheless, there would seem to be some unknown condition or conditions which must be fulfilled if a good yield of crystals is to be obtained. Attempts to duplicate successful experiments do not always produce a crystalline product. This has been noted by several workers, who have also drawn attention to the fact that different preparations of crude insulin show a marked variation in the ease with which they may be crystallised. Of further interest is the fact that it is a common belief that the chemistry of insulin is closely related to if not identical with that of proteins, and the results of an investigation concerning the exact conditions under which insulin crystallises would be of value for the isolation of other proteins in a pure form.

Insulin was first isolated in a crystalline form by Abel [1926]. The method utilised as the crystallising medium a solution which was highly buffered and consisted of several substances. In addition to ammonium acetate, it contained large quantities of brucine and pyridine. The isolation of the crystals was attributed to the fact that, by the use of such a complex system of weak bases, the acidity could be adjusted to the isoelectric point of insulin so slowly and so exactly that a supersaturated solution of insulin was obtained. The conditions then seemed to be such that the insulin separated in crystalline form. The problem of crystalline insulin was further investigated by Harington and the author [1929]. Using a buffer solution of ammonium acetate containing saponin, they were able to prepare crystals from amorphous insulin. In microscopic appearance, physiological activity and chemical analysis, these crystals were identical with those described by Abel. In this paper we stated that we were "quite unable to offer any rational explanation of the mechanism of the action of saponin in promoting crystallisation." Nevertheless two factors were apparent. One was the power of saponin to inhibit precipitation and thus permit the preparation of a supersaturated solution of insulin at the isoelectric point. The other was the fact that saponin seemed to sharpen the isoelectric point of foreign proteins in the amorphous insulin and thus permit the removal of many of the impurities at a lower acidity than that at which the insulin crystallised.

Since the publication of the saponin method, I have carried out many experiments on the crystallisation of insulin. In view of the fact that they have a distinct bearing upon the experimental work described later in the present paper, they should be mentioned briefly here. During the summer of 1929, many attempts were made to develop a method superior to the saponin procedure. It was felt that much would be gained if saponin could be replaced by some

pure substance which was highly surface-active. Of the many substances used in these experiments, two were found to be of great assistance in the crystallisation of insulin, namely potassium xanthate and thiocarbamide. Later work, however, showed that equally satisfactory results could be obtained by using definite quantities of various solvents, *e.g.* acetone, propyl alcohol or butyl alcohol. Since good yields of crystals were obtained even when working with 2-litre quantities of insulin solution, this method was an improvement on the original saponin procedure. However, when it became necessary to use a fresh phosphate buffer solution, it was found that it was no longer possible to obtain crystals. The addition of solvents or other substances previously found to be of value in the formation of crystals, did not overcome the difficulty. The fact that the fresh buffer solution was prepared in the same manner as the one which was found to be satisfactory, suggested that some change had taken place in the solutions after preparation. It seemed possible that this change might consist in the absorption of carbon dioxide from the air or the removal of some inorganic substance from the glass bottle. A great number of experiments were carried out with samples of this fresh buffer to which various amounts of carbon dioxide had been added. However, only negative results were obtained. Experiments were also conducted with this buffer solution to which varying amounts of the common ingredients of glass were added. These substances did not aid in the formation of insulin crystals. Since the distilled water used in the buffer solution which yielded crystals was obtained from a tin-lined still, it was thought that traces of tin might be the essential factor. Accordingly, experiments were conducted with the buffer solution to which various amounts of stannic chloride had been added. This substance also failed to promote crystallisation. However, after storing the buffer solution for about 3 months, that is, until January 1930, it was found that it was then satisfactory for use in the production of crystalline preparations of insulin. Again a fresh phosphate buffer solution was prepared and the same difficulties in crystallisation were encountered.

Since it was impossible to attribute successful crystallisation to any particular constituent of the phosphate buffer, the water or glass, and since it seemed likely that inorganic substances played a definite rôle in the crystallisation of insulin, there was a possibility that pancreas might contain inorganic substances which would assist in the crystallisation of insulin. In searching the literature, it was found that pancreas contains appreciable quantities of zinc [Lutz, 1926], cobalt and nickel [Bertrand, 1926]. To ascertain the effects of these metals upon the formation of insulin crystals, work was initiated in which zinc was added to the crystallising medium. The first experiment was conducted in the following manner. Two phosphate buffer solutions containing about 0.1 % insulin and 10 % acetone were adjusted to p_H 6.2. After standing for 24 hours at room temperature, the precipitates were examined microscopically, and found to be completely amorphous. To one of the beakers, a few granules of zinc were added. They were then allowed to stand for 2 more days and again examined. The beaker to which zinc had been added contained the characteristic rhombohedral insulin crystals, whereas the other precipitate remained completely amorphous. Using a solution of zinc chloride, a great number of experiments were carried out to ascertain the minimum amount of zinc necessary to promote the development of crystals in a phosphate buffer solution containing insulin. This amount was found to be about 2 mg. of zinc for each 100 ml. of phosphate buffer-insulin solution. The crystals obtained in this manner were separated from the solution and re-crystallised. A biological assay showed that they contained 25 i.u. of insulin per mg. Similar experiments were then carried out with

nickel chloride and cobaltic chloride. These substances likewise, in the same concentration as in the zinc experiments, greatly facilitated the formation of crystals. Since iron is grouped with cobalt and nickel in the periodic table, experiments were conducted with various concentrations of ferric chloride. It was found that iron was of no value in the formation of crystals. The effect of other elements in Group II of the periodic table was next determined. It was found that the addition of 2 mg. of cadmium per 100 ml. of buffer greatly facilitated crystallisation. Magnesium, however, had no such effect. During 1931, many similar experiments were conducted using small amounts of the heavy metals. None of these metals was found to aid in the separation of crystals from a phosphate buffer solution containing insulin. Thus it was shown that there were at least four metals, namely, zinc, cadmium, cobalt and nickel, which aided in the preparation of crystals from amorphous insulin. Three of these (Zn, Co and Ni) have been reported to be present in the pancreas. Zinc and cobalt were superior to the others in the promotion of the crystallisation of insulin.

Although the mechanism of the action of these metals in the formation of crystals from solutions of insulin was entirely unknown, there seemed to be three possibilities, *viz.* (1) these elements might form nuclei with the phosphate buffer which initiated the crystallisation of the insulin; (2) these elements might combine with amorphous foreign material and thus leave the insulin free to crystallise; (3) an insulin salt might be formed. In view of the fact that all these metals produce crystals having the same microscopic appearance, it was thought that the second possibility was the most plausible one.

During 1932 little or no work was done on the crystallisation of insulin in these laboratories. In 1933, however, it became necessary to prepare a large quantity of crystalline insulin. There were four methods by which this might be done: (1) the brucine-pyridine procedure, (2) the saponin method, (3) the phosphate buffer method, in which the buffer is stored for 3 months before using, (4) modifications of method (3), using small amounts of either zinc, cadmium, nickel or cobalt, which have already been outlined. On account of the difficulties which might be encountered in removing the metallic elements from the final product if any modification of the fourth method was used, it was decided to prepare the material by means of methods 2 and 3. It seemed particularly advantageous to use method 2, since there were at hand samples of saponin which had been found to be completely satisfactory for the process. After a sufficiently large quantity of crystalline insulin had been obtained, the material was recrystallised from a phosphate buffer solution. The product was thoroughly mixed and samples were removed for biological assay and chemical analysis. The results of assays using 400 mice showed that the crystals contained 25 i.u. per mg. The chemical analysis gave carbon 49.79 %, hydrogen 6.91 %, ash 1.46 %. The values for carbon and hydrogen agree remarkably well with those published by Abel, and by Harington and Scott [1929]. The ash, however, was somewhat higher than the values reported for crystalline insulin by Abel. Unfortunately, no ash values were recorded for the insulin crystals isolated by the saponin method by Harington and Scott. In some unpublished work, I have repeatedly found the ash of crystalline material to be approximately 1 %. The ash value of 1.46 %, while higher than usual, is not surprising when it is remembered that the insulin was recrystallised from a phosphate buffer solution. Thus it would undoubtedly contain a small amount of phosphate. Moreover, since one of the prime objects of this work was to obtain as large a quantity of crystalline material as possible from the insulin at hand, all the solutions were

allowed to stand for one week at 6° before removing the crystals. It is not inconceivable that, in this comparatively long time, some inorganic substances might be precipitated. These would then be included with the crystals and would increase the ash content of the final product. In view of my former findings, namely, that small quantities of zinc, cobalt and nickel facilitate the crystallisation of insulin, the ash was tested for the presence of these metals. It was found to contain phosphate and gave positive tests for iron and zinc. It was free from cobalt and nickel. The amorphous insulin from which the crystals had been prepared, also showed the presence of zinc but no cobalt or nickel. Similarly, the saponin used contained zinc and iron but showed negative tests for cobalt and nickel. Some time ago, Prof. Abel was kind enough to send to these laboratories two crystalline preparations of insulin which he had obtained by his brucine-pyridine method. The first preparation contained 1.03 % ash. This ash, to my surprise, gave a strongly positive test for zinc. The tests for cobalt and nickel were negative. This striking result was confirmed by analysis of a second sample of the same material. The second preparation gave an ash content of 0.6 % and likewise showed a positive test for zinc and negative tests for cobalt and nickel.

Attempts were next made to remove the inorganic substances contained in the crystalline material. Isoelectric precipitation was first tried but this procedure removed little or no inorganic material. Experiments were next conducted with isoelectric precipitation in various concentrations of aqueous acetone solutions. It was found that the ash content could be reduced to about 1.0 % by isoelectric precipitation in an aqueous solution containing about 20 % of acetone. If this product were dissolved in a small volume of 0.1 *N* HCl and the insulin precipitated by the addition of alcohol and ether, a material was obtained which contained 0.25 % ash. A second treatment with acid alcohol and ether yielded a product containing 0.15 % ash. This ash was tested for zinc and the test was still found to be positive. A sample of the material containing 0.15 % ash was assayed on 400 mice. It showed a potency of 24.8 i.u. per mg. Experiments were next carried out in an attempt to crystallise this product. From a freshly prepared phosphate buffer solution no crystals were obtained, whereas from a similar solution containing, in addition to the insulin, small amounts of zinc, cadmium, nickel or cobalt, the characteristic rhombohedral crystals readily formed. Since the last traces of metal apparently could not be removed by treatment with alcohol and ether, attempts were made to do so by electro-dialysis. The material so obtained contained 0.13 % ash and still gave a positive test for zinc. This product was assayed on 300 mice and showed no loss in potency. Attempts were made to crystallise this material. Crystals only formed when zinc, cobalt, nickel or cadmium was added to the phosphate buffer solution.

Thus it has been shown that zinc occurs in the crystalline insulin preparations of Abel, in the crystals obtained by the saponin procedure and also in commercial samples of insulin. In view of this apparent affinity between zinc and insulin, it was decided to prepare a quantity of crystals from a buffered solution of insulin, using the method involving the addition of small amounts of zinc chloride. Further it was intended in these experiments to attempt removal of the zinc by treatment with alcohol and ether followed by electro-dialysis, as mentioned above, and to ascertain whether this product still retained its physiological activity and the property to crystallise. Zinc was chosen in preference to cadmium, nickel or cobalt for three reasons. In the first place, it has been my experience that it promotes crystallisation slightly better than the

other elements. Secondly, the solubility of zinc chloride in alcohol and ether is such that it is more readily removed than either cobalt, nickel or cadmium. Further, zinc is a metal normally occurring in commercial preparations of insulin. These experiments are described in detail below.

EXPERIMENTAL.

A phosphate buffer solution was prepared. This contained 335 g. Na_2HPO_4 , $12\text{H}_2\text{O}$ and 55 g. KH_2PO_4 in 20 litres of freshly distilled water. The reaction of the solution was p_{H} 7.15. 250 ml. of this phosphate buffer were measured into a 600 ml. beaker, and to it was added an equal volume of distilled water together with 20 ml. of N HCl. The p_{H} was 2.3. Then 20 ml. of an insulin solution containing 600 mg. of protein were added, followed by 5 ml. of 0.5 % ZnCl_2 and 50 ml. of acetone. The solution was adjusted to p_{H} 6.2 by adding 15 ml. of N ammonium hydroxide. (The solution is generally clear. It sometimes happens, however, that if the insulin is very impure, there is a distinct haze at this acidity, and under these conditions it is preferable to adjust the reaction to p_{H} 6.5 to dissolve the precipitate, and then readjust the solution to p_{H} 6.2 with N HCl.) The inside of the beaker was then rubbed with a glass rod and allowed to stand. In about 20 minutes, crystals began to form, and in one hour the solution became quite opaque. After standing at room temperature for 6 hours, the mixture was placed in the refrigerator for 2 days. Twenty-four experiments were conducted in this manner simultaneously. At the end of 2 days, the supernatant liquids were decanted. The crystals were separated by centrifuging and dried *in vacuo*. The yield from the 24 experiments amounted to 7.5 g. Recrystallisation was then conducted in the following manner. The crystals were added to 150 ml. of water containing sufficient N HCl to dissolve most of the material. The insoluble material was removed by filtration. 750 ml. of phosphate buffer solution (p_{H} 7.15) were diluted with an equal volume of distilled water in a 2 litre beaker. This solution was heated to 55° and to it were added 150 ml. of acetone and 30 ml. of the clear insulin filtrate. Thus 1.5 g. of the material prepared by the first crystallisation were used. The solution was immediately adjusted to p_{H} 6.0 with N HCl and the inside of the beaker rubbed with a glass rod. The solution remained clear until it had cooled to about 40° . Crystals then began to form. A series of five experiments was conducted in this manner. After standing at room temperature for 6 hours, they were placed in a refrigerator. At the end of 2 days, the supernatant liquids were decanted, the precipitates combined and the crystals centrifuged in a 50 ml. centrifuge-tube. The crystals were washed three times with distilled water, using 40 ml. each time. They were then dried *in vacuo* and weighed 5.5 g. After being thoroughly mixed, 4.051 mg. were weighed (sample A) and dissolved in 10 ml. isotonic saline (p_{H} 2.5). This was further diluted and the physiological activity was determined by the mouse method of assay. The standard insulin solution used for all the biological assays in this work was prepared by dissolving 6.206 mg. of International Standard Insulin powder (8.0 units per mg.) in 107 ml. of isotonic saline (p_{H} 2.5). Samples of this stock solution were suitably diluted with isotonic saline as required for the assays. As shown by the protocols, the crystals contained 25.6 I.U. of insulin per mg. Chemical analysis showed that the crystals contained: C, 49.30; H, 6.91; N, 14.33; ash, 1.68 %.

A large proportion of the inorganic material was removed from the above recrystallised preparation by the following treatment: 0.5 g. of the crystals was dissolved in a 250 ml. centrifuge-tube containing 100 ml. of water acidified with

1.0 ml. of N HCl. After the addition of 20 ml. of acetone, the solution was adjusted to p_H 5.3 with N NaOH and placed in the refrigerator overnight. In the morning the mixture was centrifuged and the supernatant liquid discarded. The precipitate was dissolved in 10 ml. of 0.1 N HCl and the insulin precipitated immediately by the addition of 100 ml. of absolute ethyl alcohol and 140 ml. of ether. The mixture was placed in the refrigerator. The following morning it was centrifuged and the precipitate redissolved in 10.0 ml. of 0.1 N HCl. The insulin was again precipitated by adding alcohol and ether and placed in the refrigerator overnight. The following morning, the precipitate was removed by centrifuging and dried *in vacuo*. Four such experiments were conducted in this manner and the precipitates were thoroughly mixed. 6.805 mg. (sample B) were dissolved in 10 ml. of isotonic saline (p_H 2.5) and after further dilution a biological assay was made. As shown by the protocols, this material contained 26.2 i.u. of insulin per mg. The ash content was 0.12 %, and traces of zinc were still present. The product from the four experiments described above was then used in attempts to crystallise the insulin in the following manner. 100 mg. of this insulin powder were dissolved in 50 ml. of phosphate buffer solution containing 4 ml. of N HCl. To this were added 50 ml. of water and 10 ml. of acetone. 10 ml. of this solution were measured into each of a series of ten 15 ml. centrifuge-tubes. The first six tubes of this series were then adjusted with $N/3$ ammonium hydroxide to the following acidities, *viz.* p_H 5.23, 5.45, 5.56, 5.84, 5.95, 6.18. To solution No. 7 in this series, 0.1 ml. of 0.5 % $ZnCl_2$ was added, and to samples 8, 9 and 10 similar quantities of solutions of $CdCl_2$, $NiCl_2$ and $CoCl_2$ were added respectively. These last four insulin solutions, containing the metals, were then adjusted to p_H 6.2. The precipitates so produced were in striking contrast to those formed without the addition of metallic salts. Within an hour, the tubes to which zinc, cobalt or cadmium had been added contained large quantities of crystals entirely free from amorphous material. From the solution to which nickel had been added, a precipitate amorphous in appearance was obtained, but on examination the following morning this was found to be completely crystalline also. Furthermore, regardless of which salt had been added, the crystals all had the same microscopic appearance, namely, the rhombohedral form characteristic of insulin. In those tubes which did not contain these salts, the precipitates were amorphous. From an observation of the amount of precipitate in each of the first six tubes, the isoelectric point of insulin free from metals seems to be about p_H 5.5.

Since the ash of the material purified with alcohol and ether still showed traces of zinc, an attempt was made to remove this metal by electro-dialysis. This was done in the following manner. 1.0 g. of the insulin powder in a litre beaker was dissolved in 500 ml. of distilled water containing 2 ml. of N HCl. This solution was then electro-dialysed. For this purpose the type of apparatus of King and Dolan [1934] was used, in which the electrodes are placed in dialysing thimbles, through which water flows continuously. In this experiment, the electrodes were strips of platinum 1×2 in. in size. During the experiment distilled water acidified with HCl to p_H 2.5 flowed through the dialysing thimbles. The insulin solution in the beaker was stirred continuously. In spite of these conditions, a precipitate continued to form on the dialysing thimble containing the cathode. This precipitate was constantly scraped off and it redissolved in the insulin solution. Dialysis was continued for one hour using a current of 0.3 ampère. At the end of this time the insulin solution in the beaker was filtered, and N NaOH added until the first sign of a turbidity appeared. Further additions of 0.1 ml. of alkali were made at half-hour intervals until a flocculent

precipitate appeared. The beaker was then placed in the refrigerator. The following morning the precipitate was removed by centrifuging and the acidity of the supernatant liquid determined. It was found to be p_H 6.3. The precipitate was dissolved in 20 ml. of 0.1 *N* HCl. 10 ml. quantities of this solution were measured into each of two 250 ml. centrifuge-tubes, and to each were added 100 ml. of absolute ethyl alcohol and 140 ml. of ether. Both tubes were then placed in the refrigerator. The following morning, after centrifuging, the precipitates were dried *in vacuo*. These were intimately mixed and 6.99 mg. (sample C) were dissolved in 50 ml. isotonic saline p_H 2.5. An aliquot was further diluted and assayed biologically. As shown by the protocols, this material contained 23.1 i.u. of insulin, and had an ash content of 0.04 %. The ash gave a faintly positive test for zinc.

In attempts to crystallise the product obtained after dialysis, experiments were conducted in the following manner. 100 mg. were placed in a beaker and to them were added 50 ml. of phosphate buffer solution, 50 ml. of water and 4 ml. of *N* HCl. After the insulin had dissolved, 10 ml. of this solution were measured into each of a series of ten 15 ml. centrifuge-tubes. To each tube, 1.0 ml. of acetone was added. The solutions in the first six tubes of this series were adjusted with 0.25 *N* NaOH to acidities between p_H 5 and 6.5. To solutions 7, 8, 9 and 10, 0.1 ml. of 0.5 % $ZnCl_2$, $NiCl_2$, $CoCl_2$ and $CdCl_2$ was added respectively, and the acidities were adjusted to p_H 6.2. All the tubes in the series were rubbed with a glass rod. Within an hour, a flocculent precipitate formed in the first six tubes. In the solution containing the zinc chloride, crystallisation occurred readily, the precipitate settling to the bottom of the tube. In the experiments using the chlorides of nickel, cadmium and cobalt, precipitates also settled, although less firmly than in the zinc experiment. After standing at room temperature for 2 hours, the tubes were placed in the refrigerator. The following morning, the precipitates which had formed in the presence of the metals presented a striking contrast to those in the first six tubes. Without disturbing the precipitate in any tube of the series, a photograph was taken (Plate X). On microscopic examination, it was found that the precipitates in the tubes containing the metals were completely crystalline, whereas the first six in the series were amorphous. The mixtures were then centrifuged and the acidities of the supernatant liquids were accurately determined and found to be as follows: p_H 5.26, 5.43, 5.64, 5.82, 5.98, 6.18, 6.15, 6.20, 6.24, 6.16. The crystalline material in the last four tubes was separated by centrifuging, washed with distilled water and dried *in vacuo*. Samples of each were then weighed and assayed against the International Standard Insulin. The weights of the various samples were 3.316, 4.901, 3.104 and 2.652 mg. respectively. Each sample was diluted to 50.0 ml. with isotonic saline and aliquots further diluted for assay. These samples are designated in the protocols as D, E, F, G, respectively (tubes 7, 8, 9 and 10).

Since, in the above experiments, and in others like them, it was found that the presence of certain metals was of prime importance for the formation of insulin crystals, attempts were made to determine the possibility of promoting crystallisation by some other means. A series of experiments was carried out in which the acidity of the phosphate buffer-insulin solutions was adjusted with ammonium hydroxide instead of sodium hydroxide. The precipitates were amorphous. Experiments were also conducted with a phosphate buffer solution to which no acetone was added, and again amorphous products were obtained. Similar experiments were carried out with buffer solutions of ammonium acetate and sodium acetate. Although various solutions covering a wide range of acidity

Protocols of mouse tests.

International Standard Insulin				Insulin of unknown potency			
Test No.	Dilution	Amount injected per 20 g. mouse ml.	Convulsion rate	Dilution	Amount injected per 20 g. mouse ml.	Convulsion rate	Units per mg.
Sample A (insulin crystals by zinc method):							
1	1/8	0.25	12/24	1/175	0.25	13/24	26.3
2	"	"	14/24	"	"	7/24	18.5
3	"	"	12/24	"	"	12/24	25.0
4	"	"	9/24	"	"	11/24	27.0
5	"	"	14/24	"	"	15/24	26.0
6	"	"	20/24	"	"	15/24	20.0
7	"	"	7/24	"	"	9/24	27.5
8	"	"	7/24	"	"	14/24	34.3
Mean							25.6
Sample B (amorphous insulin after ether and alcohol):							
1	1/8	0.25	6/24	1/300	0.25	6/24	25.6
2	"	"	11/24	"	"	15/24	30.2
3	"	"	12/24	"	"	15/24	28.9
4	"	"	17/24	"	"	13/24	22.0
5	"	"	15/24	"	"	10/24	20.9
6	"	"	15/24	"	"	16/24	26.8
7	"	"	17/24	"	"	20/24	29.6
8	"	"	16/24	"	"	15/24	24.5
Mean							26.1
Sample C (amorphous insulin after electro-dialysis):							
1	1/8	0.25	13/24	1/57	0.25	16/24	26.7
2	"	"	10/24	"	"	5/24	18.2
3	"	"	10/24	1/60	"	12/24	27.1
4	"	"	8/24	"	"	7/24	23.6
5	"	"	7/24	"	"	5/24	22.4
6	"	"	10/24	"	"	9/24	23.9
7	"	"	13/24	"	"	7/24	18.9
8	"	"	10/24	"	"	9/24	23.9
Mean							23.1
Sample D (recrystallised insulin by zinc method):							
1	1/9	0.25	25/36	1/32	0.25	22/36	22.9
2	"	"	24/36	"	"	22/36	23.6
3	"	"	13/36	"	"	9/36	21.7
4	"	"	16/36	"	"	13/36	23.1
Mean							22.8
Sample E (recrystallised insulin by nickel method):							
1	1/9	0.25	24/36	1/50	0.25	19/36	23.1
2	"	"	15/36	"	"	13/36	25.0
3	"	"	21/36	"	"	20/36	25.8
4	"	"	17/36	"	"	17/36	26.3
Mean							25.1
Sample F (recrystallised insulin by cobalt method):							
1	1/9	0.25	19/36	1/30	0.25	25/36	25.3
2	"	"	20/36	"	"	32/36	31.3
3	"	"	25/36	1/35	"	25/36	29.0
4	"	"	15/36	"	"	16/36	29.9
Mean							28.9
Sample G (recrystallised insulin by cadmium method):							
1	1/8	0.25	17/24	1/23	0.25	14/24	22.4
2	"	"	23/24	"	"	23/24	25.2
3	1/9	"	12/24	1/26	"	8/24	21.3
4	"	"	14/24	1/23	"	19/24	28.0
Mean							24.2

were used, only amorphous precipitates were obtained. Only by adding small amounts of one of the four metals, namely, zinc, nickel, cobalt or cadmium, have we been able to crystallise this material, although it has approximately the same physiological activity as crystalline insulin.

DISCUSSION.

In the present paper, a very brief review of some work on the crystallisation of insulin has been reported. It was noted during 1930 and 1931 that small amounts of various salts (the chlorides of cobalt, cadmium, nickel and zinc) aided in the formation of crystals from amorphous insulin. It was thought at that time, since the same type of crystal was formed irrespective of the salt which was added, that these metals combined with the foreign proteins in the crude insulin preparations, leaving the insulin free to crystallise. This view was questioned, however, when it was found that crystalline insulin, prepared by the phosphate buffer method and the saponin method, contained zinc. When the ash content was reduced to 0.15 %, the insulin, although having the potency of crystalline material, failed to crystallise from a phosphate buffer solution. Crystals, however, could be formed readily when traces of zinc, cadmium, nickel or cobalt were added to the crystallising medium. Furthermore, two samples of crystalline insulin prepared by Prof. Abel by the pyridine-brucine method were found to contain zinc. There is also presented in detail a method for preparing quantities of crystalline material from amorphous insulin by adding small amounts of zinc chloride to a phosphate buffer solution of insulin. These crystals had the same physiological activity and chemical composition as the crystals prepared by other methods. The inorganic material was partially removed from the crystals by isoelectric precipitation, in 20 % aqueous acetone, and by two precipitations with acid alcohol and ether. The product contained 26.2 i.u. of insulin per mg. It contained 0.12 % ash and showed traces of zinc. This material, however, would only crystallise in the presence of zinc, nickel, cobalt or cadmium. A material having an ash content of 0.04 % and containing 23.1 i.u. of insulin per mg. was obtained by electro-dialysis. This material could only be crystallised in the presence of traces of zinc, cadmium, nickel or cobalt.

The fact that the product after electro-dialysis, containing only 0.04 % ash and giving a faintly positive test for zinc, showed certain properties not usually associated with insulin, is worthy of consideration. When acid solutions of this material were neutralised slowly with sodium hydroxide, no flocculation occurred until the solution had an acidity of p_H 6.3—a significant difference from the commonly accepted isoelectric point of insulin. Biological assays showed that the precipitate was slightly less active than crystalline insulin. This was surprising, since in the electro-dialysis experiment, reported in the introductory part of this paper, no loss in potency could be detected. The differences between these two products may be due to the fact that, in the first experiment, dialysis was continued for $\frac{1}{2}$ hour only. The material so obtained, precipitated in a more acid solution, had an ash content of 0.13 %, and gave a positive test for zinc. The discrepancies between these two experiments will be further investigated.

Perhaps, in view of the present findings, the brucine-pyridine method and the saponin method of crystallising insulin should be discussed further. The success of the saponin method can be explained readily by the fact that saponin suitable for crystallising insulin contains appreciable amounts of zinc and is highly surface-active. This latter factor is comparatively important, as very few insulin crystals are obtained when using the zinc method described in this

paper, unless acetone or some other substance which reduces surface tension is used. It is highly probable that the saponin used by Harington and Scott contained zinc or some other inorganic elements essential for the success of their process. Various saponins, upon analysis, were shown by them to contain "large amounts of inorganic matter even including traces of the heavy metals." Several samples of saponin in my laboratory when tested for zinc, all gave positive tests for this element. Further, a sample of saponin which had been found to be very satisfactory for the crystallisation of insulin, contained appreciable amounts of zinc. This fact, in view of the results of the present investigation, may explain in part the success of the saponin method. With respect to crystals prepared by the brucine-pyridine method of Abel, it might be well again to point out that two samples of crystalline insulin from his laboratory contained zinc. The most probable source of the metal is the pancreatic glands from which amorphous insulin is obtained. Lutz [1926] has reported that the pancreas of various species contains appreciable quantities of this element. Working with the pancreatic glands of the ox, Delezenne [1919] found 23 mg. of zinc per kg. of tissue. In the commercial preparation of insulin, the hormone is extracted from the pancreas with acid alcohol. The alcohol is commonly acidified with hydrochloric acid. Thus the solvent which is satisfactory for the extraction of insulin is also one which would dissolve the zinc contained in the pancreatic glands. Perhaps the most important step in the purification of insulin is isoelectric precipitation. Experiments have shown that zinc adheres to insulin protein when precipitated at the isoelectric point. It is thus most likely that all insulin preparations contain zinc. Certainly all the preparations which have been thus far tested gave a positive test for this element. However, in view of the use of slightly different methods of preparation, it is conceivable that all commercial preparations of insulin do not contain the same amount of zinc. This may account for the fact that insulin from one manufacturer sometimes crystallises more readily than that from another source. In the Abel method of crystallisation there is, of course, the added possibility that the brucine contains small amounts of this element. That a phosphate buffer solution, aged for 3-4 months, more readily promotes the crystallisation of insulin can hardly be attributed to small amounts of zinc, nickel, cobalt or cadmium absorbed from the glass, since glass contains little, if any, of these substances. It is possible, however, that some other element in glass is responsible for this phenomenon. It is also possible that traces of other inorganic substances in the glass have dissolved and are combining with the foreign proteins in insulin and thus leaving the zinc in commercial insulin free to initiate crystallisation.

The observation that crystalline insulin prepared by the brucine-pyridine method of Abel or by the saponin method of Harington and Scott contains zinc may explain the fact that solutions of crystalline insulin are slightly less stable than amorphous preparations. In the crystalline material there is a higher concentration of zinc per unit of physiological activity than in amorphous insulin. In solution, this zinc may exhibit reducing properties and thus destroy the physiological activity of the hormone. From this investigation, there arises the question of the possibility of the rhombohedral crystals being a metallic salt of insulin. There are at least two reasons for believing that such may be the case. In the first place, the addition of small amounts of zinc, cadmium, nickel or cobalt to solutions of amorphous insulin greatly facilitates the formation of crystals. Secondly, if steps are taken to remove the metals from crystalline preparations of insulin, the resulting material cannot be crystallised unless one of the metals cadmium, zinc, nickel or cobalt is added. Thus it

seems possible that a salt of insulin has been formed—probably a simple salt. The fact that the crystals prepared by the addition of the different salts have the same microscopic appearance might lend support to this view. If these metals form simple salts with insulin, other proteins might be expected to behave in a similar manner. Investigations are now being conducted along this line of endeavour. On the other hand, these crystalline preparations of insulin may be double or triple salts. In this connection, one is reminded of the fact that many diabetics who are hypersensitive to commercial preparations of insulin are also hypersensitive to crystalline insulin. If crystalline insulin is the same chemical entity as occurs in human pancreas, the above reactions would not be expected to take place. That crystalline insulin does not give an X-ray pattern under ordinary conditions, is another fact indicating that the crystals may not be a simple salt. Thus it is felt that investigations on the problem of the crystallisation of insulin, together with that of the chemical constitution of the crystals, must be continued, bearing in mind the very definite rôle played by some of the metallic elements.

SUMMARY.

It has been shown that insulin crystals, prepared by the pyridine-brucine method or by the saponin method, contain zinc. A method for obtaining insulin crystals from amorphous preparations by means of zinc chloride is described. A process for reducing the inorganic substances contained in the insulin preparation to as low as 0.04 % is given. Such material could not be crystallised by the ordinary procedures. Crystals were obtained only in the presence of small amounts of zinc, nickel, cobalt or cadmium. The significance of these findings has been discussed.

I wish to express my indebtedness to Dr H. Stantial, of the Department of Chemistry, for making the many ash estimations and qualitative tests for inorganic substances. I am also grateful to Dr J. C. FitzGerald and Dr C. H. Best for their interest in this work.

REFERENCES.

- Abel (1926). *Proc. Nat. Acad. Sci.* **12**, 132.
- Bertrand (1926). *Science*, **64**, 629.
- Delezenne (1919). *Ann. Inst. Pasteur*, **33**, 68.
- Harrington and Scott (1929). *Biochem. J.* **23**, 384.
- King and Dolan (1934). *Canad. Chem. Metal.* **18**, 7.
- Lutz (1926). *J. Ind. Hygiene*, **8**, 177.

Attempts to recrystallise insulin of low ash content.

Tubes 1-6 contain insulin precipitated from phosphate buffer solutions adjusted to various acidities. All precipitates were amorphous. In tubes 7-10 are precipitates prepared from buffered solutions of insulin containing traces of the metals as indicated. The solutions were adjusted to the same acidity as the solution in tube 6. These precipitates were crystalline.