CLXXXVI. PURINOLYTIC ENZYMES OF THE HUMAN ORGANISM.

By RICHARD TRUSZKOWSKI.

From the Biochemical Laboratory, Faculty of Medicine, Warsaw University.

(Received September 24th, 1930.)

IN spite of the importance to human pathology of elucidating whether or not the human organism is capable of decomposing uric acid no actually decisive experiments on this question have been described, and the papers published give conflicting results. Pfeiffer [1906] stated that the kidney tissue of adults has a strong uricolytic action. Schittenhelm and Schmidt [1907] found that all tissues of new-born children are able energetically to destroy uric acid, and Cannata [1910] found that the human placenta possesses similar properties. Schittenhelm expressed the opinion that the reason why uricolysis did not normally take place in the human organism was that certain inhibiting substances were present in the body fluids. Folin, Berglund and Derick [1924] came to the conclusion that uric acid can, in normal persons, be oxidised by circulating blood, but that this property is immediately lost by removal of blood from circulation. Folin's experiments indicated, further, that the blood of gouty subjects was not able to destroy uric acid, and on this observation Folin based a theory of the etiology of gout. Flatow [1926] found that shed blood contained a uricolytic substance termed by him urico-oxidase; this however appears to be haemoglobin entering solution on haemolysis of erythrocytes, as is indicated by Gomolinska's work [1928]. Rowinska (unpublished work) showed further that unhaemolysed blood does not, under aseptic conditions, destroy uric acid. Wiechowski [1909], who examined various human organ preparations, was not able to find any uricolytic action, and similar results were obtained by Künzel and Schittenhelm [1908] and by Wells and Corper [1909]. Beck and Truszkowski [1930], using glycerol-chloroform extracts of foetal tissues taken from the 8th to the 40th week of pregnancy, found no evidence of the presence of uricase in unshaken systems.

The above somewhat chaotic results must be ascribed in some cases (Wells and Corper) to a misconception as to the nature of uricase, which was previously thought to be a soluble enzyme, and which was accordingly, in certain of the above researches, sought for in the extracts of organs instead of in the residue after extraction, which according to Truszkowski's work [1930] alone contains this biocatalyst. A second source of error lies in the action of micro-organisms on uric acid in the presence of such weak antiseptics as thymol, sodium fluoride, etc.; to this we would ascribe the results of Pfeiffer, Schitten-

helm and Schmidt, and Cannata. Finally, the disturbing effect of xanthine oxidase was not allowed for in the above experiments on isolated tissues and their extracts. Thus, even if the tissues examined had possessed a feeble uricolytic action, the diminution in the uric acid content of the systems might readily be more than compensated for by the formation of fresh uric acid from purine bodies present in the organ or its extract; *i.e.* uricase cannot with certainty be stated to be absent unless xanthine oxidase is either removed or proved to be absent. As to metabolism experiments, such as those of Folin, in which it was attempted to obtain a balance between injected and excreted uric acid, these cannot be considered to be of decisive value, in view of the fact that uric acid is excreted not only with the urine, but also with the bile, and possibly also directly into the intestines, in which bacterial decomposition may, and in all probability does supervene.

The above critical survey of previous work done in this field shows that from none of the papers hitherto published can it be definitely stated whether or not human tissues are possessed of uricolytic properties either in intra- or extra-uterine life.

As to xanthine oxidase, Wells and Corper [1909] were unable to demonstrate its presence in the human foetus before the 6th month of pregnancy. Beck and Truszkowski [1930] found this enzyme in glycerol-chloroform extracts of foetal tissues from the 8th week of pregnancy up to the full term; the results of these authors are however only qualitative.

For the above reasons it was considered desirable to re-investigate the whole question, applying the methods of preparation of uricase described by the present author [1930]. The establishment of the insolubility of uricase affords a simple method for its separation from soluble xanthine oxidase, and therefore of the examination of the action of these two enzymes separately, so that the results obtained appear to be free methodically from the objections raised to those of earlier workers. The organs examined were the liver and, in some cases, the kidneys. These were taken for the greater part from embryos, and in some cases from healthy adults who had died suddenly from accidental causes. These organs were selected because in mammals other than man and the anthropoid apes either the liver or kidney or both of these organs possess the most pronounced uricolytic activity.

EXPERIMENTAL.

The liver and kidneys were removed as soon as possible after death (0.5–3 hours) and were minced, or, if too small, as in the case of foetuses born before the 5th month of pregnancy, cut up into small pieces, and ground in a mortar with chloroform. Ten times its weight (in the case of the youngest foetus 15–20 times) of water was added to the comminuted tissue, which was left with excess of chloroform during 48 hours, after which part of the suspension was centrifuged, and the uricolytic actions of the whole suspension, of the centrifugate, and of the residue were examined. In view of Schittenhelm's

theory of the presence of substances inhibiting uricolysis, the effect of adding centrifugate to systems containing bovine kidney uricase was also examined (Table I). Uric acid solutions were in all cases prepared by dissolution of uric acid in boiling saturated lithium carbonate. Uric acid was determined colorimetrically by the method of Folin and Denis [1913].

A number of 100 cc. flasks each containing 10 cc. of about 0.2 % lithium urate solution (p_H 8.5) were prepared. To certain of these flasks were added 10 cc. of water and 10 cc. of centrifugate, of a suspension of the residue after centrifuging, of the whole suspension, or of a 2 % suspension of bovine kidney uricase. In addition to these systems, flasks were prepared containing 10 cc. of lithium urate solution and 20 cc. of water, 10 cc. of centrifugate and 20 cc. of water, and finally, 10 cc. of lithium urate solution, 10 cc. of centrifugate and 10 cc. of a suspension of bovine uricase. To each flask 1 cc. of toluene was added, in order to inhibit bacterial growth. The flasks were then tightly stoppered, using rubber corks, and were placed in an Ostwald water thermostat maintained at $37^\circ \pm 0.2^\circ$, with constant shaking. It has been shown by the author that uricase is most active under these conditions, 1 g. of bovine kidney powder being capable of catalysing the oxidation of 220 mg. of uric acid in 1 hour.

Three livers, taken from a foetus born at term (*placenta praevia*), and from two adult females, aged 27 and 57 years, were examined. Death took place in both cases suddenly, without any previous history of disease, so that the results can be regarded as being typical of normal healthy organs. Systems containing preparations of the foetal liver were shaken for 14.5 hours, whilst those of adult liver were shaken for 24 hours; those systems containing bovine uricase were shaken respectively for 1, 2 and 2 hours. The results given in Table I show that in no case is any diminution in uric acid content demonstrable, except in the presence of bovine uricase. It follows that the human liver is not possessed of even the feeblest uricolytic power. The uric acid content of systems containing centrifugate or whole extract is in all cases slightly higher than that of the control systems or of those to which the residue was added—this difference is due to preformed uric acid present in the extract, as is shown by the results found for the diluted centrifugate alone. The centrifugate has a slight inhibitory influence on the action of bovine uricase; this however, as was shown by the author in a previous paper [1930], may be ascribed largely to increase in viscosity of the medium occasioned by soluble proteins. Since however the corresponding centrifugate of bovine kidney extracts has a considerably greater inhibiting action, Schittenhelm's hypothesis that human organs would be uricolytically active were it not for the presence of inhibitory substances must be abandoned in its entirety.

It appears from the above experiments that during extra-uterine life the human organism does not possess catalysts capable of decomposing uric acid. The possibility exists that such factors may exist prior to birth, at some stage of embryonic development corresponding to that of mammals other than man,

Table I. *Action of preparations of human liver on uric acid.*

No. of exp.	Material	Systems contain						Mg. uric acid		Remarks
		Uric acid cc.	Suspension of residue cc.	Centri- fugate cc.	Whole extract cc.	2 % suspension of bovine uricase cc.	Water cc.	Found	Oxidised	
1	10-month old foetus. Systems shaken 14.5 hrs. at 40°	10	10	—	—	—	10	22.8	0.0	Shaken 1 hr.
		10	—	10	—	—	10	24.3 (-1.5)	0.0	
		10	—	—	10	—	10	24.3 (-1.5)	0.0	
		—	—	10	—	—	20	1.5	—	
		10	—	10	—	10	—	17.55 (-1.5)	6.75	
		10	—	—	—	10	10	14.7	8.1	
		10	—	—	—	—	20	22.8	—	
2	Adult, ♀, aged 27. Systems shaken 24 hrs. at 37°	10	10	—	—	—	10	18.6	0.0	Shaken 2 hrs.
		10	—	10	—	—	10	21.6 (-3.0)	0.0	
		10	—	—	10	—	10	21.6 (-3.0)	0.0	
		—	—	10	—	—	20	3.0	—	
		10	—	10	—	10	—	6.35 (-3.0)	15.25	
		10	—	—	—	10	10	2.43	16.17	
		10	—	—	—	—	20	18.6	—	
3	Adult, ♀, aged 57. Systems shaken 24 hrs. at 38°	10	10	—	—	—	10	18.6	0.0	Shaken 2 hrs.
		10	—	10	—	—	10	20.1 (-0.5)	0.0	
		10	—	—	10	—	10	20.1 (-0.5)	0.0	
		—	—	10	—	—	20	0.5	—	
		10	—	10	—	10	—	4.2 (-0.5)	14.8	
		10	—	—	—	10	10	2.43	16.17	
		10	—	—	—	—	20	18.6	—	

Note. Figures in brackets refer to preformed uric acid.

which might be presumed to possess uricase. Morphologically, such a stage occurs during the early phases of development, before the 8th week of pregnancy; it is in practice not possible to obtain intact embryos from these early stages. Since however Jones and Austrian [1907], and Mendel and Mitchell [1907] found that the appearance of a different enzyme, xanthine oxidase, does not take place till a comparatively late stage of development in pig embryos, it is possible that the disappearance of uricase might be similarly retarded in the human embryo. Przyłęcki and Rogalski [1927-8] reported the appearance of uricase in embryonic chicks on the 2nd day of development and its disappearance on the 10th day.

As to xanthine oxidase, Wells and Corper [1909] came to the conclusion that this does not appear before the 6th month of pregnancy. Beck and Truszkowski [1930] on the other hand demonstrated its presence in all embryos examined, from the 8th to the 40th week of pregnancy; no attempt was however made in this research to obtain other than qualitative evidence of its presence.

The livers and kidneys of six embryos from the 12th to the 40th week of pregnancy were examined for uricase by the method described above, *i.e.* the residue obtained by centrifuging chloroform-water extracts of the finely divided organs was shaken for 24 hours at 37° with lithium urate solution. In no case was any oxidation of uric acid found to take place, so that the conclusions drawn for post-natal organs can be extended to all foetuses after the 12th week of pregnancy.

Table II. *Xanthine oxidase in human liver extracts.*

Week of pregnancy	Mg. uric acid found after leaving 20 cc. of centrifugate 48 hrs. at 25° with 10 mg. of xanthine
12	0.3
18	3.06
20	2.00
24	3.3
28	2.81
40	9.5

The centrifugates were diluted to a volume of about 22 cc., and two 10 cc. portions of each centrifugate were placed in a number of flasks. One flask of each pair was heated at 100° for 3 min. in order to destroy the enzyme. 10 cc. of a 1 % suspension of xanthine and 2 cc. of toluene were then added to each flask, and the flasks, after plugging with cotton-wool, were left for 48 hours at 25°, after which time uric acid was determined. No uric acid was found in those flasks which had previously been heated at 100°, or in those containing kidney extract centrifugates. Varying amounts were found in the systems containing liver extract centrifugates (see Table II). The activity of the respective centrifugates obviously depends not only on the enzyme content of the organ, but also on its dilution, which was necessarily greater for material taken from early pregnancy. It will be seen that little variation exists between the 18th and the 28th week of pregnancy, whilst for the 40th week practically all the xanthine added (10 mg.) had been converted into uric acid.

SUMMARY.

1. The livers of a new-born child and of two adults have been examined for uricase, which is shown to be absent.
2. The livers and kidneys of six fetuses from the 12th to the 40th week of pregnancy have been similarly examined; uricase is in all cases absent.
3. Xanthine oxidase is absent from the extracts of the kidneys of the above fetuses.
4. Xanthine oxidase has been found in all liver extracts from the 12th to the 40th week of pregnancy; Wells and Corper's view that this enzyme does not appear before the 6th month of pregnancy is therefore incorrect.

The author wishes to express his gratitude to Prof. St J. Przyłęcki for his interest in this research and to Docent H. Beck for supplying the material.

REFERENCES.

- Beck and Truszkowski (1930). *Med. Doswiad.* (Warsaw), **11**, 36.
Cannata (1910). *Fol. Clin. Microscopia*, **2**, 10.
Flatow (1926). *Münch. med. Woch.* **73**, 12.
Folin, Berglund and Derick (1924). *J. Biol. Chem.* **60**, 361.
Folin and Denis (1913). *J. Biol. Chem.* **14**, 95.
Gomolinska (1928). *Biochem. J.* **22**, 1307.
Jones and Austrian (1907). *J. Biol. Chem.* **3**, 227.
Künzel and Schittenhelm (1908). *Zentr. Phys. Path. Stoffwechsels*, **3**, 721.
Mendel and Mitchell (1907). *Amer. J. Physiol.* **20**, 97.
Pfeiffer (1906). *Beitr. chem. Physiol. Pathol.* **7**, 463.
Przyłęcki and Rogalski (1927-8). *Arch. int. Physiol.* **29**, 423.
Schittenhelm and Schmidt (1907). *Z. exp. Path. Ther.* **4**, 424.
Truszkowski (1930). *Biochem. J.* **24**, 1340, 1349, 1359.
Wells and Corper (1909). *J. Biol. Chem.* **6**, 469.
Wiechowski (1909). *Arch. exp. Path. Pharm.* **60**, 180.