REFERENCES

- Beijerinck, M. W. [1901]. *Arch. néerl. Sci. Ser. (2)*, **6**, 212.
 Butlin, K. R. [1938]. *Biochem. J.* **32**, 508, 1187.
 Callow, A. B. [1923]. *J. Path. Bact.* **26**, 320.
 Cosbie, A. J. C. Tošić, J. & Walker, T. K. [1942]. *J. Inst. Brew.* **48**, 82.
 Kluyver, A. J. [1924]. *Hoppe-Seyl. Z.* **138**, 100.
 Kolthoff, I. M. & Sandell, E. B. [1937]. *Quantitative Analysis*. New York: Macmillan.
 McLeod, J. W. & Gordon, J. [1923]. *J. Path. Bact.* **26**, 326.
 Morgulis, S., Beber, M. & Rabkin, I. [1926]. *J. biol. Chem.* **68**, 521, 535.
 Shimwell, J. L. [1936]. *J. Inst. Brew.* **42**, 585.
 Virtanen, A. I. & Karström, H. [1925]. *Biochem. Z.* **161**, 9.
 Visser't Hooft, F. [1925]. Thesis, Delft.
 — [1939]. Statement in *Bergey's Manual of Determinative Bacteriology*, 1939 ed. p. 231.

Riboflavin, Vitamin B₆ and Filtrate Factors in Wheaten Flours and Offals

By A. M. COPPING, *The Division of Nutrition, Lister Institute, Roebuck House, Old Chesterton, Cambridge*

(Received 9 October 1942)

In recent years there have been many studies of the vitamin B₁ content of wheaten flours and breads, but relatively few estimations have been made of the content of members of the vitamin B₂ complex. Some five years ago Copping & Roscoe [1937] published a survey of the water-soluble B vitamins in yeast, flour and bread, and gave values for the riboflavin and 'factors other than flavin'. These estimations were, however, made at a time when our knowledge of the members of the vitamin B₂ complex was more incomplete than it is at present, and the methods used appear far from satisfactory in the light of later work. New methods for the biological estimation of riboflavin have been described by El Sadr, Macrae & Work [1940], and on the basis of these methods others have now been evolved for estimating vitamin B₆ and filtrate factors. Almost all of the previous values for riboflavin and vitamin B₆ which have been put forward for flours have been obtained by chemical or microbiological methods.

The term filtrate factors is taken to mean all growth factors of the vitamin B₂ complex required by the rat, other than riboflavin, nicotinic acid and vitamin B₆ [El Sadr, Hind, Macrae, Work, Lythgoe & Todd, 1939; Macrae, Todd, Lythgoe, Work, Hind & El Sadr, 1939]. Filtrate factors are now known to include pantothenic acid, biotin and one or more unidentified substances. Lepkovsky [1942] reviews the subject most lucidly.

It is important at the present time to have some precise indication of the nutritive value of the flour and bread available to the people of this country, and on this account, as well as to obtain values by biological methods for comparison with those obtained by other methods, the further study of wheaten flours was undertaken.

MATERIALS

The materials used were supplied through the courtesy of the Research Association of British Flour Millers, who arranged to have suitable flours milled and to store some of the grist so that further samples could be milled if required. In January 1941 the following samples were milled and put into store in a cool place in airtight containers: (1) straight run unbleached white flour of 73% extraction; (2) whole-meal flour of 100% extraction; (3) weatings or middlings; (4) bran; (5) germ. The germ represented all the clean germ obtained in the milling of the white flour, and the bran and weatings together comprised the balance of the wheat after removal of 73% as white flour.

About 6 months later, when the Accessory Food Factors Committee of the Medical Research Council and the Lister Institute had laid down the specification for the preparation of the 85% extraction national wheatmeal [Medical Research Council, 1941], a sample of this type of flour was milled from the original grist and tested in the same way as the other samples.

Table 1. *Composition of the flours and offals (%)*

	Water*	Ash	Nitrogen	Protein
73% flour	13.6	0.411	2.042	11.64†
85% flour	12.64	0.861	2.103	11.98†
100% flour	13.53	1.545	2.190	12.48†
Weatings	14.6	4.31	3.04	17.33†
Bran	13.9	6.55	2.495	15.6†
Germ	13.98	4.18	4.76	29.75†

* Dried at 105–110°.

† N × 5.7.

‡ N × 6.25.

The vitamin estimations were made on the flours as they were received; they contained about 13 % of water (Table 1).

Since the earlier results of Copping & Roscoe [1937] had shown that there was no great difference in the amounts of vitamin B₂ factors present in flours and in the breads baked from them, it was considered unnecessary to bake breads from the flours used for the present investigation, and estimations were made only on the flours.

METHODS

In all tests the criterion was the rate of growth of young rats weaned at 21 days and given the basal diet immediately. In each test comparison was made with a standard preparation of the vitamin in question. Littermate rats, as far as possible of the same sex, were given the doses to be compared, so that for each group of four or more animals on a dose of the cereal product there was a corresponding group on a dose of the standard [Coward, 1938].

The rats received a basal diet containing 60 % carbohydrate (maize starch or cane sugar), 20 % purified casein, 15 % fat and 5 % McCollum's normal salt mixture. The fat consisted of 4 parts of hardened arachis oil and 1 part of pure lard. When maize starch was the source of carbohydrate the dry ingredients were mixed with their own weight of water and the mixture was steamed for 3–5 hr., depending on the amount made up, in order to dextrinize the starch and guard against refection [Fridericia, Freudenthal, Gudjonsson, Johansen & Schoubye, 1927]. The sugar-containing diet was not cooked.

The flours tested were made up into similar diets containing 68 % flour, 12 % casein, 15 % fat and 5 % salt mixture, and these were cooked in the steamer in the same way as the basal diet. These diets contained approximately the same proportions of carbohydrate, protein and fat as the basal diet and were given as daily doses in weighed amounts, the remainder of the animal's food requirement being made up by giving the basal diet *ad lib*. In the case of bran, difficulty was experienced in making a cohesive mixture with a high proportion of bran in the diet, and a diet containing 30 % bran, 30 % maize starch, 20 % casein, 12 % arachis oil, 3 % lard and 5 % salt mixture was used. The doses of test diets were given in amounts varying from 3 to 10 g. daily, and if any of the test diet was left an equivalent extra amount was given on the next day. As the diets were mixed with their own weight of water the intake of flour was calculated as 34 % of the diet eaten. Thus a daily dose of 10 g. of the test diet was equivalent to an intake of 3.4 g. of the flour. Wheat germ was a sufficiently rich source of all the factors tested to be given alone as a separate dose apart from the diet.

Riboflavin estimation. The method was essentially that of El Sadr *et al.* [1940]. The basal diet containing maize starch was supplemented with 0.1 ml. cod liver oil as a source of vitamins A and D, with pure vitamin B₁, and with a whole-liver extract, treated with charcoal to remove the riboflavin, as a source of all the vitamin B₂ factors except riboflavin. Norite charcoal, as used in the original method, was no longer available, but a decolorizing charcoal was substituted and, after some adjustment of quantities, was found satisfactory. In every test two doses of pure riboflavin were given as standard for comparison, usually 6 and 12 µg. per day. The test materials were fed at two levels also, and 4–8 rats were used for each dose level. In the tests with white flour the riboflavin content was so low that the response was too poor to obtain a result of any accuracy; 6 µg. riboflavin were therefore given daily in addition to the flour dose, and the value of the latter deduced by calculation from the curve of response.

Vitamin B₆ (pyridoxine) estimation. The method for the biological estimation of vitamin B₆ was based on that for riboflavin. Rats taken at weaning and weighing about 40 g. were given the vitamin B-free basal starch diet with the addition of daily doses of vitamin B₁ and cod liver oil for 7 days. They were then transferred to a basal diet containing sugar and given additional daily supplements of 40 µg. riboflavin and a fuller's earth filtrate from a liver residue extract. This filtrate supplied the filtrate factor portion of the vitamin B₂ complex; a dose of 1 ml. was equivalent to 6 g. of the original liver. It was found unsatisfactory to give a starch diet in tests for vitamin B₆, as all cereal starches contain some of this vitamin [Chick, Macrae & Worden, 1940]. After some preliminary experiments it was found satisfactory to give the basal diet and supplements other than vitamin B₆ for a standard period of 14 days before administering the test material. The rats usually grew well in the 1st week, but very poorly in the 2nd week, and showed a satisfactory response to a subsequent adequate dose of vitamin B₆. Two levels of dose of pure crystalline vitamin B₆ hydrochloride were given in every test, and the test material was fed at two levels as in the riboflavin tests. This permitted of a good comparison in most cases, since flours and offals were good sources of vitamin B₆.

Estimation of filtrate factors. The filtrate factor portion of the vitamin B₂ complex is known to contain pantothenic acid and at least one other factor [Lythgoe, Macrae, Stanley, Todd & Work, 1940]. An attempt was made to estimate pantothenic acid separately, but so far difficulty has been experienced in devising a basal diet free from pantothenic acid yet containing all other factors of the vitamin B₂ complex. It has, therefore, been possible

only to test for the whole filtrate complex. For this purpose young rats were prepared as for the vitamin B₆ tests and were given the basal sugar diet supplemented with 10 µg. crystalline vitamin B₁, 40 µg. riboflavin, 10 µg. crystalline vitamin B₆ hydrochloride, 1 mg. inositol, 1 mg. nicotinic acid, 3 mg. choline chloride and 0.1 ml. cod-liver oil daily. It was thought advisable to give the additional supplements of inositol, nicotinic acid and choline chloride when the other B₂ vitamins were supplied in the crystalline form. Choline chloride was certainly needed, for it was observed in some tests, before we had seen the reports of György & Goldblatt [1940], that the young rats occasionally died without apparent cause within a week of receiving the basal diet with vitamin B₁, riboflavin and vitamin B₆. Abnormalities of the liver and kidneys, as described by György & Goldblatt, were found in these animals at autopsy, and after choline chloride was given no further losses occurred. As in the vitamin B₆ tests

Table 2. *Summary of tests for riboflavin content of flours and offals*

Supplement	Daily dose	No. of rats	Av. weekly gain in wt. g.	Calc. content of riboflavin µg./g. fresh wt
1. White flour: 73% extraction	2.7 g. + 6 µg. riboflavin	14	14	0.85
Riboflavin	6 µg.	14	12	
	12 µg.	14	16	
2. National wheatmeal: 85% extraction	1.7 g.	8	7.5	2.0
	3.4 g.	8	10	
	2.7 g. + 6 µg. riboflavin	4	16	
Riboflavin	6 µg.	12	11	
	12 µg.	12	15	
3. Wholemeal: 100% extraction	2.0 g.	4	10	2.3
	4.0 g.	4	14	
	2.7 g. + 6 µg. riboflavin	4	17	
Riboflavin	6 µg.	8	12	
	12 µg.	8	16	
4. Weatings	1.7 g.	4	14	4.0
	2.7 g.	4	15	
Riboflavin	6 µg.	4	13	
	12 µg.	4	16	
5. Bran	1.6 g.	3	14	4.8
	2.16 g.	5	18	
Riboflavin	12 µg.	4	18	
	18 µg.	4	21	
6. Germ I	200 mg.	4	9	7.8
	400 mg.	4	14	
Riboflavin	6 µg.	4	16	
	12 µg.	4	19	
7. Germ II	200 mg.	4	12	12
	400 mg.	4	13	
Riboflavin	6 µg.	3	16	
	12 µg.	3	20	

a preliminary period of 14 days with all vitamin B supplements other than filtrate factors was found suitable as a means of preparing the rats for testing. As no source of filtrate factors in a pure form was available a dried baker's yeast was taken as standard of comparison in these tests; it was given at two levels of 100 and 200 mg. daily. The results were calculated in terms of mg.-equivalents of the standard yeast.

Table 3. *Summary of tests for vitamin B₆ content of flours and offals*

Supplement	Daily dose	No. of rats	Av. weekly gain in wt. g.	Calc. content of vitamin B ₆ hydrochloride µg./g. fresh wt.*
1. White flour: 73% extraction	1.7 g.	8	11.5	2.5
	3.4 g.	8	15.5	
Vitamin B ₆ hydrochloride	2.5 µg.	8	10.5	
	5.0 µg.	8	13	
2. National wheatmeal: 85% extraction	1.36 g.	8	14	4.2
	2.7 g.	8	18	
Vitamin B ₆ hydrochloride	2.5 µg.	8	9	
	5.0 µg.	8	13.5	
3. Wholemeal: 100% extraction	1.0 g.	8	14	c. 5.0
	2.0 g.	8	19	
Vitamin B ₆ hydrochloride	2.5 µg.	3	12	
	5.0 µg.	8	13.5	
	10 µg.	4	14.5	
4. Weatings	1.02 g.	4	20	13
	2.04 g.	4	27	
Vitamin B ₆ hydrochloride	2.5 µg.	4	9	
	5.0 µg.	4	14	
5. Germ I	200 mg.	4	12	13.6
	400 mg.	4	15	
Vitamin B ₆ hydrochloride	2.5 µg.	3	11	
	5.0 µg.	4	15	
6. Germ II	200 mg.	4	9.5	12
	400 mg.	4	15	
Vitamin B ₆ hydrochloride	2.5 µg.	4	10.5	
	5.0 µg.	4	14.5	

* These values $\times 0.82$ give the actual vitamin B₆ content of the materials.

All results given in the final columns of Tables 2, 3 and 4 are calculated by reference of the weight increase corresponding to the test doses of the unknown substances, to the straight line obtained by plotting the logarithm of the two standard doses to the growth response which they elicited. In no case was use made of a composite curve constructed for the responses to the standard, but in each separate experiment the results were calculated with reference to the line obtained from the results with the standard doses in that particular test.

Table 4. Summary of tests for filtrate factor content of flours and offals

Supplement	Daily dose	No. of rats	Av. weekly gain in wt. g.	Calc. content of filtrate factors mg.-equiv. of D.C.L. yeast per g.
1. White flour: 73% extraction	1.7 g. 2.7 g.	8 8	7.5 11	40-50
D.C.L. yeast	100 mg. 200 mg.	8 8	9 12	
2. National wheatmeal: 85% extraction	1.36 g. 2.7 g.	8 8	10.5 16.5	80
D.C.L. yeast	100 mg. 200 mg.	7 8	10 15.5	
3. Wholemeal: 100% extraction	1.02 g. 2.04 g.	11 11	11 16.4	110
D.C.L. yeast	100 mg. 200 mg.	10 11	9.7 15.5	
4. Weatings	1.02 g. 2.04 g.	4 4	15 21	190
D.C.L. yeast	100 mg. 200 mg.	4 4	8 15	
5. Germ I	250 mg. 500 mg.	4 4	6 10	200
D.C.L. yeast	100 mg. 200 mg.	4 4	11 17	
6. Germ II	250 mg. 500 mg.	4 4	5.5 9	250
D.C.L. yeast	100 mg. 200 mg.	4 4	8 14.5	

DISCUSSION

The recorded estimations of riboflavin in cereals include few carried out by biological methods with rats as the test animals. The earlier studies of Copping [1936] and Copping & Roscoe [1937] gave an approximation, but no quantitative values, since factors other than riboflavin were limiting the growth of the rats under the conditions of the tests. Most of the existing values for riboflavin have been obtained by fluorimetric or microbiological methods. Conner & Straub [1941*a*] have reported good agreement in their results obtained by a combined method for estimating vitamin B₁ and riboflavin simultaneously, and by a simple fluorimetric method for estimating riboflavin alone, as used by Hodson & Norris [1939]. Hodson & Norris compared the results obtained with their fluorimetric method for estimating riboflavin with those obtained by the microbiological method of Snell & Strong [1939] and also found good agreement.

All the values reported for wheat and wheat products by the above workers are considerably lower than those found in the present biological estima-

tions. Dr Davis and Mr Newland, of the National Institute for Research in Dairying, Shinfield, have tested the present series of flour samples by a modification of the microbiological method of Snell & Strong and have obtained results in remarkably good agreement with those obtained in the present biological tests (Table 5). Barton-Wright [1942], who used the microbiological method, reports a series of values in good agreement with those obtained by Dr Davis and Mr Newland and by the biological method. Andrews, Boyd & Terry [1942] described studies of riboflavin in cereals by the microbiological method, but their results were much lower than those obtained at Shinfield. It may be that the wheat blend used in milling the present series of flour samples was particularly rich in riboflavin, though this is unlikely. Conner & Straub [1941*a*] report values ranging from 0.89 to 1.91 $\mu\text{g.}$ per g. for 15 varieties of hard wheats and from 0.81 to 1.48 $\mu\text{g.}$ for 16 varieties of soft wheats. The generally higher results obtained by biological methods with rats, as compared with those obtained by fluorimetric or microbiological methods may be due to the difficulty in the extraction of riboflavin from cereal products which is necessary in the latter procedures, or, on the other hand, to the presence of other factors affecting the growth of rats in the biological method. The biological method for estimating riboflavin as originally described by El Sadr *et al.* [1940] has been applied to examining many materials in this laboratory and has been constantly checked by experiments with different doses of pure riboflavin. No evidence has been obtained of the need of the rat for B vitamins other than those present in the liver extract used. The good agreement between the results of biological and microbiological tests on the materials used in this study would also appear to confirm the reliability of the biological method.

Fraenkel & Blewett [1941] demonstrated differences in the riboflavin content of patent, straight run and wholemeal flours by a method in which the criterion adopted was the development of the flour beetle, *Tribolium confusum*. These workers have also demonstrated gradation of riboflavin content in the flour samples used in the present study but have not as yet been able to make a quantitative estimation by their method of riboflavin in such relatively poor sources.

For comparison the riboflavin values obtained for different wheat products by several observers are collected together in Table 5.

Values for the vitamin B₂ content of cereals have not yet been reported to any extent, though it was demonstrated qualitatively some years ago [Birch & György, 1935; Copping, 1936] that most cereals and their milling products were good sources of this nutrient. Swaminathan [1940*a, b*] developed a

Table 5

	Riboflavin $\mu\text{g./g.}$ estimated by					
Material	(1)	(2)	(3)	(4)	(5)	(6)
73 % ext. flour	0.85	0.83	0.75-1.25	0.62	0.4	—
85 % ext. flour	2.00	2.14	2.00-2.1	—	—	—
100 % ext. flour	2.30	2.78	2.65-3.7	1.00	—	0.89-2.03
Weatings (middlings)	4.00	5.90	—	3.80	1.3	—
Bran	4.80	—	5.8	2.80	1.0	3.17
Germ	7.8	—	9.7-12.85	—	—	3.78-5.56
and						
12						

- (1) Copping by biological method in present study.
- (2) Davis & Newland by microbiological method on same materials.
- (3) Barton-Wright [1942] by microbiological method.
- (4) Andrews *et al.* [1942] by microbiological method.
- (5) Hódson & Norris [1939] by fluorimetric method.
- (6) Conner & Straub [1941b] by fluorimetric method.

colorimetric method by means of which he found values of 3.0 and 7.6 $\mu\text{g.}$ per g. for white flour and whole wheat. These values are rather higher than those obtained by the biological method but of the same order. Recently, Conger & Elvehjem [1941] have described a biological method for estimating vitamin B_6 by the growth of rats which is essentially the same as that used in the present study. While these workers did not include cereals among the materials which they examined, so that no comparative values exist, it is worthy of note that the rates of weight increase of their control animals receiving vitamin B_6 hydrochloride in daily doses of 2, 5 and 10 $\mu\text{g.}$ were, respectively, 9, 14 and 19 g. weekly, and in the present tests for rats receiving 2.5 and 5 $\mu\text{g.}$ they were 10 and 14 g. weekly, respectively. This would seem to indicate that the methods in these two laboratories were directly comparable and that biological testing of vitamin B_6 can now be said to be on a sound basis.

The results of the tests for filtrate factors in flours and offals are important in so far as they give a comparison of the value of the different milling fractions and indicate the distribution of the filtrate factors through the grain. This appears to follow that of vitamin B_6 fairly closely, a considerable amount of both vitamins being found in the white flour, with a marked increase in the transition from 73 to 85% extraction and a less marked increase in the further advance to 100% extraction. A great increase in the amount in flour of 85% as compared with that in 73% extraction also held for riboflavin, but the amounts present in the flours and offals were relatively lower than the amounts of vitamin B_6 and filtrate factors. The germ and the weatings or middlings fraction of the grain proved very good sources of all three vitamins, and it is the inclusion of these

fractions in the 85% extraction national wheatmeal which raises its nutritive value with regard to all members of the vitamin B_2 complex so greatly above that of white flour.

SUMMARY

1. Estimations have been made by biological methods, with the weight increase of young rats as criterion, of the riboflavin, vitamin B_6 , and filtrate factors present in white flour, national wheatmeal and wholemeal, and in wheat germ, bran and middlings or weatings.

2. All the flours were poor sources of riboflavin, but there was a threefold increase with the rise of extraction from 73% for white flour to 85% for national wheatmeal. Wholemeal was only slightly superior to national wheatmeal in its riboflavin content. Wheat germ contained per g. more than three times as much riboflavin as wholemeal, while bran and weatings were about twice as rich in this factor as wholemeal.

3. The differences in vitamin B_6 content in white flour, national wheatmeal and wholemeal were less marked than those in riboflavin content, indicating a more even distribution of vitamin B_6 throughout the grain. Wheat germ and weatings contained per g. about three times as much vitamin B_6 as wholemeal.

4. The amounts of filtrate factors in the flours and offals showed a similar distribution to the amounts of vitamin B_6 .

5. The results are summarized in Tables 2-4, and in Table 5 riboflavin values for wheat products obtained by several workers are collected for comparison.

My thanks are due to the Medical Research Council for a part-time personal grant, to Miss A. S. Cole for her assistance during the latter part of this investigation, and to the Lister Institute Division of Nutrition, now kindly housed by Sir Charles Martin, for hospitality. Most particularly I have to thank the Research Association of British Flour Millers for their assistance in supplying the materials for study. I wish also to thank Glaxo Laboratories Ltd. for liver extracts, Roche Products Ltd. for supplies of pure riboflavin and vitamin B_1 and Merck and Co. Inc. of Rahway, N.J., for vitamin B_6 hydrochloride.

Throughout the investigations Dr H. Chick and Sq.-Ldr. T. F. Macrae have been most helpful with their advice and criticism. I acknowledge gratefully the co-operation of Dr Davis and Mr Newland, of the National Institute for Research in Dairying, Shinfield, and of Dr Fraenkel, of the Imperial College of Science and Technology, in making tests of the riboflavin content of my materials by other methods.

REFERENCES

- Andrews, J. S., Boyd, H. M. & Terry, D. E. [1942]. *Cereal Chem.* **19**, 55.
- Barton-Wright, E. [1942]. *Nature, Lond.*, **149**, 696.
- Birch, T. W. & György, P. [1935]. *Chem. Industr.* **54**, 507.
- Chick, H., Macrae, T. F. & Worden, A. N. [1940]. *Biochem. J.* **34**, 580.
- Conger, T. W. & Elvehjem, C. A. [1941]. *J. biol. Chem.* **138**, 555.
- Conner, R. T. & Straub, G. J. [1941a]. *Industr. Engng Chem. (Anal. ed.)*, **13**, 385.
- [1941b]. *Cereal Chem.* **18**, 671.
- Copping, A. M. [1936]. *Biochem. J.* **30**, 849.
- Copping, A. M. & Roscoe, M. H. [1937]. *Biochem. J.* **31**, 1879.
- Coward, K. H. [1938]. *Biological Standardization of the Vitamins*. London: Baillière, Tindall and Cox.
- El-Sadr, M. M., Hind, H. G., Macrae, T. F., Work, C. E., Lythgoe, B. & Todd, A. R. [1939]. *Nature, Lond.*, **144**, 73.
- El Sadr, M. M., Macrae, T. F. & Work, C. E. [1940]. *Biochem. J.* **34**, 601.
- Fraenkel, G. & Blewett, M. [1941]. *Nature, Lond.*, **147**, 716.
- Fridericia, L. S., Freudenthal, P., Gudjonsson, S., Johansen, G. & Schoubye, N. [1927]. *J. Hyg., Camb.*, **27**, 70.
- György, P. & Goldblatt, H. [1940]. *J. exp. Med.* **72**, 1.
- Hodson, A. Z. & Norris, L. G. [1939]. *J. biol. Chem.* **131**, 621.
- Lepkovsky, S. [1942]. *Nutr. Abstr. Rev.* **11**, 363.
- Lythgoe, B., Macrae, T. F., Stanley, R. H., Todd, A. R. & Work, C. E. [1940]. *Biochem. J.* **34**, 1335.
- Macrae, T. F., Todd, A. R., Lythgoe, B., Work, C. E., Hind, H. G. & El Sadr, M. M. [1939]. *Biochem. J.* **33**, 1681.
- Medical Research Council Accessory Food Factors Committee [1941]. *Lancet*, **1**, 703.
- Snell, E. E. & Strong, F. M. [1939]. *Industr. Engng Chem. (Anal. ed.)*, **11**, 346.
- Swaminathan, M. [1940a]. *Nature, Lond.*, **145**, 780.
- [1940b]. *Indian J. med. Res.* **28**, 427.

The Intravenous Glucose Tolerance Equation

By G. D. GREVILLE, *Biochemical Department, Runwell Hospital, near Wickford, Essex*

(Received 11 September 1942)

The intravenous glucose-tolerance test in which the blood-sugar is measured at intervals for an hour or two following a single intravenous injection of glucose has received steady but not intensive attention during the last 20 years. Recent important contributions include those of Crawford [1938], Ross [1938], Lozner, Peters, Taylor & Winkler [1941], Hamilton & Stein [1942], McKean, Myers & von der Heide [1935] and Tunbridge & Allibone [1940]. The two last-named papers contain reviews of the literature up to 1935 and 1938 respectively.

The lack of agreement among the various workers regarding procedure is remarkable. This applies not only to the dose of glucose given, but also to the assessment of the results. As the latter involves interpretation of the curve obtained for each test when the blood-sugar is plotted against the time after injection, a study of such curves seems most desirable. This paper concerns curves obtained in a series of glucose tolerance tests performed in this Hospital in 1940 in collaboration with Dr Derek Russell Davis; I undertook recently an examination of them. The following data are important for an appreciation of what follows; fuller practical details are given later in the paper.

In all, 35 tests were carried out on 13 adult psychotic patients. In 26 tests the dose of glucose was 15 g. per sq. m. body surface, in 8 it was 0.5 g. per kg. body weight, and in 1 test it was 27.6 g. [cf.

Tunbridge & Allibone, 1940]. The glucose was injected in 33½ % solution in water over a period of 2 min., and the time at which the blood specimens were taken was measured from the mid-time of the injection. 14 specimens were taken at intervals from 5 to 90 min. (both inclusive) after, and 4 specimens before the administration of glucose. The blood-sugar (in mg./100 ml.) will be denoted by y , the time (in min.) by t , and the graph obtained by plotting y against t will be called 'the curve'. Two of the curves obtained are shown in Figs. 1 and 3 (points represented by circles).

AN EQUATION FOR THE INTRAVENOUS GLUCOSE TOLERANCE CURVE

A simple relationship which might hold approximately is that the rate of fall of blood-sugar at any instant after the injection of glucose is proportional to the value of the blood-sugar at that time, i.e.

$$\frac{dy}{dt} = -Ky, \quad (i)$$

where K is a constant. This gives on integration

$$\log y = -Kt + \text{const.} \quad (ii)$$

Such a relationship has been found to hold approximately for the disappearance of foreign sugars from the blood [Fishberg, 1930], and also for the disappearance of calcium ion from the blood following