aspartic acid. This appears to be the first isolation of ornithine by acid hydrolysis of an intact protein or polypeptide.

3. The phenylalanine isolated had mainly the *d*-configuration; the other amino-acids were mainly *l*-.

4. 88–105 % of the N of tyrocidine has now been accounted for, but the variability of the analytical

figures has not permitted stoichiometric calculations. The implications of this variability, which exceeds the analytical errors, are discussed.

We are grateful to Dr R. J. Dubos and to Sir H. H. Dale, P.R.S., for making a supply of tyrothricin available to us.

We wish to thank the Director and Council of the Wool Industries Research Association and the International Wool Secretariat for permission to publish this work.

REFERENCES

- Bergmann, M. & Zervas, L. [1928]. *Biochem. Z.* **203**, 280.
 Christensen, B. E., West, E. S. & Dimick, K. P. [1941]. *J. biol. Chem.* **137**, 735.
 Christensen, H. N., Edwards, R. R. & Piersma, H. D. [1941]. *J. biol. Chem.* **141**, 187.
 Dubos, R. J. [1942]. *Ann. Rev. Biochem.* **11**, 659.
 — & Hotchkiss, R. D. [1942]. *Trans. Coll. Phys. Philad.* (4th ser.), **10**, 11.
 Dunn, M. S. [1925]. *J. Amer. chem. Soc.* **47**, 2564.
 Fearon, W. R. [1939]. *Biochem. J.* **33**, 902.
 Gordon, A. H., Martin, A. J. P. & Syngé, R. L. M. [1943a]. *Biochem. J.* **37**, 79.
 — — — [1943b]. *Biochem. J.* **37**, 86.
 — — — [1943c]. *Biochem. J.* **37**, 92.
 Gortner, R. A. & Norris, E. R. [1923]. *J. Amer. chem. Soc.* **45**, 550.
 Holm, G. E. & Gortner, R. A. [1920]. *J. Amer. chem. Soc.* **42**, 2378.
 Homer, A. [1915]. *J. biol. Chem.* **22**, 369.
 Hotchkiss, R. D. [1941]. *J. biol. Chem.* **141**, 171.
 — & Dubos, R. J. [1941]. *J. biol. Chem.* **141**, 155.
 Hunter, A. [1939]. *Biochem. J.* **33**, 27.
 Kossel, A. & Weiss, F. [1913]. *Hoppe-Seyl. Z.* **84**, 1.
 Lipmann, F., Hotchkiss, R. D. & Dubos, R. J. [1941]. *J. biol. Chem.* **141**, 163.
 Martin, A. J. P. & Syngé, R. L. M. [1941]. *Biochem. J.* **35**, 91.
 Onslow, H. [1921]. *Biochem. J.* **15**, 383.
 — [1924]. *Biochem. J.* **18**, 63.
 Van Slyke, D. D., Dillon, R. T., MacFadyen, D. A. & Hamilton, P. [1941]. *J. biol. Chem.* **141**, 627.
 Du Vigneaud, V. & Meyer, C. E. [1932]. *J. biol. Chem.* **98**, 295.
 Wada, M. [1933]. *Biochem. Z.* **257**, 1.
 Wood, J. K. [1914]. *J. chem. Soc.* **105**, 1988.

The Structure of the Galactose-phosphate Present in the Liver During Galactose Assimilation

By H. W. KOSTERLITZ, *From the Physiology Department, Marischal College, Aberdeen*

(Received 25 February 1943)

In this paper further support is provided for the view [Kosterlitz, 1937; 1938; 1939b] that in the galactose-phosphate present in the liver during galactose assimilation the phosphate group is attached to carbon atom 1. It has not yet been possible to separate this ester from the accompanying glycerophosphate, and from small quantities of glucose-1-phosphate and an unidentified ester. But the facts that reducing hexosemonophosphates can be removed by alkaline hydrolysis, and that Ba glycerophosphate is very resistant to acid hydrolysis and has a specific rotation of 0°, have made possible the approximate determination of the rate of acid hydrolysis of the galactose-phosphate and of its specific rotation. Since the ester is non-reducing, resistant to alkaline hydrolysis and very labile to acid hydrolysis, it is concluded that the first carbon atom is substituted. The velocity constant (*k*) of hydrolysis of the natural ester in 0.25*N* HCl at 25° is 0.91×10^{-3} , while that of synthetic galactose-1-phosphoric acid is 0.89×10^{-3} .

Natural Ba galactose-1-phosphate has $[\alpha]_{5461} + 113^\circ$ while the synthetic ester has $[\alpha]_{5461} + 109.5^\circ$. These values strongly suggest that the natural and synthetic esters are identical. It is of interest to note that Wolfrom & Pletcher [1941] and Wolfrom, Pletcher & Brown [1942] have succeeded in finally proving that the glucose-1-phosphate of Cori, Colowick & Cori [1937] has the structure of phosphoryl α -mono-*d*-glucopyranoside. Since the principle of the synthesis of galactose-1-phosphate is analogous to that devised by Cori *et al.* [1937] for glucose-1-phosphate, the findings of Wolfrom *et al.* [1942] give further support to the view expressed on other grounds [Kosterlitz, 1939a] that the synthetic galactose-1-phosphate is phosphoryl α -mono-*d*-galactopyranoside.

EXPERIMENTAL

The general methods were the same as those described in previous papers [Kosterlitz, 1937; 1939a]. The Ba salts of the phosphoric esters were obtained

from the livers of 38 rabbits which had been fed with galactose. The method of isolation has been described previously [Kosterlitz, 1937]. No nitrogenous substances were present in the ester mixture. The yield of Ba salts was 1.75 g. from 2.41 kg. liver.

Analysis of Ba salts of the ester mixture. The substance contained 6.87% total P and 1.88% acid-labile P (=27.4% of the total P). The initial reduction was 9.1% 'glucose'; 12 min. hydrolysis in 0.09 *N* H₂SO₄ at 100° increased the reduction by 8.9% 'glucose' or 11.7% 'galactose'. The fermentation analysis gave the following results: change after fermentation with *Saccharomyces ludwigii* (*S. ludwigii*) = +2.8% 'glucose'; change after fermentation with galactose-adapted *Saccharomyces cerevisiae* (*S. cerevisiae*) Froberg = -8.9% 'glucose'; residual reduction: 3.0% 'glucose'. The mixed Ba salts, therefore, contained 11.7% galactose as ester or 25.7% anhydrous Ba galactose-1-phosphate. Since Ba glycerophosphate was found to be more easily soluble in aqueous ethanol than Ba galactose-phosphate, a separation of the two compounds was attempted by fractional precipitation with aqueous ethanol. This attempt, however, was not very successful [Kosterlitz, 1938].

Acid and alkaline hydrolysis of the ester mixture. A sample of a composition slightly different from that described in the preceding paragraph was used for this particular experiment. 17.96 mg. Ba salt were dissolved in 15 ml. H₂O containing 0.87 ml. *N* H₂SO₄ and the precipitated BaSO₄ removed by centrifuging. The initial reduction was 7.76% 'glucose' and the total P 7.02%. The acid-labile P, determined by 3 min. hydrolysis in 0.09 *N* H₂SO₄ at 100°, was 1.84%; the alkali-labile P, determined by hydrolysis in 0.2 *N* NaOH at 100°, was 1.29% after 20 min. and 1.33% after 30 min. hydrolysis. The acid-labile P constituted

and the precipitate successively extracted with 20, 20, 10, 10 and 15 ml. H₂O. The residue was dissolved in a minimum of 0.25 *N* HCl, the solution brought back to pH 8.4 by means of Ba(OH)₂, and then centrifuged. The supernatant fluid was added to the aqueous extracts, followed by ethanol to give a concentration of 60%. The extraction of the centrifuged precipitate was repeated twice and the final precipitate dried *in vacuo* over P₂O₅. Yield: 293 mg. This material together with 23 mg. from a previous experiment (containing 3.0 mg. galactose as phosphate) was dissolved in 26 ml. H₂O, the insoluble material separated and washed by centrifuging. To the clear solution (32 ml.) an equal quantity of ethanol was added; after standing at 0° for 2 hr., the Ba salts were centrifuged off, washed with 50 and 80% ethanol and dried *in vacuo* over CaCl₂. Yield: 295 mg. The slightly turbid solution of this product in 10 ml. H₂O was centrifuged, and an equal quantity of ethanol was added to the supernatant fluid. After standing at 0° for 2 hr., the precipitate was centrifuged off and then successively extracted with 15, 10 and 15 ml. 10% ethanol. Ethanol was added to the supernatant fluid to give a concentration of 50%. After 2 hr. standing at 0°, the precipitate was centrifuged off, washed with 80% ethanol and dried *in vacuo* over CaCl₂. The Ba salts were now completely soluble in H₂O. Yield: 228 mg. containing 16.3% =37.2 mg. galactose as phosphate. The crude material contained 62.5 mg. (+3.0 mg.) galactose. Yield: 57%.

Analysis of the Ba salts of the ester mixture freed from reducing phosphates. The substance contained 7.46% total P and 2.77% acid-labile P (=37.2% of total P). $[\alpha]_{5461}^{20} = +34.4$ ($c=1.476$; $l=2$). The initial reduction was negligible (0.45% 'glucose') and 3 min. hydrolysis in 0.09 *N* H₂SO₄ at 100° increased the reduction by 12.4% 'glucose' or 16.3% 'galactose'. The fermentation analysis (Table 1) showed that most of this increase was due to galactose:

Table 1. Analysis of saccharide phosphate mixture by fermentation

	Unhydrolysed esters	Hydrolysed esters
Initial reduction	0.38% 'glucose'	12.9% 'glucose'
Change after fermentation with <i>S. ludwigii</i>	+10.9% 'glucose'	-1.53% 'glucose'
Change after fermentation with galactose-adapted <i>S. cerevisiae</i> Froberg	-10.5% 'glucose'	-10.2% 'glucose'
Residual reduction	=13.8% 'galactose'	=13.3% 'galactose'
	0.72% 'glucose'	1.15% 'glucose'

* This increase in reduction was caused by phosphatatic hydrolysis of galactose-1-phosphate [Kosterlitz, 1937; 1939a].

26.2% and the alkali-labile P 18.7% of the total P so that 55% of the total P was resistant to hydrolysis. Galactose-1-phosphate and glucose-1-phosphate were the main sources of the acid-labile P, and reducing hexosephosphates and possibly pentosephosphates those of the alkali-labile P.

Removal of reducing phosphate esters by hydrolysis in 0.2 *N* NaOH. 534 mg., containing 11.7% galactose (=62.5 mg.) as phosphate, were dissolved in 47.4 ml. H₂O and 2.58 ml. *N* H₂SO₄. The precipitated BaSO₄ was removed by centrifuging and washed with 5 ml. H₂O. To the total of 55 ml. were added 2.58+13.75 ml. *N* NaOH (final concentration=0.193 *N*). The hydrolysis was allowed to proceed at 100° for 30 min., during which time the solution turned deeply yellow, due to the action of the alkali on the reducing sugars. After cooling and addition of 16.33 ml. *N* HCl, Ba(OH)₂ was added until the pH was 8.4. The precipitated Ba phosphate was centrifuged off and washed. To the supernatant fluid, ethanol was added to give a concentration of 60%. After 2 hr. standing at 0°, the material was centrifuged, the supernatant fluid decanted,

From these results, the following conclusions may be drawn: (1) alkaline hydrolysis practically abolished the initial reduction which was caused by reducing hexosephosphates and possibly pentosephosphates; (2) the greatest part but not all of the reducing substances produced by acid hydrolysis consisted of galactose (13.5%). Besides galactose, 1.53% glucose was liberated, and 0.93% expressed as glucose of a non-fermentable reducing substance. Thus, the mixture of Ba salts contained as anhydrous salts: 29.6% Ba galactose-1-phosphate, 3.4% Ba glucose-1-phosphate, and 2.0% unidentified acid-labile ester (calculated on basis of hexosephosphate). This made a total of 35% anhydrous Ba salt or 39.8% Ba salt containing 3H₂O.

These results show up a fact which so far has escaped attention, namely, that during galactose assimilation, glucose-1-phosphate is present in the

liver besides galactose-1-phosphate. In experiments on rats, Kosterlitz & Ritchie [1943*b*] could demonstrate an increase in glucose-1-phosphate in the livers after administration of galactose as well as of glucose. Nothing can so far be said about the unknown acid-labile ester.

Hydrolysis in 0.09N H₂SO₄ at 100° of the ester mixture freed from reducing phosphates

(Concentration of acid-labile P = 0.525 m.-molar)

Time min.	Increase in reduction mg. 'glucose'/ mg. Ba salt	Increase in P mg./mg. Ba salt
2	0.125	0.0272
3	0.125	0.0277
10	0.126	0.0277
180	0.130	0.0318

The liberation of reducing substances and of P was practically complete after 2-3 min. hydrolysis; under the conditions of this experiment, galactose-1-phosphate is completely hydrolysed in 2 min. and glucose-1-phosphate in 5 min. [Kosterlitz & Ritchie, 1943*a*]. Between 10 and 180 min. there was an insignificant increase in reduction, while there was still a liberation of inorganic P; this would suggest the presence of a non-reducing difficultly hydrolysable ester (or esters) such as glycerophosphoric acid.

Isolation of the difficultly hydrolysable esters by acid hydrolysis of the ester mixture freed from reducing phosphates. 96.5 mg. Ba salts were dissolved in 29.5 ml. H₂O and 3.49 ml. N H₂SO₄ (=free esters in 0.09N H₂SO₄). This solution was hydrolysed at 100° for 10 min. After addition of Ba(OH)₂ (pH 8.4), the precipitate was centrifuged off and 2 vol. of ethanol added to the supernatant fluid. The precipitate was dissolved in the minimum amount of 0.25N HCl, the solution warmed in order to precipitate colloidal BaSO₄, and the BaSO₄ centrifuged off. Ba(OH)₂ was added to the supernatant fluid (pH 8.4), the precipitate of inorganic P centrifuged off, and 2 vol. of ethanol added. The BaSO₄ was warmed with 0.25N HCl and the HCl extract treated as described above. The combined aqueous alcoholic mixtures were allowed to stand for 2 hr. at 0°. After centrifuging, the precipitate was exhaustively extracted with H₂O and the centrifuged extracts treated with 2 vol. ethanol. After standing at 0° for 2 hr., the precipitate was centrifuged and the process of extraction repeated once with H₂O and a second time with 10% aqueous ethanol. The final precipitate was dried *in vacuo* over CaCl₂. Yield: 39 mg. Expected: 60% of 96.5 mg. = 58 mg.; yield: 67%.

Analysis of the Ba salt of the difficultly hydrolysable ester. Total P = 7.48%; inorganic P = 0; acid-labile P = 0. Initial reduction = 0.58% 'glucose'; acid-labile reduction = 0.3% 'glucose'. $[\alpha]_{5461}^{17.5} = -3.4^\circ$ ($c = 0.778$; $l = 2$).

Fermentation analysis: initial reduction = 0.61% 'glucose'; change after fermentation with *S. ludwigii* = +0.38% 'glucose'; change after fermentation with galactose-adapted *S. cerevisiae* = -0.57% 'glucose'; residual reduction = 0.42%.

This ester was practically non-reducing; it was not further analysed but was probably composed mainly of Ba α -glycerophosphate (+3H₂O = 8.6% P) which is present in liver in considerable quantities [Fiske & Subbarow, 1929].

However, the low P value and the slight laevo-rotation indicated that other unknown esters were also present.

Calculation of the specific rotation of natural Ba galactose-1-phosphate. The Ba salts of the ester mixture containing 29.6% anhydrous galactose-1-phosphate had $[\alpha]_{5461} + 34.4^\circ$. By deducting -2.0° for the 60% believed to be present of the Ba salt of the difficultly hydrolysable ester, and $+3.0^\circ$ for 3.4% anhydrous Ba glucose-1-phosphate, a value of $+33.4^\circ$ was obtained, which corresponded to 29.6% anhydrous Ba galactose-1-phosphate. Thus, natural anhydrous Ba galactose-1-phosphate has approximately $[\alpha]_{5461} = +113^\circ$, while for the anhydrous Ba salt of the synthetic ester $[\alpha]_{5461} = +109.5^\circ$. No allowance could be made for the 2% unidentified acid-labile ester.

Hydrolysis of the acid-labile fraction of the ester mixture, freed from reducing phosphates in 0.25N HCl at 25°. 28.19 mg. Ba salt were dissolved in a total of 10 ml. with addition of 1 drop sat. K₂SO₄ solution and 1.365 ml. of 0.1N H₂SO₄, in order to liberate the free acid. After centrifuging off the BaSO₄, 9 ml. 0.5N HCl were added to 9 ml. of the supernatant fluid. The concentration of acid-labile P was 1.17 m.-molar.

Hydrolysis in 0.25N HCl at 25°

Time min.	Increase in reduction (% of total acid-labile reduction)	$k \times 10^3$	Inorganic P formed (% of total acid- labile P)	$k \times 10^3$
180	29.6	0.847	29.8	0.854
255	38.3	0.764	38.6	0.775
330	45.8	0.751	46.0	0.744
Average		0.787		0.791

Calculation of the k of hydrolysis of natural galactose-1-phosphoric acid by 0.25N HCl at 25°. The acid-labile fraction of the ester mixture contained 84.6% galactose-1-phosphoric acid, 9.7% glucose-1-phosphoric acid, and 5.7% unidentified ester. The k of the acid-labile fraction was 0.79×10^{-3} . The k of glucose-1-phosphoric acid (0.2×10^{-3}) was calculated from the known k values of the glucose-1-phosphoric and galactose-1-phosphoric acids at 25°, 37° and 50° [Cori *et al.*, 1937; Kosterlitz, 1939*a*; Kosterlitz & Ritchie, 1943*a*]. No allowance could be made for the 5.7% unidentified ester. By deducting 0.02×10^{-3} for the 9.7% glucose-1-phosphate present in the acid-labile fractions a value of 0.77×10^{-3} was obtained, which corresponded to 84.6% galactose-1-phosphoric acid. Thus, the k of natural galactose-1-phosphoric acid is approximately 0.91×10^{-3} , while the k of the synthetic ester is 0.89×10^{-3} .

SUMMARY

1. The mixture of phosphoric esters isolated from the livers of rabbits assimilating galactose was subjected to alkaline hydrolysis in order to remove reducing hexosephosphates.

2. The product obtained in this way contained 29.6% anhydrous Ba galactose-1-phosphate, 3.4% anhydrous Ba glucose-1-phosphate and 2% Ba salt of unidentified acid-labile ester; the remainder consisted of a difficultly hydrolysable non-reducing ester. The Ba salts of the ester mixture freed from

reducing phosphates had $[\alpha]_{5461} + 34.4^\circ$ and the Ba salt of the difficultly hydrolysable ester $[\alpha]_{5461} - 3.4^\circ$. From these values it was calculated that the natural anhydrous Ba galactose-1-phosphate has $[\alpha]_{5461} + 113^\circ$, while the synthetic product has $[\alpha]_{5461} + 109.5^\circ$.

3. The velocity constant (k) of hydrolysis of the acid-labile fraction of the ester mixture, freed from reducing phosphates, in $0.25N$ HCl at 25° was 0.79×10^{-3} . From this value the k of the natural

galactose-1-phosphoric acid was calculated to be 0.91×10^{-3} , while the k of the synthetic product is 0.89×10^{-3} .

4. These results strongly suggest that the natural and synthetic galactose-1-phosphoric acid are identical. The ester probably is an α -pyranoside as is the natural glucose-1-phosphoric acid.

An expenses grant by the Medical Research Council which partly defrayed the cost of this investigation is gratefully acknowledged.

REFERENCES

- Cori, G. T., Colowick, S. P. & Cori, C. F. [1937]. *J. biol. Chem.* **121**, 465.
 Fiske, C. H. & Subbarow, Y. [1929]. *Science*, **70**, 381.
 Kosterlitz, H. W. [1937]. *Biochem. J.* **31**, 2217.
 — [1938]. *J. Physiol.* **93**, 34P.
 — [1939a]. *Biochem. J.* **33**, 1087.
 Kosterlitz, H. W. [1939b]. *Nature, Lond.*, **144**, 635.
 — & Ritchie, C. M. [1943a]. *Biochem. J.* **37**, 181.
 — [1943b]. Unpublished results.
 Wolfson, M. L. & Pletcher, D. E. [1941]. *J. Amer. chem. Soc.* **63**, 1050.
 — & Brown, A. E. [1942]. *J. Amer. chem. Soc.* **64**, 23.

The Apparent Dissociation Constants of Galactose-1-Phosphoric Acid

By H. W. KOSTERLITZ, *From the Physiology Department, Marischal College, Aberdeen*

(Received 25 February 1943)

Meyerhof & Suranyi [1926], Meyerhof & Lohmann [1927], and Cori, Colowick & Cori [1937] found that the dissociation constants (pK' values) of hexose-diphosphoric, hexose-6-monophosphoric, fructose-6-phosphoric, and glucose-1-phosphoric acids were smaller than those of orthophosphoric acid. In the present paper α (?) galactose-1-phosphoric acid is shown to have dissociation constants similar to those of the other hexosemonophosphates. The knowledge of the acidic properties of galactose-1-phosphoric acid is necessary for the assessment of the changes in pH which may be brought about by the phosphorylation of galactose.

EXPERIMENTAL

The pH values were measured by means of a Cambridge pH meter with dip glass electrode (portable pattern). A buffer consisting of equal parts of Na acetate and acetic acid ($0.1N$) and having pH 4.62 was used for standardizing the glass electrode. The galactose-1-phosphoric acid was prepared by treating the Ba salt (from the K_2 salt [Kosterlitz, 1939]) with the exact quantity of H_2SO_4 . The concentration of galactose-1-phosphoric acid in the final solution was determined by estimation of phosphorus. 10 ml. of $0.1015M$ (Exp. 1) and $0.1038M$ (Exp. 2) galactose-1-phosphoric acid were titrated with CO_2 -free $0.0999N$ NaOH (temp. = $18-20^\circ$). For comparison, 10 ml. of $0.1001M$ H_3PO_4 were

similarly titrated. The results were plotted after correction for the deviations from 0.1 molarity

Fig. 1. 10 ml. $0.1M$ orthophosphoric acid $\bullet-\bullet$ 10 ml. $0.1M$ galactose-1-phosphoric acid: Exp. 1 $---\times$; Exp. 2 $\times-\times$.

(Fig. 1). The dissociation constants were calculated from 3 to 4 values each; pK'_1 by means of

$$K'_a = \frac{[H]^+ (B + [H]^+)}{C - (B + [H]^+)}$$