Obituary Notice

ERNEST WALKER (1900-1942)

During August 1942, Dr E. Walker's sudden death came as a considerable shock to all who knew him and occurred while he was still a member of the Committee of the Biochemical Society.

Born in 1900, he was the son of a Liverpool corn merchant, but was left an orphan early; both parents died when he was nine years old. He was educated at Pinner School and the Liverpool Collegiate School, and left at the age of seventeen to join the Army (The Artists' Rifles). Returning from the war, in 1919 he studied chemistry at Liverpool University under the inspiration of Prof. E. C. C. Baly.

In 1920, he came to the Biochemical Laboratory, Cambridge, to work as assistant to Dr R. A. Peters upon problems of chemical constitution and physiological action in relation to Chemical Defence Research. This work was continued after 1923 in the Department of Biochemistry, Oxford, whence he accompanied Dr R. A. Peters and became a member of St John's College. This research formed the basis of his thesis for the M.Sc. Liverpool, and the D.Phil. (Oxford) degree which he took while temporarily employed at the Government Experimental Station at Porton in 1927. During this period, a fundamental contribution on the physical chemistry of mustard gas and related compounds in relation to vesication was published (*Biochem. J.* 1923, 17, 260, with R. A. Peters). He also developed earlier work by McClelland and Peters upon the toxicity of ter- and pentavalent arsenical compounds to Protozoa (*Biochem. J.* 1928, 22, 292). Some of the work of this period was inspired by that of Sir F. G. Hopkins upon sulphhydryl compounds, particularly his research work upon the fixed —SH group in muscle tissue (with M. Dixon). In especial Walker worked out a nitroprusside test for —S—S— groups, using cyanide, and located the extent of the layer in skin showing the nitroprusside reaction. It was this research which led to the important discovery (independently and about the same time as Voegtlin) of the effect of an —SH compound (monothioethylene glycol) in reversing the action of diphenyl arsenious oxide upon ciliates.

After a temporary interlude at the Experimental Station, Porton, he returned to Oxford in 1928 and proposed at first to qualify in medicine. Subsequently, however, he gave up this plan and settled down as a Demonstrator, occupying the post of Departmental and then University Demonstrator

until the end. He also undertook the secretarial work of the Department; here he devoted himself to promoting successfully its efficiency by his continual and unselfish attention to its interests as a whole, and skill in administrative detail. The students owe much to his attention to the smooth running of the classes, as also to his constant interest in teaching fundamentals. The development of the courses for the Supplementary Subject in Biochemistry for the Final Honour School fell to him, and he was responsible for teaching the Microbiological Biochemistry.

In the period from 1928 to 1940, Dr E. Walker published several papers, on a variety of subjects, several with junior pupils and colleagues, though much of his constant labours did not see the light. Becoming suspicious of the Pettenkofer reaction as an indication of the presence of bile salts in normal blood, he found that this was due to cholesterol oleate, which he isolated (*Biochem. J.* 1930, 24, 1489). With A. R. Armstrong (*Biochem. J.* 1932, 26, 143) he identified the substance in pregnancy urine giving Knoop's test, as histidine, a piece of work which seems to have been overlooked by some recent workers. Returning to sulphhydryl compounds, he studied (with A. Todrick) the effect of allyl isothiocyanate upon cysteine, isolating the compound formed; at the same time and independently of Anson and Mirsky they developed a method (differing from their own) for estimating the sulphhydryl groups in protein, which was applied to ovalbumin and albumin, also myosin (*Biochem. J.* 1937, 31, 297). In his later work he turned to microbiological biochemistry. With F. L. Warren, he worked upon the decomposition of cellulose by *Cytophaga I* (*Biochem. J.* 1938, 32, 31). Chemical advances in a difficult field were here made with this organism, which has so unique an appetite for cellulose; characteristic was the realization of the biological importance which may be played by oxycellular mucilage in the humus of the soil. His final publication with W. H. Maguigan (*Biochem. J.* 1940, 34, 804) was concerned with the sterol metabolism of yeast; this formed part of a more ambitious programme, some of which has not yet been published, and broke new ground of importance. Facts were established about the time of origin of sterol and about precursors. During this war, Walker was engaged upon problems in connexion with the disposal of sewage; but much of his time

latterly was occupied in administrative problems and in the early development of the Oxford Nutrition Survey, which owes much to his wise guidance. Appraising the work as a whole, we realize the careful chemical interest in completion, combined with a wide knowledge and appreciation of biological applications; he had learnt fully the necessary physiology and bacteriology required. It indeed seems that his studies on micro-organisms were only at an initial stage.

The Secretary has told me that as a member of the Committee of the Society, Walker displayed a quiet ability combined with a sense of humour that proved to be invaluable in finding a way through the complicated results of a discussion on procedure or the like. His complete familiarity with the rules

of the Society and of syntax alike could on occasions be a little disconcerting but extremely helpful.

Quiet and modest in manner, unfailingly honest in his judgements and dealings with men, he proved a wise, independent-minded, courageous and respected counsellor in the laboratory. He had the capacity for carrying through administrative detail without appearing flurried or perturbed; in part this was due to an unfailing sense of humour which never deserted him, and which so often proved disconcerting. One of his many hobbies was chess playing.

In 1932 he married Dr Vera Reader, whose work upon nutritional problems is well known to biochemists, and is survived by his wife and two children.

R. A. PETERS

The Enzymic Production of Levan

By S. HESTRIN, S. AVINERI-SHAPIO AND M. ASCHNER, *Chemistry Department, Cancer Laboratories, and Department of Hygiene and Bacteriology, The Hebrew University, Jerusalem*

(Received 10 April 1943)

This paper describes the preparation and study of cell-free levan-synthesizing enzyme systems (levan-sucrase) from two bacterial species [cf. Aschner, Avineri-Shapiro & Hestrin, 1942]. A particularly convenient source of synthesizing enzyme has been found in a newly isolated non-sporulating bacterium to be referred to as *Aerobacter levanicum*. Recently Hehre has achieved the cell-free synthesis of dextran by the action of an enzyme isolated from *Leuconostoc mesenteroides* [Hehre, 1941; Hehre & Sugg, 1942; Stacey, 1942]. Levan- and dextran-synthesizing enzymes are thus now available in sterile form. The two principal polysaccharides which bacteria form specifically from sucrose may henceforth be included, together with glycogen and starch, in the category of macromolecular materials whose production can be effected *in vitro* at will.

Nomenclature

Beijerinck [1910, 1912] coined the word *viscosaccharase* in generic designation of enzyme systems which mediate polysaccharide-gum production from sucrose. We propose to designate individual enzymes within this group in each case in accordance with the polysaccharide whose production is catalysed. Thus for levan-synthesizing enzyme: *levansucrase*; for dextran-synthesizing enzyme: *dextran-sucrase*. Since the *viscosucrases* are very different in specificity from phosphorylase, the enzyme whose activity mediates the production of starch or glycogen, it seems correct to retain for the dextran- and levan-synthesizing systems the nomenclature and classification proposed by Beijerinck. It should not be overlooked, however, that the

over-all processes of levan and dextran production fit ill in any recognized class of carbohydrate-mediated reactions. The assignation of the dextran- and levan-synthesizing enzyme group to the *sucrases* can therefore at best be only tentative.

The enzymes of levan *hydrolysis* are, as will be shown elsewhere, distinct from *levansucrase*. They belong by definition, together with *inulase*, to the class of *polyfructosidases*. They may conveniently be referred to, therefore, as *levanases*.

EXPERIMENTAL

Bacterial strains used. Three bacterial strains were selected for investigation after preliminary trials with a sucrose-agar medium had indicated their excellence as levan formers. The first was *Bacillus subtilis*, Marburg strain, the levan of which has been the subject of extensive chemical study [Hibbert & Brauns, 1931]. This species is typical of a large group of sporulating bacteria which induce levan deposition in sucrose-agar medium at a distance from the actual growth site of the colony. The second organism selected was an aerobic gas-producing spore former received from the National Collection of Type Cultures under the name *B. polymyxa* (Prazmowski) Migula [*B. asterosporus* (Meyer)] N.C.T.C. no. 4744. It differs morphologically in the mode of its levan production from organisms of the *B. subtilis* type, depositing levan in sucrose medium at the cell-medium interface in the form of a giant capsule. The third test organism was a lactose-fermenting non-sporulating bacterium isolated from a fruit-tree root in Jerusalem. Like *B. polymyxa* it produced levan at the cell-medium interface only. The organism is actively motile, possessing peritrichous flagella. It ferments glucose, fructose, lactose, sucrose, raffinose and mannitol with the production of acid and gas, but ferments glycerol, inulin and levan with pro-