compound such as betaine or choline [Challenger & Higginbottom, 1935; Challenger, 1942]. The formation of tetramethyl putrescine in *Hyoscyamus muticus* could be explained by assuming either that dimethylamine reacts with succindialdehyde, or that glyoxylic acid (or formaldehyde) brings about direct methylation of putrescine.

SUMMARY

1. An enzyme capable of oxidizing putrescine with the formation of ammonia and an aldehyde

has been shown to exist in the tissues of *Atropa belladonna*.

2. The enzyme is located chiefly in the root system, but it also occurs in etiolated shoots.

3. Putrescine has been isolated in small amounts from leaves and upper stems of *Atropa belladonna* and *Datura stramonium*.

4. The possible role of putrescine in the synthesis of hyoscyamine and related alkaloids is discussed.

REFERENCES

- Best, C. H. & McHenry, E. W. [1930]. *J. Physiol.* **70**, 349.
 Challenger, F. [1942]. *Chem. Ind.* **61**, 397, 413, 456.
 — & Higginbottom, C. [1935]. *Biochem. J.* **29**, 1757.
 Ciamician, G. & Ravenna, C. [1911]. *R.C. Accad. Lincei*, **20**, 614. (Quoted from *Chem. Abstr.* 1911, **5**, 3466.)
 Cromwell, B. T. [1943]. *Biochem. J.* **37**, 717.
 Gale, E. F. [1940]. *Biochem. J.* **34**, 392.
 — [1942]. *Biochem. J.* **36**, 64.
 Greenish, H. G. [1933]. *A Text-Book of Pharmacognosy*, 6th ed. London: Churchill.
 Henry, T. A. [1924]. *The Plant Alkaloids*, 2nd ed. London: Churchill.
 Robinson, R. [1917]. *J. chem. Soc.* **111**, pp. 762, 876.
 — [1936]. *J. chem. Soc.* p. 1079.
 Schöpf, C. & Lehmann, G. [1935]. *Liebigs Ann.* **518**, 1.
 Zeller, E. A. [1938]. *Helv. chim. Acta*, **21**, 1645.
 — [1940]. *Helv. chim. Acta*, **23**, 1418, 1502.
 — Birkhauser, H., Mislin, H. & Wenk, M. [1939]. *Helv. chim. Acta*, **22**, 1381.

Studies in the Biochemistry of Micro-organisms

72. GENTISYL ALCOHOL (2:5-DIHYDROXYBENZYL ALCOHOL), A METABOLIC PRODUCT OF *PENICILLIUM PATULUM* BAINIER

By J. H. BIRKINSHAW, A. BRACKEN AND H. RAISTRICK, *Division of Biochemistry, London School of Hygiene and Tropical Medicine, University of London*

(Received 17 December 1943)

Workers in this laboratory [Birkinshaw, Bracken, Michael & Raistrick, 1943] have recently described the isolation of patulin, $C_7H_8O_4$, believed to be anhydro-3-hydroxymethylene-tetrahydro- γ -pyrone-2-carboxylic acid (I), from cultures of *Penicillium patulum* Bainier grown on Raulin-Thom medium. Patulin is a strong anti-bacterial substance. It completely inhibits the growth of a large series of Gram-positive and Gram-negative bacteria at concentrations of 1:33,000 to 1:100,000. It is also a powerful anti-fungal agent, since it completely inhibits, at a concentration of 1:400,000, the growth of a number of species of *Pythium* which are plant pathogens [Anslow, Raistrick & Smith, 1943].

Birkinshaw *et al.* [1943] recorded that the ethereal mother liquors, from which patulin had been isolated, still contained other metabolic products of *P. patulum*. One of these products, which forms the subject of the present communication, has now been isolated, in yields of 0.6 g./l. of culture filtrate, and identified as *gentisyl alcohol* (2:5-dihydroxy-

benzyl alcohol) (II). Gentisyl alcohol has only slight anti-bacterial activity against *Staphylococcus aureus*.

It is a curious fact that, so far as we can ascertain, a compound of this simple chemical constitution has not been described previously, although the corresponding aldehyde, gentisic aldehyde, and the corresponding acid, gentisic acid, which is also formed by *P. patulum*, have been known for a long time.

The chemical constitution of the new metabolic product was established as follows. It has the empirical formula, $C_7H_8O_3$. On long continued reduction with Pd-norite- H_2 it is converted into toluquinol. Oxidation of the dimethyl ether of the metabolic product with $KMnO_4$ gives 2:5-dimethoxybenzoic acid which on demethylation affords gentisic acid (2:5-dihydroxybenzoic acid). Finally, gentisyl alcohol was synthesized by the partial reduction of gentisic aldehyde, the initial quick reduction being stopped after the absorption of 1 mol. of hydrogen. Synthetic gentisyl alcohol and the new mould metabolic product have the same m.p. 100° , and a mixture of the two substances melts at the same temperature.

EXPERIMENTAL

Isolation of the new metabolic product. Cultures of *Penicillium patulum* Bainier, strain Ad 77, on Raulin-Thom solution, were prepared in the usual way. After 14 days' incubation at 24° the bulked culture filtrate from 52 flasks had the following characteristics: total vol. 16-14 l.; residual glucose by polarimeter, 0.74%; pH, 5.0; anti-bacterial activity against *Staphylococcus aureus*, complete inhibition at a dilution of 1:160, partial inhibition at 1:320. The bulked filtrate was evaporated *in vacuo* to 800 ml. and extracted three times with 800 ml. each time of ethyl acetate. The extract was dried over Na_2SO_4 and the solvent was removed *in vacuo*. The residual oil (58 g.) which was beginning to crystallize was dissolved in ether and passed through a column of alumina as described by Birkinshaw *et al.* [1943].¹ Two crops of patulin, wt. 19.42 and 0.88 g., were isolated.

The ethereal mother liquors remaining after separation of the patulin were freed from solvent and the residual oil was dissolved in ether (150 ml.) and hot chloroform (1 l.) was added. The crude metabolic product separated in clusters of needles containing a little oil. The product was collected after 4 days and washed with a little ether. Wt. 6.98 g., m.p. $98-99.5^\circ$. The mother liquor, after standing in an open flask for a considerable time, deposited a second crop of needles mixed with some oil. Wt. 2.90 g., m.p. $84-96^\circ$.

Purification of the crude metabolic product presented some difficulty owing to the tendency of the substance to separate as an oil. It also appears to be somewhat sensitive to heat. No completely satisfactory solvent was found and the method finally adopted was as follows: A portion (4 g.) of the first crop was finely powdered and added to 1 l. of boiling chloroform. The mixture was refluxed for 5 min. and filtered. On cooling, 1.10 g. of almost colourless needles were obtained; m.p. 100° after sintering at 98° . The undissolved portion, 1.97 g., was reworked in subsequent crystallizations.

General properties of the metabolic product. Colourless needles, m.p. 100° , which sublime in a high vacuum at a bath temperature of $70-80^\circ$, though not as readily as might be expected of a substituted quinol. (Found: C, 59.87, 60.01; H, 5.89, 6.06; N, nil; OCH_3 , nil. $C_7H_8O_3$ requires C, 60.01; H, 5.75%. 2.53 active H atoms in pyridine, 0.41 atoms in anisole. Kuhn-Roth method gives 0.112 mol. acetic acid.)

It is readily soluble in water, ethanol and ether, slightly soluble in benzene and chloroform and insoluble in light petroleum. An aqueous solution has the following properties: neutral to litmus; instantly decolorizes $KMnO_4$; reduces $AgNO_3$ in the cold with mirror formation; gives an evanescent blue colour with aqueous $FeCl_3$, changing almost immediately to yellow; reacts with aqueous iodine with absorption of two atoms of iodine which is slowly lost on titration with thiosulphate; does not react with hydroxylamine even after 24 hr. With 2N NaOH it gives an immediate yellow colour rapidly darkening to strong reddish brown. When heated with 2N H_2SO_4 , in which it is readily soluble in the cold, it gives in a few seconds a white amorphous precipitate.

It gives nearly complete inhibition of the growth of *Staphylococcus aureus* at a concentration of 1:500, partial inhibition at 1:1000 and 1:2000 and no inhibition at 1:4000.

Catalytic reduction of the metabolic product. Formation of toluquinol. A solution of the metabolic product (0.5 g.) in ethanol (10 ml.) was hydrogenated at ordinary pressure with a palladium-norite catalyst prepared from 0.2 g. of $PdCl_2$ and 0.5 g. of norite. Hydrogenation, which was slow throughout, was discontinued after $8\frac{1}{2}$ hr. (Uptake of hydrogen, 57.8 ml. (corr.); theor. for uptake of 2H per mol., 80 ml.) After filtering from catalyst the ethanol was removed *in vacuo*. A mass (0.48 g.) of crystals containing some oil remained. This material sublimed readily in a water-pump vacuum at a bath temperature of $110-115^\circ$. The sublimate formed large colourless blades, m.p. 127° , setting on cooling and remelting at 127° . Resublimed in a high vacuum the sublimate melted at 128° , alone or in admixture with authentic, sublimed toluquinol, m.p. 128° . (Found: C, 67.39; H, 6.40. Calc. for $C_7H_8O_2$: C, 67.75; H, 6.50%.)

Methylation of the metabolic product. A solution of the metabolic product (2.0 g.) in dry ether (25 ml.) was treated with an excess of ethereal diazomethane for 3 days. On removal of the solvent a reddish oil (2.35 g.) remained, which, on distillation *in vacuo*, gave 1.46 g. of a fairly mobile yellow oil, b.p. $140^\circ/3.3$ mm. (Found: OCH_3 , 34.1. $C_9H_{12}O_3$ requires, for $2OCH_3$, 36.9%.)

Oxidation of crude dimethoxy derivative. Formation of 2:5-dimethoxybenzoic acid and demethylation

to *gentisic acid*. The above oil (1 g.) was suspended in hot water (50 ml.) containing N KOH (1 ml.) and to it was added a solution of KMnO_4 (1.5 g.) in hot water (150 ml.). After $\frac{1}{2}$ hr. heating on the water-bath no KMnO_4 remained. The solution was filtered, the filtrate evaporated to 25 ml. and acidified to Congo red with 2N HCl (3 ml.). An emulsion was formed which rapidly crystallized, forming colourless needles, wt. 0.55 g., m.p. 72–76°. After recrystallization from water the oxidation product melted at 75–76°, not depressed on admixture with authentic, synthetic 2 : 5-dimethoxybenzoic acid. (Found: C, 59.30; H, 5.60. Calc. for $\text{C}_9\text{H}_{10}\text{O}_4$: C, 59.33; H, 5.53 %.)

– The acid (0.13 g.) was heated with HI (sp. gr. 1.7; 3 ml.) in a Zeisel apparatus in a stream of CO_2 , the bath temperature being maintained at 130–140° during 1 hr. On cooling, fine brownish needles separated. These were collected and washed with conc. HCl. Wt. 0.07 g., m.p. 199–201°. The crystals gave a colourless crystalline sublimate in a high vacuum, m.p. 204° not depressed on admixture with authentic *gentisic acid*, m.p. 203°. (Found: C, 54.51; H, 4.09. Calc. for $\text{C}_7\text{H}_5\text{O}_4$: C, 54.55; H, 3.92 %.)

Synthesis of gentisyl alcohol (2 : 5-dihydroxybenzyl alcohol). *Identity with metabolic product*. *Gentisic aldehyde* (2 : 5-dihydroxybenzaldehyde) (0.5 g.),

prepared from salicylaldehyde by the method of Neubauer & Flatow [1907] and dissolved in ethanol (10 ml.), was hydrogenated at ordinary pressure with a palladium-norite catalyst prepared from 0.2 g. of PdCl_2 and 0.5 g. of norite. The uptake of hydrogen was fairly rapid. Hydrogenation was discontinued after 50 min. (Uptake of hydrogen, 82.5 ml. (corr.); theor. for uptake of 1 mol. H_2 , 81.2 ml.) After filtering from catalyst the ethanol was removed *in vacuo*. A yellow oil (0.57 g.) remained and began to crystallize overnight. The product was recrystallized from chloroform in the same way as is described above for the purification of the crude metabolic product. After two recrystallizations it was obtained as almost colourless needles, m.p. 99° after softening at 95°. It was finally sublimed in a high vacuum. Colourless crystalline sublimate, m.p. 100°, alone or in admixture with the metabolic product. It also gave the same reactions. (Found: C, 60.09; H, 5.79. Calc. for $\text{C}_7\text{H}_5\text{O}_3$: C, 60.01; H, 5.75 %.)

SUMMARY

Gentisyl alcohol, 2 : 5-dihydroxybenzyl alcohol, occurs together with patulin as a metabolic product of *Penicillium patulum*, Bainier. Its isolation, identification and synthesis are described.

REFERENCES

- Anslow, W. K., Raistrick, H. & Smith, G. [1943]. *J. Soc. chem. Ind., Lond.*, **62**, 236.
 Birkinshaw, J. H., Bracken, A., Michael, S. E. & Raistrick, H. [1943]. *Lancet*, **2**, 625.
 Neubauer, O. & Flatow, L. [1907]. *Hoppe-Seyl. Z.* **52**, 375.