absorption of the exciting light. The high quenching effect of lactate is remarkable, and requires further investigation for its explanation.

The quenching effect in *isobutanol* solutions is influenced by the absolute and relative solubilities of the quenching compounds in water and *isobutanol*. In the saturated *isobutanol* solution it is higher for most substances than in aqueous solution. For a number of substances which do not quench in aqueous solution, slight quenching occurs in *isobutanol* saturated with water, but for a number of substances, particularly chloride, bromide and iodide, the quenching in *isobutanol* is considerably greater than in aqueous solution. The effect of potassium chloride is probably due to the chloride ion; the slightly greater one of Mg and Ca may be partly due to the cation. The quenching effect seen in the *isobutanol* extracts of the aqueous solutions is governed by the partition coefficient of the quenching substance between water and *isobutanol*.

Apart from the quenching effect of the halide and the slight effect of some others, the considerable quenching by the ferric cation and the ferricyanide anion is remarkable. Potassium ferricyanide in a concentration of 0.02N has a quenching effect of 18%. Since this substance is used in the preparation of thiochrome for the assay of vitamin B₁, the concentration of this salt in the final extract may be of great influence.

The curve of the dependence of the fluorescence intensity of riboflavin on the pH shows a maximum between pH 5.9 and 7.7; between 3.8 and 5.9 the intensity is about 97% of the maximum. At pH values below 3.8 and above 7.7 there is a rapid fall in the fluorescence intensity. The curve corresponds to that published by Kuhn & Moruzzi (1934).

With thiochrome in aqueous buffer solutions the maximum fluorescence intensity is reached at the alkaline end of the range examined (pH 11). The intensity decreases slowly at pH 8.8, then more rapidly at pH 7.5 and falls steeply at pH 5. This is in accordance with the findings of Kuhn & Vetter (1935). Below this pH the blue fluorescence has completely disappeared and there is only a small amount of green fluorescence. In *isobutanol* solutions saturated with buffer solutions the effect is less marked. There is very little alteration between pH 11 and 4.5, but below 4.5 the fluorescence intensity falls steeply.

SUMMARY

The influence of changes in pH of the solution, and of the presence of a number of inorganic and a number of organic compounds on the fluorescence intensity of riboflavin and thiochrome, has been investigated.

The authors wish to thank Glaxo Laboratories Ltd. for a gift of riboflavin and Roche Products Ltd. for thiamine.

REFERENCES

- Barger, G., Bergel, F. & Todd, A. R. (1935). *Ber. dtsh. chem. Ges.* **68**, 2257.
Cohen, F. H. (1934). *Acta brev. neerl. Physiol.* **4**, 46.
Eisenbrand, D. J. (1934). Quoted in Dankworth, P. W., *Luminiszenz Analyse in filtriertem ultravioletten Licht*, 3rd ed. p. 133. Leipzig: Acad. Verlaganstalt.
Ellinger, P. & Holden, M. (1944). *J. Soc. Chem. Ind.* **63**, 115.
v. Euler, H. & Adler, E. (1934). *Hoppe-Seyl. Z.* **223**, 105.
Jansen, B. C. P. (1936). *Rec. Trav. Chim. Pays-Bas*, **55**, 1046.
Joseph, B. (1934). *Acta brev. neerl. Physiol.* **4**, 46.
Kuhn, R. & Moruzzi, B. (1934). *Ber. dtsh. chem. Ges.* **67**, 888.
— & Vetter, H. (1935). *Ber. dtsh. chem. Ges.* **68**, 2375.
Medical Research Council (1943). *Biochem. J.* **37**, 433.
Najjar, V. A. (1941). *J. biol. Chem.* **141**, 355.
Weil-Malherbe, J. & Weiss, J. (1942). *Nature, Lond.*, **149**, 471.
Weiss, J. (1938). *Trans. Faraday Soc.* **34**, 451.

A Method for the Estimation of Nicotinamide Methochloride in Urine

By R. A. COULSON, P. ELLINGER AND M. HOLDEN, *Lister Institute of Preventive Medicine, London, S.W. 1*

Received 9 February 1944)

Najjar & Wood (1940), Najjar & Holt (1941 *a, b*) and Najjar, Stein, Holt & Kabler (1941) described a fluorescent pigment 'F₂' in the urine of normal dogs and of man, the elimination of which increased after ingestion of nicotinamide and related compounds, but which was absent from the urine of

pellagrins. It was suggested by Coulson, Ellinger & Platt (1942) that information about the metabolism of these compounds could be expected from an investigation of the urinary elimination of F₂, and indeed a preliminary method of assay was developed for this specific purpose before the chemical

nature of the compound was known (Coulson, Ellinger, Glock & Platt, 1942). A large number of estimations was made with this method, which provided relative but not absolute values. The investigation of the nature of F_2 led Huff & Perlzweig (1943a) to state that the fluorescent F_2 had the same absorption spectrum as, and was identical with, nicotinamide methochloride. Ellinger & Coulson (1943) and Coulson & Ellinger (1943), however, showed that synthetic nicotinamide methochloride (1-methylpyridinium chloride-3-carboxylic acid amide) is not fluorescent and only becomes fluorescent on treatment with alkali and transfer into *isobutanol*. They showed further that the substance which is found in urine in increased amounts after ingestion of nicotinamide is also non-fluorescent until treated with alkali and *isobutanol*. The natural untreated substance was then isolated and identified as nicotinamide methochloride. It was suggested that F_2 formed *in vitro* was most probably a mixture of F_{2b} , 6-hydroxy-1:6-dihydro-1-methylpyridine-3-carboxylic acid amide, with an absorption maximum at 358 $m\mu$, and another carbinol F_{2a} with an *o*-quinoid structure and three absorption maxima at 271, 306 and 347 $m\mu$. The elucidation of the nature of nicotinamide methochloride and of the mechanism of the fluorescence development provided a means of improving the assay so that absolute values could be obtained. While this work was in progress two methods of assay were published by Huff & Perlzweig (1943 b, c). With regard to the nomenclature it should be mentioned that the non-fluorescent precursor should be called nicotinamide methochloride, since the crystalline form on which the measurement is based is a chloride. The fluorescent derivatives should be called ' F_2 ' until their structure is fully proved by further experimental evidence.

METHOD OF ASSAY

The principle of the method is based on the separation of nicotinamide methochloride from urine by adsorption and elution, transformation of the substance into fluorescent derivatives (F_2) by treatment with alkali and extraction with *isobutanol*, and estimation of the concentration of F_2 in *isobutanol* by visual fluorimetry.

The assay is carried out as follows. The lower ends of thistle funnels of 4-5 mm. internal diameter are plugged with glass wool, and 1 g. Decalso (60 mesh) (Permutit Co., Ltd.) is washed into each tube. Five ml. or less of the urine diluted to 10 ml. are filtered through the column, which is then washed with 50 ml. distilled water and eluted with 14 ml. 25% (w/v) KCl solution. The eluate is collected in a dry tube and can be stored in the cold without deterioration. The eluate is divided into two equal portions which are transferred into glass-stoppered tubes of about 12 ml. capacity. Non-fluorescent *isobutanol* (2 ml.) is added to each and 1 ml. 15% (w/v) NaOH to one to develop fluorescence. The tubes are shaken at once vigorously for 5 min., and after separation of the layers the *isobutanol* is pipetted

off into identical non-fluorescent test-tubes, each containing about 100 mg. anhydrous Na_2SO_4 .

The *isobutanol* extract of the neutral half of the KCl-eluate is used as a reagent blank. The fluorescence intensity of the extract is compared with that of standards. The extracts are stable at room temperature in the dark.

Standards are prepared by passing a series of 10 ml. portions containing 1-12 $\mu g./ml.$ of crystalline nicotinamide methochloride, prepared as described by Karrer, Schwarzenbach, Benz & Solmssen (1936), in distilled water, through 1 g. Decalso columns. The columns are washed and eluted with 14 ml. 25% KCl solution in the manner described above. Seven ml. of each of the eluates are made alkaline and extracted with 2 ml. *isobutanol*. The separated extracts, dried with Na_2SO_4 , are used as the standards. The tubes should be identical with those used for the extracts in the assay and the standards should be kept in the dark.

The *isobutanol* extracts are read in the filtered 366 $m\mu$ line beam of a high-pressure mercury lamp by naked-eye comparison of the unknown with the standards, in a dark room. With such standards no instrument for measuring fluorescence intensity is necessary apart from the high-pressure mercury lamp. To increase the accuracy of the assay the results of three estimations with different amounts of urine should be averaged.

For large-scale routine work wooden racks supporting twelve thistle funnels were used. Sixteen tubes were shaken simultaneously in a box constructed for the purpose.

Examination of the various steps involved in the assay

Stability of nicotinamide methochloride in urine. Nicotinamide methochloride in urine was found to be stable for at least a week at room temperature in the dark in spite of the decomposition of the urine. It was therefore possible to collect specimens in distant districts and assay them, without loss, in the laboratory at a convenient time. Substances generally used for preserving urines interfere with the various steps of the procedure and should not be used. Boiling of urine for 15 min. does not affect the assay.

Adsorption. Of a number of adsorbents examined Decalso was found to be the most satisfactory from the standpoint of adsorption and elution. The speed of filtration, an important factor in large-scale experiments, is satisfactory with samples of about 60 mesh. The disadvantages of this adsorbent are that it cannot be sufficiently reactivated for further use, and that different batches may have a different physical and chemical constitution and therefore different adsorptive properties. Decalso of the same batch should be used for assay and preparation of standards. Up to 1 mg. of nicotinamide methochloride in 10 ml. is completely adsorbed from aqueous solutions by 1 g. Decalso. The adsorption is affected by electrolytes in concentrations above 0.4M but not by urea or changes of pH within the range 3-8. Five ml. of urine should be used for the assay if the concentration of nicotinamide methochloride is less than 12 $\mu g./ml.$ and the hourly urine volume is greater than 100 ml. With higher concentrations and lower urine volumes, correspondingly smaller amounts of urine must be taken and diluted to 10 ml. to reduce the danger of incomplete adsorption.

Washing. Washing with distilled water is essential to remove all materials which might interfere with the assay, particularly fluorescent pigments contained in the original

urine. Fifty ml. of water are always sufficient for this purpose.

Elution. Elution by the chlorides of Na, K, Ca, Sr and Ba was investigated. KCl in a concentration of 25% was found to be the most efficient. The chlorides of the bivalent cations gave almost unbreakable emulsions when shaken with *isobutanol*. The elution of the nicotinamide methochloride is an equilibrium process and is not complete when 14 ml. 25% KCl are used for 1 g. Decalso under these conditions. Two-thirds of the total adsorbed material is always eluted, as shown in Table 1.

Table 1. *Relative fluorescence intensities of two series of isobutanol extracts of successive KCl eluates from the same Decalso columns*

No. of elution	Relative intensities of fluorescence	
	100	100
1	100	100
2	29	33
3	9.7	10
4	3.1	2.8

Conversion and extraction. The conversion of nicotinamide methochloride into fluorescent derivatives, and the extraction of these derivatives, are also equilibrium processes. In aqueous alkaline solution the equilibrium favours the quaternary ammonium base and very little of the fluorescent derivatives is present. In *isobutanol* the fluorescent substances which are readily soluble in water and in *isobutanol* are predominant (Ellinger & Coulson, 1943). In the assay the mixture of the two fluorescent derivatives can be treated as a single pigment for practical purposes. An aqueous alkaline solution of nicotinamide methochloride is unstable. In the first 10 min. after the addition of alkali the deterioration is negligible, but standing for 12 hr. completely destroys the substance. The extraction should be carried out as quickly as possible after the NaOH has been added to the eluate, to avoid the risk of loss by destruction. For optimal extraction vigorous shaking by hand or mechanical shaker for not less than 5 min. is essential. Longer shaking or high speed mechanical shaking does not increase the yield.

In order to compare the intensity of the fluorescence derived directly from nicotinamide methochloride with that of the standards which had been adsorbed on Decalso, an *isobutanol* stock solution of the fluorescent derivatives was prepared in the following way. Ten mg. nicotinamide methochloride dissolved in 2 ml. of water were diluted to 100 ml. with non-fluorescent *isobutanol*. A pellet of NaOH was added and the solution shaken for about 5 min. until no further increase in fluorescence was observed. After standing for $\frac{1}{2}$ hr. in the dark the remainder of the NaOH pellet was removed. Solutions containing 0.025–0.5 $\mu\text{g.}/\text{ml.}$ at narrow intervals were prepared by dilution of this stock solution. The diluted solutions were then fluorimetrically compared with the standards used in the assay. The results are shown in Fig. 1.

If we assume that nicotinamide methochloride in *isobutanol* is converted quantitatively into the fluorescent derivatives, an *isobutanol* solution of 0.5 $\mu\text{g.}/\text{ml.}$ corresponds in fluorescence intensity to that of the 2 ml. *isobutanol* extract derived from an aqueous or urine solution which contained 12.12 $\mu\text{g.}$ and was treated in the manner of the assay of nicotinamide methochloride. The 2 ml. *isobutanol* extract derived from the above urine would contain 1.0 $\mu\text{g.}$

of the F_2 . This represents an over-all yield of 16.5%, since only one-half of the KCl eluate is used for the assay.

Under the conditions of the assay (7 ml. 25% KCl, 1 ml. 15% NaOH and 2 ml. *isobutanol*), one single extraction

Fig. 1. Relation of fluorescence intensity obtained from 10 ml. solutions of nicotinamide methochloride that have undergone the assay (A) to that of F_2 solutions developed directly by alkaline *isobutanol* treatment of nicotinamide methochloride (B).

transfers 45% of the pigments extractable into the *isobutanol* phase. If the eluate is extracted with four successive 2 ml. portions the aqueous solution is practically exhausted. The yields of four successive extracts are shown in Table 2.

Table 2. *Relative yield of fluorescent derivatives of nicotinamide methochloride from four successive 2 ml. isobutanol extractions of 7 ml. 25% KCl eluates*

(Eluates obtained from 0.3–3.0 ml. of same 23 hr. urine sample after an intake of 500 mg. nicotinamide.)

Vol. urine used (ml.)	Fluorescence intensity (arbitrary units) of extract				Fluorescence intensity of first extract in % of the total*
	First	Second	Third	Fourth	
0.3	2.1	2.1	0.2	0.06	47.1
0.5	3.5	2.5	0.9	0.0	50.7
0.8	5.8	5.8	1.5	0.06	44.0
1.0	6.8	6.3	1.5	0.06	46.3
2.0	12.0	12.0	2.8	0.1	44.6
3.0	15.5	15.0	4.0	0.8	43.3

* Values of column 6 are obtained by dividing the values of column 2 by the sums of columns 2–5 and multiplying them by 100.

The final yield after the processes of elution, conversion and extraction is, as shown, about 16.5%. Since the loss in elution is 33%, and that in extraction 55%, then by calculation the loss in conversion is about 45%.

To render the *isobutanol* non-fluorescent it is sometimes necessary to remove any fluorescent pigments present by shaking with charcoal; norite or activated charcoal (British Drug Houses, Ltd.) were found to be suitable for this

purpose. Contact of *isobutanol* with corks that have not been exhaustively extracted with *isobutanol* must be avoided.

Measurement of fluorescence intensity. The molar fluorescence intensity of F_2 in *isobutanol* is relatively low, being only about one-third that of quinine sulphate in 0.1 N- H_2SO_4 and one-ninth that of thiochrome in *isobutanol*. The fluorescence efficiency in the 366 m μ line (intensity when 50 % of the incident light is absorbed) is low, about one-tenth that of quinine sulphate in 0.1 N- Na_2SO_4 and one-sixth that of thiochrome in *isobutanol*. Therefore only a comparatively narrow range can be measured with sufficient accuracy. Between 0.025 and 0.5 $\mu\text{g./ml.}$ of the fluorescent derivatives can be estimated with the naked eye and 0.1–4.0 $\mu\text{g./ml.}$ with the photoelectric fluorimeter of Ellinger & Holden (1944), with a 2 mm. cuvette. The fluorescence intensity in *isobutanol* is not affected by pH in the range between 9.24 and 3. Of a number of inorganic substances which are used in the assay, or which might be present in the urine, a slight quenching effect was found only with Mg^{++} , Ca^{++} , SO_4^{--} and SO_3^{--} . This quenching effect, however, is negligible with the concentrations of these substances likely to be encountered.

The measurement of the fluorescence intensity is carried out preferably by a visual method, first, on account of the higher speed and secondly, because lower concentrations are measurable. By this method it is possible to detect a difference of 0.15 $\mu\text{g./ml.}$ of nicotinamide methochloride in the range between 0.6 and 6 $\mu\text{g./ml.}$ of original aqueous solution and a difference of 0.3 up to 12 $\mu\text{g./ml.}$ of solution. If the concentration in the final extract is higher than that derived from 12 $\mu\text{g./ml.}$, accurate reading is impossible. Fairly accurate estimation can be obtained by dilution of the *isobutanol* extract until it is within the range of the standards; it is, however, preferable to repeat the assay with a smaller amount of urine to safeguard against the possibility of incomplete adsorption.

Fresh standards should be prepared each week in sealed tubes and kept cool in the dark. The fluorescent pigments are unstable in the long wave ultra-violet region of mercury or daylight. The repeated short exposures in the usual readings of one week, do not, however, materially affect the fluorescence intensity.

RESULTS

To test the accuracy of the method, various concentrations of nicotinamide methochloride in urine of known low F_2 content were estimated. The results are shown in Fig. 2.

It is seen that accurate estimations are possible for any concentration greater than 0.6 $\mu\text{g./ml.}$ Between 0.6 and 1.0 $\mu\text{g./ml.}$ the accuracy is about 10 % and from 1.0 $\mu\text{g./ml.}$ upwards the accuracy is about 5 %. Discrimination of small differences in fluorescence intensity is best in concentrations between 2 and 5 $\mu\text{g./ml.}$ From Fig. 2 it is evident that the nicotinamide methochloride estimations of aqueous solutions are more accurate (0.4 % mean error) than those of urine solutions (5 % mean error). The values from urine are consistently lower than calculated. Experiments are in progress to reduce this error and to determine whether it is due to incomplete adsorption caused by the presence of electrolytes or

to some quenching substances accompanying nicotinamide methochloride through the assay procedure.

Fig. 2. Values obtained for a series of known nicotinamide methochloride concentrations in water (+) and in urine (Δ). The straight line represents the calculated values.

DISCUSSION

The method has been evolved to make it possible to assay a large number of urine samples at the same time. Sixty single examinations can be carried out in one day by two workers. The two methods published by Huff & Perlzweig (1943 b, c) do not seem to have been developed for large-scale surveys.

With the method described, the daily output of nicotinamide methochloride was determined for normal persons and for experimental subjects who had taken nicotinamide and related compounds. Details of these experiments will be reported later. The daily output varies greatly in different individuals and in the same person at different times. From 2 to 8 mg./day are eliminated by normal persons. Somewhat lower results (0.1–3.2 mg. in 14 hr.) are reported by Sarett (1943). Variations are also observed in the nicotinamide methochloride urinary output of different subjects after the ingestion of nicotinamide. There is an average recovery of about 10–15 % after the ingestion of 100 mg. nicotinamide. Continued administration leads to a gradual increase in the yield of nicotinamide methochloride in the urine. By repeated estimation of the daily output after successive administrations of nicotinamide a 'saturation test' can be obtained.

SUMMARY

A method is described for the routine estimation of nicotinamide methochloride in urine. Diluted urine is passed through Decalso to adsorb the nicotinamide

methochloride. The Decalso is eluted with KCl solution, which is then made alkaline and extracted with isobutanol. The fluorescence produced is compared with that of standards similarly treated.

This investigation forms part of an investigation of nicotinamide deficiency carried out on behalf of the Air Ministry. We wish to thank Air Marshal Sir H. E. Whit-

tingham, K.B.E., K.H.P., Director-General of the Medical Services of the Royal Air Force, for facilities provided. Our thanks are also due to L.A.C.W. A. E. Wrigglesworth for technical assistance. One of the authors (R. A. C.) is a member of the Civilian Technical Corps (Air Ministry), attached to the Lister Institute for an investigation of nicotinamide deficiency. The authors wish to thank Roche Products, Ltd., for a generous supply of nicotinamide.

REFERENCES

- Coulson, R. A. & Ellinger, P. (1943). *Biochem. J.* **37**, *Proc.* xvii.
 ——— Glock, G. E. & Platt, B. S. (1942). *Biochem. J.* **36**, *Proc.* xix.
 ——— & Platt, B. S. (1942). *Biochem. J.* **36**, *Proc.* xii.
 Ellinger, P. & Coulson, R. A. (1943). *Nature, Lond.*, **152**, 383.
 ——— & Holden, M. (1944). *Chem. Industr.* **63**, 115.
 Huff, J. W. & Perlzweig, W. A. (1943a). *Science*, **97**, 538.
 ——— (1943b). *J. biol. Chem.* **150**, 395.
 ——— (1943c). *J. biol. Chem.* **150**, 483.
 Karrer, P., Schwarzenbach, G., Benz, F. & Solmssen, U. (1936). *Helv. chim. Acta*, **19**, 826.
 Najjar, V. A. & Holt, L. E. Jr. (1941a). *Science*, **93**, 20.
 ——— (1941b). *Proc. Soc. exp. Biol., N.Y.*, **48**, 413.
 ——— Stein, H. J., Holt, L. E., Jr. & Kabler, C. V. (1941). *J. clin. Invest.* **21**, 263.
 ——— & Wood, R. W. (1940). *Proc. Soc. exp. Biol., N.Y.*, **44**, 386.
 Sarett, H. P. (1943). *J. biol. Chem.* **150**, 159.

Enzyme Formation and Polysaccharide Synthesis by Bacteria

2. POLYSACCHARIDE FORMATION BY *RHIZOBIUM RADICICOLUM* STRAINS

By H. G. BRAY, E. SCHLÜCHTERER AND M. STACEY, *Chemistry Department, The University, Birmingham*

(Received 7 February 1944)

An investigation into polysaccharide production by various strains of *Rhizobium* growing on the basal culture medium of Cooper, Daker & Stacey (1938) is described. The strains used were provided by Dr Hugh Nicol, and it was found that his two clover strains termed 'Clover Bartel A' and 'Clover 2027' were remarkably prolific in polysaccharide production. In general the purified polysaccharides gave laevo-rotatory, viscous solutions in water, although some differences in properties and yields of the polysaccharide were apparent from strain to strain. About 200 g. of the polysaccharide were obtained from the Bartel A strain.

In order to prevent contamination of the polysaccharide with the agar medium some modifications to the normal cultural conditions were made. For example, the usual agar culture medium, distributed in large Petri dishes, was covered with a layer of sterile cellophan according to the method of Birch-Hirschfeld (1933-4) and the surface inoculated with the bacterial suspension. The polysaccharide, uncontaminated by agar, was produced on the cellophan surface as a white mucilaginous growth which could easily be washed off and purified. In another method, suggested by Prof. W. N. Haworth, F.R.S., thick porous plates were embedded in agar medium so that nutrient material could freely diffuse through and be utilized by the organisms inoculated on to

the surface of the plates. Polysaccharide production was good. In a third method the agar in the basal medium was replaced by acid-free kieselguhr, to provide a pasty solid medium which was distributed in Roux bottles and conical flasks. The organism grew well on the medium, covering the surface with a white mucilaginous material from which the polysaccharide could be obtained by extraction with water. The product made in this way generally had a much higher ash content than that made by other methods.

By an experimental procedure involving an examination of the hydrolysis products of its methylated derivative, which will be described elsewhere, part of the structure of the Clover Bartel A polysaccharide has been determined. The hydrolysis products were 2:3:6-trimethylglucopyranose (1 part), 2:3-dimethylpyranose (1 part) and 2:3-dimethylglucuronic acid (1 part). The type of structure assigned to the repeating unit is that of a polymerized aldobionic acid in which the carboxyl groups of the glucuronic acid residues are free. It has been shown previously (Cooper *et al.* 1938) that *Rhizobia* polysaccharides belong to the same class of compound as the specific polysaccharides from types II and III *Pneumococcus* (see Heidelberger & Goebel, 1927; Hotchkiss & Goebel, 1937). The structure of the latter has been worked out in some