benzidine. The amount present in the liver must be very small, however, and it is evident that the main tendency is for azobenzene to be rapidly eliminated as the water-soluble hydrazo derivative.

Azobenzene is not carcinogenic, but in work which is being carried out with the carcinogenic *p*-dimethylamino-azobenzene evidence has been obtained that similar metabolic changes take place. In this case, however, the hydrazo derivative does not appear to be so stable and undergoes the benzidine rearrangement in the body more readily, since a substance giving the colour reaction of one of the 'benzidine change' products of *p*-dimethylamino-azobenzene can be extracted directly from the neutral or alkaline urine.

SUMMARY

The urine of rats which have received azobenzene has been found to contain aniline, and a water-soluble derivative which, after acidification with dilute acid followed by addition of sodium hydroxide and subsequent extraction with ether, yields benzidine.

We desire to thank I.C.I. (Dyestuffs) Ltd. for permission to publish this work. We are also indebted to the Sir Halley Stewart Trust for a Fellowship held by one of us (F. L. W.) during this work and to the British Empire Cancer Campaign and to the Anna Fuller Fund for generous grants which have helped to support this investigation.

REFERENCES

- Andervont, H. B. & Edwards, J. E. (1941). *J. Nat. Cancer Inst.* **2**, 139.
 Boyland, E. & Brues, A. M. (1937). *Proc. Roy. Soc. B*, **122**, 429.
 Cook, J. W., Hewett, C. L., Kennaway, E. L. & Kennaway, N. M. (1940). *Amer. J. Cancer*, **40**, 62.
 Fischer, B. (1906). *Münch. med. Wschr.* **53**, 2041.
 Hayward, E. (1909). *Münch. med. Wschr.* **56**, 1836.
 Kinoshita, R. (1937). *Trans. Jap. Path. Soc.* **27**, 665.
 Maughan, G. B., Evelyn, K. A. & Browne, J. S. L. (1938). *J. biol. Chem.* **126**, 567.
 Rose, F. L. & Bevan, H. G. L. (1944). *Biochem. J.* **38**, 116.
 Yoshida, T. (1934). *Trans. Jap. Path. Soc.* **24**, 523.

Changes in the Skeleton during Gestation and Lactation in the Rat

By G. M. WARNOCK AND J. DUCKWORTH, *The Rowett Institute, Aberdeen*

(Received 3 March 1944)

Many observations show that during pregnancy mammals store Ca and P in the skeleton and that these substances are released during the ensuing lactation (Forbes, Black, Braman, Frear, Kahlenberg, McClure, Swift & Voris, 1935; Coons, Schiefelbusch, Marshall & Coons, 1935; Duckworth & Warnock, 1942-3; Goss & Schmidt, 1930). The extensive growth of cancellous bone which takes place during the pre-ovulatory stage with subsequent resorption about the time of ovulation in the pigeon suggests that a similar process occurs also in birds (Kyes & Potter, 1934; Bloom, Bloom & McLean, 1941). The classical studies of Bauer, Aub & Albright (1929) showed that cancellous bone served as a store of Ca and P and that storage and depletion were governed by the level of Ca intake. Using alizarin red as an intravital stain, they also demonstrated that mobilization of the trabeculae was caused by parathyroid hormone.

Apparently, no investigation has been specifically designed to determine if the trabeculae are the sole physiological store of Ca and P in the adequately nourished animal, or whether, in lactation, the shaft

is also drawn upon by direct resorption, or by partial demineralization without resorption, or by resorption with concomitant partial demineralization of the unresorbed portion. The importance of this fundamental question is underlined by the description given by Theiler & Green (1931-2) of the modern, high milk-producing, dairy cow: 'such an animal is pathological; oscillating between a compact bone tissue and a mildly osteoporotic framework in each reproductive cycle.' In addition no comparison seems to have been made of the behaviour of the skeleton of animals lactating normally with that of animals which, at the conclusion of gestation, are not allowed to suckle their young. Neither have comparative studies been made of the rates of replenishment of skeletal stores after lactation in pregnant and non-pregnant animals.

The present investigation was designed to examine these points. The shafts of certain long bones were separated from the ends and the two portions studied. Although this mode of approach has obvious limitations, it permitted some clarification of the problem.

EXPERIMENTAL

(a) *Groups of animals*

One hundred and fifty female hooded Lister rats were divided into twenty-five subgroups of six rats each and these subgroups were distributed among the three groups described below. All rats were transferred to the experiment on reaching 100 days of age. Litter mates were distributed evenly among the three groups, being placed in appropriate corresponding subgroups. This plan was not strictly followed, for some rats failed to conform to the standard of performance given below; they were discarded and replaced by others.

The three groups into which the animals were divided were treated as follows:

Mated-rat group. These rats were introduced to the male at 100 days of age and the occurrence of coitus was detected by vaginal smear. Of the nine subgroups three were killed during the first gestation, one at the seventh day (reckoned from the date at which either a vaginal plug was seen or remains of spermatozoa were found in the smear), one at the fourteenth day and one at parturition. Six subgroups began lactation, one being killed on the seventh day, one on the fourteenth day and one on the twenty-third and final day of lactation. The remaining three subgroups were reintroduced to the male at the end of lactation, tested for coitus and killed at the same intervals as in the first gestation.

Control Group A. Of the six subgroups in this group three served as non-lactating controls to those lactating in the mated-rat groups. The rats of these subgroups were mated as in the above group but their litters were killed at birth. One subgroup was killed on the seventh day after parturition, one on the fourteenth day and one on the twenty-third day. The rats of the remaining three subgroups suckled their young but were not re-mated; they acted as controls for the animals in the mated-rat groups during the second gestation. They were killed at times corresponding to those in the second gestation of the mated-rat group.

Control Group B. The rats in this group served as non-pregnant and non-lactating controls of the mated-rat group. They were sacrificed at the same time as their corresponding litter-mates in that group. This group also included a subgroup, the rats of which were killed at 100 days of age.

The normal stock diet of the colony was used and consisted of feeding cubes (Thomson, 1936), greenstuff (cabbage, kale or clover), 5 ml. fresh milk daily and tap water. The percentage composition of the feeding cubes was: wheat offal (fine bran), 17.76; ground wheat, 17.76; Sussex ground oats, 17.76; ground barley, 8.80; ground maize, 8.80; meat and bone meal (45% protein), 8.80; dried skimmed milk, 14.00; white fish meal (60% protein), 4.30; dried yeast (40% protein), 1.10; sodium chloride, 0.46; cod-liver oil, 0.46. With the start of the experiment the milk allowance of all rats was increased to 10 ml. daily, with a further increase to 25 ml. during lactation. The animals were housed in screen-bottomed, galvanized iron cages with wood wool as bedding. They were weighed every third day and killed at the appropriate time with coal gas. Litters were reduced to eight in number on the day following parturition and were weaned at 23 days of age.

Rats were discarded and replaced by others if: (1) their clinical examination revealed any abnormality such as respiratory disorders, (2) when mated they failed to become pregnant within two oestrus cycles, (3) their litters con-

tained less than eight living young, (4) of the eight young ones left with the mother any died during lactation.

(b) *Methods*

After dissection from the body the hind legs were immersed in boiling water for 3 min. to facilitate cleaning. The cleaned bones were then repeatedly extracted with boiling ethanol to remove fat.

Both tibiae and femora were used for analyses. One-quarter of the length of each femur was measured from each end and these portions sawn off with a jeweller's file. The loss in bone dust due to sawing is small and causes no serious errors in analysis. The tibia was divided so that one-quarter of its length, measured from the proximal end, and one-tenth of its length, measured from the distal end, could be sawn off. The shafts were opened and the cancellous bone and marrow within the cavity was added to the eight ends from the tibiae and femora in order that, as far as possible, the shafts should include only compact bone and that cancellous bone should be included in the composites subsequently referred to as 'ends'. The remainder of the bones, subsequently called 'shafts', thus represented half the length of the femora and thirteen-twentieths of the length of the tibiae.

The ends and shafts were dried in an oven at 99° to constant weight. The ashing was carried out in an electric furnace at dull red heat until weight was constant.

RESULTS

The curves in Figs. 1-3 were obtained by joining together the average values of the body weight and bones observed at death in the subgroups. In discussion they will be referred to as growth curves. It should be noted that of the six subgroups of control A rats, three served as controls during the lactation period and three as controls during the second gestation and therefore the points on the curve at the twenty-third day of lactation and the seventh day of the second gestation could not be joined.

Fig. 1 shows the degree of variability in the average body weights of the different subgroups. Comparison of groups and subgroups should be made with these variations in mind, particularly in the case of minor differences. Certain points, outside the scope of the present study and technique, are investigated more satisfactorily by the metabolic balance method.

In Fig. 2 the weights of the dry, fat-free ends and shafts of the bones, and of whole bones are recorded. The values for whole bone were obtained by combining the corresponding values for ends and shafts. During the first gestation period the weights of whole bones of the mated rats were slightly greater than the weights of those of control B rats; the rates of increase in bone-weight were similar. With the onset of lactation there was a sharp reduction in the weight of whole bones of the mated-rat groups which continued in an almost linear manner throughout lactation. In the second gestation there was a

sharp increase, again almost linear, in the bone weights of the mated-rat groups. Throughout the study the bone weights of the control B rats increased in a somewhat irregular manner owing to

Fig. 1. Average body weight of each subgroup at the time of killing, showing variation in body weight within groups. — Mated-rat group. —●— Control group A. ---- Control group B, non-pregnant, non-lactating rats.

Fig. 2. Average dry fat-free weight, in mg., of the shafts, ends and whole bone (shaft and ends) of each subgroup at different stages of gestation and lactation. The femora and tibiae were used. — Mated-rat group. —●— Control group A. ---- Control group B, non-pregnant, non-lactating rats.

differences in body weight between subgroups, as is shown in Fig. 1. It is, of course, obvious that no comparison on a body weight basis can be made of the bone data of pregnant or lactating rats with non-pregnant or non-lactating rats. The bones of the control A rats, whose litters had been killed at birth, behaved in a manner similar to those of control B rats during the interval corresponding to

lactation in the mated-rat group. The greater weights of control A rats as compared with control B rats during this period may be due to the previous pregnancy. After lactation the control A rats showed a rate of bone growth which was almost identical with that of the mated-rat groups undergoing the second gestation. The total weight of dry fat-free bone was greater in the mated-rat group at the end of the second gestation than at the end of the first gestation. From the end of the first gestation the whole bones of the control B group were consistently heavier than those of the mated-rat groups.

The changes observed in whole bone weight during lactation and the second gestation reflected the changes which occurred in the bone ends. As measured by weight, shaft growth continued during the pregnancy of the mated rats, and may have undergone a change during lactation. Compared with the B control rats, the mated rats showed during their second gestation a decrease in bone weight. This decrease, also seen in the non-pregnant A rats, may have resulted from a slackening of bone growth during lactation. There is, of course, no evidence of a resorption of shaft in any way comparable with that occurring at the bone ends.

Fig. 3. Percentage ash in dry fat-free shafts, ends and whole bones (ends and shafts) of each subgroup at different stages of gestation and lactation. The femora and tibiae were used. — Mated-rat group. —●— Control group A. ---- Control group B, non-pregnant, non-lactating rats.

The values for the weight of ash in whole bones, ends and shafts followed paths almost identical with those of dry fat-free bones. The changes in ash content of the whole bones and ends were, however, exaggerations of those found for dry fat-free bone, with the result that the percentage ash values did not remain constant, as is shown in Fig. 3.

The percentages of ash in the shafts of the rats of all groups were not significantly different, and were remarkably constant after an initial reduction followed by an increase. In the whole bone of the mated-rat group a reduction in the percentage of ash during lactation resulted from changes in the percentage of ash in the bone ends. During the interval corresponding to lactation in the mated-rat group the percentages of ash in the whole bones and ends of the control A rats followed the path of the values for control B rats. The reduction observed at the seventh day in the non-lactating control rats is difficult to explain, but since it occurred equally in ends and shafts it may have arisen from causes not connected with the experiment.

The changes in the percentage of ash in the ends of the bones from the seventh to fourteenth day of the second gestation were unexpected. They occurred in both the non-pregnant rats of control A group and in the pregnant rats of the mated-rat group. The percentage of ash in the bones of both groups followed almost identical paths. The final value for percentage ash in the ends did not again reach the high level at the termination of the first gestation although the total amount of ash in the ends was greater. The percentage ash in the whole bones, however, came close to the term value of the first gestation, because of increases in the percentage of ash in the shaft during the second gestation.

DISCUSSION

The present technique, although suitable for the purpose of this study, is not refined sufficiently for the detection of slight differences. The amount of minerals deposited in the skeleton during pregnancy in preparation for the forthcoming lactation is small in relation to the weight of the skeleton and even to the weight of long bones, in which the principal storage occurs. Although the present study indicated storage during gestation, too much reliance should not be placed in the technique used as a means of investigating such problems. It must be noted that at 100 days the skeleton of the female rat has not completed its own growth and that the observations might have been different if very young or mature breeding females had been used. However, the data do warrant the conclusion that mineral deposition during pregnancy in the rat proceeds at a rate which is at least normal and that this continued growth is not restricted to the spongiosa, but extends to the shaft as well.

The sharp reduction in the weight of the whole bone, with an even greater reduction in the total quantity of ash, that begins with and continues throughout lactation was restricted to the bone ends. There is, perhaps, some indication that shaft growth is suspended during lactation, but there was no evidence of resorption or demineralization with-

out resorption in any way comparable with the changes in bone ends.

The total evidence in this study indicates that two processes operate in the bone during lactation: resorption and demineralization of the unresorbed residue. Resorption requires no further discussion beyond calling attention to the presence of considerable quantities of cortex in the samples referred to as ends. If the corticalis in the epiphyseal and metaphyseal regions has the same composition as that in the diaphyseal zone, the changes in the spongiosa must have been correspondingly greater.

In studies of the bones of pigeons during the laying period and under the influence of toxic doses of parathyroid hormone, Bloom & Bloom (1940), Bloom *et al.* (1941) and McLean & Bloom (1940, 1941), found that bone resorption occurred and that the organic matrix and bone salt disappeared simultaneously without preliminary demineralization or the formation of osteoid borders. If histological staining techniques can detect partial demineralization processes physiologically distinct from resorption, the reductions observed in the percentages of ash during lactation could only have arisen from an accumulation of non-matrix organic matter within the bone cavity, which increases the volume as the spongiosa recedes.

On the other hand, the changes in the percentages of ash during the first and second week after the end of lactation could be explained easily if partial demineralization in addition to resorption had occurred during lactation. While the number of rats in each group is comparatively small, it should be noted that after lactation almost identical trends were found in both pregnant and non-pregnant rats. Thus, if, after lactation, minerals were deposited in the partially demineralized bone as well as being utilized for new bone growth, the rate of mineral deposition relative to the organic matter deposited would be greater in the earlier stages.

The behaviour of the bones of the non-lactating rats whose litters had been killed at birth indicates that the changes occurring in the lactating rats were a consequence of lactation and not of pregnancy or parturition. No endocrine mechanism intervenes to cause liberation of the mineral stores of pregnancy if they are not required subsequent to parturition.

The almost identical rates of replenishment of skeletal reserves in the pregnant and non-pregnant rats, during the period when the rats of the mated group were undergoing their second gestation, indicates that, within the limits of the present method, pregnancy did not exert a marked influence on storage. Since the pregnant rats were depositing minerals in the foetuses at this time, there was, of course, a greater total storage.

It is clear that the bone shafts of the lactating rat with a history of good nutrition, as in this study,

are not used as a source of minerals to sustain the mammary supply during lactation. In other words, an osteoporotic skeleton is not an obligatory consequence of lactation in this species at least. Considerable difficulties exist in the determination of these requirements by the metabolic balance technique, since this method does not indicate when mobilization of minerals from the spongiosa is accompanied by mobilization from the shaft. Further, the balance over a whole lactation is often small in relation to the total quantities of materials analyzed and may be close to the analytical errors of the methods when the diet given is rich in minerals. The present observations may find some application in the determination of certain mineral requirements of lactation. Because the losses of Ca and P are unavoidable, involving the gradual liberation of reserves laid down at an earlier period, it has been difficult to estimate the requirements for lactation on the basis of the negative values obtained by the balance technique. Since the present study shows that, in the rat, it is possible to restrict the withdrawal of bone salt to the spongiosa, leaving the shaft unaffected, it seems reasonable to suggest that lactation requirements could be defined as those intakes which, for a given output of minerals in milk, will restrict the withdrawals to the spongiosa. Thus, using a similar technique, the problem of the mineral requirements for lactation in domestic animals might become more amenable to experimental attack. This preservation of the diaphyses may be an unattainable ideal in the high-producing

dairy cow, but in sheep and swine it may be possible as in the rat. Also since Theiler (1934) pointed out that, in cattle, P deficiency had a more marked effect on the skeleton than Ca deficiency, the method might be more suitable for the determination of P than Ca requirement in the cow. While no evidence has yet been produced to show that a mild degree of osteoporosis is seriously detrimental to the individual, its avoidance would preserve the value of the skeleton from a structural standpoint.

SUMMARY

1. A technique for studying changes in the ends and shafts of the femur and tibia of the rat is described and used to follow the effect of gestation and lactation upon bone.

2. The percentage ash content and weight of dry, fat-free bones of well-fed rats were determined. During lactation and pregnancy the changes which occurred were confined to the bone ends.

3. The resorption of the spongiosa which occurred in lactating rats was not observed in non-lactating animals. The rate of replenishment of the depleted spongiosa at the end of lactation was the same, irrespective of whether the animals were then pregnant or non-pregnant.

4. The shafts of the bones of rats were little affected during lactation. The use of this finding in the determination of the Ca and P requirements of domestic animals is discussed.

The authors are indebted to Mr A. C. Dalgarno for assistance in the care of the animals.

REFERENCES

- Bauer, W., Aub, J. C. & Albright, F. (1929). *J. exp. Med.* **49**, 145.
 Bloom, W. & Bloom, M. A. (1940). *Anat. Rec.* **78**, 497.
 ——— & McLean, F. C. (1941). *Anat. Rec.* **81**, 443.
 Coons, C. M., Schiefelbusch, A. T., Marshall, G. B. & Coons, R. R. (1935). *Bull. Okla. agric. Exp. Sta.* no. 223.
 Duckworth, J. & Warnock, G. M. (1942-3). *Nutr. Abstr. Rev.* **12**, 167.
 Forbes, E. B., Black, A., Braman, W. W., Frear, D. E. H., Kahlenberg, O. J., McClure, F. J., Swift, R. W. & Voris, L. (1935). *Tech. Bull. Pa agric. Exp. Sta.* no. 319.
 Goss, H. & Schmidt, C. L. A. (1930). *J. biol. Chem.* **86**, 417.
 Kyes, P. & Potter, T. S. (1934). *Anat. Rec.* **60**, 377.
 McLean, F. C. & Bloom, W. (1940). *Anat. Rec.* **78**, 333.
 ——— (1941). *Arch. Path.* **32**, 315.
 Theiler, A. (1934). *Vel. J.* **90**, 143.
 ——— & Green, H. H. (1931-2). *Nutr. Abstr. Rev.* **1**, 359.
 Thomson, W. (1936). *J. Hyg., Camb.* **36**, 24.

Qualitative Analysis of Proteins: a Partition Chromatographic Method Using Paper

By R. CONSDEN, A. H. GORDON AND A. J. P. MARTIN, *Wool Industries Research Association, Torridon, Headingley, Leeds, 6*

(Received 13 May 1944)

Gordon, Martin & Synge (1943*b*) attempted to separate amino-acids on a silica gel partition chromatogram, but found it impracticable owing to adsorption by the silica of various amino-acids.

They obtained, however, good separations by using cellulose in the form of strips of filter paper. Following further work along these lines, the present paper describes a qualitative micro-analytical tech-